DEUTSCHES ELEKTRONEN-SYNCHROTRON DESY

DESY 76/45
August 1976

受入
76-9-139
高工研図書室

# Weak Production of Charmed Baryon Resonances

by

C. Avilez, T. Kobayashi and J. G. Körner

Abstract:

We evaluate weak neutrino production cross sections for 7 conjectured charmed baryon resonances in the four flavour quark model. Effects resulting from the mass difference of uncharmed and charmed quarks are explicitly taken into account. The quark model results are only taken at $q^2 = 0$ to determine the $q^2 = 0$ values of the invariant transition form factors. These are then continued to $q^2 \neq 0$ by suitably chosen form factors. Our numerical results are compared with the results of other calculations of weak charm production.

2 HAMBURG 52 · NOTKESTIEG 1

To be sure that your preprints are promptly included in the
HIGH ENERGY PHYSICS INDEX ,
send them to the following address ( if possible by air mail ) :

DESY
Bibliothek
2 Hamburg 52
Notkestieg 1
Germany

受入
76-9-139
高工研図書室

DESY 76/45
August 1976

# Weak Production of Charmed Baryon Resonances

by

C. Avilez, T. Kobayashi and J. G. Körner

## ERRATUM

We have corrected an error in our numerical calculation.

Replace Figs.1 and 2 by the enclosed new Figs.1 and 2.

Replace p.6, line 4 - line 11 "Our $C_0^+$ cross section ..... on their form factors." by "Our $C_0^+$ cross section is ≈ 2 times larger than the result of LS. This discrepancy is not easily accounted for. The $C_1^{*++}$ cross section is smaller than the LS result for energies $E \gtrsim 7$ GeV. This is mostly accounted for by the fact that the $q^2$-dependence of their form factors is flatter than ours, since they are not imposing the asymptotic constraints Eq.(6) on their form factors."


σ
[10^-40 cm^2]
100
10
1
MODEL I
MODEL II
5/2+; 0; 2.96
3/2−; 0; 2.83
1/2+; 0; 2.25
3/2+; 1; 2.52
1/2+; 1; 2.32
3/2−; 0; 2.83
3/2−; 1; 2.80
5/2+; 0; 2.96
5/2+; 1; 3.05
10
20
30
40
50
E [GeV]

Fig. 1




Fig. 2

# WEAK PRODUCTION OF CHARMED BARYON RESONANCES

by

C. Avilez [+,++], T. Kobayashi [+,+++] and J.G. Körner

Deutsches Elektronen-Synchrotron DESY, Hamburg

Abstract: We evaluate weak neutrino production cross sections for 7 conjectured charmed baryon resonances in the four flavour quark model. Effects resulting from the mass difference of uncharmed and charmed quarks are explicitly taken into account. The quark model results are only taken at $q^2 = 0$ to determine the $q^2 = 0$ values of the invariant transition form factors. These are then continued to $q^2 \neq 0$ by suitably chosen form factors. Our numerical results are compared with the results of other calculations of weak charm production.

[+] Alexander von Humboldt Foundation Fellow

[++] Supported in part by Consejo Nacional de Ciencia y Tecnologia, Mexico

[+++] On leave of absence from the Department of Physics, Tokyo University of Education, Tokyo, Japan



Recently some evidence for the existence of charmed particle has been reported in $e^+e^-$-experiments [1]. The dilepton events [2] in $\nu$-reactions are also considered as strong evidence for the production of charmed particles. The production of the ground state charmed baryons has been evaluated by Lee and Schrock [3] and Finjord and Ravndal [4]. In this paper we give calculations of the production cross-section for several charmed baryons including excited states in the quark model. A characteristic point of our evaluation is the introduction of the quark masses. Since the mass difference of the normal quarks (p, n) and the charmed quark (c) may be quite large, the symmetry breaking which is induced by the quark mass difference will be large and must play an important role.

## Model

The production of charmed baryons is represented as the quasi-elastic process [5,6] $\nu N \to \mu^- C$ (C = charmed baryon). We treat the excitations to the L = 0 ground state charmed baryons with $J^P = \frac{1}{2}^+, \frac{3}{2}^+$ and to the excited states $\frac{3}{2}^-$ and $\frac{5}{2}^+$ in analogy to the $D_{13}$ and $F_{15}$ seen in electroproduction:

$$\begin{array}{lll} L=0 : & C_0 \equiv (20;0;\tfrac{1}{2}^+)\,,\ C_1 \equiv (20;1;\tfrac{1}{2}^+)\,, & C_1^* \equiv (20';1;\tfrac{3}{2}^+) \\ L=1 : & (20;0;\tfrac{3}{2}^-)\,, & (20;1;\tfrac{3}{2}^-) \\ L=2 : & (20;0;\tfrac{5}{2}^+)\,, & (20;1;\tfrac{5}{2}^+) \end{array}$$

with the notation (SU (4) representation; I; $J^P$). The internal wave function of the ground state baryons composed of 3 quarks in momentum space is written as follows:

$$\Phi(\vec{p}) = \int \prod_{i=1}^{3} d^3k_i\, \delta^3(\vec{p} - \sum_{i=1}^{3} \vec{k}_i)\, \phi(\vec{k}_1, \vec{k}_2, \vec{k}_3)\, \chi_{123} \qquad (1)$$

where $\phi$ and $\chi$ stand for the space and the spin-unitary spin wave functions, respectively. We shall evaluate the hadronic matrix elements of charmed particle production in the isobar rest frame. One obtains

$$\bar{\Phi}_f(\vec{p}_f)\, \Gamma_\mu^{V,A}(0)\, \Phi_i(\vec{p}_i) = \langle C.G. \rangle \int \prod_{j=1}^{3} d^3k_j\, \delta^3(\vec{p}_i - \sum_{j=1}^{3} \vec{k}_i)\, \phi_f^+(\vec{k}_1, \vec{k}_2, \vec{k}_3 + \vec{q})\, \phi_i(\vec{k}_1, \vec{k}_2, \vec{k}_3)$$
$$\otimes\ \bar{u}_c(\vec{k}_3 + \vec{q})\, \Gamma_\mu^{V,A}\, u_n(\vec{k}_3)\,, \qquad (2)$$

where the relevant charm changing current operator is $\Gamma_\mu^V + \Gamma_\mu^A = \gamma_\mu(1-i\gamma_5)\cos\Theta_c$ with $\sin\Theta_c = 0.235 \pm 0.005$. $\langle C.G.\rangle$ is a Clebsch-Gordan type coefficient.
In the non-relativistic model the $\vec{k}$-dependence of the Dirac spinor is neglected and instead the Pauli spinors are used. Such an approximation is reliable only if $|\vec{q}|$ and the cut-off momenta for the $\vec{k}$'s in $\Phi$ are small. But in the present case the mass difference between the n and c quarks is so large that $|\vec{q}|$ is very large even at $q^2 \approx 0$. Therefore we may neglect the $\vec{k}$-dependence of the Dirac spinors, but not their $\vec{q}$-dependence. In the above approximation we reduce eq. (2) as follows:

$$\langle C.G.\rangle \bar{u}_c(\vec{q})\Gamma_\mu^{V,A} u_n(0) \int \prod_{j=1}^{3} d^3k_j\, \delta^3(\vec{p}_i - \sum_{j=1}^{3}\vec{k}_i)\, \phi_f^+(\vec{k}_1,\vec{k}_2,\vec{k}_3+\vec{q})\, \phi_i(\vec{k}_1,\vec{k}_2,\vec{k}_3) \,. \quad (3)$$

The dependence of the current matrix elements on the quark masses $m_n$ and $m_c$ is explicitly contained in the Dirac spinors $u_n$ and $u_c$. The overlap integral for the ground state in eq. (3) is a function of $q^2$ which is normalized to 1 at $q^2 = 0$. In the case of the production of excited states the overlap integral $\int \phi_f^+ \phi_i$ is proportional to $|\vec{q}|^L$ which gives a very large cross-section for large mass excited states even at $q^2 = 0$. Here we evaluate the following two cases for the excited states

Model I: $$\int \phi_f^+ \phi_i \Big|_{q^2=0} = \left(\sqrt{\tfrac{2}{\Omega}}\,|\vec{q}|\right)^L \exp\left[-\frac{m^2 q_{oc}^2}{\Omega(m^2+M^2)}\right]\Big|_{q^2=0} \quad (4)$$

Model II: $$\int \phi_f^+ \phi_i \Big|_{q^2=0} = \left(\frac{|\vec{q}|}{M}\right)^L\Big|_{q^2=0} \,,$$

where m(M) = mass of nucleon (resonance), $|\vec{q}|^2 \equiv q_c^2 = [(M+m)^2 - q^2][(M-m)^2 - q^2]/4M^2$, $q_{oc} = (M^2 - m^2 - q^2)/2M$ and $\Omega = 1.05\ \mathrm{GeV}^2$. The above forms of the overlap integrals correspond to the model presented by Feynman-Kislinger-Ravndal [7] (model I) and a simple dimensional argument for the invariant coupling strengths (model II) +).

+) It should be noted that the quark model presented by Fujimura-Kobayashi-Namiki [8] gives a suppression form $\int \phi_f^+ \phi_i \propto \left(\frac{|\vec{q}|}{\gamma}\right)^L$, where $\gamma = (M^2+m^2)/2Mm$ at $q^2 = 0$. That form is quite similar to that of the model II for an excited state with large mass.



In principle the quark model prediction for the $q^2$-dependence of the current matrix elements is contained in eq. (3) and the equivalent expressions for resonance excitation. However, beside the difficulty that the $q^2$-dependence of the overlap integral is model dependent, the $q^2$-structure of the remaining spin part tends to be unreliable. For example for the electro-production of the $D_{13}(1520)$, the quark model prediction for the $q^2$-dependence of the ratio of the $\frac{1}{2}$ to $\frac{3}{2}$ helicity amplitudes does not agree with experiment. Thus we shall use the quark model results only at $q^2 = 0$ and continue to $q^2 < 0$ by continuing a suitable set of constraint free form factors via a generalized meson dominance $q^2$-dependence. In ref. [9] it was shown that resonance excitation in electro-production can be accounted for quite well using such a continuation procedure.

## Cross-Section and Form Factors

If one neglects the lepton mass one obtains a very simple expression of the differential cross-section $d\sigma/dq^2$ in terms of the isobar rest frame helicity amplitudes. One has [5]

$$\frac{d\sigma}{dq^2} = \frac{G_c^2}{4\pi}\left[2uv|f_0|^2 + \frac{-q^2}{q_c^2}(u^2|f_+|^2 + v^2|f_-|^2)\right] \,,$$

where
$$2M^2|f_\pm|^2 = \tfrac{1}{2}\sum_\lambda |\langle N^*,\lambda\mp 1|J_\pm|N,\lambda\rangle|^2 \quad (5)$$
$$4M^2|f_0|^2 = \tfrac{1}{2}\sum_\lambda |\langle N^*,\lambda|J_0 + \tfrac{q_{oc}}{q_c}J_3|N,\lambda\rangle|^2$$

and $J_\pm = \frac{1}{2}(J_x \pm iJ_y)$. The weak hadron current is $J_\mu = J_\mu^V + J_\mu^A$. The u and $v$ are functions of $q^2$ and of E and E', the lab. energies of initial and final lepton: $2Eu = E + E' + \frac{M}{m}q_c$ and $2Ev = E + E' - \frac{M}{m}q_c$. For the weak coupling $G_c$ one has $G_c = G\sin\Theta_c$ for the weak charm production processes treated here.

The asymptotic $q^2$-dependence of the helicity form factors will be determined from the Drell-Yan threshold relation. If one has $\nu W_2 \sim (1-x)^{2c-1}$ (c = 2 corresponds to the canonical dipole case) and asymptotic suppression of the scalar cross-section $\sigma_S \propto |f_0|^2$ relative to the transverse cross section $\sigma_T \propto |f_+|^2, |f_-|^2$ vz. $\sigma_S/\sigma_T \sim q^{-2}$ one obtains

$$f_\pm \sim (q^2)^{-c+\frac{1}{2}}, \quad f_0 \sim (q^2)^{-c-\frac{1}{2}} \,. \quad (6)$$

In the case $\frac{1}{2}^+ \to \frac{1}{2}^+$ (unequal mass) this implies for the constraint free invariant form factors [9] $G_i'^{V,A} \sim (q^2)^{-1-c}$, where we define

$$\langle \tfrac{1}{2}^+ | J_\mu | \tfrac{1}{2}^+ \rangle = \bar{u}(p^*) \Big\{ \gamma_\mu [G_0'^V - i\gamma_5 G_0'^A] + \frac{i}{2M} \sigma_{\mu\nu} q_\nu [G_1'^V - i\gamma_5 G_1'^A] + \frac{1}{M^2}(q^2\gamma_\mu - \not{q} q_\mu)[G_2'^V - i\gamma_5 G_2'^A] \Big\} u(p) \quad . \tag{7}$$

For the $\frac{1}{2}^+ \to \frac{3}{2}^+$ transition one has $G_i^{V,A} \sim (q^2)^{-1-c}$ (i = 0,1,2) and $G_3^{V,A} \sim (q^2)^{-2-c}$, where we define

$$\langle \tfrac{3}{2}^+ | J_\mu | \tfrac{1}{2}^+ \rangle = \bar{u}_\beta(p^*) \Gamma_{\beta\mu} i\gamma_5 u(p) \quad , \tag{8}$$

and where

$$\Gamma_{\beta\mu} = [G_0^V - i\gamma_5 G_0^A] g_{\beta\mu} + [G_1^V + i\gamma_5 G_1^A](q_\beta \gamma_\mu - \not{q} g_{\beta\mu})/M + [G_2^V - i\gamma_5 G_2^A](q_\beta p_\mu^* - p^* q g_{\beta\mu}) + [G_3^V - i\gamma_5 G_3^A](q_\beta q_\mu - q^2 g_{\beta\mu})/M^2 .$$

For the other cases treated in this paper constraint free invariant form factors can be defined in complete analogy to eq. (7) and (8).

For the invariant form factors we shall make a generalized meson dominance ansatz in the form of a product of meson poles as predicted by the dual current model [10] for leading baryon resonance excitation [11] . We write

$$F(q^2) = \prod_{n=0}^{N(c,J)-1} \left( 1 - \frac{q^2}{m_M^2 + n\alpha'^{-1}} \right)^{-1} \quad , \tag{9}$$

where $m_M$ is the meson mass, $\alpha'$ the Regge slope and where the number of poles N(c,J) is determined by the desired large $q^2$-behaviour as discussed before.

The form factors $G_i^{V,A}$ will in general obtain contributions from mesons with $J^{PC} = 1^{--}, 1^{++}, 1^{+-}, 0^{-+}$ and $0^{++}$. In the case of charm production the mesons in (9) carry a charm unit. Since not much is known about the mass spectrum of charmed mesons yet, we shall be using one common set of mass values of the charmed mesons entering in eq. (9), namely the $D^*$ with mass 2.0 GeV [12]. For the Regge slope which determines the masses of the $D^*$-recurrences we take $\alpha' = 0.5$ GeV$^{-2}$, a value that lies between the $\alpha' \approx 1$GeV$^{-2}$ of uncharmed vector mesons and the $\alpha' \approx 0.25$ GeV$^{-2}$ of the $\Psi, \Psi'$ sequence.



Results

The model is now completely specified: the quark model is used to calculate the $q^2 = 0$ helicity amplitudes which are then projected onto the invariant amplitudes at $q^2 = 0$. The invariant amplitudes are then continued to $q^2 \neq 0$ by the GMOM form factors as described before. Except for the transition $\frac{1}{2}^+ \to \frac{3}{2}^+$ we shall assume canonical behaviour (c=2) for all form factors. For the former we take c = 3 corresponding to a faster fall-off of the N-$\Delta$ form factor as observed in electro-production experiments [13].

The model contains 4 parameters, namely the mass of the produced charmed baryon M, the mass of the charmed vector meson $m_{D^*}$, the slope of the $D^*$-trajectory $\alpha'_{D^*}$ and the ratio of the masses of the n and c quarks. For these we use $m_n \approx \frac{m}{3} = 0.313$ GeV and $m_c \approx \frac{1}{2} m_\Psi \approx 1.6$ GeV. The charmed baryon masses are estimated in a heuristic way by noting that every charmed baryon resonance (C = +1) has a strange baryon partner (S = -1) with the same values of I and $J^P$. We then estimate M(C = +1) = M(S = -1)$-m_\lambda + m_c$ with $m_\lambda \approx 0.466$ GeV. The results are not very sensitive to small variations in the masses M.

The dependence on the form factor parameters $\alpha'$ and $m_{D^*}$ is such that $\sigma$(small $\alpha'$)>$\sigma$(large $\alpha'$) and $\sigma$(small $M_{D^*}$) <$\sigma$(large $m_{D^*}$) which immediately follows from the form factors eq. (9). The variation for $0.25 < \alpha'_{D^*} < 1$ is weak but quite strong for $1.8 < m_{D^*} < 2.3$. Our subsequent discussion is based on the choice $\alpha'_{D^*} = 0.5$ GeV$^2$ and $m_{D^*} = 2.0$ GeV [12] .

An interesting point is the difference between the results of models I and II (fig. 1). Model I has

$$\sigma \text{ (ground state)} < \sigma \text{ (excited state)}$$

whereas model II gives

$$\sigma \text{ (ground state)} > \sigma \text{ (excited state)}$$

We believe therefore that the results of model II are more reliable since the results of model I would lead to unreasonably large cross-sections for high orbital excitations.

In fig. 1 we also present our predictions for the production cross sections of the re-

maining 4 I = 1 states using model II. The production cross section for $C_0^+$ off neutrons is large compared to the other cross-sections. In fig. 2 we compare our results for the L = 0 states $C_0^+$, $C_1^{++}$ and $C_1^{*++}$ with those of Lee and Shrock (LS) [3]. In the case of the $C_L^{++}$ we get similar results as LS. Our $C_0^+$ cross-section is 5 times larger than the result of LS. This discrepancy is not easily accounted for. The $C_1^{*++}$ cross section is smaller by one order of magnitude compared to the LS result. This is partly accounted for by noting that LS have divided by the initial state nucleon mass m in order to apply SU(4) to dimensionless invariant couplings. In the quark model approach these mass factors turn out to contain the charmed mass scale. Secondly, the $q^2$-dependence of their form factors is flatter than ours, since they are not imposing the asymptotic constraints eq. (6) on their form factors.

The authors would like to express their gratitude for valuable discussions with the members of the theory group at DESY. Thanks are due to M. Krammer for reading the manuscript and suggesting some improvements. Two of them (C.A. and T.K.) would also like to thank Prof. H. Joos for his hospitality.

Figure Captions

Fig. 1: Full line: Production cross-sections for 7 prominent charmed baryons and resonances using model II. Dashed line: Production cross sections for I = 0 states with $J^P = \frac{1}{2}^+, \frac{3}{2}^-, \frac{5}{2}^+$ using model I.

Fig. 2: Comparison of our results with results of LS. Mass parameters are $M_{C_0^+} = M_{C_1^{++}} = 2$ GeV and $M_{C_1^{*++}} = 3$ GeV and, in the case of LS $m_{0*} = 1.95$ GeV. For $C_1^{*++}$ we take the LS tripole result.

+) The cross-section estimates of ref. [4] are orders of magnitude lower. As remarked by LS, this is due to an imprudent choice of form factors.



References

1 Talks of V. Lüth and D. Schmitz in the Neutrino Conference at Aachen, June 1976. DASP Collaboration, DESY 76/37.

2 Talks of A. Benvenuti, B.C. Barish, V. Khovansky, E. Radermacher, A. Lutz, J. von Krogh, F.A. Nezrick, R. B. Palmer, C. Baltay and D. Carmony in the Neutrino Conference at Aachen, June, 1976.

3 R.E. Schrock and B.W. Lee, Phys. Rev. D13, (1976), 2539.

4 J. Finjord and F. Ravndal, Physics Letters 58B, (1975) 61.

5 C.H. Albright and L.S. Liu, Phys. Rev. 140 (1965), B 478.
F. Ravndal, Nuovo Cimento 18A (1973), 385.

6 T. Iwata, T. Takemura and T. Kobayashi, Prog. Theor. Phys. 42 (1969), 676.

7 R.P. Feynman, M. Kislinger and F. Ravndal, Phys. Rev. D3 (1971), 2706.

8 K. Fujimura, T. Kobayashi and M. Namiki, Prog. Theor. Phys. 43 (1970), 73 and 44 (1970), 193.

9 R.C. Devenish, T.S. Eisenschitz and J.G. Körner, Phys. Rev. D (July 1976).

10 H. Sugawara; Tokyo University of Education Report (1969); unpublished; I. Ohba, Progr. Theor. Phys. 42, 432 (1969); M. Ademollo and E. del Guidice, Nuovo Cimento 63A, 639 (1969).

11 A. Actor, I. Bender and J.G. Körner, DESY 75/57.

12 G. Goldhaber et al., preprint LBL-5309, SLAC-1762 (1976).

13 R.C. Devenish and D.H. Lyth; Nucl. Phys. B93, 109 (1975).


σ
[10^-40 cm^2]
LS
MODEL II
C_0^+
C_1^++
C_1^*++
10
1
10
20
E [GeV]

Fig.2


σ
[10^-40 cm^2]
MODEL I
MODEL II
5/2+; 0; 2.96
3/2-; 0; 2.83
1/2+; 0; 2.33
3/2-; 0; 2.83
1/2+; 1; 2.26
3/2-; 1; 2.81
5/2+; 0; 2.96
5/2+; 1; 3.06
3/2+; 1; 2.53
100
10
50
100
E [GeV]

Fig.1