

**HAMBURGER BEITRÄGE
ZUR MATHEMATIK**

Heft 140

Complete minors in $K_{s,s}$ -free graphs

D. Kühn, Hamburg, D. Osthus, HU Berlin

FEBRUAR 2002

Complete minors in $K_{s,s}$ -free graphs

Daniela Kühn

Deryk Osthus

Abstract

We prove that for a fixed integer $s \geq 2$ every $K_{s,s}$ -free graph of average degree at least r contains a K_p minor where $p = r^{1+\frac{1}{2(s-1)}+o(1)}$. A well-known conjecture on the existence of dense $K_{s,s}$ -free graphs would imply that the value of the exponent is best possible. Our result implies Hadwiger's conjecture for $K_{s,s}$ -free graphs whose chromatic number is sufficiently large compared with s .

1 Introduction

For every $r > 0$ define $p = p(r)$ to be the largest integer such that all graphs G of average degree at least r contain the complete graph K_p on p vertices as a minor. Kostochka [7] and Thomason [15] independently proved that there exists a positive constant c such that

$$p(r) \geq c \frac{r}{\sqrt{\log r}}, \quad (1)$$

which improved a bound of Mader. Random graphs show that (1) gives the correct order of magnitude. Recently, Thomason [16] showed that $p(r) = (1 + o(1))\gamma r/\sqrt{\log r}$ for an explicit constant γ .

In this paper we prove that if the graph G is locally sparse in the sense that it does not contain a fixed complete bipartite graph $K_{s,s}$ as a subgraph, then G has a K_p minor where p is asymptotically much larger than the average degree of G :

Theorem 1 *For every integer $s \geq 2$ there exists an r_s such that every $K_{s,s}$ -free graph of average degree at least $r \geq r_s$ contains a K_p minor for all*

$$p \leq \frac{r^{1+\frac{1}{2(s-1)}}}{(\log r)^3}.$$

As every graph of chromatic number k contains a subgraph of minimum degree at least $k - 1$, this implies Hadwiger's conjecture for $K_{s,s}$ -free graphs of sufficiently large chromatic number:

Corollary 2 *For every integer $s \geq 2$ there exists an integer k_s such that every $K_{s,s}$ -free graph of chromatic number $k \geq k_s$ contains a K_k minor. \square*

In Section 3 we will see that there exists an absolute constant α so that we can take $k_s := s^{\alpha s}$ in Corollary 2.

A simple observation (Proposition 14) shows that the bound on p in Theorem 1 is best possible up to the logarithmic term, provided that there exist $K_{s,s}$ -free graphs G with at least $c_s|G|^{2-1/s}$ edges. These are known to exist for $s = 2, 3$ and have been conjectured to exist also in general (see e.g. Bollobás [2, p. 362]).

In [11, Cor. 3] we showed that if G is not only $K_{s,s}$ -free but has large girth, then it contains even larger complete minors than those guaranteed by Theorem 1: for every odd integer g there exists a positive constant c such that every graph of average degree at least r and girth at least g contains a K_p minor for all $p \leq cr^{\frac{g+1}{4}}/\sqrt{\log r}$. Note that the case $g = 5$ immediately implies the case $s = 2$ of Theorem 1 because every $K_{2,2}$ -free graph G can be made into a bipartite graph (which has girth at least 5) by deleting at most half of the edges of G . Related results concerning topological minors in graphs of large girth can be found in [9, 10, 12, 13, 14]. For example, a result in [9] implies the conjecture of Hajós for all graphs of girth at least 15 and sufficiently large chromatic number.

We now turn to an application of Theorem 1 to highly connected graphs. Bollobás and Thomason [3] proved that every $22k$ -connected graph is k -linked. As is well known and easy to see, the graph obtained from K_{3k-1} by deleting k independent edges shows that the function $22k$ cannot be replaced by anything smaller than $3k - 2$. On the other hand, a result in [3] states that if a graph G is $2k$ -connected and contains a minor H with $2\delta(H) \geq |H| + 4k - 2$ then G is k -linked. Together with Theorem 1 this immediately implies the following.

Corollary 3 *For every integer $s \geq 2$ there exists an integer k_s such that for all $k \geq k_s$ every $2k$ -connected $K_{s,s}$ -free graph is k -linked. \square*

Mader [13, Cor. 1] showed that for $k \geq 2$ one cannot replace $2k$ by $2k - 1$.

Note that Theorem 1 is far from being true if we forbid a non-bipartite graph H instead of a $K_{s,s}$. Indeed, recall that there are graphs of average degree r containing no complete graph of order at least $c'r/\sqrt{\log r}$ as minor. These graphs can be made bipartite (and thus H -free) by deleting at most half of their edges. In particular, the resulting graphs G contain no complete graph as minor whose order exceeds the average degree of G . However, replacing average degree with chromatic number might help:

Problem 4 *Given an integer $s \geq 3$, does there exist a function $\omega_s(k)$ tending to infinity such that every K_s -free graph of chromatic number k contains a K_p minor for all $p \leq k \cdot \omega_s(k)$?*

In other words, the question is whether for K_s -free graphs of sufficiently large chromatic number Hadwiger's conjecture is true with room to spare. For a survey on Hadwiger's conjecture and related questions see e.g. [5].

This paper is organized as follows. In Section 2 we introduce some notation and state several results which we will need later on. Theorem 1 is then proved in Section 3. The methods are related to those in [11]. The final section is concerned with upper bounds for the size of the complete minor in Theorem 1.

2 Notation and tools

All logarithms in this paper are base e , where e denotes the Euler number. We write $e(G)$ for the number of edges of a graph G and $|G|$ for its order. We denote the degree of a vertex $x \in G$ by $d_G(x)$ and the set of its neighbours by $N_G(x)$. We denote by $\delta(G)$ the minimum degree of G , by $\Delta(G)$ its maximum degree and by $d(G) := 2e(G)/|G|$ the average degree of G . Given $A, B \subseteq V(G)$, an A - B edge is an edge of G joining a vertex in A and to a vertex in B , the number of these edges is denoted by $e_G(A, B)$. If A and B are disjoint, we write $(A, B)_G$ for the bipartite subgraph of G whose vertex classes are A and B and whose edges are all A - B edges in G . More generally, we write (A, B) for a bipartite graph with vertex classes A and B .

A graph H is a *minor* of G if for every vertex $h \in H$ there is a connected subgraph G_h of G such that all the G_h are disjoint and G contains a G_h - $G_{h'}$ edge whenever hh' is an edge in H . The vertex set of G_h is called the *branch set corresponding to h* . A *subdivision* of a graph G is a graph TG obtained from G by replacing the edges of G with internally disjoint paths. So if G contains a subdivision of a graph H then H is a minor of G .

Let us now collect some results which will be used in the proof of Theorem 1. We will need the following two easy propositions.

Proposition 5 *Every graph G contains a subgraph of average degree at least $d(G)$ and minimum degree at least $d(G)/2$.*

Proposition 6 *The vertex set of every graph G can be partitioned into disjoint sets A, B such that the minimum degree of $(A, B)_G$ is at least $\delta(G)/2$.*

Moreover, we will use the following Chernoff type bound (see e.g. [4, Cor. 2.3]).

Lemma 7 *Let X_1, \dots, X_n be independent 0-1 random variables with $\mathbb{P}(X_i = 1) = p$ for all $i \leq n$, and let $X := \sum_{i=1}^n X_i$. Then*

$$\mathbb{P}(X \leq \mathbb{E}X/2 \text{ or } X \geq 2\mathbb{E}X) \leq 2e^{-\mathbb{E}X/12}.$$

A proof of the next lemma can be found in [2, Ch. VI, Lemma 2.1].

Lemma 8 *Let (A, B) be a bipartite $K_{s,t}$ -free graph and suppose that on average each vertex in A has d neighbours in B . Then*

$$|A| \binom{d}{s} \leq t \binom{|B|}{s}.$$

Lemma 8 can be used to prove the following upper bound on the number of edges of a $K_{s,t}$ -free graph (see e.g. [2, Ch. VI, Thm. 2.3]).

Theorem 9 *Let $t \geq s \geq 1$ be integers. Then every $K_{s,t}$ -free graph G has at most $t|G|^{2-1/s}$ edges.*

Finally, we will need the following special case of [8, Cor. 19].

Lemma 10 *Let ℓ, t be integers with $\ell \geq 8t$. Let $G = (A, B)$ be a $K_{t,t}$ -free bipartite graph such that $|A| \geq \ell^{12t}|B|$ and $d_G(a) = \ell$ for every vertex $a \in A$. Then G contains a subdivision of some graph of average degree at least $\ell^9/2^{14}$.*

3 Dense Minors in $K_{s,t}$ -free graphs

Instead of proving Theorem 1, we will prove the following slightly more general result on the existence of dense minors in $K_{s,t}$ -free graphs.

Theorem 11 *For all integers $t \geq s \geq 2$ and all $r \geq (100t)^{16s}$ every $K_{s,t}$ -free graph G of average degree r contains a minor of average degree at least*

$$d := \frac{r^{1 + \frac{1}{2(s-1)}}}{10^9 t^4 (\log r)^{2 + \frac{1}{s+1}}}. \quad (2)$$

Note that asymptotically the restriction on the range of r is not too severe: if $r \leq t^s$, then (2) is already smaller than the trivial lower bound of r on the average degree of the densest minor of G .

Proof of Theorem 1. Theorem 1 immediately follows by an application of (1) to the minor obtained from the $s = t$ case of Theorem 11. \square

Furthermore, Theorem 11 shows that there exists an absolute constant α so that we can take $k_s := s^{\alpha s}$ in Corollary 2. (Indeed, given a $K_{s,s}$ -free graph G of chromatic number $r + 1$, apply Theorem 11 to a subgraph H of G of minimum degree at least r . If $r \geq s^{\alpha s}$ where α is sufficiently large compared with the constant c appearing in (1), then this shows that H contains a K_{r+1} minor, since then the value d in (2) satisfies $cd/\sqrt{\log d} \geq r + 1$.)

Our aim in the proof of Theorem 11 is to find disjoint stars in G such that a large fraction of the edges of G joins two distinct stars. If the number of these stars is not too large and if only a few edges join the same pair of stars, then the minor of G obtained by contracting the stars (and deleting all other vertices) has large average degree, as desired. We will find such stars by first choosing the set X of their centres at random and then assigning vertices $v \in G$ with distance one to X to one of the centres adjacent to v in a suitable way. For this to work we need that G is ‘almost regular’. The following lemma allows us to assume this at the expense of only a small loss of the average degree.

Lemma 12 *For all integers $t \geq 2$ and all $r \geq 10^9 t^4$ every $K_{t,t}$ -free graph G of average degree at least r either contains a subdivision of some graph of average degree at least r^3 or a bipartite subgraph H such that $\delta(H) \geq \frac{r}{400t \log r}$ and $\Delta(H) \leq r$.*

Proof. Apply Propositions 5 and 6 to obtain a bipartite subgraph G' of G of minimum degree at least $d := \lceil r/4 \rceil$. Let A be the larger vertex class of G' and delete edges if necessary to obtain a (bipartite) subgraph G'' with $d_{G''}(a) = d$ for all $a \in A$. Let B be the set of all vertices in $G'' - A$ that are not isolated and put $G^* := (A, B)_{G''}$. So $d_{G^*}(a) = d$ for all vertices $a \in A$ and thus $d(G^*) \geq d$ (since $|A| \geq |B|$). Put $N := \lceil 1 + (6t + 1) \log d \rceil$ and note that

$$\frac{d}{N} \geq \frac{d}{8t \log d} \geq 10^5 t^2. \quad (3)$$

Partition B into N disjoint sets B_1, \dots, B_N such that

$$\begin{aligned} e^{i-1} &\leq d_{G^*}(x) < e^i \quad \forall x \in B_i, \quad i = 1, \dots, N-1 \\ e^{N-1} &\leq d_{G^*}(x) \quad \forall x \in B_N. \end{aligned}$$

Then there exists an index i such that $e_{G^*}(A, B_i) \geq e(G^*)/N$. First assume that $i \leq \log d$. Then Proposition 5 implies that $(A, B_i)_{G^*}$ contains a subgraph H with $\delta(H) \geq d((A, B_i)_{G^*})/2 \geq d/2N$. As $\Delta(H) \leq \Delta((A, B_i)_{G^*}) \leq d$, H is as required in the lemma.

Next assume that $i = N$. Let A^* be the set of all those vertices in A which send at least $\lfloor \sqrt{d}/(2N)^{1/9} \rfloor =: \ell$ edges in G^* to B_N . Then

$$d|A^*| + \ell|A| \geq e_{G^*}(A, B_N) \geq \frac{e(G^*)}{N} = \frac{d|A|}{N},$$

and therefore

$$|A^*| \geq \left(\frac{d}{N} - \ell \right) \frac{|A|}{d} \geq \frac{|A|}{2N}. \quad (4)$$

Moreover, $d|A| = e(G^*) \geq e^{N-1}|B_N| \geq d^{6t+1}|B_N|$. Together with (4) this implies that

$$|A^*| \geq \frac{d^{6t}|B_N|}{2N} \geq \ell^{12t}|B_N|.$$

Let H^* be the graph obtained from $(A^*, B_N)_{G^*}$ by deleting edges if necessary such that $d_{H^*}(a) = \ell$ for all $a \in A^*$. Since $\ell \geq 8t \geq 2$ by (3), Lemma 10 implies that H^* (and hence G) contains a subdivision of some graph of average degree at least

$$\frac{\ell^9}{2^{14}} \geq r^3 \cdot \frac{d^{3/2}}{4^3 \cdot 2^{15+9N}} \stackrel{(3)}{\geq} r^3.$$

So we may assume that $\log d < i < N$. Set $k := \lfloor d/2N \rfloor$ and let A_p be a random subset of A which is obtained by including every vertex into A_p with probability $p := 2k/e^{i-1}$ independently of all other vertices. Then for every vertex $b \in B_i$ we have

$$2k \leq d_{G^*}(b)p = \mathbb{E}(|N_{G^*}(b) \cap A_p|) \leq 2ek \leq d/2. \quad (5)$$

Let us call a vertex $b \in B_i$ *bad* if $|N_{G^*}(b) \cap A_p| \leq k$ or $|N_{G^*}(b) \cap A_p| \geq d$. So (3), (5) and Lemma 7 together imply that the probability that a given vertex $b \in B_i$ is bad is at most $2e^{-k/6} \leq 1/24$. So the expected number of bad vertices in B_i is at most $|B_i|/24$. Hence Markov's inequality implies that

$$\mathbb{P}(\geq |B_i|/6 \text{ vertices of } |B_i| \text{ are bad}) \leq 1/4. \quad (6)$$

Moreover

$$2k|A| \leq \frac{d|A|}{N} \leq e_{G^*}(A, B_i) \leq e^i|B_i|,$$

and so $|A| \leq e^i|B_i|/2k$. Hence

$$\mathbb{E}(|A_p|) = p|A| \leq \frac{pe^i|B_i|}{2k} = e|B_i|.$$

Thus Markov's inequality shows that

$$\mathbb{P}(|A_p| \geq 4|B_i|) \leq e/4.$$

Together with (6) this implies that with probability at least $1 - 1/4 - e/4 > 0$ there exists an outcome A_p such that $|A_p| \leq 4|B_i|$ and at most $|B_i|/6$ vertices of B_i are bad. Let H' be the subgraph of G^* induced by A_p and those vertices in B_i that are not bad. Then $\Delta(H') \leq d$ and $e(H') \geq 5k|B_i|/6$. Moreover, $|H'| \leq |A_p| + |B_i| \leq 5|B_i|$, and so the average degree of H' is at least $k/3$. By Proposition 5, H' has a subgraph H with

$$\delta(H) \geq \frac{k}{6} \stackrel{(3)}{\geq} \frac{d}{100t \log d} \geq \frac{r}{400t \log r}.$$

So H is as required in the lemma. \square

Proof of Theorem 11. Apply Lemma 12 to G to obtain (without loss of generality) a bipartite subgraph H with $\Delta(H) \leq r$ and

$$\delta := \delta(H) \geq \frac{r}{400t \log r}. \quad (7)$$

Define ε by

$$r^\varepsilon = \frac{r^{\frac{1}{2(s-1)}}}{32t(r/\delta)^{\frac{1}{s+1}}}. \quad (8)$$

Put $\ell := r^{1-\varepsilon}$ and let X be a random subset of $V(H)$ which is obtained by including each vertex into X with probability $p := 2\ell/\delta$ independently of all other vertices. The branch sets of our minor of large average degree will consist of stars whose centres are precisely the vertices in X . Since $r \geq (100t)^{16(s-1)}$, for every vertex $v \in H$ we have

$$\begin{aligned} \mathbb{P}(v \in X) = p &= \frac{2r}{r^\varepsilon \delta} \leq \frac{2 \cdot 400t \log r}{r^\varepsilon} \\ &\leq \frac{2 \cdot 32t \cdot (400t)^2}{r^{\frac{1}{4(s-1)}}} \cdot \frac{(\log r)^2}{r^{\frac{1}{4(s-1)}}} \leq 1/20. \end{aligned} \quad (9)$$

Call a vertex $v \in H$ *good* if it satisfies the following two conditions.

- (i) $v \notin X$.
- (ii) $|N_H(v) \cap X| \geq \ell$.

We will now show that with large probability a given vertex $v \in H$ is good. First note that

$$\mathbb{E}(|N_H(v) \cap X|) = d_H(v) \cdot p \geq 2\ell.$$

As $\ell \geq \sqrt{r}$, Lemma 7 implies that

$$\mathbb{P}(|N_H(v) \cap X| < \ell) \leq 2e^{-\sqrt{r}/6} \leq 1/20.$$

Together with (9) this implies that the probability that a given vertex $v \in H$ is not good is at most $1/10$. Call an edge $uv \in H$ *good* if both u and v are good. So the probability that a given edge $uv \in H$ is not good is at most $1/5$ and therefore

$$\mathbb{E}(\text{number of edges which are not good}) \leq e(H)/5.$$

So Markov's inequality implies that

$$\mathbb{P}(\geq e(H)/2 \text{ edges are not good}) \leq 2/5.$$

Using Markov's inequality once more, we see that

$$\mathbb{P}(|X| \geq 2p|H|) \leq 1/2.$$

Thus with probability at least $1 - 2/5 - 1/2 > 0$ there is an outcome X with $|X| \leq 2p|H|$ and for which at least half of the edges of H are good. Let U be the set of good vertices of H . So $e_H(U, U)$ is precisely the number of good edges of H . For every $x \in X$ put $U_x := U \cap N_H(x)$. Note that, since H is bipartite, $H[U_x]$ consists of isolated vertices. Given a vertex $u \in U$, let $X_u := X \cap N_H(u)$. So condition (ii) implies that $|X_u| \geq \ell$.

For every vertex $u \in U$ choose a vertex $x_u \in X_u$ uniformly at random, independently of all other vertices in U . For all $x \in X$, let S_x be the set of all those $u \in U_x$ with $x_u = x$. Note that the S_x are disjoint and their union is U . Moreover, every good edge of H joins vertices in distinct S_x . We will now show that with positive probability the minor M of H whose branch sets are the $S_x \cup \{x\}$ ($x \in X$) has large average degree. For this, we will show that with positive probability a large fraction of good edges joins different pairs S_x, S_y and thus corresponds to different edges of M . As $|X|$ (i.e. the number of vertices of M) is relatively small, this will imply that M has large average degree. Thus, given a good edge $uv \in H$, we say that

- uv is of *type I* if there exists a good edge $ab \neq uv$ joining S_{x_u} to S_{x_v} such that ab and uv are disjoint,
- uv is of *type II* if there exists a good edge $ab \neq uv$ joining S_{x_u} to S_{x_v} such that a is an endvertex of uv and $|N_H(a) \cap U_{x_w}| \leq \ell/30$, where w is the endvertex of uv distinct from a (Fig. 1),
- uv is of *type III* if there exists a good edge $ab \neq uv$ joining S_{x_u} to S_{x_v} such that a is an endvertex of uv and $|N_H(a) \cap U_{x_w}| > \ell/30$, where w is the endvertex of uv distinct from a .

Note that for all distinct $x, y \in X$ the graph $H[U_x \cup U_y]$ does not contain a $K_{s-1, t}$ (since this would form a $K_{s, t}$ together with either x or y). So Theorem 9 implies that

$$e_H(U_x, U_y) \leq 4tr^{2-\frac{1}{s-1}}. \quad (10)$$

Recall that the S_x - S_y edges are precisely those U_x - U_y edges uv (with $u \in U_x$ and $v \in U_y$) for which u has chosen x and v has chosen y , i.e. for which $x = x_u$ and




Figure 1: The two possibilities for an edge uv to be of type II

$y = x_v$. Since the probability that $x = x_u$ and $y = x_v$ is $|X_u|^{-1}|X_v|^{-1} \leq \ell^{-2}$, it follows that

$$\begin{aligned} \mathbb{P}(\text{there is a good } S_x\text{-}S_y \text{ edge}) &\leq e_H(U_x, U_y) \cdot \left(\frac{1}{\ell}\right)^2 \\ &\stackrel{(10)}{\leq} 4tr^{2\varepsilon - \frac{1}{s-1}} \stackrel{(8)}{\leq} \frac{1}{60}. \end{aligned} \quad (11)$$

So given a good edge uv we have

$$\begin{aligned} \mathbb{P}(uv \text{ is of type I}) &= \sum_{x \in X_u, y \in X_v} \mathbb{P}(uv \text{ is of type I and } x = x_u \text{ and } y = x_v) \\ &= \sum_{x \in X_u, y \in X_v} \mathbb{P}(\text{there is a good } S_x\text{-}S_y \text{ edge disjoint from } uv) \cdot \frac{1}{|X_u|} \cdot \frac{1}{|X_v|} \\ &\stackrel{(11)}{\leq} \frac{1}{60}. \end{aligned}$$

Moreover, given x_u and x_v , in the definition of a type II edge uv there are at most two possibilities for a and at most $\ell/30$ candidates for b and $\mathbb{P}(x_b = x_w) \leq 1/\ell$. Thus

$$\begin{aligned} \mathbb{P}(uv \text{ is of type II}) &= \sum_{x \in X_u, y \in X_v} \mathbb{P}(uv \text{ is of type II and } x = x_u \text{ and } y = x_v) \\ &\leq \sum_{x \in X_u, y \in X_v} 2 \cdot \frac{\ell}{30} \cdot \frac{1}{\ell} \cdot \frac{1}{|X_u|} \cdot \frac{1}{|X_v|} = \frac{4}{60}. \end{aligned}$$

Hence

$$\mathbb{E}(\text{number of good edges which are type I or II}) \leq e_H(U, U)/12,$$

and so Markov's inequality implies that

$$\mathbb{P}(\geq e_H(U, U)/4 \text{ good edges are of type I or II}) \leq 1/3. \quad (12)$$

It remains to show that also with only small probability a large fraction of the good edges is of type III. This trivially holds for $s = 2$. Indeed, as $\ell/30 \geq t$,

the vertices a and x_w in the definition of a type III edge form a $K_{2,t}$ together with any t vertices in $N_H(a) \cap U_{x_w}$. Thus there are no good edges of type III in this case. So suppose that $s \geq 3$. Given a vertex $y \in X$, let V_y be the set of all those vertices in U which have at least $\ell/30$ neighbours in U_y . So $V_y \subseteq U \setminus U_y$. As H is $K_{s,t}$ -free, Lemma 8 implies that

$$|V_y| \binom{\ell/30}{s} \leq t \binom{|U_y|}{s}.$$

Thus

$$|V_y| \leq \left(\frac{32}{\ell}\right)^s t \cdot |U_y|^s \leq (32r^\varepsilon)^s t. \quad (13)$$

Given distinct good edges uv and ub and vertices $x, y \in X$, we say that the ordered quadruple uv, ub, x, y forms a *configuration of type III* if $u \in U_x$, $v, b \in U_y$ and if u has at least $\ell/30$ neighbours in U_y . So each configuration of type III can be obtained by first selecting a vertex $v \in U$, then selecting a vertex $y \in X_v$, then selecting a neighbour u of v which lies in V_y (i.e. which lies in U and sends at least $\ell/30$ edges to U_y), then we select a vertex $x \in X_u$ and finally we select a neighbour b of u in $U_y \setminus v$. We say that a configuration of type III *survives* if u has chosen x and both v and b have chosen y , i.e. if $x = x_u$ and $y = x_v = x_b$. Thus the probability that it survives is precisely $|X_u|^{-1}|X_v|^{-1}|X_b|^{-1} \leq |X_u|^{-1}|X_v|^{-1}/\ell$. Hence

$$\begin{aligned} & \mathbb{E}(\text{number of good edges which are of type III}) \\ & \leq \mathbb{E}(\text{number of surviving configurations of type III}) \\ & \leq \sum_{v \in U} \sum_{y \in X_v} \sum_{u \in N_H(v) \cap V_y} \sum_{x \in X_u} \sum_{b \in N_H(u) \cap U_y \setminus v} \frac{1}{|X_u||X_v|\ell} \\ (13) \quad & \leq \frac{|H|(32r^\varepsilon)^s t r}{r^{1-\varepsilon}} \\ (8) \quad & \stackrel{=}{=} |H| 32^s t \left(\frac{r^{\frac{1}{2(s-1)}}}{32t}\right)^{s+1} \cdot \frac{\delta}{r} \\ (s \geq 3) \quad & \stackrel{\leq}{=} \frac{\delta|H|}{32} \leq \frac{e(H)}{16} \leq \frac{e_H(U, U)}{8}. \end{aligned}$$

Hence Markov's inequality implies that also for $s \geq 3$

$$\mathbb{P}(\geq e_H(U, U)/4 \text{ good edges are of type III}) \leq 1/2.$$

Together with (12) this shows that for every $u \in U$ there exists a choice of x_u such that at most $e_H(U, U)/2$ good edges are of type I, II or III. Let F be the set of all good edges which are not of type I, II or III.

Consider the minor M of H whose branch sets are the sets $S_x \cup \{x\}$ (for all $x \in X$). As H is bipartite, every edge in F joins distinct branch sets and, by definition of F , no two edges in F join the same pair of branch sets. Thus

$e(M) \geq |F|$ and so

$$\begin{aligned}
d(M) &\geq \frac{2|F|}{|X|} \geq \frac{e_H(U, U)}{|X|} \geq \frac{e(H)}{2 \cdot 2p|H|} \geq \frac{\delta}{8p} = \frac{\delta^2}{16r^{1-\varepsilon}} \\
&\stackrel{(8)}{=} \frac{r^{1+\frac{1}{2(s-1)}}}{16 \cdot 32t} \cdot \left(\frac{\delta}{r}\right)^{2+\frac{1}{s+1}} \\
&\stackrel{(7)}{\geq} \frac{r^{1+\frac{1}{2(s-1)}}}{16 \cdot 32t \cdot (400t \log r)^{2+\frac{1}{s+1}}} \geq \frac{r^{1+\frac{1}{2(s-1)}}}{10^9 t^4 (\log r)^{2+\frac{1}{s+1}}},
\end{aligned} \tag{14}$$

as required. \square

Note that for regular graphs G the logarithmic term in (2) is not necessary. Indeed, we only have to replace the graph H in the proof of Theorem 11 with a bipartite subgraph obtained from G by an application of Proposition 6, and then (14) shows that this subgraph contains a minor of the required average degree. Moreover, for non-regular graphs the exponent $2 + \frac{1}{s+1}$ of the logarithmic term can be reduced to $1 + \frac{1}{2(s-1)}$. However, we do not give the details as we conjecture that (as in the case $s = 2$, see [11, Thm. 12]) the logarithmic term in (2) can be removed altogether.

4 Upper bounds

In this section we observe that the truth of the following well-known conjecture about the existence of dense $K_{s,t}$ -free graphs would imply that for fixed s and t Theorems 1 and 11 are best possible up to the logarithmic term.

Conjecture 13 *For all integers $t \geq s \geq 2$ there exists a positive constant $c = c(s, t)$ such that for all integers n there is a $K_{s,t}$ -free graph G of order n with at least $cn^{2-1/s}$ edges.*

(See e.g. Bollobás [2, p. 362] for the case $s = t$ which of course would already imply the general case.) In other words, the conjecture states that the upper bound on the number of edges of a $K_{s,t}$ -free graph in Theorem 9 gives the correct order of magnitude. Conjecture 13 is known to be true for all $t \geq s$ with $s = 2, 3$ (see [2, Ch. VI]). Furthermore, Alon, Rónyai and Szabó [1] proved the conjecture for all $t \geq s \geq 2$ with $t > (s-1)!$ by modifying a construction of [6]. The following proposition immediately implies that Theorems 1 and 11 are best possible up to the logarithmic term, provided that Conjecture 13 holds.

Proposition 14 *For every $c > 0$ and every $s \geq 2$ there exists a constant $C = C(c, s)$ such that whenever G is a graph with $e(G) \geq c|G|^{2-1/s}$ then every minor H of G satisfies*

$$d(H) \leq C \cdot d(G)^{1+\frac{1}{2(s-1)}}.$$

Proof. Put $n := |G|$, $r := d(G)$ and $d := d(H)$. For every vertex $h \in H$ let $V_h \subseteq V(G)$ be the branch set corresponding to h . Then

$$nr = 2e(G) \geq \sum_{h \in H} \sum_{v \in V_h} d_G(v) \geq \sum_{h \in H} d_H(h) = 2e(H) \geq d^2,$$

and so $d \leq \sqrt{nr}$. But $r \geq 2cn^{1-1/s}$, i.e. $n \leq (r/2c)^{\frac{s}{s-1}}$. Therefore

$$d \leq \frac{r^{\frac{1}{2}(1+\frac{s}{s-1})}}{(2c)^{\frac{s}{2(s-1)}}},$$

as required. □

In general, for $s \geq 4$ the best known lower bound on the maximum number of edges of a $K_{s,s}$ -free graph G is $c|G|^{2-2/(s+1)}$ (see e.g. [2, Ch. VI, Thm. 2.10]). Together with Proposition 14, this still yields an upper bound of $c'r^{1+\frac{1}{s-1}}$ for the size of the complete minor in Theorem 1 and the average degree of the minor in Theorem 11.

References

- [1] N. Alon, L. Rónyai and T. Szabó, Norm-graphs: variations and applications, *J. Combin. Theory B* **76** (1999), 280–290.
- [2] B. Bollobás, *Extremal Graph Theory*, Academic Press 1978.
- [3] B. Bollobás and A. Thomason, Highly linked graphs, *Combinatorica* **16** (1996), 313–320.
- [4] S. Janson, T. Łuczak and A. Ruciński, *Random graphs*, Wiley-Interscience 2000.
- [5] T.R. Jensen and B. Toft, *Graph Coloring Problems*, Wiley-Interscience 1995.
- [6] J. Kollár, L. Rónyai and T. Szabó, Norm graphs and bipartite Turán numbers, *Combinatorica* **16** (1996), 399–406.
- [7] A. V. Kostochka, A lower bound for the Hadwiger number of graphs by the average degree, *Combinatorica* **4** (1984), 307–316.
- [8] D. Kühn and D. Osthus, Induced subdivisions in $K_{s,s}$ -free graphs of large average degree, submitted.
- [9] D. Kühn and D. Osthus, Topological minors in graphs of large girth, submitted.
- [10] D. Kühn and D. Osthus, Subdivisions of K_{r+2} in graphs of average degree at least $r + \varepsilon$ and large but constant girth, submitted.
- [11] D. Kühn and D. Osthus, Minors in graphs of large girth, submitted.

- [12] D. Kühn and D. Osthus, Large topological cliques in graphs without a 4-cycle, in preparation.
- [13] W. Mader, Topological subgraphs in graphs of large girth, *Combinatorica* **18** (1998), 405–412.
- [14] W. Mader, Subdivisions of a graph of maximal degree $n + 1$ in graphs of average degree $n + \varepsilon$ and large girth, preprint.
- [15] A. Thomason, An extremal function for contractions of graphs, *Math. Proc. Cambridge Philos. Soc.* **95** (1984), 261–265.
- [16] A. Thomason, The extremal function for complete minors, *J. Combin. Theory B* **81** (2001), 318–338.

Daniela Kühn
Mathematisches Seminar
Universität Hamburg
Bundesstraße 55
D - 20146 Hamburg
Germany
E-mail address: `kuehn@math.uni-hamburg.de`

Deryk Osthus
Institut für Informatik
Humboldt-Universität zu Berlin
Unter den Linden 6
D - 10099 Berlin
Germany
E-mail address: `osthus@informatik.hu-berlin.de`