


DESY 77/76  
November 1977



Weak Decays of Charmed Particles and Heavy Leptons

by

T. F. Walsh

*Sektion Physik  
Universität Wuppertal*

and

*Deutsches Elektronen-Synchrotron DESY, Hamburg*

To be sure that your preprints are promptly included in the  
**HIGH ENERGY PHYSICS INDEX** ,  
send them to the following address ( if possible by air mail ) :

DESY  
Bibliothek  
2 Hamburg 52  
Notkestieg 1  
Germany

WEAK DECAYS OF CHARMED PARTICLES  
AND HEAVY LEPTONS

T.F. Walsh  
Sektion Physik, Universität Wuppertal  
and  
Deutsches Elektronen-Synchrotron DESY, Hamburg +  
Invited Talk

1977 International Symposium on Lepton and  
Photon Interactions at High Energies

I. Introduction

We now believe that charm (1) is in broad outline correct. You have heard the evidence in the experimental talks. Despite the successes of the theory, weak decays of charm may still surprise us. If charm is to fit into a larger world including  $\tau$  (2) and  $\eta$  (3), it is unlikely that the 1970 charm theory is correct in all particulars. So we should examine the basis of charm and ask where new post-1970 physics may turn up. Much of this talk is therefore concerned with what we do not know about charm: the chirality of the  $\Delta C = 1$  current, the Cabibbo suppressed decays, and  $\Delta C = 1$  nonleptonic decays generally.

With the  $\tau$  we are faced with basic issues: what is the neutral particle "v" appearing in  $\tau$  decay? Does  $\tau$  decay via the old weak current or a new one? Where does  $\tau$  fit into the lepton world? There are partial answers to the first two questions (with one disturbing element:  $\tau \rightarrow \nu \pi$  is not seen). The last question will surely occupy us for years, and we may face dramatic surprises. Perhaps we already have one, in the two mysterious HPWF super trimonon events.

II A. Charm's Chirality

Direct lepton production in  $\nu N, \bar{\nu} N$  and  $e^+ e^-$  annihilation is evidence for charm. However, we still have no solid evidence for the chirality of the  $\Delta C = 1$  current (1). Where should we look?

+ permanent address

I. Distributions

In dimuon production by  $\nu_\mu$  off d or s quarks (or  $\bar{\nu}_\mu s$ ) (4)

$$\nu_\mu d + \bar{\nu}_\mu c \rightarrow \mu^+ \nu_\mu s \text{ or } d \quad (\propto \sin^2 \theta_c); \nu_\mu s + \bar{\nu}_\mu c \rightarrow \mu^+ \nu_\mu s \text{ or } d \quad (\propto \cos^2 \theta_c) \quad (1)$$

$\nu_\mu$  is left handed and so is d or s in the 1970 GIM theory. The  $\mu^-$  distribution is thus isotropic in the  $\nu_\mu d$  rest frame. Boosted to the lab frame this gives a flat distribution in fractional muon energy loss or fractional hadron shower energy  $y = 1 - E_\mu^-/E_\nu = E_{had}^-/E_\nu$ . If the struck quark were right handed, angular momentum shows that it would go backward in the rest frame and have a lower lab energy, the  $\mu^-$  carrying off the greater part (the y distribution is, of course,  $(1-y)^2$ ).

Experimentally we have the claim of the CDHS group that the overall dimuon distribution is compatible with being flat (5), as expected for a V-A  $\Delta C = 1$  current (V+A is thus ruled out). However, the group does not give upper limits on the amount of  $s_R$  (from low x events) or  $d_R$  (from high x or valence quark events) allowed in the  $\Delta C = 1$  current.

There is another application of these ideas. Namely in

$$e^+ e^- \rightarrow D^+ D^+ \dots \rightarrow e^+ \nu_e^+ \left\{ \begin{array}{l} K \text{ (V current)} \\ \bar{K} \text{ (V,A currents)} \end{array} \right. \quad (2)$$

the kinematic configuration with maximum e energy is forbidden by angular momentum for a left handed  $s_L$  (see the figure). So we expect soft e with  $K$ . For  $s_L$  and  $s_R$  appearing symmetrically in the current (6), the maximum e energy is allowed, and a harder spectrum will result.

Experimentally, it does not yet appear possible to decide the issue; we need more data on D decays at the  $\psi(3772)$ .

2. Polarization

Neither of the above tests is satisfying; it would be better to infer the s polarization from the polarization of a  $\Lambda^0$  containing it.

Consider neutrino production of the charmed baryon  $C_0^+(cud)$



$$\nu_n + \bar{\mu} \rightarrow \bar{c}^+ \rightarrow e^+ \nu_e \Lambda^0 \quad (3)$$

studied by G. Köpp et al. (8) In general  $\bar{c}^+$  (c ud) will be polarized (in the production plane, by T-invariance). Because  $d_L + c_L$  is proportional to  $\sin\theta_c$ , this polarization is very sensitive to any admixture in the current.

Turning to the  $\Lambda^0$  polarization, the authors of (8) find nearly complete longitudinal polarization, using the 1970 GIM theory. If transitions  $c \rightarrow s_L$  and  $c \rightarrow s_R$  were equally strong and in phase, the current would be pure vector, and  $\langle p_{\Lambda^0}^+ \rangle = 0$ .

Another place to look for longitudinal  $\Lambda^0$  polarization:

$$e^+ e^- \rightarrow \bar{c} c^+ \rightarrow e^+ \nu_e \Lambda^0 + \dots \quad (4)$$

(or equivalently  $e^+ \Lambda^0 + \dots$ , where C is now a generic charmed baryon). Because of  $\gamma$  exchange, C is now unpolarized. A nonzero longitudinal polarization is thus, of course, evidence for parity violation.

Cross sections for  $\bar{c} c^+ \dots$  as large as several percent of  $\sigma_{tot}$  are credible theoretically and experimentally. With  $B(C \rightarrow e \nu \Lambda^0 + \dots) \sim 10\%$ , about 1 in  $10^3$   $e^+ e^-$  events above  $E_{CM} \sim 4.5$  GeV will contain  $e^+ \Lambda^0$ . Of course  $e^+ \Lambda^0$  from an exclusive channel like  $C^+ C^0$  will be less frequent. Because  $\bar{c} c^+ \dots$  is an inclusive process, we should not expect  $\langle p_{\Lambda^0}^+ \rangle$  as large as in (3). Of course, observation of any nonzero polarization would be of great significance.

Note that in  $e^+ \Lambda^0$  analogous to (4),  $e^+$  tags a semileptonic decay of C, whereas  $\Lambda^0$  comes from a (mostly nonleptonic) decay of  $\bar{c}$ . As a result, it will also be interesting to examine  $\langle p_{\Lambda^0}^+ \rangle_{e^+ \Lambda^0}$ . Maybe it will tell us something about chirality in  $\Delta C = 1$  nonleptonic decays.

### II B. Cabibbo's Angle

The dominance of kaons in charm decays is a dramatic confirmation of the  $\Delta C = \Delta S = 1$  rule. Now we need to look for violations of this rule. The 1970 theory predicts that  $\Delta C = 1, \Delta S = 0$  decays will take place at a rate  $\sim \tan^2 \theta_c$  times the rate of  $\Delta C = \Delta S = 1$  decays. Deviations from this prediction will be evidence for something new in charm decay.

### I. Semileptonic Processes

Tests of the charm current are cleanest in semileptonic processes, since the hadronic current appears once, multiplying a known leptonic

current. The leading  $\Delta C = \Delta S = 1$  piece has  $\Delta I = 0$ , leading to  $\Gamma(D^0 \rightarrow \ell^+ \nu \bar{K}) = \Gamma(D^+ \rightarrow \ell^+ \nu \bar{K}^0)$ . It may be possible to test this by looking for decays in emulsions. This consequence of the  $\Delta I = 0$  rule may also be useful in normalizing the relative total  $D^0$  and  $D^+$  decay rates as seen in  $e^+ e^-$  annihilation.

decay	amplitude	$\Delta S$	$\Delta I$
$e^+ \Lambda^0 \nu_e$	$\cos\theta_c$	1	0
$e^+ \Lambda^0 \nu_d$	$-\sin\theta_c$	0	1/2

The specific Cabibbo structure of the current appears in

$$\frac{\Gamma(D^0 \rightarrow e^+ \nu \pi^-)}{\Gamma(D^+ \rightarrow e^+ \nu K^-)} = \frac{BR(D^0 \rightarrow e^+ \nu \pi^-)}{BR(D^+ \rightarrow e^+ \nu K^-)} = \left[ \frac{1.6}{2.0} \tan^2 \theta_c \right] \quad (5)$$

which can be measured in  $e^+ e^-$  annihilation. (The numbers in the bracket are calculated using  $F_+(Q^2) = F_+(0)$ , giving the 1.6, and  $F_+(Q^2) = (1 + Q^2/M_c^2)^{-1} F_+(0)$ , giving 2.0 SU<sub>3</sub> for  $F_+(0)$  has also been used, from the Ademollo-Gatto theorem (12). Significant violation of (5) would indicate a new piece in the hadronic  $\Delta C = 1$  current.

Notice that (5) only checks whether d and s are mixed as the 1970 theory predicts - not whether the decaying c quark is entirely the 1970 charmed quark. Suppose  $c''$  entered (5) rather than c, where

$$\begin{pmatrix} c'' \\ t'' \end{pmatrix} = \begin{pmatrix} \cos\phi & \sin\phi \\ -\sin\phi & \cos\phi \end{pmatrix} \begin{pmatrix} c \\ t \end{pmatrix} \quad (6)$$

(t being a new  $e = 2/3$  quark, probably with its own charge  $-1/3$  partner b) (13). Now (5) is unchanged, but c is now no longer the promised 100% charmed quark (charm is distributed among c and t). We can check this by comparing

$$\mu^+ d \rightarrow \bar{\mu} c \rightarrow \mu^+ \nu_s \text{ or } d \quad \text{and} \quad e^+ e^- \rightarrow c^+ \bar{c} \rightarrow \mu^+ \nu_s \text{ or } d \quad (7)$$

For the latter,  $\sigma_{charm}$  is known, and DASP gives us the semileptonic branching ratio  $B_{charm} = 11 - 3\%$ . The former is from dimuon data at large x where valence d quarks dominate) gives  $B \sin^2 \theta_c \cos^2 \phi$ . The CDHS group have not analyzed their data with

decays promptly in  $\sim 10^{-13}$  sec. to mesons and possibly leptons. There is no time for mixing with  $D^0$ . The mixing must compete with  $\Delta C = -\Delta S$  decays at a fractional level of order  $\theta \sim 10^{-6}$ .

A very clean way of looking for mixing or a  $\Delta C = -\Delta S$  decay is through production and decay of  $\psi(3772)$ ,

$$e^+ e^- \rightarrow \psi(3772) \rightarrow D^0 + \bar{D}^0 \rightarrow \begin{cases} K^+ K^- + \dots \\ \pi^+ \pi^- + \dots \\ e^+ e^- + \dots \end{cases} \quad (9)$$

The final state with  $K^+ K^-$  arises through mixing and  $\Delta C = -\Delta S$  decay:  $e^+ e^-$  solely through mixing provided the semileptonic  $\Delta Q = \Delta C$  rule is valid. Experimentally we know that  $\int_0^1 e^+ e^-$  annihilation a  $D^0$  decays as a  $D^0$  less than 18% of the time (20) (same sign dimuon data) appear to lead to a similar but more model dependent number.

While this data excludes the existence of an order  $G_F \Delta C = 0$  neutral current (which would give complete mixing) (15), we are still 2 orders of magnitude away from the prediction of the 1970 theory. Activity in this unexplored range would be evidence for something new. Notice in particular that we have no bounds or data on final states (9) with same sign electrons. DELCO should be able to deliver on this.

II C. Nonleptonic Decays: Models

Can we expect to improve our understanding of nonleptonic weak interactions by studying charm decays? This is not obvious: final state multiplicities are large and information may be washed out by final state interactions. (This does not apply to selection rules respected by the strong interactions, of course.) Probably we should only accept as potential weak interaction physics those things which deviate significantly from what would be expected in a statistical or thermodynamic model (21). The relative decay rates into two body channels seems an ideal place to look. Remember that at Frascati energies  $\sigma(e^+ e^- \rightarrow \pi^+ \pi^-) \sim \sigma(e^+ e^- \rightarrow K^+ K^-)$  (evidence for the SU(3) classification of the photon), but that inclusively few heavy particles are produced (information on SU(3) is lost) (23). Here  $H_{NL}$  will play the role  $\gamma$  has in  $e^+ e^-$  annihilation. Let us look at some options.

1. Sextet Dominance

$H_{NL}^{quark}$  ( $\Delta C = 1$ ), broken down into SU(4) and SU(3) representations is shown in the table. It is the sum of two pieces (24):

- (i) an SU(3) V-spin singlet in an SU(3) 6 which in turn is in an SU(4) 20 and
- (ii) a V-spin triplet in an SU(3) 15, buried in an SU(4) 84

this in mind. They do claim that a credible s to d ratio,  $(S) 5\% B^e$  and  $\sin^2 \theta_c = 0.05$  (no factor  $\cos^2 \theta$ ) agrees with their data. Probably  $\cos^2 \theta$  as small as  $\sim 1/2$  is not yet excluded. This should be looked at more carefully.

2.  $\Delta C = 1$  Nonleptonic Decays

The nonleptonic decay Hamiltonian transforms as a sum of the symmetric 20 and 84 representations of SU<sub>4</sub> (see section IIIC). An immediate consequence of this and SU<sub>3</sub> invariance is

decay	amplitude	$\Delta S$	$\Delta I$
$c \rightarrow u\bar{s}$	$\cos^2 \theta_c$	1	1
$c \rightarrow u\bar{d}$	$-\cos \theta_c \sin \theta_c$	0	1/2, 3/2
$c \rightarrow u\bar{s}s$	$\cos \theta_c \sin \theta_c$	0	1/2
$c \rightarrow u\bar{s}d$	$-\sin^2 \theta_c$	-1	0, 1

that for certain nonleptonic decays only one reduced matrix element contributes, and the ratio of Cabibbo suppressed to allowed decays is predicted to be

$$\frac{\Gamma(D^0 \rightarrow \pi^+ \pi^-) \Gamma(D^+ \rightarrow \pi^0 \pi^+)}{\Gamma(D^0 \rightarrow K^+ \pi^-) \Gamma(D^+ \rightarrow K^0 \pi^+)} = \frac{\Gamma(F^+ \rightarrow K^0 \pi^+)}{2 \Gamma(F^+ \rightarrow \eta \pi^+)} \quad (8)$$

$$= 2 \frac{\Gamma(F^+ \rightarrow K^+ \pi^0)}{\Gamma(F^+ \rightarrow K^0 K^+)} = R.M.F. \tan^2 \theta_c$$

(the above uses Isgur's wave function  $\eta = \frac{1}{\sqrt{2}}(u\bar{u} + d\bar{d} - \sqrt{2}s\bar{s})$  (16), and R.M.F. stands for relative momentum factors--phase space times barrier penetration factors or form factors). Simple S-wave phase space gives  $< 20\%$  corrections to (8); this is probably the order of magnitude of SU<sub>3</sub> violations also). Large violations of (8) would mean

- (i) There is a new piece in the  $\Delta C = 1$  weak current, or
  - (ii) The still poorly understood nonleptonic decay dynamics (17) induces representations other than 20 and 84 in  $H_{NL}$ .
- Experimentally, we only know that  $\Gamma(D^0 \rightarrow \pi^+ \pi^-) / \Gamma(D^0 \rightarrow K^+ \pi^-) < 8\%$ .

3.  $D^0 - \bar{D}^0$  Mixing and  $\Delta C = -\Delta S$  Decays

The 1970 theory predicts small  $D^0 - \bar{D}^0$  mixing (1), (18). A produced  $D^0$

$\Delta C = 1$	$SU_4$	$SU_3$	V spin	comment
$(\bar{c}c)(\bar{u}d) - (\bar{u}c)(\bar{c}d)$	$20$	$6$	0	$\Delta C = 0$ is pure octet
$(\bar{c}c)(\bar{u}d) + (\bar{u}c)(\bar{c}d)$	$84$	$15$	1	$\Delta C = 0$ is octet plus $27$

Many authors observed that the  $84$  contains a  $\Delta C = 0$  piece with  $\Delta I = 3/2$ , whereas the  $20$  has a pure octet  $\Delta C = 0$  ( $\Delta I = 1/2$ ) piece. (24) Accepting exact  $SU(4)$ , they concluded that the octet enhancement of nonleptonic strange particle decays implied an enhancement of the  $SU(4)$   $20$  piece (or perhaps suppression of the  $84$ , or both simultaneously). Then  $H_{NL}$  effective ( $\Delta C = 1$ ) transforms as an  $SU(3)$   $6$  and charm decays satisfy a  $\Delta V = 0$  rule. Thus follows

$$\Gamma(D^+ \rightarrow \bar{K}^0 \pi^+) = 0 \quad (10)$$

(among other relations) (24).

Actual enhancement of the nonleptonic  $6$  would naturally lead to a large nonleptonic decay rate and a small semileptonic branching ratio (and perhaps a difference in  $D^0$  and  $D$  lifetimes).

None of the above has come to pass. From the DASP group we have  $BR(D \rightarrow e \nu + \dots) = 11 \pm 3\%$  (confirmed within errors by DELCO and SPEAR experiment SP-26). This is not small. SP-26 has reported the first clean evidence for  $D \rightarrow K \pi$  at this conference. They quote  $BR(D^+ \rightarrow \bar{K}^0 \pi^+) = 1.5 \pm .6\%$  and  $BR(D^0 \rightarrow K \pi^+) = 2.2 \pm .6\%$ . It does not seem likely that  $6$  dominance will play a role in the future. We must look elsewhere.

## 2. Quark Comics

Let us turn to a simple alternative: quark diagrams and quark as a description for  $\Delta C = 1$  two body decays. These comic book pictures are familiar by now and are shown in the tables: amplitudes calculated using color factors and exact  $SU(3)$  and ignoring gluons (25). The resulting picture for  $\Delta C = 1$  decays is certainly too naive, but it is clear, uncomplicated and predictive. Seen this way, a D meson is a  $c\bar{q}$  atom in which  $q$  is either captured by  $c$  or  $c$  decays ( $c\bar{q} \rightarrow q\bar{q}$  or  $c \rightarrow q\bar{q}$ ,  $q$  being  $u, d$  or  $s$ ). In the limit of zero  $q$  mass, quark capture vanishes by helicity: a  $V-A$  interaction gives  $q_L$  and  $q_R$  and a  $S$   $c\bar{q}$  state cannot go to  $q_L q_R$  (spin 1) (26). Charm decays simply involve the decay of the  $c$  nucleus of the  $c\bar{q}$  atom. The basic test of this is

picture	contributes to	comment
	$D^0 \rightarrow \pi^+ \pi^-$ $D^+ \rightarrow \pi^+ \pi^0$	Cabibbo suppressed; zero in free quark model; ignore.
	$D^0 \rightarrow \bar{K}^0 \pi^0$ $D^+ \rightarrow \bar{K}^0 \pi^+$	vanishes by helicity for $m_q = 0$ ; ignore
	$F^+ \rightarrow \pi^+ \pi^+ \pi^-$	vanishes by helicity for $m_q = 0$ ; absence of $3\pi$ mode tests idea

quark capture,  $c\bar{q} \rightarrow q\bar{q}$

picture	contributes to	comment
	$D^+ \rightarrow \bar{K}^0 \pi^+$ $D^0 \rightarrow \bar{K}^0 \pi^+$ $F^+ \rightarrow \pi^+ \pi^+$	color factor 3 in amplitude enhances rate; inclusively gives ratios e vs: $\bar{p}vs: \bar{u}ds \sim 1:1:3$
	$D^+ \rightarrow \bar{K}^0 \pi^+$ $D^0 \rightarrow \bar{K}^0 \pi^0$ $F^+ \rightarrow \bar{K}^0 \pi^+$	no color factor in amplitude; rates not enhanced (alternatively, they are suppressed relative to e.g. $D^0 \rightarrow \bar{K}^0 \pi^+$ )

quark decay  $c\bar{q} \rightarrow q\bar{q}q$

III. Tau

Persuasive evidence for a heavy lepton  $\tau$  with pointlike  $e^+e^-$  production cross section has been presented at this conference and ably reviewed by Perl (2). The  $\tau$  mass lies in the range 1.8 - 1.95 GeV, and lepton spectra and all branching ratios except  $\tau \rightarrow \nu_\pi \pi$  agree with what is expected for a  $V-A$   $\tau$  with its own massless neutrino. In particular,  $BR(\tau \rightarrow e\nu\nu) \approx BR(\tau \rightarrow \mu\nu\nu) \approx 15-20\%$ .

I have only two comments by way of introduction:

Firstly, there is a decisive test for a pointlike  $j = 1/2 \tau$ . Measure  $e^+e^- \rightarrow \tau^+ \tau^- \rightarrow e^+ + l$  prongs, where


$$\frac{d\sigma}{d\Omega} \approx 1 + \alpha_e [\cos^2 \theta_e + p^2 \sin^2 \theta_e \cos^2 \phi_e] \quad (14)$$

( $P$  is the  $e^-$  polarization,  $\theta_e$  the polar angle and  $\phi_e$  the azimuth measured from the ring plane). Now, Pais and Trieman pointed out that  $\langle \alpha \rangle$  ( $\alpha$  integrated over  $P_e$ ) is independent of the  $\tau$  decay interaction. It depends only on  $\beta_e = p_\tau/E_B$  for a pointlike  $j = 1/2 \tau$  and can be calculated exactly (e.g.  $\langle \alpha \rangle = .25$  for  $E_{beam} = 3.5$  GeV).


In addition  $\alpha$  ( $\alpha = P_e/E_B, E_B$ ) can be calculated (31) and turns out to be insensitive to the  $V, A$  structure of the  $\tau$ - $\nu_\tau$  vertex. (See the figure, from Ref. (31).)

It would be useful (if perhaps anticlimactic) to check this prediction.

Second remark: we know that charm is produced singly by the weak charged current in  $\nu$  interactions, and pairwise from  $l\bar{l}$  in  $e^+e^-$  annihilation. By contrast,  $\tau^+\tau^-$  is produced in  $e^+e^-$ , but not by a weak current striking a nucleon. This gives a consistency check on the belief that  $\tau$  is a heavy lepton (and not, for example a liberated quark): the lepton spectra in



(15)



$$\frac{\Gamma(F^+ \rightarrow \pi^+ \pi^+ \pi^-)}{\Gamma(F^+ \rightarrow K^+ K^+ \pi^-)} \ll 1 \quad (11)$$

(see the table;  $\Gamma(F^+ \rightarrow \rho^+ \pi^+) \ll \Gamma(F^+ \rightarrow \bar{K}^0 K^+)$  might prove a better test, as it involves two body decays). If (11) fails, the simple quark picture must be discarded.

Suppose (11) succeeds. We can then proceed. Notice that the first amplitudes in the quark decay table get a factor 3 from the sum over three colors of quark in one of the outgoing mesons. Because the colors must match in the second set of amplitudes, they receive no such enhancement. We immediately find many relations for Cabibbo allowed decays,

$$\Gamma(D^0 \rightarrow K^+ \pi^-) = 2\Gamma(F^+ \rightarrow \eta \pi^+) = (3/4)^2 \Gamma(D^+ \rightarrow \bar{K}^0 \pi^+) \quad (12)$$

$$= 18 \Gamma(D^0 \rightarrow \bar{K}^0 \pi^0) = 9\Gamma(F^+ \rightarrow K^+ \bar{K}^0)$$

(it is easy to extend this to Cabibbo suppressed decays). Now, we see that  $D^0 \rightarrow K^+ \pi^-$  is no longer suppressed, but  $F^+ \rightarrow K^+ \bar{K}^0$  is (satisfying our condition that there must be deviations from a statistical model, for which  $F^+ \rightarrow K^+ \bar{K}^0$  is unsuppressed).

If  $F^+ \rightarrow K^+ \bar{K}^0$  is indeed small, it would explain why  $F^+$  has not so far been seen in this channel (a better charged mode for an  $F^+$  search would be  $K^+ K^+ \pi^-$ ). Further, (12) is consistent with the fact that  $F^+ \rightarrow \eta \pi^+$  has been seen.

Quark decay naturally leads to equal semileptonic branching ratios for  $D^0, D^+$  and  $F^+$  (c knows nothing of the flavor of  $q$  in a  $q\bar{q}$  state). The DELCO group has presented data bearing on this (14). They note that the semileptonic branching ratios from  $2e$  events/le events and le events/all charm at  $\psi(3770)$  are equal. But it is easy to see that if  $\sigma(D^0 D^0) = \sigma(D^+ D^+)$ ;  $\sigma(F^+ F^+) = 0$ ,

$$\frac{B_{2e}}{B_{le}} - 1 = \left( \frac{B_e(D^+) - B_e(D^0)}{B_e(D^+) + B_e(D^0)} \right)^2 \quad (13)$$

and  $B_{2e} = B_{le}$  implies  $B_e(D^+) = B_e(D^0)$ . (At present the errors are rather big; hopefully we will soon have separate values for  $B_e(D^+)$  and  $B_e(D^0)$ .)

If this simple model turns out to have anything to do with real charm decays, it might help us on the way to a general understanding of non-leptonic decays. It may also be of some use in myopic futurism: guessing how final states will look above the threshold for producing the objects of which T(9.4) is made (if it is a quark-antiquark bound state).

should be the same (the  $\tau$  contribution to the former can be eliminated by demanding high multiplicities). The comparison depends on the assumption that a charmed quark fragments the same way in  $e^+e^-$  and  $\nu N$ . It should be carried out at the same value of the parent quark momentum in both processes or, accepting scaling, by plotting  $e^+e^-$  and  $\nu N$  data versus a common scaling variable (e.g.  $z = p_e/p_{\text{quark}}$ ).

Data on inclusive muons in multiprong  $e^+e^-$  events has been published by the SLAC-LBL group for mean CM energy 6.9 GeV (33). This is easily plotted versus a scaling variable. Unfortunately, the  $\mu$  spectra in  $\nu N \rightarrow \mu \mu^+ \dots$  have not yet been plotted this way, which makes comparison difficult. However, Odorico (34), in a contributed paper, points out that the energy distribution of the "soft" muon in dimuon events is well accounted for if one assumes a  $D \rightarrow e \nu K$  decay and a charmed quark to D meson fragmentation function  $D(z) = \text{constant}$ . Coincidentally, T.C. Yang and the present author have independently noted that the SLAC-LBL data on  $e^+e^- \rightarrow \mu^+ \mu^-$  multiprongs (35) be accounted for using the same constant fragmentation function.

There is thus no evidence for a second "tau-like" component to the soft muon spectrum in dimuon events, and we have no reason from this corner to suspect that  $\tau$  is anything but a heavy lepton. However, a direct comparison of data would be more satisfying.

To sum up: there are still unexplored issues, but every reason to believe that the heavy lepton  $\tau$  exists. The remainder of this talk is about the physics issues arising from this fact.

### III A. What is " $\nu_\tau$ "?

We know little about the missing neutral in decays like  $\tau^- \rightarrow e^- \nu_e \nu_\tau, \mu^- \nu_\mu \nu_\tau$ . From PLUTO we have the mass limit  $m_{\nu_\tau} < 540 \text{ MeV}$  (36) and from the existence of  $\mu e$  events and nonexistence of  $\mu e$  plus further tracks it seems plausible that the lifetime is  $\tau_{\nu_\tau} > 10^2 \text{ cm}$ .

It is not yet even established that " $\nu_\tau$ " does not interact strongly with matter.

Let us assume that " $\nu_\tau$ " is a light spin 1/2 weakly interacting neutral.

What can we say about the possibility that it is a known neutrino?

#### 1. " $\nu_\tau$ " is $\bar{\nu}_e$ or $\bar{\nu}_\mu$

Suppose " $\nu_\tau$ " is the same as the right-handed antineutrino  $\bar{\nu}_e$  or  $\bar{\nu}_\mu$  appearing at the  $e$  or  $\mu$  vertex in  $\tau^- \rightarrow \nu_\tau e^- \bar{\nu}_e$  or  $\tau^- \rightarrow \nu_\tau \mu^- \bar{\nu}_\mu$ . There is then a statistics factor 2 in the rate for  $\tau^- \rightarrow \nu_\tau e^- \bar{\nu}_e$  or  $\tau^- \rightarrow \nu_\tau \mu^- \bar{\nu}_\mu$ , independent of the  $\tau$  lifetime (37), (38). There is

evidence against such a factor for " $\nu_\tau$ " =  $\bar{\nu}_e$  and weaker evidence against such a factor for  $\bar{\nu}_\mu$ .  
The fact that the reaction (39)



is not seen argues against  $\bar{\nu}_\mu$ , provided the weak interaction in this process is order  $G_F$ .

Notice that neither of these arguments would apply if one supposed that " $\nu_\tau$ " were a new left-handed antineutrino with the same lepton number as  $\nu_\tau$  or  $\bar{\nu}_e$ , but produced only in  $\tau$  decay. In the following we follow Perl and (tentatively) discard " $\nu_\tau$ " =  $\bar{\nu}_e$  or " $\nu_\tau$ " =  $\bar{\nu}_\mu$ .

#### 2. " $\nu_\tau$ " is $\nu_{eL}$ or $\nu_{\mu L}$

Data presented at this conference bears on this question - in particular of the possibility that " $\nu_\tau$ " =  $\nu_{\mu L}$  (the left-handed neutrino in  $K^0$  or  $\pi^+ \rightarrow \mu^+ \nu_\mu$ ). PLUTO has presented a  $2\sigma$  upper bound on the  $\tau$  lifetime,  $\tau_\tau < 1 \times 10^{-11} \text{ sec}$ . (36) In the unit

$$\tau_0 = \frac{1}{5} (1.90 \text{ GeV}/m_\mu)^5 \tau_\mu = 2.3 \times 10^{-13} \text{ sec}$$

this is

$$\tau_\tau / \tau_0 < 43 \quad (17)$$

In addition, both FNAL and CERN bubble chamber groups have looked for  $\nu_{\mu L} + N \rightarrow \tau + \dots$  with  $\tau$  decaying to  $e + \dots$ . Their limits, expressed in the same unit, are

$$\tau_\tau(\nu_\mu) / \tau_0 > \begin{cases} 12 & \text{FNAL} \\ 15 & \text{CERN} \end{cases} \quad (18)$$

( $\tau_\tau(\nu_\mu)$  is the inverse of the rate to  $\nu_\mu + \dots$ ).

Evidently, events of this sort (soft  $e^-$  in a  $\nu$  beam) will be seen soon, or " $\nu_\tau$ " =  $\nu_{\mu L}$  ruled out. It is obviously much harder to rule out " $\nu_\tau$ " =  $\nu_{eL}$ , because of the low intensity of  $\nu_e$  beams. In order to go



latter corroborated by SP-26 (2), and that with the exception of the branching ratio for  $\tau \rightarrow \nu_\tau \pi$  (14), the predicted final states in  $\tau$  decay are at about the level expected for a sequential heavy lepton

We can test quantitatively whether  $\tau$  decays via the conventional current. We assume

- (i) The conventional weak current is solely responsible for  $\tau$  decay
- (ii) The weak  $\tau$ - $\nu_\tau$  vertex is any mixture of pointlike V and A (but these alone)
- (iii) The mass of the  $\tau$  associated neutrino is not too large ( $m_{\nu_\tau} < .7$  GeV).

Eliminating the uncertain total  $\tau$  decay rate by taking a ratio of branching ratios, we can then derive lower bounds for the quantities

$$\frac{BR(\tau^- \rightarrow \nu_\tau H^-)}{BR(\tau^- \rightarrow \nu_\tau e^- \nu_e)} > \begin{cases} 0.50 & H = \pi^- \\ 1.10 & H = \rho^- \end{cases} \quad (19)$$

(The derivation is a straightforward application of formulae in refs. (37), (42)). The bound for  $\rho$  is uncertain to 10-20% due to finite-width effects and errors on the experimental quantity  $f_0^2/4\pi$  (the  $\rho^0$ - $\gamma$  coupling). By contrast, the bound for  $\pi$  is good to a few percent (the pion decay constant  $f_\pi$  is well known. We heard from Yamada that DASP finds no  $\tau \rightarrow \nu_\tau \pi$  at a level 2-3 $\sigma$  below the lower bound (18). The discrepancy is 3 $\sigma$  for a leptonic branching ratio 20%, and 1.5 $\sigma$  for a 15%  $\tau \rightarrow e\nu_\tau$  branching ratio; the DASP value is 20-3%. If this is confirmed,  $\tau$  cannot decay solely via the conventional weak current. An extraordinary result, since apparently  $\tau \rightarrow \nu_\tau \rho$  and  $\tau \rightarrow \nu_\tau A_1$ , and the branching ratios  $\tau \rightarrow \nu_\tau \mu\nu$ ,  $\nu_{e\nu}$  are as predicted, and - a vital point in view of the nearby charm threshold - very few kaons appear in  $\tau$  decay.

Is there a simple and plausible explanation for a small or vanishing  $\tau \rightarrow \nu_\tau \pi$ ? Some exploratory options:

- (i) The  $\tau$ - $\nu_\tau$  vertex is V and A but has structure. This contradicts our belief in a pointlike  $\tau$ .
- (ii) Besides the old current, a new pseudoscalar piece is present which interferes destructively with the old current in  $\tau \rightarrow \nu_\tau \pi$ . This strains credibility. And why then is the ratio  $\pi \rightarrow e\nu/\pi \rightarrow \mu\nu$  small (it agrees with V - A theory)?

beyond these considerations, we will have to turn to models (in section III C, D). From now on " $\nu_\tau = \nu_\tau$ "; we assume a new neutrino.

III B. Testing the Current

We want to know whether or not the  $\tau$  decays via the "known" vector and axial vector weak current, ignoring at first the possibility of S P T contributions to the vertex

1. V, A Structure

The V, A structure of the  $\tau$ - $\nu_\tau$  vertex shows up in the decay spectrum  $\tau \rightarrow \nu_\tau \nu$  and in the angular correlation of  $e\nu$  events (41)(42). Theoretically, a V - A  $\tau$ - $\nu_\tau$  vertex is most popular. It may be worth recalling that this is expressed in terms of the Michel parameter  $\rho = 0$  for  $(V - A)_\tau (V + A)_e$  and  $\rho = 3/4$  for  $(V - A)_\tau (V - A)_e$  (the subscripts refer to the vertex:  $\tau$  for  $\tau$ - $\nu_\tau$  and  $e$  or  $\mu$  for  $e$ - $\nu_e$  or  $\mu$ - $\nu_\mu$ ). Consistency of the decay spectrum with  $\rho = 3/4$  only implies a  $\bar{V} - A$   $\tau$ - $\nu_\tau$  vertex given the conventional  $\nu_e R$  or  $\nu_\mu R$  with  $e$  or  $\mu$ . This amounts to assuming in advance that the old V - A current is responsible for  $\tau$  decay. A new current coupling  $\tau$  to  $\nu_e$  and  $e$  or  $\mu$  to new left-handed antineutrinos would reverse the chirality of the  $\tau$ - $\nu_\tau$  vertex if we assume that  $\rho = 3/4$ . A polarization measurement would be needed in order to disentangle this.

It would be well to keep in mind that we have at present little evidence about the V, A structure of the  $\tau$  decay interaction.

2. The Hadronic Current in  $\tau$  Decay

The decay of  $\tau$  to specific channels is usually calculated from the known weak current

- (i) whose leptonic part is the V - A theory with two component left handed neutrinos
- (ii) whose vector hadronic part satisfies CVC, is of first class, and obeys Cabibbo universality
- (iii) whose axial part is responsible for  $\pi \rightarrow \mu\nu$  decay, is also first class and universal.

(Of course, the current-current Hamiltonian or W boson exchange is also used). While the decay rates to specific channels may be predicted accurately, branching ratios are more uncertain. This is because multibody decays are model dependent, and they enter branching ratios through the total rate. It is therefore important to emphasize our qualitative knowledge, and those places where the theory can be precisely tested.

Qualitatively, we know that  $e$ - $\mu$  universality is not violated by as much as a factor two, that vector and axial vector currents appear to be present (this depends on the impressive evidence presented by DASP and PLUTO for  $\tau \rightarrow \nu_\tau \rho$  and  $\tau \rightarrow \nu_\tau A_1$ , respectively). The

(iii) There is a new current in  $\tau$  decay (e.g.  $V + A$ , mediated by a new  $W$  boson), and its axial piece is conserved or nearly conserved. This is not consistent with the axial quark current  $\psi \gamma_5 \psi$ . Its divergence is the same for the old and new currents. It looks hard to escape  $\tau \rightarrow \nu \pi$  if  $\tau \rightarrow \nu A_1$  is present.

We emphasize that it is not logically inconsistent to have  $\tau \rightarrow \nu \pi$  but not  $\tau \rightarrow \nu A_1$ , but clearly dramatic steps will be necessary if the  $\pi$  decay is really small or zero. If  $\tau$  is really very near in mass to  $D$  and refuses to decay as our preconceptions would have it, we could even be precipitated into a "D- $\tau$  puzzle".

It is very important to confirm the DASP result or to find the  $\pi$  decay.

From now on we assume that  $\tau \rightarrow \nu \pi$  will be found at the expected level.

III C. A New Lepton Number for  $\tau$ ?

If  $\tau$  is a sequential heavy lepton with its own massless neutrino, it automatically has a new lepton number. Then we already know a substantial fraction of what we will ever learn about  $\tau$ . (Of course,  $\tau$ -number violating decays  $\tau \rightarrow (84)$  or  $\mu \gamma$  and  $\tau \rightarrow (e \text{ or } \mu) + (e \text{ or } \mu \mu)$  may occur at some level (44).)

Even if  $\tau$  has a new lepton number, interesting things can happen.

I. A Massive Neutral Partner  $N_\tau$  for  $\tau$

Suppose that  $\tau$  follows the pattern (45)  $\nu_e, e$  and  $\nu_\mu, \mu$  except that its neutral partner is heavy,  $M_{N_\tau} > M_\tau$ . In general,  $N_\tau$  will mix with  $\nu_e, \nu_\mu$  (which need not be massless) and the  $V - A$  doublet involving  $\tau$  will be

$$\begin{pmatrix} N_\tau + \epsilon_\tau \nu_e + \epsilon_\mu \nu_\mu \\ \tau \end{pmatrix}_L \quad (20)$$

(plus mixed doublets involving  $e$  and  $\mu$ ).

Cabibbo universality (46) and CVC are then generally assumed to give  $\epsilon_e, \epsilon_\mu < .1$ . (45), (46) The result is a long-lived  $\tau$ ,

$$\frac{\tau}{\tau_0} = \frac{1}{\epsilon_e^2 + \epsilon_\mu^2} > 50 \quad (21)$$

which modestly disagrees with the  $\tau$  lifetime limit; any improvement would exclude this model with  $\epsilon_e, \epsilon_\mu < .1$ . If neutrino experiments are able to exclude  $\epsilon_\mu = 0$  (see Sec. IIIA) this model is presumably already defunct.

An interesting temporary escape from (21) has been noted by Glasgow. (45) If the mixing in the lepton sector is the same as in the quark sector (by the Cabibbo angle  $\theta$ ), (21) is weakened to  $\tau/\tau_0 = 1/2\theta^2 = 9.5$ . Then we expect  $\tau$  production in  $\nu N$  only a factor  $\sim 2$  below the present FNAL and CERN limits.

2. Massive Neutrals and  $V + A$  Currents

Another possibility, suggested by Fayet (47) and applied to muon number nonconservation by Cheng and Li (48), is to add to each  $V - A$  lepton doublet  $\begin{pmatrix} \nu_i \\ \ell_i \end{pmatrix}_L$  a  $V + A$  doublet  $\begin{pmatrix} N_i \\ \ell_i \end{pmatrix}_R$  involving a heavy neutral

lepton. (The neutral leptons will in general mix with one another). This model is theoretically interesting because of possible lepton number nonconservation (44). It is experimentally interesting because it predicts no parity violation in heavy atoms (47). It is also a useful paradigm in studying the production and experimental signature of heavy neutral leptons which might be seen in  $e \rightarrow \nu N + \nu N$  (48) in  $\tau \rightarrow e N \nu_\tau$  (if  $M_N < M_\tau$ ), followed by decay of  $N_i$  into  $e^+ e^-$ .

III D. An Old Lepton Number for  $\tau$ ?

It seems prodigal to introduce a new lepton number for each new lepton. An attractive alternative would be to give (50) an old lepton number. This idea goes back to Konopinski and Mahmoud (50) who put  $\mu, \nu, e$  in a triplet with the same lepton number. Following up this possibility for  $\tau$  within the framework of gauge theories opens some curious and exciting possibilities. We illustrate these possibilities by discussing some selected models:

I. Neutrino Swapping in  $SU_2 \times U_1$

One can explore an option where  $\mu$  and  $e$  share their neutral partners  $\nu_\mu, \nu_e$  with the  $\tau$ , which has no partner of its own. This offensive possibility is excluded in  $SU_2 \times U_1$  gauge models (one  $W$ , one  $Z$ ) because  $\tau$  is then wholly unsure of its own lepton number, resulting in a bound (51)

$$BR(\tau \rightarrow e \ell^+ \ell^- + \mu \ell^+ \ell^-) > 5\% \quad (1 = e \text{ or } \mu) \quad (22)$$

contradicting the PLUTO (SLAC-MBL) limits which require this branching ratio to be below (1%) (.6%).

The interesting possibilities appear when we allow for the possibility that there are many weak bosons (gauge group bigger than  $SU_2 \times U_1$ ).

### 2. $SU_3 \times SU_3$ Model of Bjorken and Lane

A theoretically respectable model which shows clearly what happens when there are few lepton multiplets but many weak bosons has been invented by Bjorken and Lane (52). (The model uses a lepton classification due to Guresey and Sikivie (53)). This model has great phenomenological utility, whatever its experimental fate. The model has two octets of gauge bosons mediating the weak and electromagnetic interactions. The familiar  $W^+$ ,  $W^-$  are in the first octet (together with other new charged and neutral bosons). The second octet contains no "known" charged boson. The neutral weak current is distributed among both octets. There is, of course, only one massless boson (the  $\gamma$ ). Existing experiments require that many of the unseen weak bosons must be at least several times heavier than the  $W^-$ .

The leptons in this model are in  $(\bar{3}, 3)$  representations with  $e$  and  $\mu$  number. There are new neutrinos as well as new heavy leptons,

$$E = \begin{pmatrix} \tau^0 & e^+ & \tau^+ \\ e^- & \nu_e & \nu_E \\ \tau^- & \nu_\tau & \tau^0 \end{pmatrix} L \quad M = \begin{pmatrix} M^0 & M^+ & \mu^+ \\ M^- & M^0 & \nu_M^+ \\ \mu^- & \nu_\mu & \nu_M \end{pmatrix} L \quad (23)$$

The notation is as follows. The familiar  $W^+$  causes transitions between the first two leptons in each row, and a new  $V$  between the first and third leptons. A new neutral boson mediates the nondiagonal transitions between the second and third leptons. A second octet of bosons acts similarly on the leptons in columns.

The leptonic decay of  $\tau$  takes place through the usual weak current ( $\tau \leftrightarrow W \nu_\tau$ ) and for the final state with an electron via a new current (e.g.  $\tau^- \leftrightarrow W^0 e^-$ ). This new boson contributes incoherently and

$$\frac{BR(\tau^- \rightarrow \nu_e \nu_\tau)}{BR(\tau^- \rightarrow \nu_\mu \nu_\tau)} > 1 \quad (24)$$

- i.e.  $e-\mu$  universality is violated in  $\tau$  decay because  $\tau$  has the lepton

number of  $e^-$ . The amount of violation is proportional to  $M_W^4/M_U^4$ .

The model also has  $M_{\tau^0} = M_\tau$ ;  $\tau^0$  might be seen via  $F^+ \rightarrow \tau^0 e^+$  followed by  $\tau^0 \rightarrow e^- \pi^+$  at a low rate (54). It might be better to look for  $\tau^0$  (or any other heavy neutral lepton with  $e$  or  $\mu$  number) through the semileptonic decay of a heavy quark  $Q$  to a light one  $q$ , mediated by  $W^+$  or a new gauge boson. In this model the situation is complicated. However,

$$Q \rightarrow \bar{\nu}_E \tau^0 q \quad (25)$$

$$\rightarrow \nu_\tau \tau^0 q$$

are allowed if the neutral gauge boson is not very massive. Above the threshold for heavy quarks of mass 5 GeV in  $e^- e^+$  annihilation a crude estimate of the number of  $Q$  decays containing  $\tau^0 \rightarrow e^- \pi^+$  is  $1-10$  in  $10^4$ .  $\tau^0$  will also be produced via  $W$  exchange in  $e^- + \nu_\tau \rightarrow e^- + \nu_\tau$ , and  $e^- e^+ \rightarrow \gamma \rightarrow M^+ M^-$  will provide the muonic partner of  $\tau^0$ .

Muon neutrino beams will produce  $M^0$  by exchange of one of the new neutral weak bosons. Since this coupling involves a light and a heavy lepton, a heavy quark will be produced too,

$$\nu_\mu q \rightarrow M^0 Q \quad (26)$$

The signature is, of course, a multimuon event generated by the leptonic decay of  $M^0$  and the semileptonic decay of  $Q$ .

We now turn to a more ambitious (if theoretically less respectable) scheme which exhibits the same phenomenological features.

### 3. Octet Leptons

The idea that all leptons might be put together into an octet has been around for some time (56). There is a contributed paper by Achiman on an  $SU_3 \times SU_3$  model of this type. We will discuss here a simpler model

$$\begin{matrix} \bar{N}_e & e^+ & \bar{\nu}_e & e^+ \\ \tau^- & \nu_\tau, N_\sigma & \sigma^+ & \tau^- & \nu_\tau, N_\sigma & \sigma^+ \\ \mu^- & \nu_\mu & \mu^- & \mu^- & N_\mu & \mu^- \end{matrix}$$

left-handed right-handed

due to Horn and Ross (56). See the figure for the classification of left and right handed leptons. This time there are only 8 gauge bosons. This model also shows the phenomenological features we can expect when we attempt to conserve on lepton numbers:

- (i)  $e-\mu$  universality is violated in  $\tau$  decay due to the presence of new neutral bosons. (Constraints on weak boson or neutral lepton masses also arise from  $\mu$ -decay and  $\beta$ -decay)
- (ii) There is a heavy neutral lepton produced together with  $\nu$  via the weak interactions in  $e^+e^-$  annihilation. It may also appear in semileptonic decays of a heavy quark.  $\sigma^+ \sigma^-$  will be produced via  $\nu$  in  $e^+e^-$ .
- (iii) When new heavy quarks are added, they will be produced together with a new lepton by neutrino beams,

$$\nu_\mu q \rightarrow N \mu$$

The signature here is again multimion production.

We emphasize again that these models are illustrative and not exhaustive; other models exist which make predictions for  $\tau$  decay (58). The lesson these models pass on is that if the lepton world has a rich structure but few (or no) new lepton numbers, then we expect that  $e, \mu$  and  $\nu$  communicate with new leptons and new quarks. "Old" lepton beams will then prove to be a good source of "new" leptons.

Within the next few years we may learn from  $e^+e^-$  and neutrino experiments whether these ideas have anything to do with reality or not, and perhaps where the  $\tau$  fits in the lepton world.

#### IV. Conclusion

Studies of the production and decay of charm and  $\tau$  should be based on the conviction that they are only part of a larger reality. We don't understand why  $\tau$  has appeared (with nearly the D mass!), and in view of the discovery of T, we ought to consider it surprising that charm seems to behave as was predicted it should. We have a lot to learn.

#### Acknowledgements

I wish to thank H. Joos, N. Kawamoto, M. Kugler, A. Sanda, B. Wiik and T.C. Yang for their comments. This talk was prepared during a stay at Rutherford Laboratory, and I want to thank R.J.N. Phillips and G. Ringland for their hospitality.

#### References

- (1) Y. Hara, Phys. Rev. 134, 701 (1964)  
J.D. Bjorken and S.L. Glashow, Phys. Lett. 11, 255 (1964)  
S.L. Glashow, J. Illiopoulos and L. Maiani, Phys. Rev. D2, 1285 (1970)  
gauge theory calculations can be traced through:  
M.K. Gaillard, B.W. Lee and R.E. Schrock, Phys. Rev. D13, 2674 (1976)  
V.A. Novikov, M.A. Shifman, A.I. Vainshtein and V.I. Zakharov, Phys. Rev. D16, 223 (1977)  
Phenomenology and references:  
M.K. Gaillard, B.W. Lee and J.L. Rosner, Rev. Mod. Phys. 17, 277 (1975)  
experimental discovery:  
G. Goldhaber et al., Phys. Rev. Lett. 37, 255 (1976)  
I. Peruzzi et al., Phys. Rev. Lett. 37, 569 (1976)  
R. Brandelik et al., Phys. Lett. 70B, 132 (1977)  
For information on charmed baryons, see the talk of W. Lee
- (2) M. Perl, talk at this conference
- (3) L. Lederman, talk at this conference
- (4) C. Rubbia, XVII International Conference on High Energy Physics, London 1974, p. IV-117  
A. Benvenuti et al., Phys. Rev. Lett. 34, 419 (1975)  
phenomenology:  
B.W. Lee "Dimuon Events", FNAL preprint, October, 1975  
L.M. Sehgal and P.M. Zerwas, Phys. Rev. Lett., 36, 399 (1976)  
Nucl. Phys. B108, 483 (1976)  
V. Barger, R.J.N. Phillips and T. Weiler, Phys. Rev. D13, 2511 (1976)
- (5) P. Bloch, talk at this conference
- (6) A. Ali and T.C. Yang, Phys. Lett. 65B, 275 (1976)  
V. Barger and R.J.N. Phillips, Phys. Rev. D10, 80 (1976)  
M. Gronau et al., Nucl. Phys. B123, 47 (1977)  
F. Bletzacker, H.T. Nieh and A. Soni, Phys. Rev. D16, 732 (1977)  
See also contributed papers no. 161 and no. 162.  
A right handed  $c_\alpha (1 + \gamma_5)$ 's coupling was suggested by  
H. Fritzsch, M. Gell-Mann and P. Minkowski, Phys. Lett. 59B, 256 (1975)  
F. Wilczek et al., Phys. Rev. D12, 2768 (1975)  
Adding this to the GIM current results in a pure vector  $\Delta C = 1$  current (for  $\theta = 0$ ). I've checked using  $c \rightarrow e \nu_s$  that the V spectrum differs noticeably from V-A (for  $m_c = m_D, m_s = m_K$  the V spectrum is near V+A).  
See also F. Bletzacker et al., ref. (6).
- (7) Semileptonic decay spectra may be found in R. Brandelik et al., Phys. Lett. 70B, 387 (1977) and in the talk of Yamada. For the state of the  $\psi(3772)$  see the talks by J. Kirkby and A. Barbaro-Galtieri.

(21) M. Peshkin and J.L. Rosner, Nucl. Phys. B122, 144 (1977)  
 J. Rosner, Proc. Orbis Scientiae (Jan. 1977)  
 C. Quigg and J.L. Rosner, FNAL preprint 77/60 THY (July, 1977)

(22) M. Bernardini et al., Phys. Lett. 46B, 261 (1973)

(23) e.g. H.L. Lynch, SLAC-PUB-1643 (August, 1975)  
 (data at  $\sqrt{s} = 3.0$  GeV)

(24) G. Altarelli, N. Cabibbo and L. Maiani, Nucl. Phys. B88, 285 (1975)  
 R.L. Kingsley et al., ref. (16)  
 M.K. Gaillard, et al., ref. (1)  
 M.B. Einhorn and C. Quigg, Phys. Rev. D12, 2015 (1975)

For theory arguments against  $\bar{b}$  dominance, see  
 J. Ellis, M.K. Gaillard and D.V. Nanopoulos, Nucl. Phys. B100,  
 313 (1975)

(25) After preparing this part of the talk, I received a nice paper  
 by D. Fakirov and B. Stech (contributed paper no. 164) which  
 discussed these matters in detail, including short distance gluon  
 corrections,  $SU_3$  breaking and also pseudoscalar vector decays not  
 mentioned here. The reader should consult this paper for further  
 information. For theoretical background, see J. Ellis et al.,  
 ref. (24).

(26) S. Mussinov, Phys. Rev. D15, 2025 (1977)  
 J. Ellis, et al., ref. (24)

(27) Suppression of  $F^+ \rightarrow K^0 \bar{K}^0$  was noted by J. Ellis et al., in ref.  
 (24) and attributed to a cancellation of  $20$  and  $84$  amplitudes.  
 I prefer color.

(28) J. Kirkby (ref. (7)) has pointed out that data at  $\psi(3772)$  com-  
 bined with that at 4.028 can be used to extract  $B(D^0)$  and  $B(D^+)$ .  
 (The model predicts not only that these are equal but that  $B_c$  for  
 D and a heavy lepton of the same mass should be equal. This is  
 roughly consistent with data on the  $\tau$ . See Peri, ref. (2)).

(29) T.F. Walsh, Discussion Meeting on PETRA Experiments, Frascati,  
 March 1-5, 1976 and DESY 76/47.  
 R.N. Cahn and S.D. Ellis, University of Michigan preprint  
 UM-HE-76-45  
 J. Ellis, M.K. Gaillard, D.V. Nanopoulos and S. Rudaz, CERN TH 2346  
 (July, 1977)  
 A. Ali, CERN preprint in preparation

(30) A. Pais and S.B. Treiman, Phys. Rev. D14, 293 (1976)  
 These authors also discuss tests for e.g. second class currents  
 in  $\tau$  decay.

(31) T. Hagiwara, S-Y Pi and A.I. Sanda, Ann. Phys. 106, 134 (1977);  
106, 171 (1977)

(8) G. Köpp, L.M. Sehgal and P.M. Zerwas, Nucl. Phys. B123, 61  
 (1977); B123, 77 (1977)  
 background:  
 A. Buras, Nucl. Phys. B109, 373 (1976)  
 A. Buras and J. Ellis, Nucl. Phys. B111, 341 (1976)

(9) Persistent theory folklore; exploited by  
 A. De Rújula, H. Georgi and S.L. Glashow, Phys. Rev. Lett. 37,  
 785 (1976)  
 Attempts to calculate  $e^+e^- \rightarrow C\bar{C}$  yield cross sections which are  
 even too large compared to present limits (J. Körner, private  
 communication)  
 New results on  $\bar{p}$  and  $\Lambda$ ,  $\bar{\Lambda}$  production may be found in the talk of  
 A. Barbaro-Galtieri.

(10) J. von Krogh et al., Phys. Rev. Lett. 36, 710 (1976)  
 Burmester et al., Phys. Lett. 64B, 369 (1976)  
 and ref. (1); see also the talk of D. Cline

(11) A. Pais and S.B. Treiman, Phys. Rev. D15, 2529 (1977)  
 A. Pais and O. Nachtmann, submitted paper no. 70 (Phys. Rev. D16,  
 630 (1977))

(12) R.E. Marshak, Riazuddin and C.P. Ryan, Theory of Weak Interactions  
 in Particle Physics. (Wiley-Interscience, 1969)  
 M.K. Gaillard et al., ref. (1); F. Bletzacker et al., ref. (6)

(13) e.g. M. Kobayashi and T. Maskawa, Prog. Theor. Phys. 49, 652 (1973)  
 H. Harari, Stanford Lepton-Photon Symposium (Standord, 1975)  
 see also the talk of Prof. Maiani.

(14) See the talks at this conference by  
 S. Yamada (DASP), A. Barbaro-Galtieri (SP-26) and J. Kirkby (DELCO).

(15) R.L. Kingsley et al., Phys. Rev. D11, 1919 (1975)

(16) G. Karl, private communication

(17) e.g. A. Pais and V. Rittenberg, Phys. Rev. Lett. 34, 707 (1975)

(18) M.K. Gaillard et al., ref. (1)  
 L.B. Okun, V.I. Zakharov and B.M. Pontecorvo, Lett. Nuovo Cim.  
13, 218 (1975)  
 A. Pais and S.B. Treiman, Phys. Rev. D12, 2744 (1975)  
 R.L. Kingsley, Phys. Lett. 63B, 329 (1976)  
 M. Goldhaber and J.L. Rosner, Phys. Rev. D15, 1254 (1977)

(19) J. Kirkby and A. Barbaro-Galtieri in ref. (7)

(20) G. Goldhaber, Conference on Weak Interactions (Bloomington, Ind.)  
 May 1977 LBL preprint no. 6481  
 See also G.S. Abrams et al., "Summary of Charmed Particle Studies"  
 unnumbered preprint.

- (32) F. Bletzacker and H.-T. Nieh, Phys. Rev. D14, 1251 (1976)
- (33) G.J. Feldman, et al., Phys. Rev. Lett. 38, 117 (1977)
- (34) R. Odorico, contributed paper
- (35) T.C. Yang and T.F. Walsh, DESY internal note, (Feb. 1977, unpublished).  $D(z) = \text{const.}$  is one of the forms suggested by M. Gronau et al., ref. (6). (They favored  $D(z) \propto (1-z)^2$ , which accounts for  $\nu_N$  and  $e e$  data less well than  $D(z) = \text{const.}$ )
- (36) G. Knies, talk at this conference
- (37) J.D. Bjorken and C.H. Llewellyn Smith, Phys. Rev., D7, 887 (1973)  
C.H. Llewellyn Smith, Oxford preprint 33/76 (1976)
- (38) A. Ali and T.C. Yang, Phys. Rev. D11, 3052 (1976)
- (39) M. Perl (ref. (2)), quotes a mass limit for  $\nu + M^+$  of  $M_M^+ > 10 \text{ GeV}$
- (40) M. Murtagh (FNAL) and K. Schultze (CERN), talks at this conference
- (41) Y.S. Tsai, Phys. Rev. D4, 2821 (1971)
- (42) K. Fujikawa and N. Kawamoto, Phys. Rev. D14, 59 (1976)  
A.B. Berkov, E.D. Zhizhin and Yu. Nikitin, JETP Lett. 23, 408 (1976)  
R. Gatto and I. Vendramin, submitted paper no. 92  
V. Linke, E. Tränkle and I. Bender, submitted papers no. 116-120
- (43) H.B. Thacker and J.J. Sakurai, Phys. Lett. 36B, 103 (1971)
- (44) See the talk of S. Weinberg at this conference. Notice that in gauge models  $\text{BR}(\tau \rightarrow e\gamma \text{ or } \mu\gamma)$  can be orders of magnitude larger than  $\text{BR}(\mu \rightarrow e\gamma)$ ; the search for  $\tau \rightarrow e\gamma, \mu\gamma$  should continue.
- (45) H. Fritzsch, Phys. Lett. 67B, 451 (1977)  
B.W. Lee, S. Pakvasa, R.E. Schrock and H. Sugawara, FNAL preprint 77/20-THY  
S. Glashow, Harvard preprint HUTP-77/007
- (46) For more details on universality, see  
R. Gatto and G. Sartori, Phys. Lett. 67B, 467 (1977)  
B.W. Lee and R.E. Schrock, FNAL preprint 77/21-THY (February, 1977)
- (47) P. Fayet, Nucl. Phys. B78, 14 (1974)
- (48) J.D. Bjorken and C.H. Llewellyn Smith, ref. (37)  
A. Ali, Phys. Rev. D10, 2801 (1974)  
K. Fujikawa, Phys. Lett. 62B, 176 (1976) and INS report 286 (Feb. 1977)  
B. Humpert and P.N. Scharbach, Phys. Lett. 68B, 71 (1977) and SLAC-PUB-1941 (May 1977)  
F. Bletzacker and H.T. Nieh, Stony Brook preprint ITP-SB-77-27 (March 1977)

- (49) J.L. Rosner, Nucl. Phys. B126, 124 (1977)  
T.F. Walsh, DESY Internal Note (March, 1977; unpublished)  
These authors conclude from present data that  $M_N > 1 - 1.1 \text{ GeV}$ .
- (50) J. Konopinski and H. Mahmoud, Phys. Rev. 99, 1065 (1953)  
R.E. Marshak et al., ref. (12)
- (51) G. Altarelli, N. Cabibbo, L. Maiani and R. Petronzio, Phys. Lett. 67B, 463 (1970)  
D. Horn and G.G. Ross, Phys. Lett. 67B, 460 (1977)
- (52) J.D. Bjorken, Summary Talk, "Neutrino '77" (SLAC-PUB-1996, August, 1977).
- (53) F. Gürsey, P. Ramond and S. Sikivie, Phys. Lett. 60B, 177 (1976)
- (54) See J. Rosner, ref. (49); J.D. Bjorken, private communication
- (55) M. Barnett, talk at the European Conference on Particle Physics (Budapest, July, 1977) and SLAC-PUB-1961 (June, 1977)
- (56) For references prior to 1975, see  
C.H. Albright, C. Jarlskog and M.O. Tjia, Nucl. Phys. B86, 535 (1975)  
more recent work:  
H. Fritzsch and P. Minkowski; Phys. Lett. 63B (1976)  
D. Horn and G.G. Ross, Phys. Lett. 69B, 364 (1977)
- (57) Y. Achiman, submitted paper no. 143  
(Phys. Lett. 70B, 187 (1977) )
- (58) T.C. Yang, Phys. Lett. 70B, 239 (1977)