

DESY 78/04  
January 1978



MORE ABOUT THE S MATRIX OF THE CHIRAL SU(N) THIRRING MODEL

by

E. Abdalla and B. Berg

*II. Institut für Theoretische Physik der Universität Hamburg*

P. Weisz

*Institut für Theoretische Physik der Freien Universität Berlin*

NOTKESTRASSE 85 · 2 HAMBURG 52

To be sure that your preprints are promptly included in the  
HIGH ENERGY PHYSICS INDEX,  
send them to the following address ( if possible by air mail ) :

**DESY**  
**Bibliothek**  
**Notkestrasse 85**  
**2 Hamburg 52**  
**Germany**

E.Abdalla and B.Berg  
II. Institut für Theoretische Physik  
Universität Hamburg

P.Weisz  
Institut für Theoretische Physik  
Freie Universität Berlin, Berlin

More about the S matrix of the chiral SU(N) Thirring model

Abstract: We fix the bound state poles of the S matrix of the chiral SU(N) Thirring model by general arguments. Avoiding an infrared problem by using a modified 1/N-expansion, the result is confirmed in leading order.

The exact S matrix of the chiral SU(N) invariant Thirring model described by the Lagrangian

$$\mathcal{L} = i\bar{\Psi}\gamma\partial\Psi - \frac{1}{2}(\sigma^2 + \pi^2) + g\Psi(\sigma + i\gamma^5\pi)\Psi \quad (1)$$

was proposed in ref. /1/.

Let us define  $2 \rightarrow 2$  particle scattering amplitudes involving the scattering of particles  $P_\alpha$  belonging to the fundamental representation of SU(N) and the antiparticles  $A_\alpha$  by

$$\langle P_\beta(p_1') A_\delta(p_2') | P_\alpha(p_1) A_\gamma(p_2) \rangle^{in} = \alpha_\delta F_{\beta\delta}(\theta) \delta_{\alpha\gamma} - \alpha_\gamma B_{\delta\beta}(\theta) \bar{\delta}_{\alpha\gamma} \quad (2a)$$

$$\langle P_\beta(p_1') P_\delta(p_2') | P_\alpha(p_1) P_\gamma(p_2) \rangle^{in} = \alpha_\delta S_{\beta\delta}(\theta) \delta_{\alpha\gamma} - \alpha_\gamma S_{\delta\beta}(\theta) \bar{\delta}_{\alpha\gamma} \quad (2b)$$

where

$$\begin{aligned} \delta_{\alpha\gamma} &\equiv \delta(p_1' - p_1) \delta(p_2' - p_2) \\ \bar{\delta}_{\alpha\gamma} &\equiv \delta(p_1' - p_2) \delta(p_2' - p_1) \end{aligned} \quad (2c)$$

and

$$\begin{aligned} \alpha_\delta F_{\beta\delta}(\theta) &= t_1(\theta) \delta_{\alpha\beta} \delta_{\gamma\delta} + t_2(\theta) \bar{\delta}_{\alpha\gamma} \delta_{\beta\delta} \\ \alpha_\gamma B_{\delta\beta}(\theta) &= r_1(\theta) \bar{\delta}_{\alpha\delta} \delta_{\gamma\beta} + r_2(\theta) \bar{\delta}_{\alpha\gamma} \delta_{\beta\delta} \\ \alpha_\delta S_{\beta\delta}(\theta) &= u_1(\theta) \bar{\delta}_{\alpha\beta} \delta_{\gamma\delta} + u_2(\theta) \delta_{\alpha\delta} \bar{\delta}_{\gamma\beta} \end{aligned} \quad (2d)$$

with  $\text{ch } \theta = \frac{p_1 p_2}{m^2}$ . Crossing implies  $u_1(\theta) = t_1(i\pi - \theta)$ ,  $u_2(\theta) = t_2(i\pi - \theta)$  and  $r_1(\theta) = r_2(i\pi - \theta)$ .

The main result of ref./1/ was the proposal that S matrix of the chiral SU(N) Thirring model factorises and belongs to the class II of the SU(N) S matrices which were computed previously /2/. This S matrix has vanishing backward scattering and the amplitudes are given by the following formulas

$$t_1(\theta) = (\text{pole - factor}) \cdot t_1^{(0)}(\theta), \quad (3a)$$

$$u_2(\theta) = -\frac{2}{N} \frac{i\pi}{\theta} u_1 \quad \text{and} \quad r_1(\theta) = r_2(\theta) = 0 \quad (3b)$$

with

$$t_1^{(0)}(\theta) = \frac{\Gamma\left(\frac{1}{2} + \frac{\theta}{2i\pi}\right) \Gamma\left(\frac{1}{2} + \frac{1}{N} - \frac{\theta}{2i\pi}\right)}{\Gamma\left(\frac{1}{2} - \frac{\theta}{2i\pi}\right) \Gamma\left(\frac{1}{2} + \frac{1}{N} + \frac{\theta}{2i\pi}\right)} \quad (3c)$$

the minimal solution with no poles and zeros in the physical sheet

$$\text{and pole - factor} = \prod_{i=1}^L \frac{\text{sh } \frac{1}{2}(\theta + \theta_i)}{\text{sh } \frac{1}{2}(\theta - \theta_i)} \prod_{j=1}^{L'} \frac{\text{ch } \frac{1}{2}(\theta - \theta_j')}{\text{ch } \frac{1}{2}(\theta + \theta_j')} \quad (3d)$$

For the identification of this S matrix with the chiral Thirring model a variety of classical arguments and a check of the amplitudes  $r_1$ ,  $r_2$ ,  $u_2$  and  $t_2$  in leading order of the usual  $1/N$ -expansion were presented in ref. /1/. Furthermore it was noted that the multiparticle S matrix factorises in leading order  $1/N$  in terms of two particle amplitudes /3,1/ (assuming that the infrared divergent parts make sense).

There was, however, an infrared problem involved in the evaluation of the amplitudes  $u_1(\theta)$ ,  $t_1(\theta)$  within the ordinary  $1/N$ -expansion. Knowledge of these amplitudes is crucial if one wishes to check the precise form of the pole-factor (3.d) in leading order  $1/N$ . Therefore using an ad-hoc regularisation procedure and relying on consistency arguments with semiclassical calculations /4/ only speculations were made on the form of  $t_1$ . These speculations are however inconsistent.

In the present note we apply a general argument of Karowski /5/ to this model, which (under the assumption of a reasonable non-relativistic limit) leaves only one possibility for the pole-factor (3.d) open. The result is confirmed within an improved  $1/N$ -expansion which avoids the infrared problem in computing  $t_1(\theta)$  to lowest order.

The semiclassical analysis /4/ of the model indicates a rich spectrum of multifermion and boson boundstates. As in the treatment of the Gross-Neveu model by Zamolodchikov and Zamolodchikov /6/ the spectrum has to be "fitted" by an appropriate choice of the pole-factor (3d). Recently Karowski /5/ has clarified the question when an insertion of a pole-factor (with correct residuum in an appropriate channel) has a boundstate interpretation. It turns out that there are only a very limited number of possibilities to insert a boundstate pole. Poles appearing in several channels would correspond to ghosts in channels with the wrong residuum. The consistence requirement is that no boundstate-ghost transition

is allowed. This is for example fulfilled in the massive Thirring model where at points of the coupling corresponding to vanishing backward scattering redundant poles exist /7/. In the present model redundant poles are not allowed because they are in contradiction with the consistence requirements. For example if we would try to fit the boson part of the semiclassical spectrum by introducing a pole in  $t_1(\theta)$  at  $\theta = i\pi\alpha/N$  this implies poles in both positivity +, - adjoint channels  $M_{2\pm} = t_1$  only one of which has the correct residuum /1,5/. Consistence requires

$$0 = \text{out} \langle M_- \rangle \langle M_+ \rangle \sum_{\alpha} \lambda_{\beta\alpha}^i \left\{ \langle \alpha' \bar{\beta}' \gamma' | S_{23} S_{13} | \alpha \bar{\beta} \gamma \rangle - \langle \bar{\beta}' \alpha' \gamma' | S_{23} S_{13} | \bar{\beta} \alpha \gamma \rangle \right\} \lambda_{\alpha\beta}^i \quad (4)$$

leading to the equations

$$t_2' u_1 - u_1' t_2 = t_1' u_2 - u_2' t_1 = t_3' u_3 - u_3' t_3 \quad \text{for } N=2 \quad (5a)$$

and

$$\frac{u_1}{u_1'} = \frac{t_1}{t_1'} = \frac{u_2}{u_2'} = \frac{t_2}{t_2'} \quad \text{for } N \geq 3 \quad \text{where} \quad (5b)$$

$$u = u \left( \theta + \frac{i\alpha\pi}{2N} \right)$$

$$u' = u \left( \theta - \frac{i\alpha\pi}{2N} \right)$$

which are in disagreement with the S matrix (2). In fact the Semiclassical boson boundstates are explained by noting  $t_1(0) = 1$ . In the massive Thirring model one has  $t(0) = 0$  except at those points of the coupling where a boson boundstate is just at threshold where one has  $t(0) = 1$ . Therefore we claim that in the exact solution of the present model the poles interpreted in the semiclassical treatment as boson boundstates remain at threshold.

By the above argument one is left with precisely one bound state pole possible in the chiral Thirring model in the antisymmetric channel

$$\text{pole-factor} = \frac{\text{ch } \frac{1}{2} (\theta - 2\pi i/N)}{\text{ch } \frac{1}{2} (\theta + 2\pi i/N)} \quad (6)$$

This pole is so arranged that it is not present in the symmetric channel (cf. ref./1/). It implies a spectrum in exact agreement with the semiclassical multifermion spectrum

$$m_n = m \frac{\sin(\pi n/N)}{\sin(\pi/N)} \quad (7)$$

All other possible poles are CDD-poles in the real sense of the word, i.e. they have no possible boundstate interpretation. It has been noted very early that CDD poles are excluded in the S-matrixes of field theories which have a reasonable non-relativistic limit in the sense of potential scattering /8/.

Another interesting contribution to the better understanding of our S-matrix has been recently made by Kurak and Swieca /9/. These authors claim that since the physical fermions have lost not only the chiral U(1) but also the charge U(1) symmetry, they transform according to pure SU(N). They propose an interpretation of the antiparticles as boundstate of n-1 particles and perform consistency checks at the S - matrix level. The condition of vanishing backward scattering is an immediate consistency requirement. At present we are not able to comment on this approach in general. We remark, however, the fact that the fermions have lost their U(1) charge has already been demonstrated in leading order 1/N in ref. /1/. There it was

shown that the matrix element of the chiral current  $\tilde{j}_\mu^0 = \epsilon_\mu^\nu j_\mu^0$  between the physical particles vanishes in leading order 1/N. So far we have found no inconsistencies concerning the opinion /10,1/, that the 1/N-expansion is reliable as long as no infrared problems are involved. There remains only the question of the relevance of the abnormal statistics mentioned in /9/.

Now we come to the check of the bound state factor (6) within a new 1/N expansion.

There are several ways of formulating the infrared improved 1/N

expansion/10/. The method we present, using bosonization has certain advantages. Introducing fields  $\rho, \theta$  defined through:

$$\sigma + i\pi = \rho e^{i\theta} \quad (8)$$

in the Lagrangian we obtain:

$$\mathcal{L} = \bar{\Psi} i \not{\partial} \Psi - \frac{1}{2} \rho^2 + g \rho \bar{\Psi} e^{i\theta \gamma_5} \Psi \quad (9)$$

The fermions in (9) can be bosonized by the usual formulae/10/ and we obtain

$$\mathcal{L} = \sum \frac{1}{2} (\partial_\mu \varphi_i)^2 - \frac{1}{2} \rho^2 + g \rho \sum \cos(\theta + \varphi_i \sqrt{4\pi}) \quad (10)$$

we can use the bosonization formulae once more and introduce the field  $\tilde{\Psi}$  by

$$\bar{\tilde{\Psi}}_i i \not{\partial} \tilde{\Psi}_i = \frac{1}{2} \left[ \partial_\mu \left( \varphi_i + \frac{\theta}{\sqrt{4\pi}} \right) \right]^2, \text{ etc}$$

to obtain

$$\mathcal{L} = \bar{\tilde{\Psi}} i \not{\partial} \tilde{\Psi} - \frac{1}{2} \rho^2 + g \rho \bar{\tilde{\Psi}} \tilde{\Psi} + \frac{1}{2} \partial_\mu \theta \bar{\tilde{\Psi}} \gamma^\mu \gamma_5 \tilde{\Psi} + \frac{N}{8\pi} (\partial_\mu \theta)^2 \quad (11)$$

The important point is that the massless  $\theta$  field only appears in the Lagrangian as a derivative  $\epsilon_{\mu\nu} \partial^\nu \theta = A_\mu$  thus infrared divergence problems are manifestly improved. In the process the  $\Psi$  field describing infra-particles in the sense of Schroer/11/ has been replaced by the field  $\tilde{\Psi}$  which is an interpolating field for the physical particles.

The  $1/N$  perturbation theory is now set up in the standard way-expanding the action around the stationary point  $\rho=m, A_\mu=0$ , with the dynamically generated mass. The  $\rho$  and  $A_\mu$  propagators are given by:

$$D^\rho = -\frac{2\pi i}{N} \frac{\text{tr } \phi/2}{\phi} \quad (12a)$$

$$D_{\mu\nu}^A = -\frac{2\pi i}{N m^2} \frac{1}{\phi \text{tr } \phi/2} (k_\mu k_\nu - g_{\mu\nu} K^2) \underset{k^2 \rightarrow 0}{\sim} \frac{4\pi i}{N} \frac{k_\mu k_\nu - g_{\mu\nu} K^2}{K^2} \quad (12.b)$$

for  $K^2 = -4m^2 \text{sh}^2 \phi/2$

and the vertices are

$$\begin{array}{c} \diagup \\ \diagdown \end{array} \rho = -i \quad \begin{array}{c} \diagup \\ \diagdown \end{array} A_\mu \text{---} = -\frac{i \gamma_\mu}{2}$$

It is clear that to lowest order the results of ref /1/



are reproduced for amplitudes  $u_2, t_2, r_1, r_2$  which are free of infrared problems. The advantage now is that the amplitudes  $t_1, u_2$  can be calculated without any difficulty. The  $\mathcal{P}$ -exchange diagram yields for  $u_1$  a term  $\frac{\pi i}{N \sinh \theta}$  as before. The  $A_\mu$  exchange yields:

$$\frac{(-i/2)^2}{4 m \sinh \theta} \bar{u} \gamma^\mu u \bar{u}' \gamma_\mu u' \left( \frac{-4\pi i}{N} \right) = \frac{\pi i}{N} \coth \theta \quad (13)$$

The fact that the  $K_\mu K_\nu$  in the  $A_\mu$  propagator gives zero contribution can be seen by taking the particles on shell successively.

We thus obtain:

$$u_1 = 1 + \frac{i\pi}{N} \coth \frac{\theta}{2} + \dots \quad (14)$$

in agreement with the assumed form of the S-matrix.

We also have checked that the multiparticle scattering amplitudes factorize again in terms of two particle amplitudes in leading order  $-1/N$ .

Acknowledgement: One of us (E.A.) wishes to thank Dr.M.Lüscher for the introduction in this field, and Prof.H.Lehmann for the hospitality at II.Institut für Theoretische Physik. Financial support was given by Alexander von Humboldt Stiftung and Fundação de Amparo à Pesquisa do Estado de Sao Paulo.

Two of the authors (B.B. and P.W.) are indebted to M.Karowski for valuable comment. They also acknowledge usefull discussion with V.Popov (on the infrared problem) and with B.Schroer (on the Kurak-Swieca preprint).

## References

- (1) B.Berg and P.Weisz, Preprint, FU-Berlin HEP 78/18  
to be published in Nucl.Phys.B
- (2) B.Berg, M.Karowski, W.Kurak and P.Weisz, Nucl.Phys.B134  
(1978), 125
- (3) S.Meyer and R. Trostl, Phys.Lett. B (to be published)
- (4) S.Shei, Phys.Rev. D 14 (1976) 535
- (5) M.Karowski, Preprint, FU-Berlin HEP 78/
- (6) A.B.Zamolochikow and Al.B.Zamolodchikow, Phys.Lett. 72B  
(1978) 481
- (7) B.Berg, M.Karowski, W.R.Theis and H.J.Thun, Phys.Rev. D17  
(1978) 1172
- (8) R.Haag, Nuovo Cim. 5 (1957) 203
- (9) V.Kurak and J.A.Swieca, Preprint, Pontificia Universidade  
Catolica, Rio de Janeiro
- (10) E.Witten, Preprint Harvard, HU 7 P-78/Ao 27
- (11) B.Schroer, Fortsch.Phys. 11 (1963) 1