

DESY 78/33
July 1978



Light Quark Mass Spectrum in Quantum Chromodynamics

by

Paul Langacker

Deutsches Elektronen-Synchrotron DESY, Hamburg, Germany

Heinz Pagels

Princeton University, Princeton, NJ 08540

To be sure that your preprints are promptly included in the
HIGH ENERGY PHYSICS INDEX,
send them to the following address (if possible by air mail) :

DESY
Bibliothek
Notkestrasse 85
2 Hamburg 52
Germany

Light Quark Mass Spectrum in Quantum Chromodynamics

Paul Langacker ⁺

Deutsches Elektronen-Synchrotron DESY, Hamburg, Germany

Heinz Pagels ⁺⁺

Princeton University, Princeton, NJ 08540

⁺ Permanent address: University of Pennsylvania, Philadelphia, PA 19174.

⁺⁺ Permanent address: Rockefeller University, New York, N.Y. 10021

Abstract

Quantum Chromodynamics (QCD) has placed the problem of hadronic symmetry breaking on a rational basis. The current quark mass ratios can be shown to be renormalization group invariants up to small and controllable corrections from flavor interactions. We calculate the mass ratios of the light u, d and s quarks using the pseudoscalar meson mass spectrum, the baryon mass spectrum and $\eta \rightarrow 3\pi$ decay. The main theoretical assumptions are that low lying resonance and Born terms correctly estimate the photonic contribution to isotopic mass splittings and that chiral perturbation theory - equivalently kaon PCAC - correctly estimates chiral symmetry breaking. Taking account of all leading order chiral corrections to the meson spectrum and from the baryon spectrum and $\eta \rightarrow 3\pi$ decay we obtain $m_u/m_d = 0.38 \pm 0.13$ and $m_d/m_s = 0.045 \pm 0.011$. We conclude that while a vanishing up quark mass is not rigorously ruled out it is unattractive from the standpoint of the presently consistent phenomenology of hadronic symmetry breaking.

I. Current Quark Masses and Quantum Chromodynamics

In any discussion of the quark mass spectrum it is important to distinguish between current quark masses and constituent quark masses. To provide a framework for our discussion we will adopt quantum chromodynamics (QCD) as our strong interaction theory ⁽¹⁾. The Lagrangian for the $SU_c(3)$ gauge theory with N quark flavors is

$$\mathcal{L}^{\text{QCD}} = -\frac{1}{4} F_{\mu\nu}^a F_{\mu\nu}^a + \bar{q}_i \not{D} q_i + \Delta \mathcal{L}^{\text{QCD}} + \text{gauge terms}$$

$$\Delta \mathcal{L}^{\text{QCD}} = \sum_{i=1}^N m_i^0 \bar{q}_i q_i \quad (1.1)$$

where $F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + g F^{abc} A_\mu^b A_\nu^c$ is the Yang-Mills field strength and D_μ is the covariant derivative. Here m_i^0 is the bare current quark mass of the i^{th} flavored quark. An important feature of QCD in the absence of weak (flavored) interactions is that the current quark mass term $\Delta \mathcal{L}^{\text{QCD}}$ is the only term that can break the chiral $SU(N) \times SU(N)$ symmetry of \mathcal{L}^{QCD} and maintain renormalizability.

Constituent quark masses are the physical masses of the quarks as they make their appearance in hadrons. This constituent quark mass is not the same as the current quark mass since QCD presumably undergoes a PCAC phase transition. This implies that even in the absence of explicit flavor breaking ($m_i^0 = 0$) the chiral \mathcal{Y}_5 symmetry of the vacuum is broken. All the quarks then can acquire a common mass M by the Nambu-Jona Lasinio mechanism ⁽²⁾.

The problem with discussions of constituent quark masses is that the constituent

quark mass, to the authors' knowledge, has never been given a precise definition independent of a specific application and its dynamical details. If quarks could exist as free physical particles then the constituent quark mass could be precisely defined as the mass of that particle. However if quarks are permanently confined inside hadrons the constituent quark mass depends on the details of the dynamics and a precise definition has not yet been given. In spite of this difficulty the use of the concept of a constituent, physical quark mass in specific phenomenological applications, especially nonrelativistic potential models of permanently bound quarks, is quite useful.

By contrast the current quark masses can be given a precise phenomenological meaning. The reason is that the divergences of the currents that are measured in electro and neutrino production experiments are given by $m_i^0 \bar{q}_i q_i$ and $m_i^0 \bar{q}_i \gamma_5 q_i$. These operators make their appearance in the operator product expansion for the current correlation function near the light cone. For example the W_4 and W_5 structure functions measured in neutrino production from nucleons vanish in the chiral limit as $m_i^0 \rightarrow 0$ ⁽³⁾. J. Collins has shown that moments of these structure functions give direct informations about current quark masses ⁽⁴⁾. While in principle it is possible to determine the current quark masses in this way it is nearly impossible in practice since the distributions are multiplied by a lepton mass and hence are very small for the light leptons.

Actually what one obtains from such experiments that probe the light cone is not the bare mass $m_i^0(\Lambda)$ which depends on the cutoff Λ but the cutoff independent quantity $m_i(\mu)$ where μ is the renormalization mass. The relation of $m_i(\mu)$ to the bare mass $m_i^0(\Lambda)$ is

$$m_i(\mu) = Z(\Lambda/\mu) m_i^0(\Lambda) \quad (1.2)$$

The important feature of equation (1.2), valid in QCD, is that the renormalization $Z(\Lambda/\mu)$ is flavor (i) independent. This is a special feature of QCD and is a consequence of the fact that in QCD the flavor and color $SU_c(3)$ groups commute so the strong interaction is flavor independent. Equation (2) implies

$$R_{ij} = \frac{m_i(\mu)}{m_j(\mu)} = \frac{m_i^0(\Lambda)}{m_j^0(\Lambda)} \quad (1.3)$$

so that the ratios R_{ij} of the current quark masses are cutoff and μ independent. The renormalization group invariant quantities R_{ij} are unambiguous pure numbers that can, in principle, be extracted from experiment. It is the determination of these ratios that will be the subject of this article.

The μ dependence of $m_i(\mu)$ is established ⁽⁵⁾ from the renormalization group equation

$$\left(\mu \frac{\partial}{\partial \mu} + \beta(g) \frac{\partial}{\partial g} + \gamma(g) \right) m_i(\mu) = 0, \quad (1.4)$$

where $\beta(g) = -(b_0/2)g^3 + \dots$ is the Callan-Symanzik function and $\gamma(g) = \gamma_0 g^2 + \dots$ the anomalous dimension of the operator $m_i^0 \bar{q}_i q_i$ which is independent of the specific flavor, i. Equation (1.4) can be solved for

$m_i(\mu)$ up to an integration constant m_i

$$m_i(\mu) = m_i F(\mu) \quad (1.5)$$

Clearly $R_{ij} = m_i/m_j$ and for large μ asymptotic freedom implies

$$F(\mu) \xrightarrow{\mu \rightarrow \infty} (\ln \mu)^{-b_0/b_0} \quad (1.6)$$

We can define m_i to be the renormalized current quark mass.

The above discussion shows that current quark mass ratios R_{ij} are unambiguous in QCD if we ignore flavor breaking interactions due to weak interaction gluons (QFD). However if we examine isospin breaking the photon interaction must be taken into account. We will however ignore the usual weak interaction for $Q^2 \ll m_W^2 \sim (60 \text{ GeV})^2$. As a consequence of the electromagnetic flavor breaking interactions the operator $m_i^0 \bar{q}_i q_i$ is no longer universally renormalized and there are ambiguities in the quark mass ratios as emphasized by J. Collins (4). However, for quarks of the same electric charge, since the electromagnetic interaction couples only to the charge, the renormalization is still universal. Hence the quark mass ratios

$$R_{ij} = \frac{m_i^0(1)}{m_j^0(1)}, \quad q_i = q_j = \pm \frac{1}{3}, \pm \frac{2}{3} \quad (1.7)$$

for quarks of the same charge are unambiguous.

For quarks with different charges the mass ratios $m_i(\mu)/m_j(\mu)$ are not renormalization group invariant. The renormalization group equation in the presence of electromagnetism is

$$\left(\mu \frac{\partial}{\partial \mu} + \beta_g(g, e) \frac{\partial}{\partial g} + \beta_e^i(e, g) \frac{\partial}{\partial e} + \gamma_i(g, e) \right) m_i(\mu) = 0 \quad (1.8)$$

where $\beta_g(g, e) = -(b_0/2)g^3 + O(g^3e^2)$, $\beta_e^i(e, g) = (e^2/12\pi^2)q_i^2 + O(e^2g^2)$, $\gamma_i(g, e) = \gamma_0 g^2 + \gamma_1 q_i^2 e^2 + \dots$

Although β_g is flavor independent β_e^i and γ_i are not. Expanding in powers of $\alpha = e^2/4\pi$ and defining $m_i = m_i(\mu_0)$, where μ_0 is arbitrary, one finds ⁽⁴⁾ using (8)

$$R_{ij}(\mu) = \frac{m_i(\mu)}{m_j(\mu)} = \frac{m_i}{m_j} \left(1 + \alpha A (q_i^2 - q_j^2) \ln(\mu/\mu_0) \right) \quad (1.9)$$

where A is a numerical constant. This equation exhibits the ambiguity if $q_i^2 \neq q_j^2$ since $R_{ij}(\mu)$ is μ dependent.

Although this ambiguity exists it is numerically unimportant. If we compare the mass ratio $R_{ij}(\mu)$ at two different values of μ (μ_1 and μ_2) the difference is small compared to the mass ratio itself;

$$\frac{R_{ij}(\mu_1) - R_{ij}(\mu_2)}{R_{ij}} = \alpha A (q_i^2 - q_j^2) \ln(\mu_1/\mu_2) \quad (1.10)$$

$\ll 1$

providing μ_1/μ_2 is not of $O(e^{-1/\alpha})$. This is certainly the case for experimentally available energies. Equivalently the cutoff dependence in $m_i^0(\Lambda)/m_j^0(\Lambda)$ is small compared to the ratio itself and we can ignore it. We conclude that quark mass ratios in QCD, even in the presence of electromagnetism, are unambiguous up to very small and controllable corrections of $O(\alpha)$.

Having reached this conclusion there remains the problem of actually extracting the values of the quark mass ratios from experimental data. The remainder of this article will be devoted to this task. The technique we will use is the chiral perturbation theory previously developed by us (6,7). The basic observation of chiral perturbation theory is that many strong interaction amplitudes approach their chiral $SU(N) \times SU(N)$ values as $m_i \rightarrow 0$ in a non-analytic fashion such as $m_i \ln m_i$ or $m_i^{1/2}$. Chiral perturbation theory is a systematic calculational technique for calculating the coefficients of all such non-analytic terms which are usually the leading order terms in chiral symmetry breaking. The main assumptions are that these nonanalytic terms are the dominant contribution to symmetry breaking and that one can do perturbation theory in the chiral symmetry breaking parameters, the renormalized quark masses m_i . The current quark mass m_i is not a dimensionless parameter. A typical dimensionless parameter that makes its appearance in chiral perturbation theory is $\delta_i = \mu_i^2 / 16\pi^2 F_\pi^2$ where μ_i is a pseudoscalar mass that vanishes as $m_i \rightarrow 0$ and $F_\pi = 93$ MeV is the pion decay constant. As long as this parameter is small we have confidence that chiral perturbation theory is applicable.

For the lightest quarks, u and d, perturbation theory about the $SU(2) \times SU(2)$ limit is very good since $\delta_u \approx \delta_d \approx 0.014$. For the s quark $\delta_s \approx 0.1$ so

there is reason for less confidence in perturbation theory about chiral $SU(3) \times SU(3)$. However we remind the reader that in the Gell-Mann-Oakes-Renner⁽⁸⁾ scheme, which is consistent with all available information on chiral symmetry breaking⁽⁷⁾, the breaking of chiral $SU(3) \times SU(3)$ and ordinary $SU(3)$ are comparable, both breakings being generated by the strange quark mass m_s . Consequently the corrections to kaon PCAC, equivalent to chiral perturbation theory is m_s , and ordinary $SU(3)$ are comparable. Conversely any attempt to give up kaon PCAC will have to account for the success of $SU(3)$ symmetry in a new way. This seems to us to be an unattractive and unnecessary option although it is not strictly ruled out.

For heavy quarks like the c we have no reason to believe that chiral perturbation theory is applicable since $\gamma_c \approx 1$. It is difficult to suppose that the D mesons are approximate Nambu-Goldstone bosons like the π and K . In fact it is possible that as one increases a current quark mass there is a phase transition at a critical value $m = m_{c,r,T}$ beyond which there is no remnant of the Nambu-Goldstone states for that flavor. In any case we have no reliable way of determining the ratio of light (u,d,s) quark masses to heavy (c,t,b,\dots) quark masses with the methods available at this time. For this reason we restrict ourselves to examining the light quark mass ratios.

The origin of the quark mass spectrum is a deep problem. This problem, it is evident, is connected with the problem of establishing mixing angles like the Cabibbo angle and other mixing angles⁽⁹⁾ which arise upon diagonalizing the general quark mass matrix m_{ij} . A solution to this problem may come from the interplay of QCD and QFD in a unified theory and from the systematics of the quark vs. lepton spectrum. In the present state of the art of unified gauge

theories such connections remain speculative.

II. Quark Mass Ratios and Chiral Perturbation Theory

What are the motivations for examining the current quark mass spectrum?

There are at least two reasons: (a) In most weak interaction models (QFD) the current quark mass term is generated by spontaneous symmetry breaking of a local flavor group. A knowledge of mass ratios is useful in providing constraints on model building. (b) The vacuum of QCD will in general violate P and T invariance since one can add to \mathcal{L}^{QCD} the term $\Theta(g^2/32\pi^2)F_{\mu\nu} \times F_{\mu\nu}$ without altering renormalizability ⁽¹⁰⁾. This term gives a non-trivial contribution to the action from instantons if $\Theta \neq 2\pi n$ and will violate P and T symmetry. But this interaction term is not invariant under a chiral transformation on a single quark field, such as $u_R \rightarrow e^{i\alpha} u_R$, and could be rotated away if $\mathcal{L}^{\text{QCD}} + \Delta\mathcal{L}^{\text{QCD}}$ were invariant under this chiral transformation. This is possible providing $m_u^0 = 0$. While a massless up quark is not the only solution to the problem of potentially strong P and T violations ⁽¹¹⁾ it is of interest to see if this solution is phenomenologically viable.

Our plan is as follows. First we describe the problem of determining the $\Delta I = 1$ quark mass differences and then give the formulas for determining the quark mass ratios in terms of meson and baryon masses. We give a careful treatment of symmetry breaking which is controlled by chiral perturbation theory. We will discuss separately the isosinglet-isosinglet mass ratio $(m_d + m_u)/2m_s$ and the isovector-isosinglet mass ratio $(m_d - m_u)/2m_s$.

The ratio $(m_d + m_u)/2m_s$ of SU(2) x SU(2) to SU(3) x SU(3) chiral breaking

can be determined in chiral perturbation theory to lowest order to be given by $\mu_\pi^2/2\mu_K^2$. Including the complete leading order corrections from chiral perturbation theory we find

$$\frac{m_d + m_u}{2m_s} = 0.031 \quad (2.1)$$

The ratio $(m_d - m_u)/2m_s$ of isospin to SU(3) breaking is far more difficult to determine in a reliable way. The reason for this is that electromagnetism is known to violate isospin and require divergent counterterms in the quark mass matrix. Because these counterterms are divergent (in most QFD-theories) in the presence of electromagnetism there is a $m_u - m_d$ mass difference which is a free parameter. In practice the $m_u - m_d$ mass difference is determined from (i) the groundstate pseudoscalar meson mass difference $K^+ - K^0$ (ii) the $\Delta I = 1$ baryon mass differences (iii) the $\eta \rightarrow 3\pi$ decay. Below we will review the theoretical problem of defining the $m_u - m_d$ mass difference and the phenomenological problem of extracting it from the data.

From the meson mass spectrum we obtain

$$\frac{m_d - m_u}{2m_s} = 0.0175 \quad (2.2)$$

and from the baryon spectrum the consistent value

$$\frac{m_d - m_u}{2m_s} = 0.0105 \pm 0.003 \quad (2.3)$$

The extraction of this ratio from the $\eta \rightarrow 3\pi$ decay is sensitive to the details of $SU(3) \times SU(3)$ symmetry breaking (an extrapolation on the order of the η mass is required). So the $\eta \rightarrow 3\pi$ decay is inherently less reliable. If one ignores $SU(3) \times SU(3)$ symmetry breaking one obtains from the measured rate a ratio which is consistent with (2.2) and (2.3) obtained from the hadron mass spectrum. Depending on the treatment of symmetry breaking one can obtain results that improve or spoil the agreement with (2.2) and (2.3).

From (2.1) and (2.2) or (2.3) we have

$$\frac{m_u}{m_d} \approx 0.38$$

(2.4)

$$\frac{m_d}{m_s} \approx 0.045$$

which rules out a vanishing up quark mass. The main ways to avoid this conclusion are (i) The contribution of the low lying states to the Cottingham formula for the electromagnetic mass shifts of the hadrons is underestimated by a factor ≈ 4 (ii) Kaon PCAC or equivalently chiral $SU(3) \times SU(3)$ perturbation theory fails. If the latter is the case then we also lose the rationale for an approximate $SU(3)$ symmetry. Further it is not clear that if an alternative approach is adopted that it will lead to a consistent account of all symmetry breaking. We conclude that while a vanishing up quark mass is not rigorously ruled out it is unattractive from the standpoint of the presently consistent phenomenology of hadronic symmetry breaking.

A. Computational Procedure for $\Delta I = 1$ Mass Terms

The symmetry breaking part of QCD that we will consider is

$$\Delta \mathcal{L}^{\text{QCD}} = m_u^0 \bar{u}u + m_d^0 \bar{d}d + m_s^0 \bar{s}s \quad (2.5)$$

to which is to be added the conventional electromagnetic effective action

$$\mathcal{L}^{\text{EM}} = \frac{\alpha}{(2\pi)^3} \int d^4x D^{\mu\nu}(x) T^* (V_\mu^Q(\frac{x}) V_\nu^Q(-\frac{x})) \quad (2.6)$$

where V_μ^Q is the electromagnetic current and

$$D_{\mu\nu}(x) = - \int \frac{d^4q}{q^2 + i\epsilon} e^{iq \cdot x} (g_{\mu\nu} + \text{gauge terms}) \quad (2.7)$$

Before proceeding with calculations it is necessary to discuss the extraction of the electromagnetic mass shifts of quarks (tadpole) from the conventional electromagnetic interaction. This has been discussed in detail by J. Gasser and H. Leutwyler ⁽¹²⁾, J.F. Gunion ⁽¹³⁾ and more recently, from the viewpoint of the renormalization group, by J. Collins ⁽¹⁴⁾. We will simply review the conclusion of these articles.

First we note that $\Delta \mathcal{L}^{\text{QCD}}$ has no $\Delta I = 2$ part. On the other hand \mathcal{L}^{EM} , if

one examines the operator product expansion of the T^* product, has finite matrix elements for its $\Delta I = 2$ part. Therefore the Cottingham integral is convergent for $\Delta I = 2$ mass shifts like $\mu_{\pi^+}^2 - \mu_{\pi^0}^2$ and $m_{\Sigma^+} + m_{\Sigma^-} - 2m_{\Sigma^0}$. As is well known saturation of the Cottingham formula with low lying states for these combination yields excellent agreement with experiment (15). The $\Delta I = 2$ mass shifts are evidently well understood but they give no information on the quark mass spectrum.

By contrast the $\Delta I = 0, 1$ parts of \mathcal{L}^{EM} can be shown from the operator product expansion to give divergent contributions. These divergences are cancelled by the mass counterterms in $\Delta \mathcal{L}^{QCD}$ to yield finite results for symmetry breaking. While the subtraction of the logarithmically divergent part is unambiguous in a renormalizable theory like QCD there can remain a finite term after the subtraction. This finite term in the quark masses is ambiguous because it depends on the subtraction procedure used; different subtraction procedures treat finite parts differently. However this set of finite rescalings is precisely that which is controlled by the renormalization group and the ambiguity is due to the fact that the renormalization subtraction can be carried out at different renormalization masses. This has been discussed in detail by J. Collins (14).

To see how this work in practice let us regulate \mathcal{L}^{EM} by the introduction of a cutoff Λ in (2.7) according to the replacement $1/q^2 \rightarrow 1/q^2 - 1/(q^2 - \Lambda^2)$. Then if we consider the contribution of $\Delta \mathcal{L}^{QCD} + \mathcal{L}^{EM}$ to a physical mass shift ΔM it can be written as

$$\Delta M = (\Delta M)_{\Lambda}^{QCD} + (\Delta M)_{\Lambda}^{EM} \quad (2.8)$$

where the two terms correspond to the matrix elements of $\Delta\mathcal{L}^{\text{QCD}}$ and \mathcal{L}^{EM} respectively. Now as $\Lambda \rightarrow \infty$ we have

$$(\Delta M)_{\Lambda}^{\text{EM}} \xrightarrow{\Lambda \rightarrow \infty} \alpha C \ln(\Lambda/\mu) \quad (2.9)$$

with C a constant and μ the renormalization mass. By the renormalizability of QCD and QED ΔM must be Λ independent so that

$$(\Delta M)_{\Lambda}^{\text{QCD}} \xrightarrow{\Lambda \rightarrow \infty} -\alpha C \ln(\Lambda/\mu) \quad (2.10)$$

Defining the finite quantities

$$\begin{aligned} (\Delta M)^{\text{EM}} &= (\Delta M)_{\Lambda}^{\text{EM}} - \alpha C \ln(\Lambda/\mu) \\ (\Delta M)^{\text{TAD}} &= (\Delta M)_{\Lambda}^{\text{QCD}} + \alpha C \ln(\Lambda/\mu) \end{aligned} \quad (2.11)$$

we can split up the contributions to the physical mass shifts as

$$\Delta M = (\Delta M)^{\text{EM}} + (\Delta M)^{\text{TAD}} \quad (2.12)$$

where TAD means tadpole contribution. Here $(\Delta M)^{\text{EM}}$ represents the contribution of the low lying states, Born terms and resonances, before the

onset of scaling behavior in the structure functions in the Cottingham integral. The scaling part of the structure functions which gives the divergence in the Cottingham formula ⁽¹⁶⁾ has been explicitly subtracted out of the definition of $(\Delta M)^{EM}$ and is not to be included in numerical estimates.

The finite ambiguity mentioned above is simply reflected in the fact that we can change the numerical value of $(\Delta M)^{EM}$ or $(\Delta M)^{TAD}$ by a different choice of cutoff Λ or equivalently a different choice of renormalization point μ . This ambiguity we see from (2.12) is given by

$$\delta(\Delta M)^{TAD} = \alpha C \ln(\Lambda_1/\Lambda_2) \quad (2.13)$$

where Λ_1 and Λ_2 are two choices of cutoff. The constant C has been computed by Collins ⁽¹⁴⁾ with the result

$$C = \frac{(\Delta M)^{TAD}}{12\pi} \left(\frac{4m_u - m_d}{m_u - m_d} \right) \quad (2.14)$$

where $m_{u,d}$ are the renormalized quark masses. To estimate the ambiguity we choose $m_u/m_d \approx 1/3$, $\Lambda_1/\Lambda_2 = 100$ so that $\delta(\Delta M)^{TAD} = 4.4 \times 10^{-4} (\Delta M)^{TAD}$. The ambiguity due to change in cutoff by a factor of 100 is 10^{-4} compared to the tadpole term we are computing. Our conclusion is that we can extract the tadpole contribution from physical $\Delta I = 1$ mass differences if we subtract just the non-scaling Born and resonance terms in the Cottingham integral from the physical mass difference up to a small and negligible ambiguity.

B. Pseudoscalar Mesons

The groundstate pseudoscalar mesons are a good source of information about the current quark mass ratios. The η and η' can mix and this introduces an additional parameter rendering this system useless for our problem of determining quark mass ratios. This leaves the π and K system.

We use the standard method invoking the PCAC relation for the axial currents $A_\mu^a(x)$, $a=1, \dots, 8$.

$$\partial_\mu A_\mu^a(x) = i [{}^5Q^a, \mathcal{L}] \quad (2.15)$$

where $\mathcal{L} = \mathcal{L}^{\text{QCD}} + \Delta \mathcal{L}^{\text{QCD}} + \mathcal{L}^{\text{EM}}$ with $\Delta \mathcal{L}^{\text{QCD}}$ and \mathcal{L}^{EM} given by (2.5) and (2.6). \mathcal{L}^{QCD} is chiral invariant so $[{}^5Q^a, \mathcal{L}^{\text{QCD}}] = 0$. We will take the matrix element of (2.15) between the vacuum and the physical π and K states.

With the definitions

$$\begin{aligned} \langle 0 | A_\mu^{\pi^+}(0) | \pi^+(k) \rangle &= i \sqrt{2} F_{\pi^+} k_\mu \\ \langle 0 | \bar{d} \gamma_5 u | \pi^+ \rangle &= \sqrt{2} Z_{\pi^+}^{1/2}, \text{ etc.} \end{aligned} \quad (2.16)$$

(in the chiral limit the Z's are proportional to $Z(1/\mu)$ of (1.2))

we obtain the equations

$$\begin{aligned} F_{\pi^+} \mu_{\pi^+}^2 &= Z_{\pi^+}^{1/2} (m_d^0 + m_u^0) + F_{\pi^+} (\delta \mu_{\pi^+}^2)^{\text{EM}} \\ F_{\pi^0} \mu_{\pi^0}^2 &= Z_{\pi^0}^{1/2} (m_d^0 + m_u^0) + F_{\pi^0} (\delta \mu_{\pi^0}^2)^{\text{EM}} \\ F_{K^+} \mu_{K^+}^2 &= Z_{K^+}^{1/2} (m_s^0 + m_u^0) + F_{K^+} (\delta \mu_{K^+}^2)^{\text{EM}} \\ F_{K^0} \mu_{K^0}^2 &= Z_{K^0}^{1/2} (m_s^0 + m_d^0) + F_{K^0} (\delta \mu_{K^0}^2)^{\text{EM}} \end{aligned} \quad (2.17)$$

where $(\delta\mu^2)^{EM}$ is the contribution of \mathcal{L}^{EM} and is explicitly of order α . We have dropped the π^0 - η mixing terms in the π^0 equation, which are second order in isospin breaking. Otherwise, (2.17) is exact. The quantities $z^{1/2}$, m^0 , $(\delta\mu^2)^{EM}$ are all divergent. After the renormalizations and ratios are taken everything will be finite as we have previously discussed. To extract information from (2.17) we will make the safe assumption of (i) dropping terms second order in isospin breaking relative to terms first order in isospin breaking, (ii) dropping terms first order in isospin breaking relative to terms first order in SU(3) breaking, and (iii) using $\delta\mu_{\pi^+}^2/\mu_{\pi^+}^2 = 3\alpha \ln 2 m_0^2/4\pi\mu_{\pi^+}^2 + O(\alpha \ln \mu_{\pi^+}^2)$ we find we can drop $\delta\mu_{\pi^+}^2$ relative to $\mu_{\pi^+}^2$. For the quark mass ratios we can replace the bare with the renormalized values and we obtain

$$\frac{m_d + m_u}{2m_s} = \frac{\mu_{\pi^+}^2 (1-\varepsilon)}{2\mu_K^2 - \mu_{\pi^+}^2 (1-\varepsilon)} \quad (2.18)$$

$$\frac{m_u - m_d}{2m_s} = \frac{\mu_{K^+}^2 - \mu_{K^0}^2 - (\Delta\mu_K^2)^{EM}}{2\mu_K^2 - \mu_{\pi^+}^2 (1-\varepsilon)} - \frac{\mathcal{J}_K}{2}$$

and

$$\mu_{\pi^+}^2 - \mu_{\pi^0}^2 = (\Delta\mu_{\pi^+}^2)^{EM} + \mathcal{J}_{\pi} \mu_{\pi^+}^2 \quad (2.19)$$

where

$$\begin{aligned}
 \delta_{\pi} &= 1 - \frac{Z_{\pi^0}^{1/2} F_{\pi^+}}{Z_{\pi^+}^{1/2} F_{\pi^0}} \\
 \delta_K &= 1 - \frac{Z_{K^0}^{1/2} F_{K^+}}{Z_{K^+}^{1/2} F_{K^0}} \\
 \varepsilon &= 1 - \frac{Z_K^{1/2} F_{\pi}}{Z_{\pi}^{1/2} F_K}
 \end{aligned}
 \tag{2.20}$$

and

$$\begin{aligned}
 (\Delta M_K^2)^{EM} &= (\delta M_{K^+}^2)^{EM} - (\delta M_{K^0}^2)^{EM} \\
 (\Delta M_{\pi}^2)^{EM} &= (\delta M_{\pi^+}^2)^{EM} - (\delta M_{\pi^0}^2)^{EM}
 \end{aligned}
 \tag{2.21}$$

are the conventional photonic contribution to the meson mass differences from just the low lying states in the Cottingham formula. In equation (2.20) $\delta_{\pi, K}$ are of the order of isospin violation while ε is of the order of SU(3) violations. If we also drop terms of the order of isospin violation times chiral SU(2) x SU(2) violation relative to isospin violations (2.19) implies the well known result

$$\mu_{\pi^+}^2 - \mu_{\pi^0}^2 = (\Delta M_{\pi}^2)^{EM}
 \tag{2.22}$$

To determine the quark mass ratios from (2.18) we need to know $(\Delta M_K^2)^{EM}$, δ_K and ε . These quantities cannot be obtained directly from experiment and one is required to make theoretical assumptions.

The contribution from the low lying states to the kaon mass difference

$(\Delta M_K^2)^{EM} \approx 2M_K (\Delta M_K)^{EM}$ was estimated by R. Socolow ⁽¹⁷⁾ in 1965 with the result

$$(\Delta M_K)^{EM} \approx 3 \text{ MeV (Socolow)}. \quad (2.23)$$

Recently use has been made of the Dashen sum rule ⁽¹⁸⁾ which implies

$$(\Delta M_K^2)^{EM} = (\Delta M_\pi^2)^{EM} \quad (2.24)$$

which upon using (2.22) implies

$$(\Delta M_K)^{EM} \approx 1.5 \text{ MeV (Dashen Sum Rule)}$$

However it has been emphasized by us ⁽¹⁹⁾ that the Dashen sum rule is obtained in perturbation theory. The corrections to the Dashen sum rule are as large as the terms one is retaining and hence the assumption upon which it was obtained is false. This could be the explanation for the discrepancy between the Socolow calculation and the Dashen sum rule. In view of this we will use Socolow's estimate (2.23) for $(\Delta M_K)^{EM}$.

Since the quantities f_K and ϵ involve the ratio $Z_K^{1/2}/Z_\pi^{1/2}$, etc. they cannot be obtained from any experiment. However if we assume the validity of chiral perturbation theory then these quantities can be calculated exactly, independent of any model, to leading order in the chiral symmetry breaking. The results of these calculations are ^(6,7)

$$\frac{F_K}{F_\pi} = 1 + \frac{3(M_K^2 - M_\pi^2)}{64\pi^2 F_\pi^2} \ln(M^2/\mu^2)$$

$$\frac{Z_K^{1/2}}{Z_\pi^{1/2}} = 1 + \frac{(M_K^2 - M_\pi^2)}{192\pi^2 F_\pi^2} \ln(M^2/\mu^2)$$

$$\frac{Z_{K^+}^{1/2}}{Z_{K^0}^{1/2}} = 1 + \frac{(\Delta M_K^2)^{TAD}}{192\pi^2 F_\pi^2} \ln(M^2/\mu^2) \quad (2.25)$$

$$\frac{F_{K^+}}{F_{K^0}} = 1 + \frac{3(\Delta M_K^2)^{TAD}}{64\pi^2 F_\pi^2} \ln(M^2/\mu^2)$$

so

$$\delta_K = - \frac{(\Delta M_K^2)^{TAD}}{24\pi^2 F_\pi^2} \ln(M^2/\mu^2)$$

$$\varepsilon = \frac{(M_K^2 - M_\pi^2)}{24\pi^2 F_\pi^2} \ln(M^2/\mu^2)$$

where $(\Delta M_K^2)^{TAD} = M_{K^+}^2 - M_{K^0}^2 - (\Delta M_K^2)^{EM}$. Here $F_\pi = 93$ MeV, M is a cutoff we will take to be the nucleon mass and $\mu^2 = \text{mean (mass)}^2$ of the groundstate pseudoscalars ≈ 0.17 (GeV)². Using those numbers and (2.23) we obtain from (2.25)

$$\frac{F_K}{F_\pi} = 1.21 \quad \frac{Z_K^{1/2}}{Z_\pi^{1/2}} = 1.023 \quad \varepsilon = 0.18 \quad (2.26)$$

$$\frac{F_{K^+}}{F_{K^0}} - 1 = -6.3 \times 10^{-3} \quad \frac{Z_{K^+}^{1/2}}{Z_{K^0}^{1/2}} - 1 = -7.0 \times 10^{-4} \quad \delta_K = 5.6 \times 10^{-3}$$

The above calculation in chiral perturbation theory can be checked against experiment for the quantity F_K/F_π . From $K_{\ell 3}$ decay one finds

$$\frac{F_K}{F_+(0)F_\pi} = 1.26 \pm 0.02 \quad (2.27)$$

with $f_+(0)$ the $K_{\ell 3}$ decay vector form factor. From the nonrenormalization theorem⁽²⁰⁾ we know that $f_+(0)$ is near 1.0. An explicit calculation of the exact first order correction in chiral symmetry breaking gives⁽²⁰⁾ $f_+(0) = 0.97$ so that we obtain

$$\frac{F_K}{F_\pi} = 1.22 \pm 0.02 \quad (2.28)$$

in excellent agreement with the calculated ratio (2.26). This gives us confidence that our other estimates in (2.26) are roughly correct even though they cannot be directly obtained from experiment.

Using the results of chiral perturbation theory (2.26) and Socolow's estimate of the kaon electromagnetic mass difference (2.23) the quark mass ratios can be gotten from (2.18) with the result

$$\frac{m_d + m_u}{2m_s} = 0.031 \quad (0.038) \quad (2.30)$$

$$\frac{m_d - m_u}{2m_s} = 0.0175 \quad (0.015)$$

where the numbers in parenthesis are what would have been obtained if we ignored the corrections of chiral perturbation theory and set $\xi = 0$, $\delta_K = 0$. Equation (2.30) implies

$$\frac{m_u}{m_d} = 0.28 \quad (0.44) \quad (2.31)$$

so the up quark has nonvanishing mass.

Some attempts have been made to obtain $m_u = 0$. Since $Z_K^{1/2}/Z_\pi^{1/2}$ is not obtainable from experiment some authors ⁽²¹⁾ have supposed $Z_K^{1/2}/Z_\pi^{1/2} \approx 2-3$, over 100 % different from the result of chiral perturbation theory. Such an assumption implies huge violations of ordinary SU(3) symmetry. Dominguez ⁽²²⁾ has pointed out that the ratio $Z_K^{1/2}/Z_\pi^{1/2}$ makes its appearance in the non-renormalization theorem for $f_+(0)$ and a value very different from 1.0 would wreck the agreements of Cabibbo theory. Further if $Z_K^{1/2}/Z_\pi^{1/2}$ was wildly different from unity there is no good reason to suppose that a similar ratio of wavefunction renormalization constants that makes its appearance in the baryon mass difference calculations should be near unity. Then one loses the Gell-Mann Okubo formula. It seems to us unlikely that if $Z_K^{1/2}/Z_\pi^{1/2}$ differs greatly from the result of chiral perturbation theory that the impressive agreement of broken SU(3) with experiment could be maintained.

Soper and Deshpande ⁽²³⁾ have obtained $m_u = 0$ by assuming $\delta_K = 0.036$. This assumption represents a huge violation of isospin. The value of δ_K is almost an order of magnitude greater than what we have calculated from chiral perturbation theory $\delta_K = 0.0056$. We think that the successes of chiral perturbation theory do not warrant such a large value of δ_K although it

is not impossible that there is some special dynamical enhancement of δ_K to which the chiral perturbation theory is blind. In the absence of a demonstration of such enhancement it is not possible to justify the large value of δ_K .

We finally remark that the quark mass ratios we obtain are not very sensitive to the precise values of the chiral perturbation theory corrections ε , δ_K providing they are within 50 % of the values we obtain and not different by orders of magnitude.

C. Baryons

From the baryon mass spectrum it is possible to establish an estimate of the ratio $(m_d - m_u) / 2m_s$. To do so one must extract the tadpole contribution from the observed isotopic splittings according to

$$(\Delta M)_{ij} = (\Delta M)_{ij}^{EM} + (\Delta M)_{ij}^{TAD}$$

$$(\Delta M)_{ij} \equiv m_i - m_j$$

The low lying resonance and Born term contribution to $(\Delta M)_{ij}^{EM}$ from the Cottingham formula were calculated by S. Coleman and H. Schnitzer (24) for all the baryons. There have been many calculations of $(\Delta M)_{pn}^{EM}$ and these generally agree with each other (15). The results of these calculations are listed in table I along with the experimental mass differences and the calculated tadpole terms $(\Delta M)_{ij}^{TAD}$.

In order to calculate the ratio $(m_d - m_u)/2m_s$ we will (i) drop second order SU(3) violations relative to first order (ii) drop SU(2) x SU(2) violations relative to SU(3) violations so $m_s \gg m_{d,u}$. Then by considering the matrix elements of the divergence of the vector currents $\partial_\mu V_\mu^a = i[Q^a, \mathcal{L}]$ between baryon states one obtains for the $\Delta I = 1$ mass differences

$$\begin{aligned} (\Delta M)_{pn}^{TAD} &= M \left(1 + \frac{F}{D}\right) \\ (\Delta M)_{\Sigma^+ \Sigma^-}^{TAD} &= 2M \frac{F}{D} \\ (\Delta M)_{\Xi^0 \Xi^-}^{TAD} &= M \left(\frac{F}{D} - 1\right) \end{aligned} \quad (2.32)$$

where F/D is the F/D ratio for the baryon mass matrix.

The three $\Delta I = 1$ mass differences in (2.32) are not all independent in this model of symmetry breaking. First one finds from (2.32) the Coleman-Glashow formula

$$(\Delta M)_{pn}^{TAD} + (\Delta M)_{\Xi^0 \Xi^-}^{TAD} + (\Delta M)_{\Sigma^+ \Sigma^-}^{TAD} = 0 \quad (2.33)$$

which is satisfied within experimental errors. Second one obtains the F/D ratio from the tadpole splittings as

$$\frac{F}{D} = \frac{(\Delta M)_{\Sigma^+ \Sigma^-}^{TAD}}{(\Delta M)_{\Xi^0 \Xi^-}^{TAD} + (\Delta M)_{pn}^{TAD}} = -2.7 \quad (2.34)$$

which should be equal to the F/D from the medium strong splittings

$$\frac{F}{D} = \frac{2}{3} \frac{M_{\Xi} - M_N}{M_{\Lambda} - M_{\Sigma}} = -3.3 \quad (2.35)$$

since it is the same octet that is responsible for both splittings. These ratios differ by 18 % in (2.34) and (2.35) and this gives us an idea of the uncertainties in our calculations of the tadpole terms.

Finally we can compute the constant \mathcal{M} in terms of the ratio of quark masses with the result

$$\mathcal{M} = \frac{3}{2} \left(\frac{m_d - m_u}{2m_s} \right) (M_{\Sigma} - M_{\Lambda}) \quad (2.36)$$

Using this and the Gell-Mann Okubo formula, $2(M_N + M_{\Xi}) = 3M_{\Lambda} + M_{\Sigma}$, we obtain from (2.36) and (2.32) the result

$$\frac{m_d - m_u}{2m_s} = \frac{1}{2} \frac{(\Delta M)_{np}^{TAD} + (\Delta M)_{\Xi^0 \Xi^-}^{TAD}}{M_{\Xi} + M_N - 2M_{\Sigma}} \quad (2.37)$$

From our previous analysis of the pseudoscalar mesons we recognize the left hand side of (2.37) as $(\Delta\mu_{K^0 K^+}^2)^{TAD} / 2\mu_K^2$ to leading order in chiral breaking so that

$$\frac{(\Delta M_{K^0 K^+})^{TAD}}{M_K^2} = \frac{(\Delta M)_{np}^{TAD} + (\Delta M)_{\Xi^0 \Xi^-}^{TAD}}{M_{\Xi} + M_N - 2M_{\Sigma}} \quad (2.38)$$

which is just the hybrid sum rule first obtained by S. Coleman and S. Glashow⁽²⁵⁾.

The point of these remarks is that if the quark mass matrix model of symmetry breaking is correct it is equivalent to the old tadpole model and so no really independent estimate of $m_d - m_u / 2m_s$ is obtained from the baryons. However we get an idea of the uncertainties in our calculations by seeing if the baryon mass spectrum gives a consistent answer for the ratio $m_d - m_u / m_s$. From (2.37) we find

$$\frac{m_d - m_u}{2m_s} = 0.0105 \pm 0.003 \quad (2.38)$$

with an allowed 30 % uncertainty since the F/D ratio comparison is good only to 18 % and SU(3) is good to only ≈ 15 %. This result is not inconsistent with our analysis of the mesons which gave, $(m_d - m_u) / 2m_s = 0.0175$.

Allowing for experimental and calculational uncertainties we conclude from our analysis of the mesons and baryons

$$\frac{m_d - m_u}{2m_s} = 0.014 \pm 0.003$$

$$\frac{m_d + m_u}{2m_s} = 0.031 \pm 0.007$$

(2.39)

From this we obtain

$$\frac{m_u}{m_d} = 0.38 \pm 0.13$$

$$\frac{m_d}{m_s} = 0.045 \pm 0.011 \tag{2.40}$$

as our estimate of the quark mass ratios.

To improve this analysis we suggest that new estimates of the photonic contributions to kaon and baryon isotopic splittings be undertaken to check the calculations done over ten years ago (17,24). Although the chiral perturbation theory corrections to the pseudoscalar mesons have been done and provide important $\approx 20\%$ corrections to the symmetric estimates a similar calculation taking into account second order SU(3) breaking in the baryon spectrum has not been done. This might improve the agreement between the meson and baryon estimate of $m_d - m_u / 2m_s$.

D. The decay $\eta \rightarrow 3\pi$

The purely photonic contributions to the $\eta \rightarrow 3\pi$ width vanishes in the chiral SU(2) x SU(2) limit (Sutherland's theorem (20)) and for broken SU(2) x SU(2) has been estimated (27) to be two orders of magnitude smaller than the experimental width. The decay can, however, be induced by the isospin violating quark mass terms (which also induce $\pi^0 - \eta$ mixing) and these are required in any case as counterterms to the usual electromagnetic interaction in QCD (28).

The amplitude ⁽²⁹⁾ for the $\eta \rightarrow \pi^+\pi^-\pi^0$ decay is parametrized as

$$T_{+-0} = A + BE_0 \quad (2.41)$$

From the central value of the decay width ⁽³⁰⁾ $\Gamma_{+-0} \approx 203$ eV and various experimental values for B/A one finds

$$0.57 \leq |A| \leq 0.70 \quad (2.42)$$

Assuming the decay is due to the term $(m_u^c - m_d^c)(\bar{u}u - \bar{d}d)$ and using current algebra in the SU(2) x SU(2) limit one finds $B/A \rightarrow 1/\mu_\eta$ in good agreement with the observed slope $B/A \approx -2.1/\mu_\eta$. Using current algebra in the SU(3) x SU(3) limit one obtains ⁽³¹⁾

$$A = \left(\frac{m_u - m_d}{2m_s} \right) \frac{8\mu_K^2}{3\sqrt{3}F_\pi^2} \quad (2.43)$$

Assuming the sign of A is negative we obtain from the experimental range (2.42) the result

$$\frac{m_d - m_u}{2m_s} = 0.0145 \pm 0.0015 \quad (2.44)$$

in good agreement with our previous estimate (2.39) based on the meson and baryon spectrum.

The range represented in (2.44) only takes into account the experimental range (2.42) and not uncertainties associated with $SU(3) \times SU(3)$ symmetry breaking. The effects of symmetry breaking in this amplitude can be considerable since extrapolations on the order of μ_η^2 are required. In a previous publication⁽²⁹⁾ we have found a chiral symmetry correction factor $\approx 34\%$ of $O(\mu_K^2 \ln \mu_K^2)$ which suggests that corrections of $O(\mu_K^2)$ can be large also. So it is not inconceivable that symmetry breaking effects could dramatically alter the estimate (2.44) which is consistent with the meson and baryon spectrum. Another analysis of symmetry breaking in $\eta \rightarrow 3\pi$ based on extended PCAC has been done by C. Dominguez and A. Zepeda⁽³²⁾. They find that quark mass ratios so obtained are compatible with those obtained from the meson and baryon spectrum and have results consistent with ours.

We conclude that the estimate of quark mass ratios from the $\eta \rightarrow 3\pi$ decay is not inconsistent with that obtained from the meson and baryon spectrum but that the estimate is sensitive to potentially large symmetry breaking corrections which are difficult to take into account reliably.

III. Conclusions

The advent of QCD as a theory of the strong interactions has placed the problem of strong interaction symmetry breaking on a rational basis. Current quark mass ratios are renormalization group invariants up to small and controllable terms coming from QFD interactions. Further we see that the old wisdom of subtracting just the lowest lying Born and resonance (non-scaling) contributions to the Cottingham formula from the observed isotopic mass difference and identifying the remainder as the tadpole can be justified

using the renormalization group.

To actually extract the quark mass ratios from experimental data we have used (i) the meson spectrum (ii) the baryon spectrum (iii) the $\eta \rightarrow 3\pi$ decay. Beyond this one must make theoretical assumptions. The main theoretical assumptions are that (i) the low lying states in the Cottingham formula contributing to $(\Delta M)^{EM}$ have been correctly estimated (ii) chiral perturbation theory - equivalently kaon PCAC - correctly estimates the leading order corrections to the symmetric results. These assumptions, we emphasize, lead to a completely consistent and coherent account of all hadronic symmetry breaking.

We have done a careful analysis of the groundstate pseudoscalar mass spectrum taking into account all leading order chiral corrections to the quark mass ratios $m_d + m_u/2m_s$ and $m_d - m_u/2m_s$. The baryon mass spectrum and $\eta \rightarrow 3\pi$ yielded values for these ratios consistent with those from the meson spectrum and we concluded

$$\frac{m_u}{m_d} = 0.38 \pm 0.13$$

$$\frac{m_d}{m_s} = 0.045 \pm 0.011$$

These ratios are consistent with those recently obtained by S. Weinberg ⁽³³⁾, C. Dominguez and A. Zepeda ⁽³²⁾ and earlier by J. Gasser and H. Leutwyler ⁽¹²⁾ and inconsistent with those estimates which obtain a zero mass up quark.

Acknowledgements

One of us (HP) would like to thank J. Collins for making the results of his work on the renormalization group and the quark mass problem available before publication and the Aspen Center for Physics, where this work was completed, for its hospitality. The other (PL) would like to thank the theoretical groups at DESY and CERN for their hospitality, and the Department of Energy, contract EY-76-C-07-3071, for support.

Mass Difference	(ΔM)	$(\Delta M)^{EM}$	$(\Delta M)^{TAD} = (\Delta M) - (\Delta M)^{EM}$
$p - n$	- 1.3	1.1	- 2.4
$\Sigma^+ - \Sigma^-$	- 8.0	- 0.7	- 7.3
$\Xi^0 - \Xi^-$	- 6.4	- 1.3	- 5.1
$\Sigma^+ + \Sigma^- - 2\Sigma^0$	1.8	2.1	0

Table I. Baryon Mass Differences in MeV (to nearest 0.1 MeV)

References

1. QCD as a theory of the strong interaction has been reviewed by W. Marciano and H. Pagels, Physics Rep. 36C, 137 (1978).
2. A description of this mechanism in QCD is given in D.G. Caldi, Phys. Rev. Lett. 39, 121 (1977) and C. Callen, R. Dashen and D.J. Gross, Phys. Lett. 66B, 375 (1977); Phys. Rev. D17, 2717 (1978).
3. R. Jaffe and C. Llewellyn-Smith, Phys. Rev. D7, 2506 (1973).
4. J. Collins, Princeton University Preprint (1978).
5. S. Weinberg, Phys. Rev. D8, 3497 (1973), *ibid* 605, 4482 (1973) G. 't Hooft, Nucl. Phys. B61, 455 (1973), H.D. Politzer, Nucl. Phys. B117, 397 (1976), H. Georgi and H.D. Politzer, Phys. Rev. D14, 1829 (1976).
6. P. Langacker and H. Pagels, Phys. Rev. D8, 4595 (1973).
7. Chiral perturbation theory is reviewed in H. Pagels, Physics Rep. 16C, 219 (1975).
8. M. Gell-Mann, R. Oakes and B. Renner, Phys. Rev. 175, 2195 (1968).
9. An early version of obtaining the Cabibbo angle from the quark mass matrix is in H. Pagels, Phys. Rev. D11, 1213 (1975) and more recent work based on gauge theories is S. Weinberg, reference 33; F. Wilczek and A. Zee, Phys. Lett. 70B, 418 (1977), 72B, 504 (1978); H. Fritzsch, Phys. Lett. 73B, 317 (1978)
10. G. 't Hooft, Phys. Rev. Lett. 37, 8 (1976); Phys. Rev. D14, 3432 (1976)
11. R.D. Peccei and H.R. Quinn, Phys. Rev. Lett. 38 1440 (1970); S. Weinberg, Phys. Rev. Lett. 40, 223 (1978); F. Wilczek, Phys. Rev. Lett. 40, 279 (1978).

12. J. Gasser and H. Leutwyler, Nucl. Phys. B94, 269 (1975).
13. J.F. Gunion, Phys. Rev. D8, 517 (1973).
14. J. Collins, "Renormalization of the Cottingham Formula", Princeton University preprint 1978.
15. The problem of electromagnetic mass differences has been reviewed by A. Zee, Phys. Rep. 3C, 129 (1972).
16. See Appendix 0 of reference 15 and references therein.
17. R. Socolow, Phys. Rev. 137B, 1221 (1965).
18. R. Dashen, Phys. Rev. 183, 1245 (1969); see also M. Weinstein, Phys. Rev. D4, 2544 (1972).
19. P. Langacker and H. Pagels, Phys. Rev. D8, 4620 (1973)
20. The non-renormalization theorem was proved in SU(3) by M. Ademollo and R. Gatto, Phys. Rev. Lett. 13, 264 (1965) and by R.E. Behrends and A. Sirlin, Phys. Rev. Lett. 4, 186 (1960).
It was extended to chiral SU(3) x SU(3) by P. Langacker and H. Pagels, Phys. Rev. Lett. 30, 630 (1972); S. Wada, Phys. Lett. 49B, 175 (1974); M. Bace and D.T. Cornwell, Phys. Rev. D10, 2694 (1974).
21. A. Zepeda, Centro de Investigacion, (Mexico City) preprint (1978).
22. C.A. Dominguez, Centro de Investigacion, (Mexico City) preprint 1978.
23. D. Soper and N.G. Deshpande, University of Oregon preprint 1978.
24. S. Coleman and H. Schnitzer, Phys. Rev. 136, B 223 (1964).
25. S. Coleman and S. Glashow, Phys. Rev. 134, B671 (1964).

26. D.G. Sutherland, Phys. Lett. 23, 384 (1966); J.S. Bell and D.G. Sutherland, Nucl. Phys. 4B, 315 (1968).
27. W.A. Bardeen, L.S. Brown, B.W. Lee and H.T. Nieh, Phys. Rev. Lett. 18, 1170 (1967); A.D. Dolgov, A.I. Vainshtein and V.I. Zakhanov, Phys. Lett. 24B, 425 (1967).
28. The original suggestion of a u_3 term is due to S.K. Bose and A.H. Zimmermann, Nuovo Cimento 43A, 1165, (1966); see also K. Wilson, Phys. Rev. D2, 1478 (1970); 179, 1499 (1963).
29. P. Langacker and H. Pagels, Phys. Rev. D10, 2904 (1974) and references therein.
30. A. Browman et al., Phys. Rev. Lett. 32, 1067 (1974).
31. This result can be gotten from the calculated value $A = -4\varepsilon_3/3\sqrt{3}F_\pi^3$ in reference 29 and the relations $\varepsilon_3 = (\sqrt{3}\varepsilon_0 - (2/\sqrt{3})\varepsilon_8)(m_u - m_d)/2m_s$, $\varepsilon_0 = -\sqrt{2/3}F_K/M_K^2$, $\varepsilon_8/\varepsilon_0 = -\sqrt{2}$ valid to lowest order in SU(3) x SU(3) breaking.
32. C.A. Dominguez and A. Zepeda, Centro de Investigacion (Mexico City) preprint (1978).
33. S. Weinberg, in Festschrift for I.I. Rabi, ed. by Lloyd Motz (N.Y. Academy of Sciences, N.Y. 1977).