

19-11-14  
高工研圖書室

# DEUTSCHES ELEKTRONEN-SYNCHROTRON **DESY**

DESY 79/62  
October 1979

LONGITUDINAL CROSS SECTION AND ASYMMETRIES

FOR JETS IN LEPTOPRODUCTION

by

Ch. Rumpf and G. Kramer

*II. Institut für Theoretische Physik der Universität Hamburg*

NOTKESTRASSE 85 2 HAMBURG © 52

To be sure that your preprints are promptly included in the  
HIGH ENERGY PHYSICS INDEX,  
send them to the following address ( if possible by air mail ) :

DESY  
Bibliothek  
Notkestrasse 85  
2 Hamburg 52  
Germany

Longitudinal Cross Section and Asymmetries  
for Jets in Leptoproduction +)

Ch. Rumpf and G. Kramer

II. Institut für theoretische Physik der  
Universität Hamburg

So far most of the theoretical predictions on jet properties based on quantum chromodynamics (QCD) are for  $e^+e^-$  annihilation [1]. In leptoproduction previous theoretical work is mainly on one-particle inclusive processes [2]. From this it is known that first order contributions in the QCD coupling  $\alpha_s(Q^2)$  lead to angular asymmetries of singly produced hadrons relative to the lepton scattering plane. But these predictions depend on poorly known quark and gluon fragmentation functions. Furthermore, nonperturbative effects, in particular the effect of a nonvanishing transverse momentum of the incoming partons, produce similar asymmetries. From this point of view, direct predictions on global jet properties, whether as distributions in the Sterman-Weinberg cut-off variables  $(\epsilon, \delta)$  [3] in the summed transfer momentum of hadrons variable  $\pi_T = \sum_{hadrons} |\vec{p}_T|$  [4] or in the jet variables thrust  $\mathcal{T}$  and/or sphericity  $\mathcal{S}$  [5] seem more useful. In particular, by making cuts with these variables, it is easier to eliminate from the distributions unwanted two-jet contributions, which, if dressed by nonperturbative effects, dominate the distributions for small  $(\epsilon, \delta)$ ,  $\pi_T$ ,  $1-\mathcal{T}$  or  $\mathcal{S}$ . These distributions become narrower and n-arrows with increasing  $W$ , where  $W$  is the total available energy in the final state. This way one can hope that genuine QCD effects due to gluon emission and absorption become visible if  $W$  is large enough.

**Abstract:**  
We have calculated the longitudinal and other polarization dependent cross sections for jet production in deep inelastic electron-proton scattering up to order  $\alpha_s$  of the quark-gluon coupling constant and compared them with estimates of the non-perturbative contributions.

Recently experimental results for  $d\sigma/d\Omega$  and  $d\sigma/d\Omega dS$  have been reported for deep inelastic neutrino scattering for  $W \approx 10 \text{ GeV}$  [6]. These data are still in the nonperturbative region and the shape of these distributions can be very well described by known fragmentation models [7]. Data with larger center-of-mass energies  $W$  may become available with higher neutrino energies in the near future or with ep storage rings now under discussion [8].

In general, with unpolarized target and beam, the inclusive single jet cross-section can be decomposed into four separate terms which can be classified by the polarization of the absorbed virtual photon. This is well-known from the more familiar inclusive single particle cross-section [9]. These four terms can be disentangled by varying the azimuthal angle of the detected jet and the degree of the polarization of the virtual photon.

+ ) Supported by Deutsche Forschungsgemeinschaft  
Grant Kr 671/1

In this letter we study these four cross sections as a function of the thrust variable  $T$  up to the first order of the quark-gluon coupling  $g = (\frac{4\pi}{3} \alpha_s(Q^2))^{1/2}$ . We hope that the separate determination of all four cross-sections will give a further handle for the verification of QCD in the spacelike region.

First we consider the three-jet contribution proportional to  $\alpha_s(Q^2)$ , represented by the diagrams in Fig. 1b, c and d. The lowest order diagram, Fig. 1a, will be considered later. The three final state jets are the partons, quarks, antiquarks or gluons with momentum  $p_1$  and  $p_2$  and the spectator jet with momentum  $p_3$ . We use the final hadronic rest frame  $\vec{p} + \vec{p} = 0$  and use as variables  $W^2 = (p_1 + p_2)^2$ ,  $Q^2$  and  $X_i = 2p_{i0}/W$  ( $i=1,2,3$ ) with the constraint  $X_1 + X_2 + X_3 = 2$  (see Fig. 1e for the notation of momenta). As usual we define  $y = \frac{p_T}{p_0}$  and the angle  $\varphi$  as the azimuthal angle of the scattered electron relative to the plane defined by the final parton momenta. The dependence of the cross-section on  $\varphi$  and  $y$  for electron-nucleon scattering (in the one-photon exchange approximation) is of the following form<sup>†</sup>

$$\frac{d^5}{dQ^2 dW^2 d\varphi dx_1 dx_2} = T \left\{ \frac{d^2\sigma_U}{dx_1 dx_2} + \frac{2(1-y)}{1+(1-y)^2} \frac{d^2\sigma_T}{dx_1 dx_2} + \frac{2(1-y)}{1+(1-y)^2} \cos 2\varphi \frac{d^2\sigma_T}{dx_1 dx_2} - \frac{(1-y)^2(2-y)}{1+(1-y)^2} \cos\varphi \frac{d^2\sigma_T}{dx_1 dx_2} \right\} \quad (1)$$

$T$  is the well-known equivalent photon spectrum

$$T = \frac{\alpha W^2}{4\pi^2 Q^2 (W^2 - Q^2)^2} (1 + (1-y)^2) \quad (2)$$

The term  $d\sigma_U$  is the cross section for unpolarized transverse virtual photons,  $d\sigma_T$  is due to the transverse linear polarization of the photon,  $d\sigma_L$  accounts for the longitudinal polarized photons and  $d\sigma_I$  describes the interference of the longitudinal and transverse matrix element.

†) We consider  $ep$  scattering only. The case of neutrino scattering is studied in a separate paper.

The total cross-section is written in the form

$$\frac{d^2\sigma}{dQ^2 dW^2 d\varphi} = T \sigma_U \quad (3)$$

The  $\sigma_U$  in zeroth order is

$$\sigma_U^0 = \frac{\pi e^2 \alpha_s}{W^2} f_a(X_B, Q^2) \quad (4)$$

Here  $f_a(X_B, Q^2)$  is the distribution of parton a inside the nucleon with the fraction of momentum  $X_B = Q^2/(Q^2 + W^2)$  and  $Q_a$  is the charge of parton a. The derivation of the three-jet cross-section is straightforward. For the diagrams in Fig. 1b we obtain:<sup>†</sup>

$$(\sigma_U^0)^{-1} \frac{d^2\sigma_U}{dx_1 dx_2} = C_1 \frac{W^2}{X_{11} X_{13} X_3} \left\{ \frac{W^2}{\eta(W^2 + Q^2)} X_3^2 (X_{11}^2 + X_{12}^2) + X_{11} X_3 (X_{12} + X_3) + \frac{Q^2}{W^2} (X_{12}^2 + X_3^2) \right\}$$

$$(\sigma_U^0)^{-1} \frac{d^2\sigma_L}{dx_1 dx_2} = 2(\sigma_U^0)^{-1} \frac{d^2\sigma_T}{dx_1 dx_2} = \frac{4 C_1 Q^2 X_{12}}{\eta(W^2 + Q^2) X_3}$$

$$(\sigma_U^0)^{-1} \frac{d^2\sigma_I}{dx_1 dx_2} = -\frac{4 C_1}{X_3} \left( \frac{X_{12} X_{13} Q^2}{X_{11} W^2} \right)^{1/2} \left\{ (X_2 - X_1) \frac{W^2}{\eta(W^2 + Q^2)} + \frac{X_{12}}{X_3} \right\} \quad (5)$$

where  $X_{1i} = 1 - X_i$  ( $i=1,2,3$ ) and  $\eta$  is a factor universal to all four partial cross-sections:

$$C_1 = \frac{W^2}{\eta(W^2 + Q^2) X_3} \cdot \frac{f_a(\eta, Q^2)}{f_a(X_B, Q^2)} C_1 \frac{dS}{d^2\pi} \quad (6)$$

$\eta = \frac{Q^2 + (1-X_3)W^2}{Q^2 + W^2}$  is the fraction of the proton momentum carried away by the ingoing parton, i.e.  $p = \eta P$ . In these formulas we assumed that  $p_T = 0$ . For diagram Fig. 1b, where the parton a is the gluon, the corresponding formulae are:

†) These cross-sections can be derived from the formulas in ref. 10 with the substitution  $Z = 1 - X_{11}/X_3$ , where  $Z = p_{T1}/p_T$

$$(\sigma_u^0)^{-1} \frac{d^2\sigma_u}{dx_1 dx_2} = \frac{C_3}{x_{11} x_{12}} \frac{W^4 x_{12}^2 + Q^4}{\eta^2 (W^4 + Q^2)^2} (x_{11}^2 + x_{12}^2)$$

$$(\sigma_u^0)^{-1} \frac{d^2\sigma_L}{dx_1 dx_2} = 2(\sigma_u^0)^{-1} \frac{d^2\sigma_T}{dx_1 dx_2} = \frac{8 C_3 Q^2 W^2 x_{13}}{\eta^2 (W^2 + Q^2)^2}$$

$$(\sigma_u^0)^{-1} \frac{d^2\sigma_I}{dx_1 dx_2} = 4 C_3 \frac{Q^2 W^2}{\eta^2 (W^2 + Q^2)^2} \left( \frac{Q^2 x_{13}}{W^2 x_{11} x_{12}} \right)^{1/2} (x_1 - x_2) \left( \frac{W^2}{Q^2} x_{13} - 1 \right) \quad (7)$$

$C_3$  deviates from  $C_2$  just by the colour factors:  $C_3 = \frac{C_3}{C_1} C_1$  where  $C_3 = \frac{1}{2}$ ,  $C_1 = \frac{3}{2}$ . The contribution of the diagrams in Fig.1c is obtained by making the substitution  $x_1 \leftrightarrow x_2$ . We remark that in the final expression where all three contributions of Fig.1b,c and d are added the normalization is given by (4) with a sum over the quark and antiquark terms.

The two-jet contribution is calculated on the basis of the diagrams in Fig.1a. It is well known that, if there is no transverse momentum of the partons confined within the proton, the contribution of Fig.1a to  $\sigma_L$ ,  $\sigma_T$  and  $\sigma_I$  vanishes. To obtain a realistic estimate of the distortion of nonperturbative terms to  $\sigma_L$  and the azimuthal asymmetries we shall incorporate this primordial transverse momentum into the parton model. Let  $p_T$  be the transverse momentum (equal to the transverse momentum of the emitted parton). Then we get the following contribution to the various cross-sections (considering ingoing and outgoing quarks on zero-mass shell) [11]:

$$(\sigma_u^0)^{-1} \frac{d^2\sigma_L}{dx_1 dx_2} = 2(\sigma_u^0)^{-1} \frac{d^2\sigma_T}{dx_1 dx_2} = \frac{4 p_T^2}{Q^2} \delta(1-x_1) \delta(1-x_2) \quad (8)$$

$$(\sigma_u^0)^{-1} \frac{d^2\sigma_I}{dx_1 dx_2} = \frac{4 p_T}{(Q^2)^{1/2}} \delta(1-x_1) \delta(1-x_2)$$

For increasing  $Q^2$  both terms vanish like  $1/Q^2$  and  $1/(Q^2)^{1/2}$  compared to  $\alpha_s(Q^2) \sim 1/\ln Q^2$ , which determines the QCD contributions. The contribution to  $\sigma_I$ , which vanishes only like  $(Q^2)^{-1/2}$  and is comparable in magnitude to the QCD term even at  $Q^2 \approx 100 \text{ GeV}^2$ , does not bother us, since we do not distinguish the current jet from the target jet in our final result. In this case the  $\cos\varphi$  term

drops out in the two-jet contribution.

The variables  $x_1, x_2$  and  $x_3$  are transformed into jet variables. For example, we can take thrust  $T$  and sphericity  $S$ . They are related to  $x_1, x_2$  and  $x_3$  by [12]

$$T = \max(x_1, x_2, x_3) \quad (9)$$

$$S = 64 \pi^{-2} T^{-2} (1-x_1)(1-x_2)(1-x_3)$$

For  $x_1 > x_2 > x_3$  we have

$$dx_1 dx_2 = \frac{\pi^2 T dT dS}{64(1-T)(1-\pi^2 S/16(1-T))^{1/2}} \quad (10)$$

In order to calculate  $\frac{d^2\sigma}{dT dS}$  we sum over the 6 regions of phase space  $x_1 > x_2 > x_3$  etc. consisting of the 6 permutations of 1,2,3. Then the double differential distributions for the various photon polarization states are integrated over  $S$ .  $\frac{d^2\sigma}{dT dS}$  For different values of  $W^2$  and  $Q^2$  the resulting distributions  $\frac{d\sigma_L}{dT}$  and  $\frac{d\sigma_I}{dT}$  are shown in Fig. 2 and Fig. 3. In Fig. 2a, b we have  $W^2 > Q^2$  ( $Q^2 = 625 \text{ GeV}^2$  and  $W^2 = 10^4 \text{ GeV}^2$  and  $W^2 = 2500 \text{ GeV}^2$ ). In this case  $\frac{d\sigma_L}{dT} = 2 \frac{d\sigma_T}{dT}$  and  $\frac{d\sigma_I}{dT}$

have a rather steep fall-off with decreasing  $T$ . For both choices of  $W$  it should be possible to determine the  $\cos\varphi$  term  $\frac{d\sigma_T}{d\sigma_u}$  which is around 15% for  $T = 0.8$  which is outside the non-perturbative region.  $\frac{d\sigma_I}{d\sigma_u}$  is roughly of the same magnitude in the case  $W = 50 \text{ GeV}$ , but smaller for  $W = 100 \text{ GeV}$ . The results in Fig. 3a,b are for  $W=50 \text{ GeV}$  and two  $Q$  values:  $Q=100 \text{ GeV}$  and  $Q=50 \text{ GeV}$ . Here  $\frac{d\sigma_L}{dT} = 2 \frac{d\sigma_T}{dT}$  is somewhat flatter as a function of  $T$  and smaller than  $\frac{d\sigma_I}{dT}$  in the relevant region near  $T = 0.8$ . In these two cases the determination of the  $\cos\varphi$  term should be easier.

The nonperturbative distributions due to the limited transverse momentum of hadrons from quark and gluon fragmentation are also shown in Fig. 2a, b and Fig. 3a,b. These distributions are calculated from

$$(\sigma_u^0)^{-1} \frac{d^2\sigma_u}{dT} = \frac{2(1-T)}{(\Delta T)^2} e^{-\frac{(1-T)^2}{(\Delta T)^2}} \quad (11)$$

The usual assumption for the  $p_T$  distribution of the particles in a jet is  $e^{-4p_T^2}$ , which is gaussian. This has lead us to assume (similar to ref. 12) also a gaussian form for the  $\mathcal{T}$  distribution. The  $(1-T)$  factor is for the fact that massive particles cannot be produced collinearly. The width parameter  $4T$  has been determined from recent jet production experiments in  $\nu$ -nucleon scattering at  $W = 10$  GeV [6]. For the higher  $W$  values the  $4T$  was scaled down with the factor  $W^{-1}$ , which gives us  $4T = 0.048$  for  $W = 50$  GeV. We see from Figs. 2 and 3 that  $T \lesssim 0.8$  is a safe region to test the QCD cross sections. Concerning the asymmetry terms  $d\sigma_T/dT$  and  $d\sigma_{\bar{T}}/dT$  one could go even to larger  $T$  values, since the nonperturbative contribution is really small for  $d\sigma_T/dT$  in this region. This latter distribution was computed from (11) with the factor given in (8) where  $p_T$  was chosen 0.4 GeV. Further tests concerning the transverse-longitudinal cross section  $d^2\sigma/d^2T$  are possible in case one distinguishes the final jets: current jet (quark, antiquark or gluon) and the target jet. Of course, then a nonperturbative contribution is present. Results for such cases will be presented elsewhere.

Finally we mention that all our results have been calculated with  $Q^2$  dependent quark and gluon distribution functions. For the  $Q^2$  dependence we adopted the functional form of Buras and Gaemers [13] with parameters taken from the recent analysis of deep inelastic lepton scattering data by Glück and Reya [14].

In conclusion we state that the polarization dependent cross sections  $\sigma_L$ ,  $\sigma_T$  and  $\sigma_I$  for jet production in deep inelastic lepton scattering can serve as more sophisticated tools to test QCD in the space-like region of  $Q^2$ .

Acknowledgement: We are grateful to G. Schierholz for helpful discussions.

### References

- 1 G. Sterman and S. Weinberg, Phys. Rev. Lett. 39 (1977) 1436  
 H. Georgi and M. Machacek, Phys. Rev. Lett. 39 (1977) 1236  
 E. Farhi, Phys. Rev. Lett. 39 (1977) 1587  
 C.L. Basham, L.S. Brown, S.D. Ellis and T.S. Love, Phys. Rev. D17 (1978) 2298  
 A. De Rújula, J. Ellis, E.G. Floratos and M.K. Gaillard, Nucl. Phys. B138 (1978) 387  
 S.-Y. Pi, R.L. Jaffe and F.E. Low, Phys. Rev. Lett. 41 (1978) 142  
 G. Kramer, G. Schierholz and J. Willrodt, Phys. Lett. 79B (1978) 249  
 erratum *ibid.* 80B (1979) 433
- 2 H. Georgi and H.D. Politzer, Phys. Rev. Lett. 40 (1978) 3  
 J. Cleymans, Phys. Rev. D18 (1978) 954  
 G. Köpp, R. Maciejko and P. Zerwas, Nucl. Phys. B144 (1978) 123  
 P. Mazzanti, R. Odorico and V. Roberts, Phys. Lett. 81B (1979) 219  
 A. Méndez, Nucl. Phys. B145 (1978) 199  
 A. Méndez, A. Raychaudhuri and V.J. Stenger, Nucl. Phys. B148 (1979) 499
- 3 P.M. Stevenson, Nucl. Phys. B150 (1979) 357
- 4 H. Georgi and J. Sheiman, Harvard preprint HUTP-78/A043  
 A. Méndez and T. Weiler, Phys. Lett. 83B (1979) 221
- 5 J. Ranft and G. Ranft, Phys. Lett. 82B (1979) 129  
 P. Binétruy and G. Girardi, Nucl. Phys. B155 (1979) 150  
 C. Sachrajda, Proceedings of the Study of an ep Facility for Europe, DESY, Hamburg, Editor: U. Amaldi, DESY report, DESY 79/48, p. 259
- 6 K.W.J. Barnham et al., Phys. Lett. 85B (1979) 300
- 7 R.D. Field and R.P. Feynman, Nucl. Phys. B136 (1978) 1
- 8 Proceedings of the Study of an ep Facility for Europe, DESY Hamburg, April 2 and 3, 1979, Editor: U. Amaldi, DESY report, DESY 79/48

Figure Captions

9 N.S. Craigie, G. Kramer and J. Körner, Nucl. Phys. B68 (1974) 509

10 A. Méndez, see ref. 2

11 R.N. Cahn, Phys. Lett. 78B (1978) 269

12 A. De Rújula et al., see ref. 1

13 A. Buras and K.J.P. Gaemers, Nucl. Phys. B132 (1978) 249

14 M. Glück and E. Reya, DESY preprint, DESY 79/13

Fig. 1: a) Zeroth order parton diagrams for lepton-parton scattering  
 b,c,d) First order parton diagrams for lepton-parton scattering  
 e) Kinematic diagram for three-jet production in lepton-proton scattering

Fig. 2: Partial cross sections  $U: d\sigma_U/dT, L: d\sigma_L/dT, I: d\sigma_I/dT$   
 and  $T: d\sigma_T/dT = \frac{2}{3} d\sigma_I/dT$  for  
 jet production in electron-proton scattering. The dashed curves  
 are the nonperturbative cross sections,  $\bar{\sigma}$

- a)  $W = 100 \text{ GeV}, Q = 25 \text{ GeV}, \Delta T = 0.028$
- b)  $W = 50 \text{ GeV}, Q = 25 \text{ GeV}, \Delta T = 0.048$

Fig. 3: Same as Fig. 2 for other  $W$  and  $Q$

- a)  $W = 50 \text{ GeV}, Q = 100 \text{ GeV}, \Delta T = 0.048$
- b)  $W = 50 \text{ GeV}, Q = 50 \text{ GeV}, \Delta T = 0.048$





Fig.2a






Fig. 3a

2984J




Fig. 2b

2984J




Fig. 3b