

# DEUTSCHES ELEKTRONEN-SYNCHROTRON DESY

DESY 80/35  
May 1980



MEASUREMENT OF THE REACTION  $e^+e^- \rightarrow \gamma\gamma$  AT CMS ENERGIES FROM 9.4 TO 31.6 GeV

by

*PLUTO Collaboration*

NOTKESTRASSE 85 · 2 HAMBURG 52

**DESY behält sich alle Rechte für den Fall der Schutzrechtsteilung und für die wirtschaftliche  
Verwertung der in diesem Bericht enthaltenen Informationen vor.**

**DESY reserves all rights for commercial use of information included in this report, especially  
in case of apply for or grant of patents.**

To be sure that your preprints are promptly included in the  
**HIGH ENERGY PHYSICS INDEX**,  
send them to the following address ( if possible by air mail ) :

**DESY  
Bibliothek  
Notkestrasse 85  
2 Hamburg 52  
Germany**

MEASUREMENT OF THE REACTION  $e^+e^- \rightarrow \gamma\gamma$  AT CMS ENERGIES FROM 9.4 TO 31.6 GeV

PLUTO Collaboration

Ch. Berger, H. Genzel, R. Grigull, W. Lackas, F. Raupach  
I. Physikalisches Institut der RWTH Aachen<sup>1</sup>, Germany

A. Klovning, E. Lillestöl, E. Lilletethun, J.A. Skard  
University of Bergen<sup>2</sup>, Norway

H. Ackermann, G. Alexander<sup>3</sup>, F. Barreiro, J. Bürger, L. Criegee, H.C. Dehne,  
R. Devenish<sup>4</sup>, A. Eskreys<sup>5</sup>, G. Flügge<sup>6</sup>, G. Franke, W. Gabriel, Ch. Gerke,  
G. Knies, E. Lehmann, H.D. Mertens, U. Michelsen, K.H. Pape, H.D. Reich,  
M. Scarr<sup>13</sup>, B. Stella<sup>7</sup>, T.N. Ranga Swamy, U. Timm, W. Wagner, P. Waloschek,  
G.G. Winter, W. Zimmermann

Deutsches Elektronen-Synchrotron DESY, Hamburg, Germany

O. Achterberg, V. Blobel<sup>9</sup>, L. Boesten, V. Hepp<sup>14</sup>, H. Kapitza, B. Koppitz  
B. Lewendel, W. Lührsen, R. van Staa, H. Spitzer  
II. Institut für Experimentalphysik der Universität Hamburg<sup>1</sup>, Germany

C.Y. Chang, R.G. Glasser, R.G. Kellogg, K.H. Lau, R.O. Polvado, B. Sechi-Zorn,  
A. Skuja, G. Welch, G.T. Zorn  
University of Maryland<sup>10</sup>, College Park, USA

A. Bäcker<sup>11</sup>, S. Brandt, K. Derikum, A. Diekmann, C. Grupen, H.J. Meyer,  
B. Neumann, M. Rost, G. Zech  
Gesamthochschule Siegen<sup>1</sup>, Germany


T. Azemoon<sup>12</sup>, H.J. Daum, H. Meyer, O. Meyer, M. Rössler, D. Schmidt, K. Wacker<sup>11</sup>  
Gesamthochschule Wuppertal<sup>1</sup>, Germany

(Submitted to Physics Letters B)  
30. April 1980

Abstract

The differential cross section for the reaction  $e^+e^- \rightarrow \gamma\gamma$  has been measured in the CMS energy range between 9.4 and 31.6 GeV. The results are found to be in agreement with the predictions of quantum electrodynamics up to momentum transfers- $q^2$  of 900 GeV<sup>2</sup>. The data set lower limits of about 40 GeV on QED cut-off parameters. We have searched for the decay  $T(9.46) \rightarrow \gamma\gamma$  and obtain an upper limit  $\Gamma(T \rightarrow \gamma\gamma)/\Gamma(T \rightarrow \text{all}) < 1.4\% \text{ (95\% c.l.)}$ .

- 1) Supported by the BMFT, Germany
- 2) Partially supported by the Norwegian Research Council for Science and Humanities
- 3) On leave from Tel Aviv University, Israel
- 4) Now at Oxford University, England
- 5) On leave from Institute of Nuclear Physics, Krakow, Poland
- 6) Now at Universität und Kernforschungszentrum Karlsruhe
- 7) On leave from University of Rome, Italy; partially supported by INFN
- 8) Now at Tata Institute, Bombay, India
- 9) Now at CERN, Geneva, Switzerland
- 10) Partially supported by Department of Energy, USA
- 11) Now at Harvard University, Cambridge, Mass., USA
- 12) Now at University College, London, England
- 13) On leave from University of Glasgow, Scotland
- 14) On leave from Heidelberg University



We report on a study of the reaction  $e^+e^- \rightarrow \gamma\gamma$  at center of mass energies between 9.4 and 31.6 GeV. The experiment was performed at the  $e^+e^-$  colliding beam rings DORIS and PETRA at DESY. The two-photon annihilation process is particularly well suited for testing the validity of quantum electrodynamics at high energies, since contributions from electro-weak interference are expected to be absent in first order perturbation theory<sup>1</sup>. The analysis of  $e^+e^- \rightarrow \gamma\gamma$  in the  $\tau$  region is of interest, because the  $J^P = 1^-$  assignment of the  $\tau$  forbids the decay into two photons.

The experiment was carried out with the PLUTO detector which has been described in earlier publications<sup>2,3</sup>. The data analysis is closely linked to the study of Bhabha scattering<sup>3</sup> and will be described in detail only when relevant to the  $\gamma\gamma$  final state. The essential parts of the PLUTO detector for this study are:

- barrel and endcap shower counters, endcap proportional chambers and 880 helix tubes for the cluster analysis and
- 13 cylindrical proportional wire chambers for pattern recognition of charged tracks.

Data were taken at the  $\tau$  region and at center of mass energies of 9.4, 12, 13, 17, 22, 27.6 and 30 - 31.6 GeV. For the selection of  $\gamma\gamma$  events we required two back-to-back showers, containing more than 25 % of the beam energy and being collinear within 20 degrees. The angle  $\theta$  between the  $\gamma$ -rays and the electron beam was restricted to  $|\cos\theta| < 0.75$ .

In a preselection the majority of Bhabha scattering events, eey and hadronic final states were removed by rejecting events

- where at least one track could be associated with each of the clusters\*
- where more than 5 tracks were observed in the inner detector or
- where the angle between any two reconstructed tracks exceeded  $50^\circ$ .

All events surviving these cuts were scanned visually. The subsample of events with no track in the central detector (~ 60 % of the total) was found to be essentially free of background, in agreement with a Monte Carlo study.

\* This cut also excluded  $\gamma\gamma$  final states with two converted photons.

The remaining events consisted of i)  $\gamma\gamma$  final states with one converted photon (~ 30 % of the expected  $\gamma\gamma$  rate), ii) Bhabha scattering with one electron starting an early shower, iii) eey final states and iv) a negligible fraction of hadronic events. A total of 1034 events with an estimated background of ( $1 \pm 1$ ) % was accepted for the final analysis.

The fraction of events excluded by the cuts was estimated with a Monte Carlo program<sup>4</sup> to be ( $5.5 \pm 1.1$ ) %. The visual inspection showed that an additional 1.3 % of events were lost due to incorrectly analysed shower patterns. Combining both values we obtain a detection efficiency of ( $93 \pm 2$ ) % which was applied as an overall correction.

The differential cross sections were evaluated using the integrated luminosities as measured by the rate of large angle Bhabha scatters. Fig. 1 shows the angular distributions together with the QED expectations. The theoretical distributions (full curves) were folded with the angular resolution. The radiative corrections were estimated in the standard way<sup>5</sup> up to orders of  $\alpha^3$  and are incorporated into the theoretical curves. They range between 0 % and -3 % in the angular region under study. The agreement between the data and the QED predictions is good. The systematic uncertainty of the cross sections at 9.4 and 30 - 31.6 GeV is 3.5 % (2 % from event identification and 3 % from the luminosity determination). The luminosity at the remaining energies has slightly higher statistical errors (up to 5 %).

In fig. 2 the s-dependence of the cross section integrated over the measured angular range is displayed in a log-log plot. The straight line is the QED expectation and is seen to give a reasonable fit to our data.

The  $e^+e^- \rightarrow \gamma\gamma$  cross section measured on the  $\tau$  resonance at 9.46 GeV was compared with data at adjacent energies in order to look for evidence of an  $\tau \rightarrow \gamma\gamma$  decay which is forbidden for a  $J^P = 1^-$  vector meson. No enhancement in the  $\gamma\gamma$  signal was found. The upper limit for the branching ratio  $\Gamma(\tau \rightarrow \gamma\gamma)/\Gamma(\tau \rightarrow \text{all})$  is estimated to be 1.4 % (95 % c.l.).

A test of QED was made using our data at the highest PETRA energies. We have applied two possible modifications of the standard QED cross section, which have been suggested in the literature:

- based on a vertex modification (sea-gull graph<sup>7,8</sup>) the differential cross section can be written as<sup>6</sup>

$$\frac{d\sigma_V}{d\Omega} = \frac{\alpha^2}{2s} \left\{ \frac{q'^2}{q^2} |F(q^2)|^2 + \frac{q^2}{q'^2} |F(q'^2)|^2 \right\} (1 + \delta_{rad})$$

$$= \frac{\alpha^2}{2s} \left\{ \frac{q'^2}{q^2} + \frac{q^2}{q'^2} \pm \frac{4}{\Lambda_V^4} \frac{q^2 q'^2}{(1 + \delta_{rad})} \right\} (1 + \delta_{rad})$$

$$\text{with } F(q^2) = 1 \pm \frac{q^4}{\Lambda_{V\pm}^4}, \quad F(q'^2) = 1 \pm \frac{q'^4}{\Lambda_{V\pm}^4}$$

Here  $\delta_{rad}$  is the radiative correction,  $q^2 = -s \cos^2 \theta/2$  and  $q'^2 = -s \sin^2 \theta/2$ .  $\Lambda_{V\pm}$  is a cut-off parameter.

- On the other hand if one assumes the exchange of a hypothetical heavy electron  $e^*$  g then:

$$\frac{d\sigma_{e^*}}{d\Omega} = \frac{\alpha^2}{2s} \left\{ \frac{q'^2}{q^2} + \frac{q^2}{q'^2} \pm \frac{2 s^2 - 4 q^2 q'^2}{\Lambda_{e^*}^4} \right\} (1 + \delta_{rad}) \quad (2)$$

The value of  $\Lambda_{e^*}$  can be interpreted as the mass of a heavy electron assuming its coupling strength is the same as that of the electron. The parameter  $\Lambda_{e^*}$  is theoretically less motivated. In both cases the relative modification is largest and of same magnitude at  $\theta = 90^\circ$ .

Both expressions were fitted to the experimental differential cross sections at  $30 \leq E_{cm} \leq 31.6$  GeV. The fit selected the sign of  $\Lambda^4$  in eq. (1) and (2) required to describe the data. The fitted values of  $s^2/\Lambda^4$  are given in table 1. Lower bounds for the cut-off parameters  $\Lambda_{V+}$ ,  $\Lambda_{V-}$ ,  $\Lambda_{e^*}$ ,  $\Lambda_{e^*+}$  were obtained with standard  $\chi^2_{min}$  techniques.

In both parameterisations we find lower limits for the cut off parameters of about 40 GeV (95 % c.l.). The results are compared in table 1 with those from other recent measurements. It can be seen that the new results of PLUTO and JADE at the highest PETRA energies have raised the lower bounds on  $\Lambda$  by a factor of 4 to 5.

In conclusion, our results show that the two-photon annihilation process is very well described by quantum electrodynamics up to  $q^2 \approx -900$  GeV<sup>2</sup>. The absence of  $\tau + \gamma\gamma$  has been checked down to the 1 % level.

#### Acknowledgements

We wish to thank professors H. Schopper, G. Voss, E. Lohrmann and Dr. G. Söhngen for their valuable support. We are indebted to the PETRA machine group and the DESY computer center for their excellent support during the experiment under sometimes difficult circumstances. We gratefully acknowledge the efforts of all the engineers and technicians of the collaborating institutions who have participated in the construction and maintenance of the apparatus. We thank Dr. F. Gutbrod for an illuminating discussion.

References

- 1) M. Capdequi-Peyranere, G. Grunberg, F.M. Renard, M. Talon, Preprint PM/78/8 (1978), Département de Physique Mathématique, Montpellier
- 2) PLUTO Collaboration, Ch. Berger et al., Phys. Letters 81B (1979) 410
- 3) PLUTO Collaboration, Ch. Berger et al., DESY Report 80/01 (1980), to be published in Zeits. f. Physik C;
8. Koppitz, Thesis, Internal Report DESY PLUTO 80/05 (1980), unpublished
- 4) F. Rapuano, private communication
- 5) F.A. Berends, R. Gastmans, Nucl. Phys. B61 (1973) 414
- 6) R. Gatto, Proceedings of the Intern. Symposium on Electron and Photon Interactions at High Energies, Hamburg 1965, Vol. 1, p. 106, Deutsche Physikalische Gesellschaft, 1965
- 7) N.M. Kroll, Nuovo Cim. XLV (1966) 65
- 8) K. Ringhofer, H. Salecker, Contribution to the 1975 Internat. Symposium on Lepton and Photon Interactions at High Energies, Stanford University, unpublished
- 9) A. Little, Harvard University, Ph. D. Thesis (1970), unpublished
- 10) C. Bacci et al., Lett. Nuovo Cim. 2 (1971) 73
- 11) G. Hansson et al., Lett. Nuovo Cim. 7 (1973) 587
- 12) B.L. Beron et al., Phys. Rev. Lett. 33 (1974) 663
- 13) E. Hilger et al., Phys. Rev. D15 (1977) 1809
- 14) JADE Collaboration, W. Bartel et al., DESY Report 80/14 (1980)

**Table 1** Fitted values of  $s^2/\Lambda^4$  and lower limits (95 % c.l.) of the QED cut-off parameters for the reaction  $e^+e^- \rightarrow \gamma\gamma$ . Everywhere  $\epsilon = \pm 1$  represents the fitted sign in eqs. (1) and (2) and  $\Lambda_\pm$  corresponds to  $\epsilon = \pm 1$ . The statistical and systematic uncertainties of the luminosity determination have been taken into account in the fit.

ref.	$E_{cm}$ (GeV)	$\epsilon \frac{s^2}{\Lambda_4}$	$\epsilon \frac{s^2}{\Lambda_V}$	$\frac{s^2}{\Lambda_{e^*}}$	$\Lambda_{V^+}$ (GeV)	$\Lambda_{V^-}$ (GeV)	$\Lambda_{e^*}$ (GeV)
10	1.4 - 2.4				> 2.6	> 2.4	-
11	4				-	-	> 3.9
12	5.2				> 6.2	> 6.9	-
13	6.2 - 7.4				> 10.7	> 9.0	-
14	27.7 - 31.6				-	-	> 45
this exp.	30 - 31.6	-0.169 ±0.185	-0.083 ±0.145	> 4.6	> 3.6	> 4.6	

Figure captions

- Fig. 1** Differential cross section  $d\sigma/d\cos\theta$  of the reaction  $e^+e^- \rightarrow \gamma\gamma$  at different cms energies as indicated in the figure. The curves are the QED predictions including the effects of radiation and angular resolution.

- Fig. 2** The cross section for reaction  $e^+e^- \rightarrow \gamma\gamma$  for  $|\cos\theta| < 0.75$ . The curve is the QED prediction including the effects of radiation and angular resolution.




Fig. 1




Fig. 2