

BSM Higgs Physics at the LHC in the Forward Proton Mode *S. HEINEMEYER^{1†}, V.A. KHOZE^{2,3‡}, M.G. RYSKIN^{2,4§},
M. TASEVSKY^{5¶} AND G. WEIGLEIN^{6||}¹*Instituto de Física de Cantabria (CSIC-UC), Santander, Spain*²*IPPP, Department of Physics, Durham University, Durham DH1 3LE, U.K.*³*School of Physics & Astronomy, University of Manchester, Manchester M13 9PL, U.K.*⁴*Petersburg Nuclear Physics Institute, Gatchina, St. Petersburg, 188300, Russia*⁵*Institute of Physics, 18221 Prague 8, Czech Republic*⁶*DESY, Notkestraße 85, D-22607 Hamburg, Germany***Abstract**

We review the prospects for Central Exclusive Diffractive (CED) production of Higgs bosons in the SM with a fourth generation of fermions at the LHC using forward proton detectors installed at 220 m and 420 m distance around ATLAS and / or CMS. We discuss the determination of Higgs spin-parity and coupling structures at the LHC and show that the forward proton mode would provide a critical information on the \mathcal{CP} properties of the Higgs bosons.

*talk given by S.H. at the *Physics at the LHC 2010*, May 2010, DESY Hamburg, Germany

† email: Sven.Heinemeyer@cern.ch

‡ email: V.A.Khoze@durham.ac.uk

§ email: Ryskin@MR11084.spb.edu

¶ email: Marek.Tasevsky@cern.ch

|| email: Georg.Weiglein@desy.de

BSM Higgs Physics at the LHC in the Forward Proton Mode

Sven Heinemeyer^{1*}, V.A. Khoze^{2,3}, M.G. Ryskin^{2,4}, M. Tasevsky⁵, Georg Weiglein⁶

¹Instituto de Física de Cantabria (CSIC-UC), Santander, Spain

²IPPP, Department of Physics, Durham University, Durham DH1 3LE, U.K.

³School of Physics & Astronomy, University of Manchester, Manchester M13 9PL, U.K.

⁴Petersburg Nuclear Physics Institute, Gatchina, St. Petersburg, 188300, Russia

⁵Institute of Physics, 18221 Prague 8, Czech Republic

⁶DESY, Notketräße 85, 22607 Hamburg, Germany

DOI: will be assigned

We review the prospects for Central Exclusive Diffractive (CED) production of Higgs bosons in the SM with a fourth generation of fermions at the LHC using forward proton detectors installed at 220 m and 420 m distance around ATLAS and / or CMS. We discuss the determination of Higgs spin-parity and coupling structures at the LHC and show that the forward proton mode would provide a critical information on the \mathcal{CP} properties of the Higgs bosons.

1 Introduction

In the recent years there has been a growing interest in the possibility to complement the standard LHC searches a Higgs boson by the options offered by forward and diffraction physics. These assume the installation of near-beam proton detectors in the LHC tunnel installed at 220 m and 420 m around ATLAS and / or CMS, see Refs. [1–6] and references therein. The combined detection of the centrally produced system and both outgoing protons can provide valuable information on the Higgs sector of MSSM and other popular BSM scenarios [3, 7–10]. Another simple example of physics beyond the SM is a model which extends the SM by a fourth generation of heavy fermions (SM4), see, for instance, [11–13]. Here it is assumed that the masses of the 4th generation quarks are (much) heavier than the mass of the top-quark. In this case, the effective coupling of the Higgs boson to two gluons is three times larger than in the SM, and all branching ratios change correspondingly.

The central exclusive diffractive (CED) processes are of the form

$$pp \rightarrow p \oplus H \oplus p, \quad (1)$$

where the \oplus signs denote large rapidity gaps on either side of the centrally produced state. However, proving that a detected new state is, indeed, a Higgs boson will be far from trivial. In particular, it will be of great importance to determine the spin and \mathcal{CP} properties of a new state and to measure precisely its mass, width and couplings.

*Speaker

Following [8] we consider four luminosity scenarios: “60 fb⁻¹” and “600 fb⁻¹” refer to running at low and high instantaneous luminosity, respectively, using conservative assumptions for the signal rates and the experimental sensitivities; possible improvements of both theory and experiment could allow for the scenarios where the event rates are higher by a factor of 2, denoted as “60 fb⁻¹ eff×2” and “600 fb⁻¹ eff×2”.

2 The Higgs boson in the SM4

A simple example of physics beyond the SM is a model, “SM4”, which extends the SM by a fourth generation of heavy fermions, see, for instance, Refs. [11,12,14]. In particular, the masses of the 4th generation quarks are assumed to be (much) heavier than the mass of the top-quark (whereas the masses of the 4th generation leptons, which do not play a role here, are less restricted). In this case, the effective coupling of the Higgs boson to two gluons is three times larger than in the SM. No other coupling, relevant to LEP and Tevatron searches, changes significantly. Essentially, only the partial decay width $\Gamma(H \rightarrow gg)$ changes by a factor of 9 and, with it, the total Higgs width and therefore all the decay branching ratios, see for instance Ref. [13,15]. The new total decay width and the relevant decay branching ratios can be evaluated as,

$$\Gamma_{\text{SM}}(H \rightarrow gg) = \text{BR}_{\text{SM}}(H \rightarrow gg) \Gamma_{\text{tot}}^{\text{SM}}(H), \quad (2)$$

$$\Gamma_{\text{SM4}}(H \rightarrow gg) = 9 \Gamma_{\text{SM}}(H \rightarrow gg), \quad (3)$$

$$\Gamma_{\text{tot}}^{\text{SM4}}(H) = \Gamma_{\text{tot}}^{\text{SM}}(H) - \Gamma_{\text{SM}}(H \rightarrow gg) + \Gamma_{\text{SM4}}(H \rightarrow gg). \quad (4)$$

The Higgs boson searches at LEP [16,17] have been re-interpreted by `HiggsBounds` [18] in the SM4. The bound on the SM Higgs boson at LEP of $M_{H^{\text{SM}}} \geq 114.4$ GeV at the 95% C.L. is modified to $M_{H^{\text{SM4}}} \geq 112$ GeV. On the other hand Higgs boson searches in the SM4 at the Tevatron [19] have been performed. The range $130 \text{ GeV} \lesssim M_{H^{\text{SM4}}} \lesssim 210 \text{ GeV}$ is found to be excluded. Combining the two analyses leaves us with a window of allowed Higgs masses in the SM4 of $112 \text{ GeV} \lesssim M_{H^{\text{SM4}}} \lesssim 130 \text{ GeV}$.




Figure 1: Significances reachable CED Higgs production in the SM4 in the $H \rightarrow b\bar{b}$ (left plot) and $H \rightarrow \tau^+\tau^-$ (right plot) channel for effective luminosities of “60 fb⁻¹”, “60 fb⁻¹ eff×2”, “600 fb⁻¹” and “600 fb⁻¹ eff×2”. The regions excluded by LEP appear as blue/light grey for low values of $M_{H^{\text{SM4}}}$ and excluded by the Tevatron as red/dark grey for larger values of $M_{H^{\text{SM4}}}$.

We have evaluated the significances that can be obtained in the channels $H \rightarrow b\bar{b}$ and $H \rightarrow \tau^+\tau^-$. The results are shown in Fig. 1 as a function of $M_{H^{SM4}}$ for the four luminosity scenarios. The regions excluded by LEP appear as blue/light grey for low values of $M_{H^{SM4}}$ and regions excluded by the Tevatron appears as red/dark grey for larger values of $M_{H^{SM4}}$. The $b\bar{b}$ channel (left plot) shows that even at rather low luminosity the remaining window of $112 \text{ GeV} \lesssim M_{H^{SM4}} \lesssim 130 \text{ GeV}$ can be covered by CED Higgs production. Due to the smallness of $\text{BR}(H^{SM4} \rightarrow b\bar{b})$ at $M_{H^{SM4}} \gtrsim 160 \text{ GeV}$, however, the CED channel becomes irrelevant for the still allowed high values of $M_{H^{SM4}}$, and we do not extend our analysis beyond $M_{H^{SM4}} \leq 200 \text{ GeV}$. The $\tau^+\tau^-$ channel (right plot) has not enough sensitivity at low luminosity, but might become feasible at high LHC luminosity. At masses $M_{H^{SM4}} \gtrsim 220 \text{ GeV}$ it might be possible to exploit the decay $H \rightarrow WW, ZZ$, but no detailed analysis has been performed up to now.

3 Coupling structure and spin-parity determination

The determination of the spin and the \mathcal{CP} properties of Higgs bosons using the standard methods rely to a large extent on the coupling of a relatively heavy SM-like Higgs to two gauge bosons. The first channel that should be mentioned here is $H \rightarrow ZZ \rightarrow 4l$. This channel provides detailed information about spin and \mathcal{CP} -properties if it is open [20]. Within a SM-like set-up it was analyzed how the tensor structure of the coupling of the Higgs boson to weak gauge bosons can be determined at the LHC [21–23]. One study for $M_{H^{SM}} = 160 \text{ GeV}$ was based on Higgs production in weak vector boson fusion with the subsequent decay to SM gauge boson. It was shown that the discrimination between the two extreme scenarios of a pure \mathcal{CP} -even (as in the SM) and a pure \mathcal{CP} -odd tensor structure at a level of 4.5 to 5.3 σ using only about 10 fb^{-1} . A discriminating power of two standard deviations at $M_{H^{SM}} = 120 \text{ GeV}$ in the tau lepton decay mode requires an integrated luminosity of 30 fb^{-1} [23].

For $M_H \approx M_A \gtrsim 2M_W$ the lightest MSSM Higgs boson couples to gauge bosons with about SM strength, but its mass is bounded from above by $M_h \lesssim 135 \text{ GeV}$ [24], i.e. the light Higgs stays below the threshold above which the decay to $WW^{(*)}$ or $ZZ^{(*)}$ can be exploited. On the other hand, the heavy MSSM Higgs bosons, H and A , decouple from the gauge bosons. Consequently, the analysis for $M_{H^{SM}} = 160 \text{ GeV}$ cannot be taken over to the MSSM. This shows the importance of channels to determine spin and \mathcal{CP} -properties of the Higgs bosons without relying on (tree-level) couplings of the Higgs bosons to gauge bosons. CED Higgs production can yield crucial information in this context [2, 7, 8].

The $M_{H^{SM}} = 120 \text{ GeV}$ analysis, on the other hand, can in principle be applied to the SUSY case. However, in this case the coupling of the SUSY Higgs bosons to tau leptons, does not exhibit a (sufficiently) strong enhancement as compared to the SM case. Consequently, no improvement over the 2σ effect within the SM can be expected. The same would be true in any other model of new physics with a light SM-like Higgs and heavy Higgses that decouple from the gauge bosons.

The CED production channels may provide crucial information on the \mathcal{CP} properties of Higgs-like states detected at the LHC, for instance via the J_z selection rule [25]. Thanks to this selection rule in the CED case we already know that the observed new object has even parity ($\mathcal{P} = +$) and the projection of its spin is $J_z = 0$. This knowledge will greatly simplify the determination of the spin of new boson.

As discussed in [7,8] it will be challenging to identify a \mathcal{CP} -odd Higgs boson, for instance the

A boson of the MSSM, in the CED processes. The strong suppression, caused by the P -even selection rule, effectively filters out its production. However in the semi-inclusive diffractive reactions the pseudoscalar production is much less suppressed. As shown in a recent study [5] there are certain advantages of looking for the \mathcal{CP} -odd Higgs particle in the semi-inclusive process $pp \rightarrow p + gAg + p$ with two tagged forward protons and two large rapidity gaps. The amplitude of \mathcal{CP} -odd A boson production can be of the same order as the \mathcal{CP} -even boson amplitude if events with relatively hard gluons, whose energy is comparable with the energy of the whole gAg system, are selected.

4 Acknowledgments

Work supported in part by the European Community's Marie-Curie Research Training Network under contract MRTN-CT-2006-035505 'Tools and Precision Calculations for Physics Discoveries at Colliders' (HEPTOOLS).

References

- [1] M. Albrow and A. Rostovtsev, arXiv:hep-ph/0009336.
- [2] V.A. Khoze, A.D. Martin and M. Ryskin, *Eur. Phys. J. C* **23** (2002) 311.
- [3] M. Albrow, T. Coughlin and J. Forshaw, arXiv:1006.1289 [hep-ph].
- [4] A. De Roeck, V.A. Khoze, A. Martin, R. Orava and M. Ryskin, *Eur. Phys. J. C* **25** (2002) 391.
- [5] V. A. Khoze, A. Martin, M. Ryskin and A. Shuvaev, *Eur. Phys. J. C* **68** (2010) 125.
- [6] M. Albrow et al. [FP420 R&D Collaboration], *JINST* **4** (2009) T10001.
- [7] A. Kaidalov, V.A. Khoze, A.D. Martin and M. Ryskin, *Eur. Phys. J. C* **33** (2004) 261.
- [8] S. Heinemeyer, V. A. Khoze, M. G. Ryskin, W. J. Stirling, M. Tasevsky and G. Weiglein, *Eur. Phys. J. C* **53** (2008) 231.
- [9] B. Cox, F. Loebinger and A. Pilkington, *JHEP* **0710** (2007) 090.
- [10] M. Chaichian, P. Hoyer, K. Huitu, V. A. Khoze and A. Pilkington, *JHEP* **0905** (2009) 011.
- [11] P. Frampton, P. Hung and M. Sher, *Phys. Rept.* **330** (2000) 263.
- [12] B. Holdom et al., *PMC Phys. A* **3** (2009) 4 [arXiv:0904.4698 [hep-ph]].
- [13] G. Kribs, T. Plehn, M. Spannowsky and T. Tait, *Phys. Rev. D* **76** (2007) 075016.
- [14] J. Erler and P. Langacker, *Phys. Rev. Lett.* **105** (2010) 031801.
- [15] C. Anastasiou, R. Boughezal and E. Furlan, *JHEP* **1006** (2010) 101.
- [16] G. Abbiendi et al. [ALEPH, DELPHI, L3, OPAL Collaborations and LEP Working Group for Higgs boson searches], *Phys. Lett. B* **565** (2003) 61.
- [17] S. Schael et al. [ALEPH, DELPHI, L3, OPAL Collaborations and LEP Working Group for Higgs boson searches], *Eur. Phys. J. C* **47** (2006) 547.
- [18] P. Bechtle, O. Brein, S. Heinemeyer, G. Weiglein and K. Williams, *Comput. Phys. Commun.* **181** (2010) 138; arXiv:0909.4664 [hep-ph]; see: www.ipp.dur.ac.uk/HiggsBounds .
- [19] TEVNP Working Group Collaboration for the CDF Collaboration and D0 Collaboration, CDF note 10101, D0 note 6039.
- [20] V. Buescher and K. Jakobs, *Int. J. Mod. Phys. A* **20** (2005) 2523.
- [21] T. Plehn, D. Rainwater and D. Zeppenfeld, *Phys. Rev. Lett.* **88** (2002) 051801.
- [22] V. Hankele, G. Klamke, D. Zeppenfeld and T. Figy, *Phys. Rev. D* **74** (2006) 095001.
- [23] C. Ruwiedel, N. Wermes and M. Schumacher, *Eur. Phys. J. C* **51** (2007) 385.
- [24] G. Degrassi, S. Heinemeyer, W. Hollik, P. Slavich and G. Weiglein, *Eur. Phys. J. C* **28** (2003) 133.
- [25] V.A. Khoze, A. Martin and M. Ryskin, *Eur. Phys. J. C* **19** (2001) 477 [Erratum-ibid. *C* **20** (2001) 599].