

DEUTSCHES ELEKTRONEN-SYNCHROTRON **DESY**

DESY 74/10
March 1974



Quark Fragmentation Functions in Hadronic and Deep Inelastic
Single Particle Inclusive Processes

by



B. Flume-Gorczyca

To be sure that your preprints are promptly included in the
HIGH ENERGY PHYSICS INDEX,
send them to the following address (if possible by air mail) :

DESY
Bibliothek
2 Hamburg 52
Notkestieg 1
Germany

Quark Fragmentation Functions in Hadronic and Deep Inelastic
Single Particle Inclusive Processes

by

B. Flume-Gorczyca
Deutsches Elektronen-Synchrotron DESY, Hamburg

Abstract

In the framework of Feynman's quark-parton model for inclusive electroproduction and quark model of Ezawa, Maharana and Miyazawa for hadronic inclusive processes, relations between invariant cross sections for electroproduction of pions and purely hadronic production of pions in the beam fragmentation regions are derived. They are satisfied by data.

The ratio of parton fragmentation functions $\eta = D_{\text{u}}^{\pi^+}(z)/D_{\text{d}}^{\pi^+}(z)$ is estimated from purely hadronic processes independently from the parton model.

The purpose of this paper is to relate the inclusive electroproduction of pions in the parton fragmentation region with the inclusive production of pions in purely hadronic processes in the beam fragmentation region.

For electroproduction we use the framework of Feynman's quark parton model [1], [2], and for hadronic inclusive processes the quark model of Ezawa, Maharana and Miyazawa [3].

In Feynman's quark-parton model [1] $N_{ep}^{\pi}(x, z) = \frac{1}{\sigma_{\text{tot}}} \frac{d\sigma}{dx dz}$ is written as the product of probability to hit a certain type of quark in the proton times the probability that this quark decays into a pion plus anything: *)

$$\begin{aligned} N_{ep}^{\pi}(x, z) f_1^{\text{ep}}(x) &= \frac{4}{9} u(x) D_u^{\pi}(z) + \frac{4}{9} \bar{u}(x) D_{\bar{u}}^{\pi}(z) \\ &+ \frac{1}{9} d(x) D_d^{\pi}(z) + \frac{1}{9} \bar{d}(x) D_{\bar{d}}^{\pi}(z) \\ &+ \frac{1}{9} s(x) D_s^{\pi}(z) + \frac{1}{9} \bar{s}(x) D_{\bar{s}}^{\pi}(z) , \end{aligned} \quad (1)$$

where

$$\begin{aligned} f_1^{\text{ep}}(x) &= \frac{4}{9} (u(x) + \bar{u}(x)) + \frac{1}{9} (d(x) + \bar{d}(x)) \\ &+ \frac{1}{9} (s(x) + \bar{s}(x)) \end{aligned}$$

The variable x is equal to $1/\omega$ and the variable z is the energy of the pion measured in terms of the energy of the photon in the laboratory system. These D functions depend neither on q^2 nor on x [1], [2]. Six functions $u(x)$, $d(x)$, give us the average number of quarks u , d , s , ... , in the proton with a fraction x of the proton momentum. The six $D_q^{\pi}(z)$ are the parton (quark) fragmentation functions. Here the produced π carries a fraction z of the quark momentum.

Recently Ezawa, Maharana and Miyazawa (EMM) [3] proposed a simple quark model describing inclusive cross sections for hadron-hadron interactions:

$a + b \rightarrow c + \text{anything}$.

In this scheme one quark in hadron a and one in b collide and start a multiple production: quark + quark $\rightarrow c + \text{anything}$. This process contributes mainly to

the central region. Other spectator quarks are shaken off and decay into hadrons: quark $\rightarrow c + \text{anything}$. Processes of the second type contribute to the fragmentation regions of hadrons a and b .

We shall be concerned with the description of the fragmentation regions in this model. In the fragmentation region particle c is produced as a fragment of a quark. It is expected that distribution of the particle c is isotropic in the quark's rest frame and is a function of \vec{p}_c^2 (\vec{p}_c being the momentum of particle c in quark rest frame), It is easy to show [3] that $\vec{p}_c^2 = f(z, p_\perp^2)$, where z is a finite fraction of the quark momentum taken by particle c , p_\perp is its transverse momentum. Thus the fragmentation process of the quark should be described by the quark fragmentation function $D(z)$.*)

Now we confine ourselves to the inclusive meson M' production in the meson M fragmentation region. In this case we can write for $N(x_F) = 1/\sigma_{\text{tot}} d\sigma/dx_F$ the following expression

$$N_{MT}^{M'}(x_F) = \frac{1}{2} \int_0^1 f(x') \left(D_q^{M'} \left(\frac{x_F}{x'} \right) + D_{\bar{q}}^{M'} \left(\frac{x_F}{x'} \right) \right) dx', \quad (2)$$

where x_F is the Feynman's variable, x' denotes a fraction of momentum of meson M carried by quark (q or \bar{q}) and $f(x')$ describes the momentum distribution of a given quark within the meson. T denotes a target. EMM model corresponds to the specific assumption of $f(x') = \delta(x' - \frac{1}{2})$, i.e. half of the meson momentum is carried by each quark. In this case:

$$N_{MT}^{M'}(x_F) = \frac{1}{2} [D_q^{M'}(z) + D_{\bar{q}}^{M'}(z)], \quad (3)$$

where $2x_F = z$.

This assumption (claiming that both quarks in the meson take equal amount of the initial meson momentum) restricts description of the fragmentation process to the region $0 \leq x_F \leq \frac{1}{2}$. In the pion fragmentation region ($x_F > 0$) the EMM model predicts the following:**) (it is assumed that D functions are related among themselves by the charge symmetry and charge conjugation)

$$N_{\pi^+p}^{\pi^+}(x_F) = N_{\pi^-p}^{\pi^-}(x_F) = D_u^{\pi^+}(z), \quad (4)$$

$$N_{\pi^+p}^{\pi^-}(x_F) = N_{\pi^-p}^{\pi^+}(x_F) = D_d^{\pi^+}(z) \quad , \quad (5)$$

$$N_{\pi^0p}^{\pi^-}(x_F) = \frac{1}{2} (D_u^{\pi^+}(z) + D_d^{\pi^+}(z)) \quad , \quad (6)$$

where $z = \frac{P_c}{P_{\text{quark}}}$.

If we look for fragments of the beam then single inclusive distributions in hadronic processes are described just by the same quark fragmentation functions which are used for description of the single inclusive spectra in ep scattering.

Cleymans and Sehgal [4] have considered arguments supporting the assumption that in the quark parton model "functions $\bar{u}(x)$, $\bar{d}(x)$, $s(x)$ and $\bar{s}(x)$ may be neglected in comparison to $u(x)$ and $d(x)$ in an interval $x_0 \leq x \leq 1$, where x_0 is an unspecified, but presumably small number". Using this assumption they were able to extract $u(x)$ and $d(x)$ from the inelastic electron scattering data. So they have for π^\pm inclusive distributions in electron-proton scattering (excluding the small x region) the following relations:

$$N_{ep}^{\pi^+}(x, z) \approx \left[\frac{4}{9} u(x) D_u^{\pi^+}(z) + \frac{1}{9} d(x) D_d^{\pi^+}(z) \right] / f_1^{\text{ep}}(x) \quad , \quad (7)$$

$$N_{ep}^{\pi^-}(x, z) \approx \left[\frac{4}{9} u(x) D_d^{\pi^+}(z) + \frac{1}{9} d(x) D_u^{\pi^+}(z) \right] / f_1^{\text{ep}}(x) \quad , \quad (8)$$

where $f_1^{\text{ep}}(x) \approx \frac{4}{9} u(x) + \frac{1}{9} d(x)$.

Knowing the probabilities of hitting different kinds of partons (functions $u(x)$ and $d(x)$) and invariant cross sections $N_{ep}^{\pi^\pm}$ Cleymans and Rodenberg [5] extracted the quark fragmentation functions $D_u^{\pi^+}(z)$ and $D_d^{\pi^+}(z)$. Another result of this paper [5] is that now if $N_{ep}^{\pi^\pm}(x, z)$ is known (e.g. experimentally) for one particular value of x , it's easy to calculate it for all values of x excluding the small x region).

Combining relations (4-6) and (7-8) we can write down the following sum rules:

$$N_{ep}^{\pi^+}(x, z)(4u(x) + d(x)) = 4u(x)N_{\pi^+p}^{\pi^+}(x_F) + d(x)N_{\pi^+p}^{\pi^-}(x_F) \quad , \quad (9)$$

$$N_{ep}^{\pi^-}(x, z)(4u(x) + d(x)) = 4u(x)N_{\pi+p}^{\pi^-}(x_F) + d(x)N_{\pi+p}^{\pi^+}(x_F). \quad (10)$$

The comparison should be made at an experimental point where the momentum carried by the quarks is the same in ep and in πp .

Moreover, from purely hadronic inclusive cross sections in the fragmentation region in the framework of EMM model we can calculate the ratio $\eta(z) = D_u^{\pi^+}(z)/D_d^{\pi^+}(z)$. So it can be done independently from the parton model. This ratio η is an important variable as in the framework of quark parton model one needs it to calculate the charge ratio $N_{ep}^{\pi^+}/N_{ep}^{\pi^-}$. Dakin and Feldman [6] have pointed out that in the McElhaney and Tuan [7] fit to the Kuti-Weisskopf model η (average over z) gives the upper limit of $N_{ep}^{\pi^+}/N_{ep}^{\pi^-}$ for $x = 1$ (similar conclusion can be drawn from Refs.[4-5] as from graphs given there for $u(x)$ and $d(x)$ for $0.1 < x < 0.8$ one can expect that $\lim_{x \rightarrow 1} (d(x)/u(x)) = 0$).

If we want to check experimentally relations (9-10) or to calculate $\eta(z)$ from hadronic processes we meet one difficulty, namely the quark fragmentation function $D_u^{\pi^+}$ is always connected with the inclusive processes for which the leading particle effect [8], [9] plays important role (in above relations processes $\pi^+ \xrightarrow{P} \pi^+$ and $\pi^0 \xrightarrow{P} \pi^-$). Here the result of the paper given by Cleymans and Sehgal [4] is very useful. Namely for $x \geq 0.8$ we can assume that $d(x) \approx 0$. It gives us the interesting test of the relation (10):

$$N_{ep}^{\pi^-}(x \geq 0.8; z) = N_{\pi+p}^{\pi^-}(x_F) (= N_{\pi-p}^{\pi^+}(x_F)) \quad (11)$$

because in the process $\pi^- \xrightarrow{P} \pi^+$ (or $\pi^+ \xrightarrow{P} \pi^-$) π^+ (or π^-) is non-leading particle.

If we want to compare with data relation (9) or to calculate the ratio η we must avoid the contamination of the pure fragmentation mechanism by the leading particle effect in other way. To reach it we use the data for inclusive reaction $\gamma p \rightarrow \pi^- + \text{anything}$ with elastic ρ^0 production excluded [10]. To relate the invariant cross section for the process $\gamma p \rightarrow \pi^- + \text{anything}$ (with ρ^0 elastic excluded) with hadronic inclusive production of pion we use the relation obtained by combining vector meson dominance with the quark model [11]. In this way we have:

$$N_{\gamma p}^{\pi^-} = N_{\gamma p}^{\pi^+} = \frac{\gamma \xrightarrow{P} \pi^-}{\sigma_{\text{tot}}(\gamma p)} = \frac{\pi^0 \xrightarrow{P} \pi^-}{\sigma_{\text{tot}}(\pi^0 p)} \quad (12)$$

and next from the equation (6) we get

$$N_{\gamma p}^{\pi^-}(x_F) = \frac{1}{2} (D_u^{\pi^+}(z) + D_d^{\pi^+}(z)) \quad (13)$$

Now we can calculate the ratio $\eta(z) = D_u^{\pi^+}(z)/D_d^{\pi^+}(z)$ e.g. from the relation

$$N_{\gamma p}^{\pi^-}(x_F)/N_{\pi^+ p}^{\pi^-}(x_F) = \frac{1}{2} (1 + \eta(z)). \quad (14)$$

Moreover from relations (7), (8) and (13) we obtain an interesting sum rule:

$$N_{ep}^{\pi^+}(z) + N_{ep}^{\pi^-}(z) = 2 N_{\gamma p}^{\pi^-}(x_F) \quad (\text{with } \rho^0 \text{ excluded}) \quad (15)$$

Comparison with the data

a) For the relation (11):

$$N_{ep}^{\pi^-}(x = 0.8; z) = N_{\pi^+ p}^{\pi^-}(x_F)$$

The comparison of the relation (11) with the data is presented in Fig.1.

$$N_{ep}^{\pi^-}(x = 0.8; 0.3 < z < 0.7) \quad (***) \text{ is taken from the graph in Ref. [5]}$$

and

$$N_{\pi^+ p}^{\pi^-}(x_F) \text{ from Ref. [12] (at } p_{\pi^+} = 3.7 \text{ GeV).}$$

We have chosen these data as in this case the momentum carried by quarks is the same in ep and πp ($\approx 1,25$ GeV). We see that relation (11) is well attest in the region $0.3 < x_F < 0.7$.

b) For the relation (15):

$$N_{ep}^{\pi^+}(z) + N_{ep}^{\pi^-}(z) = 2N_{\gamma p}^{\pi^-}(x_F) \quad (\text{with } \rho^0 \text{ excluded}).$$

To compare the relation (15) with data we use the fit from Refs.[5], [13] namely (at $Q^2 = 1.20$ and 2.02 GeV², where W average over these two points is equal 2.4 GeV):

$$2N_{ep}^{\pi^-}(x = 0.25; z) = N_{ep}^{\pi^+}(x = 0.25; z) = \frac{\pi}{Bz} e^{-3.17z + 0.45} \quad ***)$$

In Fig.2 the points put on the experimental data for $N_{\gamma p}^{\pi^-}$ (at $E_\gamma = 2.8$ GeV and $\sigma_{\text{tot}} = 133 \mu\text{b}$) [10] are given by above fit to the data for $\frac{1}{2}(N_{ep}^{\pi^+} + N_{ep}^{\pi^-})$ with $B = 5.23 \pm 0.68$ ($0.5 < x' \approx z < 0.7$). These points fit the photoproduction data satisfactorily in the region $0.2 < x_F < 0.7$. We would like to point out that above fit in principle can be used only to check the relation (15) for the experimental point $x_F = 0.5$.

Because we see good agreement even for $x_F > 0.5$ for both relations (11) and (15) we conclude that the approximation $f(x') = \delta(x' - \frac{1}{2})$ is not far from reality.

c) Calculation of the $\eta(z)$ from the relation (14):

$$N_{\gamma p}^{\pi^-}(x_F)/N_{\pi^+ p}^{\pi^-}(x_F) = \frac{1}{2} (1 + \eta(z)) \quad .$$

We use data from Ref.[10] for $\gamma p \rightarrow \pi^- + \text{anything}$ (at $E_\gamma = 2.8$ and 4.7 GeV) and data from Ref.[12] for $\pi^+ p \rightarrow \pi^- + \text{anything}$ (at $E_{\pi^+} = 3.7$ GeV).

In this way we get $\eta = 3.0$ with the error of 20% for $0.4 \leq z \leq 0.8$. This number is identical with one obtained by Dakin and Feldman [6] and similar with one obtained by Cleymans and Rodenberg ($\eta = 2.59$ for $0.4 < z < 0.8$) [5].

Conclusions

In the framework of Feynman's parton-quark model and EMM's quark model for inclusive hadronic processes, relations between inclusive electroproduction of pions and purely hadronic inclusive production of pions in the beam fragmentation region are found.

We have considered in detail the case of pion production. We have obtained relations between $N_{ep}^{\pi^\pm}$ and $N_{\pi^\pm p}^{\pi^\pm}$, $N_{\pi^\pm p}^{\pi^\pm}$ or $N_{\gamma p}^{\pi^\pm}$ for beam fragmentation region but restricted to the region $x_F \leq 0.5$. We found that these relations are satisfied by data in the broader region of x_F , namely $0.2 \leq x_F \leq 0.7$. So we conclude that the approximation $f(x') = \delta(x' - \frac{1}{2})$ is not far from reality. The next attempt ought to be done to find more general relations for the whole region of x_F , it means one ought to look for more sophisticated function $f(x')$

different from $\delta(x' - \frac{1}{2})$.

Moreover we calculated from purely hadronic processes the ratio $\eta(z) = D_u^{\pi^+}(z)/D_d^{\pi^+}(z)$. The value obtained in this way agrees with the one obtained by other authors from data on electroproduction analyzed in the framework of parton model.

Acknowledgements

The author would like to thank Professors H. Joos, H. Schopper and G. Weber for the kind hospitality extended to her at DESY. It is a pleasure to thank T. F. Walsh and especially S. Kitakado for helpful and critical discussions. Also thanks are due to D. Schildknecht for reading the manuscript.

Footnotes

- *) In the following we will omit p_{\perp}^2 dependence of D 's. It is understood that all relations are integrated over p_{\perp}^2 .
- **) We won't quote here predictions for inclusive production of kaons (or pions) by pion, kaon or proton beams, but one can use them (additionally with help of SU(3) relations, and with presence of a strange quark fragmentation function D_s^{π}) also to test the consistency of the EMM model with Feynman's quark parton model.
- ***) The variable z is essentially identical with the variable x_F for considered kinematical region of electroproduction.

Figure Captions

Fig.1: Experimental check of the relation (11), see the main text for details.

Fig.2: Experimental check of the relation (15), see the main text for details.

References

- [1] R. P. Feynman; Photon-Hadron Interactions,
(W. A. Benjamin, Reading, Massachusetts, 1972).
- [2] M. Gronau, F. Ravndal and Y. Zarmi;
Nucl. Phys. B51, 611 (1973).
- [3] Z. F. Ezawa, J. Maharana and H. Miyazawa;
Description of Inclusive Reactions in Quark Model,
Preprint Tokyo University, UT-205, July, 1973.
- [4] J. Cleymans and L. M. Sehgal;
Nuclear Phys. B61, 479 (1973).
- [5] J. Cleymans and R. Rodenberg;
Physical Rev., to be published.
- [6] J. T. Dakin and G. J. Feldman, SLAC preprint,
SLAC-PUB-1273, July 1973.
- [7] J. Kuti and V. Weisskopf; Phys. Rev. D4, 3418 (1971),
R. McElhaney and S. F. Tuan; Preprint Hawaii, UH-511-156-73, May 1973.
- [8] H. Satz; Physics Lett. 25B, 220 (1967).
- [9] D.R.O. Morrison; Proc. of the 4th Intern. Conf. Oxford, 1972, p.253.
- [10] Stanford-Berkeley-Tufts Collaboration;
Phys. Rev. D5, 1603 (1972).
- [11] H. Satz; Physics Letters, 25B, 27 (1967).
- [12] M. Alston-Garnjost et al.;
Physics Letters 39B, 402 (1972).
- [13] Harvard-Cornell Collaboration, C. J. Bebek et al.;
Phys. Rev. Letters 30, 624 (1973).





Fig.1

$$F(x) = \frac{1}{\pi} \int_0^\infty \frac{E^*}{p_{\max}^*} \frac{d^2\sigma}{dx dp_I^2} dp_I^2 (\mu b)$$



o $F(\gamma p \rightarrow \pi^- + \text{anything})$

$$\text{I} \frac{\chi}{2\pi} \sigma_{\text{tot}}^{\gamma p} (N_{ep}^{\pi^+} + N_{ep}^{\pi^-})$$

Fig. 2