

# DEUTSCHES ELEKTRONEN-SYNCHROTRON **DESY**

DESY 75/46  
November 1975



Identification of a Possible  $\chi(3.55)$

by

G. Daboul and H. Krasemann

2 HAMBURG 52 · NOTKESTIEG 1

To be sure that your preprints are promptly included in the  
HIGH ENERGY PHYSICS INDEX,  
send them to the following address ( if possible by air mail ) :

DESY  
Bibliothek  
2 Hamburg 52  
Notkestieg 1  
Germany

# Identification of a Possible $\chi(3.55)$

by

J. Daboul<sup>+</sup> and H. Krasemann

Deutsches Elektronen-Synchrotron DESY, Hamburg

## Abstract

We argue in favor of an identification of the  $\chi(3.55)$  as the paracharmonium  $2^1S_0$ ,  $\eta'_c$ . We predict the masses of the charmonium  $2^{++}$  and  $1^{+-}$  states and give estimates for the radiative decays of the  $\psi'$  via  $\eta_c$  and  $\eta'_c$ .

<sup>+</sup> Minerva Fellow, permanent address: Physics Department,  
Ben Gurion University of Negev, Beer Sheva, Israel.

Studying the  $\psi' \rightarrow \psi + 2\gamma$  chain decays, the DASP Collaboration<sup>1</sup> at DESY detected two narrow intermediate states, denoted as  $P_c$ , with masses of 3.41 (or 3.35) and 3.51 (or 3.26) GeV. The LBL-Group at SLAC<sup>2</sup>, measuring the decays  $\psi' \rightarrow \gamma + \text{hadrons}$ , found two resonances in the same mass region, a narrow one at 3.41 GeV and a broader one at  $3.53 \pm .02$  GeV, denoted as  $\chi$ . The existence of such intermediate states was predicted by the charmonium picture<sup>3</sup>. It is plausible to assume that the  $\chi(3.41)$  and the  $P_c(3.41)$  are identical. If the mass of the second  $P_c$  is 3.26 GeV, we would have the three states 3.26, 3.41 and 3.53 GeV. But the more attractive alternative is, that the broad bump  $\chi(3.53)$  is a superposition of two narrow resonances, the  $P_c(3.51)$  and a new narrow resonance  $\chi(3.55)$ <sup>2,4</sup>.

Within the charmonium model it is almost established that the  $P_c/\chi(3.41)$  and the  $P_c/\chi(3.51)$  are the  $j^{PC}=0^{++}$  and the  $j^{PC}=1^{++}$   $c\bar{c}$  states<sup>2,4</sup>. So far the possible  $\chi(3.55)$  has been presumed to be the  $j^{PC}=2^{++}$  state, expected roughly at this mass<sup>3</sup>.

In this letter we want to argue in favor of an identification of the  $\chi(3.55)$  as the paracharmonium  $2^1S_0$  state  $\eta_c'$ . The  $\chi(3.55)$  would then be the radial excitation of the already observed  $1^1S_0$  state  $\eta_c(2.8)$ <sup>5</sup>:

i) Assuming that the S-wave ortho-para splitting can be described mainly by a term in the Hamiltonian<sup>6</sup>

$$c \int^3 (\vec{r}_1 - \vec{r}_2) \vec{S}_1 \cdot \vec{S}_2 + V_T \cdot S_{12} \quad ; \quad V_T \ll c \quad (1)$$

with the constant  $c$ , we are led to the following ratio

$$\frac{M(2^3S) - M(2^1S)}{M(1^3S) - M(1^1S)} \approx \left| \frac{\Psi(2^3S; r=0)}{\Psi(1^3S; r=0)} \right|^2 = \left( \frac{3.7}{3.1} \right)^2 \frac{\Gamma_{e^+e^-}(\Psi')}{\Gamma_{e^+e^-}(\Psi)} = 0.65 \quad (2)$$

where the last equality follows from the experimental values<sup>7</sup>

$\Gamma_{e^+e^-}(\Psi) = 4.8$  keV and  $\Gamma_{e^+e^-}(\Psi') = 2.2$  keV. Using (2) and taking  $M(\Psi) - M(\eta_c) = 0.3$  GeV as input, we obtain

$$M(\eta_c') = M(\Psi') - 0.65 \cdot 0.3 \text{ GeV} = 3.5 \text{ GeV} \quad (3)$$

In view of the crude estimate it is not unreasonable to identify the  $\chi(3.55)$  with the  $\eta_c'$ .

ii) Our identification would imply that the DASP Group cannot observe the  $\chi(3.55)$  in the decay  $\Psi' \rightarrow \Psi + 2\gamma$ . The radiative decay of  $\eta_c' \rightarrow \Psi + \gamma$  is highly suppressed, since this would be a magnetic dipole transition between orthogonal wave functions. In contrast, an identification of  $\chi(3.55)$  as  $2^{++}$  meson would lead to a ratio of  $\frac{\Gamma(\Psi' \rightarrow 2^{++} + \gamma)}{\Gamma(\Psi' \rightarrow 1^{++} + \gamma)} = \frac{5}{3} \left( \frac{k'}{k} \right)^3 \approx 1$  and together with the assumption of comparable branching ratios of  $2^{++} \rightarrow \Psi + \gamma$  and  $1^{++} \rightarrow \Psi + \gamma$  one should see the  $2^{++}$  and the  $1^{++}$  with roughly the same probability in  $\Psi' \rightarrow \Psi + 2\gamma$ .

iii) The ratio of the P level splittings

$$R_1 := \frac{M(2^{++}) - M(1^{++})}{M(1^{++}) - M(0^{++})} \quad (4)$$

depends on the strength of the tensor forces. Without tensor forces, i.e.  $V_T = 0$  in (1), the pure  $\vec{L}\vec{S}$  term gives  $R_1 = 2$ . If tensor forces are deduced from Coulomb-like potentials, similar to the Breit approximation in electrodynamics, one obtains  $R_1 = 0.8$  as in the positronium.<sup>g</sup> This lower bound would give a mass of 3.59 GeV for the  $2^{++}$ , which would still be higher than the assumed  $\chi(3.55)$ . If one applies the same approximation to <sup>the</sup> harmonic oscillator potential one obtains no tensor forces and therefore  $R_1 = 2$ .

We would like to assume in this letter that  $\psi'$  is an almost pure  $2^3S_1$ . Therefore, it is necessary to assume small tensor forces, which lead to  $R_1 \lesssim 2$ . Another argument which supports our assumption of  $R \lesssim 2$  follows from the analogy to the ordinary P wave mesons,  $A_2(1310)$ ,  $A_1(1100)$  and  $\delta(976)$ , where

$$R_1 = \frac{M(A_2) - M(A_1)}{M(A_1) - M(\delta)} \approx 1.7 \quad , \quad (5a)$$

or, if we use a quadratic equation,

$$R_1' = \frac{M^2(A_2) - M^2(A_1)}{M^2(A_1) - M^2(\delta)} \approx 2 \quad (5b)$$

These splittings, especially (5b), are described better by an ordinary  $\vec{L} \cdot \vec{S}$  term than by the analogy to the positronium. Employing the same ratios as in Eqs. (5a,b), with  $P_c(3.41) = 0^{++}$  and  $P_c(3.51) = 1^{++}$ , we get the following predictions for the  $c\bar{c} 2^{++}$  meson

$$M(2^{++}) = 3.68 \text{ GeV} \quad (6a)$$

$$M(2^{++}) = 3.70 \text{ GeV} \quad (6b)$$

In both cases the transitions  $\Psi' \rightarrow 2^{++} + \gamma$  ( or  $2^{++} \rightarrow \Psi' + \gamma$  ) are suppressed by phase space.

We can also use the above analogy to  $A_2, A_1, \delta$  to predict the mass of the  $1^1P_1(j^{PC} = 1^{+-})$  state which corresponds to the B(1235) meson. Using

$$\frac{M^i(1^{+-}) - M^i(0^{++})}{M^i(2^{++}) - M^i(0^{++})} = \frac{M^i(B) - M^i(\delta)}{M^i(A_2) - M^i(\delta)}, \quad i=1,2$$

we get for  $i=1$

$$M(1^{+-}) = 3.62 \text{ GeV} \quad (7a)$$

and for  $i=2$

$$M(1^{+-}) = 3.63 \text{ GeV} \quad (7b)$$

We note that the center of gravity (c.o.g.) of the  $2^3S$  and  $2^1S$  states (which corresponds to  $c=0$  in eq. (2) ) is 3.62 GeV. The c.o.g. of the  $2S$  and the  $1P$  states are almost degenerate, just as in the case of a Coulomb potential. However, a spin-independent Coulomb potential would in a non-

relativistic calculation give a ratio of 1/8 instead of 0.65 in eq. (3).

The M1 transition  $\Psi' \rightarrow \eta_c + \gamma$  is suppressed, because of the orthogonality of the radial wave functions. However, the M1 transition  $\Psi' \rightarrow \eta_c' + \gamma$  is allowed, and we would like to estimate the radiative decay widths of  $\Psi'$  via  $\eta_c'$ . In the absence of a reliable theory, we shall use rough analogy arguments to obtain ratios of these decay widths relative to  $\Gamma(\Psi \rightarrow \eta_c + \gamma \rightarrow 3\gamma)$ . In spite of its narrowness, the cascade  $\Psi' \rightarrow \eta_c + \gamma \rightarrow 3\gamma$  has been observed experimentally, and one finds  $\Gamma(\Psi' \rightarrow \eta_c + \gamma \rightarrow 3\gamma) \approx 10 \text{ eV}^5$ . Therefore, the comparison with this width should be useful as a guide for the feasibility of measuring the radiative decay modes of  $\Psi'$  via  $\eta_c'$ . The above ratios depend crucially on the quantity

$$R := \Gamma_{\text{tot}}(\eta_c) / \Gamma_{\text{tot}}(\Psi) \quad . \quad (8)$$

Experimentally, it has only been possible to ascertain that the width of  $\eta_c$  is smaller than the mass resolution of the DESY experiments<sup>5</sup>, i.e.

$\Gamma_{\text{tot}}(\eta_c) \leq 100 \text{ MeV}$ . This gives the upper bound  $R \leq 5000$ . It seems safe, very safe, to assume a lower bound of  $R \geq 1$ ; according to asymptotic-freedom arguments,  $R$  is roughly equal to  $100^{10}$ .

And now to our analogy arguments: Assuming  $\Gamma(\eta_c' \rightarrow 2\gamma) \approx \Gamma(\eta_c \rightarrow 2\gamma)^{11}$ , we get<sup>12</sup>

$$\begin{aligned} \frac{\Gamma(\Psi' \rightarrow \eta_c' + \gamma \rightarrow 3\gamma)}{\Gamma(\Psi \rightarrow \eta_c + \gamma \rightarrow 3\gamma)} &\approx \frac{\Gamma(\Psi' \rightarrow \eta_c' + \gamma)}{\Gamma(\Psi \rightarrow \eta_c + \gamma)} \cdot \frac{\Gamma_{\text{tot}}(\eta_c)}{\Gamma_{\text{tot}}(\eta_c')} \approx \\ &\approx \left(\frac{145}{300}\right)^3 \cdot \frac{2}{3} \cdot \frac{R}{1+R} \approx 4 - 8 \% \end{aligned} \quad (9)$$



Similarly, assuming  $\Gamma(\chi_c' \rightarrow \chi_c + 2\pi) \approx \Gamma(\psi' \rightarrow \psi + 2\pi)$ , gives <sup>12</sup>

$$\frac{\Gamma(\psi' \rightarrow \chi_c' + \gamma \rightarrow 2\pi + \chi_c + \gamma \rightarrow 2\pi + 3\gamma)}{\Gamma(\psi \rightarrow \chi_c + \gamma \rightarrow 3\gamma)} = \frac{\Gamma(\psi' \rightarrow \chi_c' + \gamma)}{\Gamma(\psi \rightarrow \chi_c + \gamma)} \cdot BR(\chi_c' \rightarrow \chi_c + 2\pi) \approx$$

$$\approx \left(\frac{145}{300}\right)^3 \cdot BR(\psi' \rightarrow \psi + 2\pi) \cdot \frac{\Gamma_{tot}(\psi')}{\Gamma_{tot}(\chi_c')} \approx$$

(10)

$$\approx \left(\frac{145}{300}\right)^3 \cdot \frac{1}{2} \cdot \frac{2}{R+1} \approx 0.002 - 6 \%$$

Note that  $R \gg 1$  corresponds to the upper bound in (9), but to the lower bound in (10). Note also that the width

$$\Gamma(\psi' \rightarrow \chi_c' + \gamma \rightarrow 2\pi + \chi_c + \gamma \rightarrow all + \gamma) = \frac{\Gamma(\psi' \rightarrow \chi_c' + \gamma \rightarrow 2\pi + \chi_c' + \gamma \rightarrow 2\pi + 3\gamma)}{BR(\chi_c \rightarrow 2\gamma)}$$

will also be suppressed <sup>10</sup> if  $R \gg 1$ .

To conclude, if the SLAC experiment with better statistics will split the broad  $\chi(3.53)$  into two resonances  $\chi(3.51)$  and  $\chi(3.55)$  and if the DESY experiment  $\psi' \rightarrow \psi + 2\gamma$  even with better statistics will fail to see the  $\chi(3.55)$ , then it will be worth to look for the  $\chi(3.55)$  in  $\psi' \rightarrow 3\gamma$  or  $\psi' \rightarrow 3\gamma + 2\pi$ . If our assignment of the  $\chi(3.55)$  proves to be correct, the charmonium spectrum will look as in Fig. 1. If, however, it turns out experimentally, that no  $\chi_c'$  exists in the mass region 3.5-3.6 GeV, it would be necessary to explain why. One possibility would be the existence of strong tensor forces <sup>6</sup>, i.e.  $V_T$  is no longer small compared to  $c$  in (1), which would in turn imply that  $\psi'$  is not a pure S wave but a mixture of  $^3S_1$  and  $^3D_1$ .

References and Footnotes

1. DASP Collaboration, Phys. Lett. 57B, 407 (1975)
2. SLAC-LBL Group, Phys. Rev. Lett. 35, 821 (1975)
3. C.G. Callan et al., Phys. Rev. Lett. 34, 52 (1975)  
T. Appelquist et al., *ibid.*, 365 (1975)  
E. Eichten et al., *ibid.*, 369 (1975)
4. Talks given by G. Goldhaber and F. Vanucci at DESY, Hamburg
5. J. Heintze, DESY 75/34 (1975)  
B.H. Wiik, DESY 75/37 (1975)
6. A strong tensor interaction  $V_T(\tau) S_{12}$  where  $S_{12} := 3(\vec{S}_1 \cdot \hat{r})(\vec{S}_2 \cdot \hat{r}) - \vec{S}_1 \cdot \vec{S}_2$  would lead to an S-D mixing for the  $\Psi'$ , and also to a reduction of the ratio  $R_1$  of the  $\bar{P}$  level splittings (see eq. (4)) from the value  $R_1 = 2$  expected for zero tensor force. In this paper however we want to assume that  $\Psi'$  is a pure or almost pure  $2^3S_1$ , so  $V_T(\tau) \ll c$ .
7. V. Lüth, Invited paper presented at the Int. Conf. on High Energy Physics, Palermo (1975) and the review given by G. Feldman, SLAC-PUB 1647
8. Assuming  $\Gamma(2^{++}) \approx \Gamma(1^{++})$ , we get  
$$\frac{BR(2^{++} \rightarrow \psi + \gamma)}{BR(1^{++} \rightarrow \psi + \gamma)} \approx \frac{\Gamma(2^{++} \rightarrow \psi + \gamma)}{\Gamma(1^{++} \rightarrow \psi + \gamma)} \cdot \left(\frac{0.46}{0.41}\right)^3 \approx 1.4$$
9. See, for example, H.J. Schnitzer, Brandeis Preprint (July 1975)
10. Because of charge conjugation,  $\eta_c$  and  $\eta_c'$  can decay into hadrons made up of non-charmed quarks via 2 gluons, whereas the  $\Psi$  and  $\Psi'$  need at least 3 gluons. Using Table I of Appelquist *et al.*<sup>3</sup>, we get  $R \approx 100$  and  $BR(\eta_c \rightarrow 2\gamma) \approx 0,02$
11. This assumption holds, if the increase in phase space is roughly compensated by the spreading of the wave function.
12. Assuming  $\Gamma_{\eta_c}(\eta_c') := \Gamma(\eta_c' \rightarrow \eta_c + 2\pi) \approx \Gamma_{\psi}(\Psi')$  and  $\Gamma_{\text{hadr}}(\eta_c')/\Gamma_{\text{hadr}}(\eta_c) \approx \Gamma_{\text{hadr}}(\Psi')/\Gamma_{\text{hadr}}(\Psi)$  gives together with the experimental result<sup>7</sup>  
 $\Gamma_{\psi}(\Psi') \approx \Gamma_{\text{hadr}}(\Psi')$   
$$\frac{\Gamma_{\text{tot}}(\eta_c')}{\Gamma_{\text{tot}}(\Psi')} \approx \frac{\Gamma_{\eta_c}(\eta_c') + \Gamma_{\text{hadr}}(\eta_c')}{\Gamma_{\text{tot}}(\Psi')} \approx \frac{\Gamma_{\psi}(\Psi') + \Gamma_{\text{hadr}}(\eta_c')}{2 \Gamma_{\text{hadr}}(\Psi')} \approx \frac{1 + R}{2}$$

Using this result and the experimental ratio  $\frac{\Gamma(\psi')}{\Gamma(\psi)} \approx 3$   
we get  $\frac{\Gamma(\psi_c)}{\Gamma(\psi_c')} \approx \frac{\Gamma(\psi_c)}{3\Gamma(\psi)} \frac{\Gamma(\psi')}{\Gamma(\psi_c')} \approx \frac{R}{3} \cdot \frac{2}{R+1}$




Figure 1. Masses and not suppressed radiative transitions of charmonium