

# DEUTSCHES ELEKTRONEN-SYNCHROTRON DESY

DESY 77/14  
February 1977



Inclusive  $K_S^0$ -Production in  $e^+e^-$  Annihilation

at Energies of 3.6 to 5.0 GeV

by

*PLUTO-Collaboration*

J. Burmester, L. Criegee, H. C. Dehne, K. Derikum, R. Devenish,  
J. D. Fox, G. Franke, G. Flügge, Ch. Gerke, G. Horlitz, Th. Kahl,  
G. Knies, M. Rössler, R. Schmitz, U. Timm, H. Wahl,  
P. Waloschek, G. G. Winter, S. Wolff, W. Zimmermann

*Deutsches Elektronen-Synchrotron DESY, Hamburg*

V. Blobel, H. Jensing, B. Koppitz and E. Lohrmann

*Lehrstuhl für Experimentalphysik der Universität Hamburg*

A. Bäcker, J. Bürger, C. Grupen, M. Rost and G. Zech

*Gesamthochschule Siegen*

and

H. Meyer and K. Wacker

*Gesamthochschule Wuppertal*

NOTKESTRASSE 85 · 2 HAMBURG 52

To be sure that your preprints are promptly included in the  
HIGH ENERGY PHYSICS INDEX ,  
send them to the following address ( if possible by air mail ) :

DESY  
Bibliothek  
2 Hamburg 52  
Notkestieg 1  
Germany

Inclusive  $K_s^0$ -production in  $e^+e^-$  annihilation at energies of 3.6 to 5.0 GeV

PLUTO-Collaboration

J. Burmester, L. Criegee, H.C. Dehne, K. Derikum, R. Devenish, J. D. Fox,  
G. Franke, G. Flügge, Ch. Gerke, G. Horlitz, Th. Kahl, G. Knies, M. Rössler,  
R. Schmitz, U. Timm, H. Wahl\*, P. Walešček, G. G. Winter, S. Wolff,  
W. Zimmermann

Deutsches Elektronen Synchrotron DESY  
Hamburg

V. Blobel, H. Jensing, B. Koppitz and E. Lehrmann\*\*

II. Institut für Experimentalphysik der Universität Hamburg

A. Bäcker, J. Bürger, C. Grupen, M. Rost and G. Zech  
Gesamthochschule Siegen

and

H. Meyer and K. Wacker  
Gesamthochschule Wuppertal

The total cross section for  $e^+e^-$  annihilation into hadrons ( $\sigma_h$ ) normalized to the muon pair cross section ( $\sigma_{\mu\mu}$ ) shows a clear step of about two units at 4 GeV center of mass energy. In addition the total cross section is not flat above 4.0 GeV but shows considerable structure <sup>1, 2</sup>. This is usually attributed to the production of pairs of charmed quarks, adding  $3 \times (2/3)^2$  units of  $R = \sigma_h/\sigma_{\mu\mu}$  to the total cross section. The charmed quarks then fragment into a pair of charmed hadrons which subsequently decay by weak interaction preferentially into strange particles. A sizeable increase in kaon production is then expected above threshold for charmed particle production. Furthermore the resonance like structures in the total cross section at 4.03 and 4.415 GeV are assumed to have strong coupling to charmed hadrons. This should be detectable by measuring the strange particle yield in this energy region.

We have measured the  $K_s^0$  production in  $e^+e^-$  annihilation into hadrons using the magnetic detector PLUTO at the  $e^+e^-$  storage ring facility DORIS at DESY. Results on  $K_s^0$ -e correlations<sup>3</sup> as evidence for the semileptonic decay mode of charmed particles and on the total cross section<sup>2</sup> have been reported. Reference should be made to those papers for details of the experimental procedure. Briefly, the detector uses a superconducting solenoid to produce a field of 20 kG. The field volume is filled with 14 cylindrical proportional chambers used for triggering and reconstruction of events. The luminosity is measured by small angle  $e^+e^- \rightarrow e^+e^-$  scattering, as determined by four counter telescopes. The detection efficiency for hadronic events is ~ 80 %.

We have searched for  $K_s^0$  production by calculating the effective mass of oppositely charged pairs of particles assuming they are pions. To suppress background from normal pion pairs a minimum distance of 15 mm from the decay vertex to the  $e^+e^-$  interaction point was required. A vertex fit is made constraining the momentum vector of the pion pair to the interaction point. The mass distribution for pion pairs satisfying the above criteria is shown in Fig. 1. The background below the  $K_s^0$  mass has been estimated from side bins and subtracted for all the following plots. The efficiency to find a  $K_s^0 \rightarrow \pi^+\pi^-$  with this method is momentum dependent and has been determined by a Monte Carlo study. The production angular distribution was assumed to be isotropic. The detection efficiency is essentially zero

Abstract

We have measured the production cross section for  $K_s^0$  in  $e^+e^-$  annihilation from 3.6 to 5.0 GeV center of mass energy. A substantial increase of the  $K_s^0$  yield is observed around 4 GeV in qualitative agreement with the charm hypothesis.

\* on leave from CERN

\*\* now at CERN

below 150 MeV/c. It rises smoothly, and goes through a maximum of 21% at 1 GeV/c. We cut at 200 MeV/c, and apply no correction for the loss of lower momentum kaons. The loss can be estimated by extrapolating the observed momentum dependence to zero momentum, and amounts to ~10% of the observed yield. Trigger losses not accounted for in the Monte Carlo calculations are estimated to be less than 10%. The data have been corrected for the unobserved decay mode  $K_s^0 \rightarrow \pi^0 \pi^0$ .

Sizeable  $K_s^0$  production is found at all energies. The cross section for inclusive  $K_s^0$  production is shown in Fig. 2a. A prominent peak at 4.03 GeV is observed. For higher energies the  $K_s^0$  cross section shows little structure but has a higher level than at 3.6 GeV.

The increase in  $K_s^0$  production is dominantly due to kaons with energies less than half the beam energy. This is evident from Fig. 3a, b, where we show the invariant cross section  $(s/B) \cdot d^2\sigma/dx$  versus  $x = E(K_s^0)/E(\text{beam})$ , at  $\sqrt{s} = 4.03$  GeV and  $\sqrt{s} > 4.03$  GeV each compared to the data at  $\sqrt{s} = 3.6$  GeV. At  $x > 0.5$  the spectra are seen to be fairly similar, as expected in quark fragmentation models.

#### Discussion

The excess  $K_s^0$  which we observe at  $\sqrt{s} = 4.03$  GeV as compared to  $\sqrt{s} = 3.6$  GeV are concentrated at  $x < 0.5$ . This supports the hypothesis that they come from the decay of intermediate charmed pairs. Since the  $K_s^0$  increase is largest for the very small  $x$  values we infer a dominance of multibody decays in agreement with previous indications<sup>3) 4) 5)</sup>.

We next determine the kaon yield per annihilation event above threshold for charmed particle production. We do so in the spirit of the quark model for  $e^+e^-$  annihilation into hadrons. The basic mechanism is assumed to be the production of a pair of quarks which subsequently fragments into hadrons. Above  $\sqrt{s} = 4$  GeV, we use the following expression for the normalized total cross section:

$$(1) \quad \sigma_H/\sigma_{\mu\mu} = R(\text{hadron}) = R_{q\bar{q}} + R_{c\bar{c}} + R_{L\bar{L}}$$

The symbol  $q$  refers to  $n, p, \Lambda$  quarks and  $c$  refers to the charm quark. A contribution  $R_{L\bar{L}}$  due to the production of heavy leptons has been included. Evidence for this lepton has been reported by the SLAC-LBL group<sup>6)</sup> and has been confirmed by a recent PLUTO experiment<sup>7)</sup>.

Equation (1) can also be written for the inclusive  $K_s^0$  cross section.

$$(2) \quad \sigma(K_s^0/\sigma_{\mu\mu}) = R(K_s^0) = R_{q\bar{q}}(K_s^0) + R_{c\bar{c}}(K_s^0) + R_{L\bar{L}}(K_s^0)$$

$R_{q\bar{q}}$  and  $R_{q\bar{q}}(K_s^0)$  have been determined at  $\sqrt{s} = 3.6$  GeV and are assumed to be independent of  $\sqrt{s}$ . The new part  $\Delta R$  due to the charm and heavy lepton contribution is then given by

$$\Delta R = R - R_{q\bar{q}} \quad (\sqrt{s} = 3.6)$$

The ratio  $\Delta R(K_s^0)/\Delta R$  is the number of  $K_s^0$  per new event. Using our measured total cross section\* and the  $K_s^0$  cross section in Fig. 2a we calculate  $\Delta R(K_s^0)/\Delta R$  shown in Fig. 2b (open points). No radiative corrections have been applied since they should approximately cancel in the ratio. The contribution due to the heavy lepton is expected to yield very few kaons<sup>8)</sup>. With the assumptions  $R_{L\bar{L}}(K_s^0) = 0$  and  $M(L) = 1.95$  GeV we determine the number of  $K_s^0$  per charm event (full points in Fig. 2b). Assuming  $2 \cdot N(K_s^0) = N(K^0 + \bar{K}^0) = N(K^+ + K^-)$  as supported by a recent experiment<sup>9)</sup>, we expect  $N(K_s^0)/\text{charm event} = 0.5$  (neglecting the Cabibbo angle). The data points, taking into account the heavy lepton contribution (full points), are slightly less than this prediction.

In the total cross section resonance like structures have been observed at 4.03 and 4.415 GeV<sup>1) 2)</sup>. At both energies we have taken fairly large data sets. Comparing the normalized  $K_s^0$  yields at the two energies (see the arrows in Fig. 2b), we have an indication of substantially less  $K_s^0$  production at 4.415 GeV.

In conclusion we find a significant increase of  $K_s^0$  yield going from  $\sqrt{s} = 3.6$  GeV to  $\sqrt{s} > 4.0$  GeV. This is very well compatible with the charm hypothesis. Just above threshold, at  $\sqrt{s} = 4.03$  GeV, we find a stronger  $K_s^0$  production than at higher energies, and there is an indication of reduced  $K_s^0$  production at the position of the 4.415 GeV resonance.

#### Acknowledgements

We thank the operation group of the storage ring for their continuous and successful effort, we are grateful to our technicians, particularly those of the cryogenic team, for their competent service during the experiment. We are indebted to the operators of the computer center for their efficient cooperation. The non-DESY members of the PLUTO-group want to thank the DESY directorate for the kind hospitality extended to them.

\* Values are taken from Ref. 2) and from more recent data (unpublished).

References

1. J. Siegrist et al., Phys. Rev. Lett. 36, 700 (1976)
2. PLUTO Collaboration, J. Burnester et al., DESY 75/53 and to be published in Phys. Lett.
3. PLUTO Collaboration, J. Burnester et al., Phys. Lett. 64B, 369 (1976)
4. DASP Collaboration, W. Braunschweig et al., Phys. Lett. 63B, 471 (1976)
5. G. Goldhaber et al., Phys. Rev. Lett. 37, 255 (1976)
6. I. Peruzzi et al., Phys. Rev. Lett. 37, 569 (1976)
7. M. Perl et al., Phys. Rev. Lett. 35, 1489 (1975); 38, 117 (1977)
8. PLUTO Collaboration to be published; H. Meyer, Talk given at the Coral Gables Meeting, January 1977
9. Y. S. Tsai, Phys. Rev. D4 2821 (1971)
10. DASP Collaboration, R. Brandelik et al., DESY 77/12 and to be published in Phys. Lett.

Figure Captions

Fig.1 Mass distribution of  $\pi^+\pi^-$  pairs with a vertex outside the  $e^+e^-$  interaction point (see text)

Fig.2 (a) The cross section for  $e^+e^- \rightarrow K_s^0 + \text{anything}$  as function of the total center of mass energy. No radiative corrections have been applied.

(b) The number of  $K_s^0$  per new event  $\Delta R(K_s^0)/\Delta R$  ( $\phi$ ); ( $\phi$ ) accounting for a heavy lepton contribution.

Fig.3 (a, b) The invariant cross section  $(s/B) d\sigma/dx$  versus  $x$  for 3.6 GeV ( $\phi$ ), 4.03 GeV ( $\phi$ ) and ( $\phi$ )  $> 4.03$  GeV.




Fig.1





Fig.3





Fig.2