

DESY 77/32
June 1977



SO(4)-Symmetric Solutions of Minkowskian Yang-Mills Field Equations

by

M. Lüscher

II. Institut für Theoretische Physik der Universität Hamburg

Abstract:

We construct all solutions to the SU(2) Yang-Mills field equations in Minkowski space that are invariant under an SO(4) subgroup of the conformal group. They are real, regular and have finite energy and action. A connection with the instanton solution is pointed out.

To be sure that your preprints are promptly included in the
HIGH ENERGY PHYSICS INDEX ,
send them to the following address (if possible by air mail) :

DESY
Bibliothek
Notkestraße 85
2 Hamburg 52
Germany

SO(4)-Symmetric Solutions of Minkowskian Yang-Mills Field Equations

M. Lüscher

II. Institut für Theoretische Physik der Universität Hamburg

I. The similarity between the four dimensional SU(2) Yang-Mills field and the two dimensional O(3) nonlinear σ -model [3] with regard to pseudo-particles [4] suggests that the Minkowskian classical Yang-Mills field theory is completely integrable, too. However, so far we know of only one non-trivial, real finite energy solution to the Yang-Mills field equations in Minkowski space. This one solution was found (in a complex form) by De Alfaro, Fubini and Furlan [1] and was later shown to be gauge equivalent to a real solution by Cervero, Jacobs and Nohl [2].

In this letter we exploit the invariance of the Yang-Mills field equations under an SO(4)-subgroup of the Minkowskian conformal group. First we rewrite the field equations in a manifestly SO(4)-covariant way.

Then, upon mixing isospin with one of the SO(4)-spins, we obtain all SO(4)-symmetric solutions. They are real, regular, non-abelian and have finite energy and action.

II. We consider an SU(2) Yang-Mills field $A_\mu = A_\mu^a \frac{\sigma^a}{2i}$ in Minkowski space * . The action is

$$(1) \quad S = -\frac{1}{4g^2} \int d^4x F_{\mu\nu}^a F^{\mu\nu a}$$

where

$$(2) \quad F_{\mu\nu} = F_{\mu\nu}^a \frac{\sigma^a}{2i} = \partial_\mu A_\nu - \partial_\nu A_\mu + [A_\mu, A_\nu]$$

* Greek indices μ, ν, ρ, \dots run from 0 to 3 and are subject to the metric $g_{\mu\nu} = \text{diag}(+, -, -, -)$. Greek indices $\alpha, \beta, \delta, \dots$ resp. latin indices a, b, \dots , k, l, \dots take on values from 1 to 4 resp. 1 to 3. Repeated indices are always summed over.

Abstract: We construct all solutions to the SU(2) Yang-Mills field equations in Minkowski space that are invariant under an SO(4) subgroup of the conformal group. They are real, regular and have finite energy and action. A connection with the instanton solution is pointed out.

Correspondingly, the field equations are

$$(3) \quad \partial_\mu F^{\mu\nu} + [A_\mu, F^{\mu\nu}] = 0$$

They are conformally invariant, i.e. if A_μ is a solution and $x^\mu \rightarrow x'^\mu$ is a conformal mapping in Minkowski space then

$$4) \quad A'_\mu(x) = A_\nu(x'(x)) \cdot \frac{\partial x'^\nu}{\partial x^\mu}$$

solves the equations (3), too.

The conformal group in Minkowski space is isomorphic to $O(4,2)$. The action of an element $\Lambda \in O(4,2)$ on a point x is conveniently written down by using projective coordinates for x , viz. we identify x with a ray in the cone [5]

$$C_{4,2} = \{ \zeta^A, A=0, \dots, 5 \mid (\zeta^0)^2 - (\zeta^1)^2 - (\zeta^2)^2 - (\zeta^3)^2 - (\zeta^4)^2 + (\zeta^5)^2 = 0 \}$$

$$(5) \quad x^\mu = \frac{\zeta^\mu}{\zeta^4 + \zeta^5}$$

The conformal mapping $x^\mu \rightarrow x'^\mu$ corresponding to Λ is then given by:

$$x'^\mu = \frac{\zeta'^\mu}{\zeta'^4 + \zeta'^5} ; \quad \zeta'^A = \Lambda^A_B \zeta^B$$

Let us now focus on the $SO(4)$ subgroup of $O(4,2)$, which acts on $\zeta^\alpha, \alpha=1, \dots, 4$, and leaves ζ^0 and ζ^5 fixed. An $SO(4)$ adapted parametrization of the rays in $C_{4,2}$ is:

$$(6) \quad \zeta = r (\sin \tau, r^1, r^2, r^3, r^4, \cos \tau)$$

This amounts to a parametrization of space-time points according to:

$$(7) \quad x^0 = \frac{\sin \tau}{\cos \tau + n^4} ; \quad x^k = \frac{n^k}{\cos \tau + n^4}$$

The space $\tilde{M} = \{(\tau, n) \mid \tau \in \mathbb{R}, n \in S^3\}$ will be called "superworld". The mapping (7) is one-to-one on the subspace $M = \{(\tau, n) \mid |\tau| < \pi, (\cos \tau + n^4) > 0\}$. Therefore, we may identify Minkowski space with M .

The superworld \tilde{M} has previously been used in conformal quantum field theory. Its outstanding features are firstly that the universal covering of the conformal group acts on it in a differentiable manner and secondly that it allows for a conformally invariant causal ordering of its points. For details, see [6].

In order to rewrite the field equations (3) in (τ, n) - coordinates it is convenient to introduce new vector potentials B_μ :

$$(8) \quad A_\mu dx^\mu = B_\mu \omega^\mu \quad \text{where} \quad \omega^0 = d\tau, \quad \omega^k = \eta^k_{\alpha\beta} n^\alpha dn^\beta$$

$\eta^k_{\alpha\beta}$ denote 't Hooft's ([7], appendix) $SO(4)$ - covariant η -symbols, i.e.:

$$(9) \quad \eta^k_{\alpha\beta} = -\eta^k_{\beta\alpha}, \quad \eta^k_{\ell 4} = \delta^k_\ell, \quad \eta^k_{\ell j} = \epsilon_{kj\ell}$$

Thus, under the conformal transformations

$$\tau' = \tau ; \quad n'^\alpha = \Lambda^\alpha_\beta n^\beta, \quad \Lambda \in SO(4)$$

the B-field transforms as:

$$(10) \quad B_0^1(\tau, n) = B_0(\tau, \Lambda n) ; \quad B_k^1(\tau, n) = B_k(\tau, \Lambda n) R(\Lambda)^k_k$$

$R(\Lambda)$ is a three dimensional orthogonal representation of $SO(4)$:

$$(11) \quad R(\Lambda)^k_\epsilon \eta_{\alpha\beta}^\epsilon = \eta_{\delta\gamma}^k \Lambda^\delta_\alpha \Lambda^\gamma_\beta$$

Let us also define a new field tensor $G_{\mu\nu}$ by

$$(12) \quad F_{\mu\nu} dx^\mu \wedge dx^\nu = G_{\mu\nu} \omega^\mu \wedge \omega^\nu ; \quad G_{\mu\nu} = -G_{\nu\mu}$$

In terms of the vector potential this becomes:

$$(13) \quad G_{0k} = \nabla_0 B_k - \nabla_k B_0 + [B_0, B_k] \\ G_{\ell k} = \nabla_\ell B_k - \nabla_k B_\ell + 2 \epsilon_{\ell k j} B_j + [B_\ell, B_k]$$

where we have introduced covariant derivatives:

$$(14) \quad \nabla_0 = \frac{\partial}{\partial \tau} , \quad \nabla_k = \frac{1}{2} \eta_{\alpha\beta}^k \left(n^\alpha \frac{\partial}{\partial n^\beta} - n^\beta \frac{\partial}{\partial n^\alpha} \right)$$

They are dual to the differential forms ω^A , i.e. for any function $f(\tau, n)$ we have:

$$df = \omega^\mu \nabla_\mu f$$

Note also that ∇_k and ∇_ℓ do not commute:

$$[\nabla_k, \nabla_\ell] = -2 \epsilon_{k\ell j} \nabla_j$$

From eq. (1) we now obtain the action in the new language:

$$(15) \quad S = - \frac{1}{2g^2} \int_{-\tilde{K}}^{\tilde{K}} d\tau \int d^4n \delta(1 - n^\alpha n^\alpha) \theta(\cos \tau + n^4) G_{\mu\nu}^a G^{a\mu\nu}$$

As expected, the integration runs over the subspace N of the superworld \tilde{M} , i.e. over a region of finite volume. Hence, any regular field B_μ on \tilde{M} has finite action.

Finally, the action principle yields the field equations:

$$(16) \quad \nabla_\ell G_{\ell 0} + [B_\ell, G_{\ell 0}] = 0 \\ \nabla_0 G_{0k} + [B_0, G_{0k}] - \nabla_\ell G_{\ell k} - [B_\ell, G_{\ell k}] + \epsilon_{k\ell j} G_{\ell j} = 0$$

III. We now seek $SO(4)$ -symmetric solutions to the field equations (16). Thus, we are concentrating on potentials B_μ^a satisfying:

$$(17) \quad B_0^a(\tau, \Lambda n) = \pi(\Lambda)^a_b B_0^b(\tau, n) \\ B_k^a(\tau, \Lambda n) = \pi(\Lambda)^a_b B_k^b(\tau, n) R(\Lambda^{-1})^l_k$$

for all $\Lambda \in SO(4)$. Here, $\pi(\Lambda)$ denotes some three dimensional orthogonal representation of $SO(4)$. A careful analysis shows that all such fields are gauge equivalent to

$$(18) \quad B_0^a = 0 , \quad B_k^a(\tau, n) = q(\tau) \delta_k^a , \quad \pi(\Lambda) = R(\Lambda)$$

where q is an arbitrary (real) function of τ .

Let us insert the Ansatz (18) into the field equations:

$$(19) \quad G_{0k}^a = \dot{q} \delta_k^a \quad ; \quad G_{kl}^a = q(q+2) \epsilon_{akl}$$

$$(20) \quad \ddot{q} + 2q(q+1)(q+2) = 0$$

(the dots denote derivation with respect to τ). This is the equation of motion of a particle moving in the double well potential

$$V(q) = \frac{1}{2} q^2 (q+2)^2.$$

There are two types of solutions of eq. (20): when the "energy"

$$(21) \quad \epsilon = \frac{1}{2} \{ \dot{q}^2 + q^2 (q+2)^2 \}$$

is smaller than the bump of $V(q)$ at $q = -1$, we find

$$(22) \quad q = -1 \pm (1 + \sqrt{2\epsilon})^{1/2} \operatorname{dn} \{ (1 + \sqrt{2\epsilon})^{1/2} (\tau - \tau_0) ; k_1 \}$$

$$k_1^2 = \frac{2\sqrt{2\epsilon}}{1 + \sqrt{2\epsilon}} \quad ; \quad \epsilon \leq 1/2$$

whereas when $\epsilon > 1/2$ the solution is:

$$(23) \quad q = -1 + (1 + \sqrt{2\epsilon})^{1/2} \operatorname{cn} \{ (\delta\epsilon)^{1/2} (\tau - \tau_0) ; k_2 \}$$

$$k_2^2 = \frac{1 + \sqrt{2\epsilon}}{2\sqrt{2\epsilon}} \quad ; \quad \epsilon > 1/2$$

(dn and cn denote Jacobian elliptic functions, [8] § 8.14).

There is also a τ -independent, but unstable solution, namely:

$$(24) \quad q = \text{constant} = -1$$

When translated back to x -coordinates one recovers the solution of De Alfaro et al. in a real gauge.

All the solutions above are regular functions on \tilde{M} . Their restriction to M therefore provides a set of regular, finite action solutions of the original equations (3). A simple expression for the energy-momentum tensor

$$(25) \quad \Theta_{\mu\nu} = \frac{1}{g^3} \left\{ -F_{\mu\lambda}^a F_{\nu}^a{}^\lambda + \frac{1}{4} g_{\mu\nu} F_{\rho\sigma}^a F^{\rho\sigma} \right\}$$

is obtained at $x^0 = 0$:

$$(26) \quad \Theta_{\mu\nu} = 0 \quad \text{for } \mu \neq \nu \quad ; \quad \Theta_{00} = \frac{3\epsilon}{g^3} \frac{16}{(1+x^2)^4} \quad , \quad \Theta_{kl} = \delta_{kl} \frac{\epsilon}{g^3} \frac{16}{(1+x^2)^4}$$

Hence, the energy of these solutions is finite, too.

For the sake of completeness we finally reproduce the solutions in their x -coordinate form:

$$(27) \quad A_0^a = 4q(\tau(x)) \delta^2 x^0 x^a \quad ; \quad \delta = [(1+x^\mu x_\mu)^2 + 4x^2]^{-1/2}$$

$$A_k^a = -4q(\tau(x)) \delta^2 \left\{ \frac{1}{2} (1+x^\mu x_\mu) \delta_k^a + \epsilon_{akj} x^j + x^a x^k \right\}$$

where $q(\tau)$ is one of the functions (22), (23) or (24) and $\tau = \tau(x)$ must be calculated from

$$\sin \tau = 2x^0 \delta \quad ; \quad \cos \tau = (1-x^\mu x_\mu) \delta \quad ; \quad |\tau| < \pi$$

IV. There is a simple connection between the mechanics of SO(4)-symmetric fields described here and the one (anti-) instanton solutions in euclidean space. In terms of (τ, \mathbf{r}) -coordinates the euclidean equations of motion are obtained by performing the following substitutions:

$$(28) \quad \begin{aligned} \sigma &= i\tau, & \mathbf{B}_0^E &= (-i) \mathbf{B}_0, & \mathbf{B}_k^E &= \mathbf{B}_k \\ G_{0k}^E &= (-i) G_{0k}, & G_{jk}^E &= G_{jk} \end{aligned}$$

The variable σ ranges from $-\infty$ to $+\infty$, i.e. the euclidean action is:

$$(29) \quad S^E = \frac{1}{2g^2} \int_{-\infty}^{\infty} d\sigma \int d^4x \delta(1 - \mathbf{r}^2 \mathbf{r}^2) G_{\mu\nu}^a G_{\mu\nu}^a$$

For SO(4) symmetric fields (eq. (18)) this becomes

$$(30) \quad S^E = \frac{6\pi^4}{g^2} \int d\sigma \left\{ \frac{1}{2} (\dot{q}^E)^2 + V(q^E) \right\}$$

and the field equations reduce to

$$(31) \quad \ddot{q}^E - 2q^E (q^E + 1)(q^E + 2) = 0$$

The potential $V(q)$ has two minima, one for vanishing fields and one at $q = -2$, i.e. for

$$(32) \quad \mathbf{B}_k = \mathbf{u}^{-1} \nabla_k \mathbf{u}, \quad \mathbf{u} = \mathbf{n}^k - i \mathbf{n}^k \sigma^k$$

which is a pure gauge. The mapping $n \rightarrow u(n)$ has winding number one [9].

In quantum mechanics tunneling takes place between the two separated vacua. In leading order of \hbar the amplitude for this process is given by the contribution to the euclidean path integral of those histories which interpolate between the vacua in question and that minimize the action (30) [9,10]. These configurations are:

$$(33) \quad q^E(\sigma) = -2 \left[1 + e^{\pm 2(\sigma - \sigma_0)} \right]^{-1}$$

In euclidean x-space these solutions turn out to be precisely the one (anti-) instanton solutions (in a singular gauge).

Acknowledgement: I thank H. Lehmann for a critical reading of the manuscript.

References

- 1 - V. De Alfaro, S. Fubini, G. Furlan: Phys. Lett. 65B, 163 (1976)
- 2 - J. Cervero, I. Jacobs, C. Nohl: Harvard preprint, CTTP 614 (1977) (submitted to Phys. Lett.)
- 3 - A. Belavin, A. Polyakov: JETP Lett. 22, 245 (1975)
- 4 - K. Pohlmeier: Comm. Math. Phys. 46, 207 (1976)
- 5 - A. Belavin, A. Polyakov, A. Schwartz, Y. Tyupkin: Phys. Lett. 59B, 85 (1975)
- 6 - R. Jackiw, C. Nohl, C. Rebbi: Phys. Rev. D 15, 1642 (1977)
- 7 - G. Mack, A. Salam: Ann. Phys. (N.Y.) 53, 174 (1969)
- 8 - M. Lüscher, G. Mack: Comm. Math. Phys. 41, 203 (1975)
- 9 - G. 't Hooft: Phys. Rev. D 14, 3432 (1976)
- 10 - I.S. Gradshteyn, I.P. Ryzhik: "Table of integrals, series and products", Academic Press (1965)
- 11 - R. Jackiw, C. Rebbi: Phys. Rev. Lett. 37, 172 (1976)
- 12 - D. McLaughlin: J. Math. Phys. 13, 1099 (1972)

