

Strictification of weakly equivariant Hopf algebras

Jennifer Maier, Thomas Nikolaus and Christoph Schweigert

*Fachbereich Mathematik, Universität Hamburg
Bereich Algebra und Zahlentheorie
Bundesstraße 55, D-20146 Hamburg*

Abstract

A weakly equivariant Hopf algebra is a Hopf algebra A with an action of a finite group G up to inner automorphisms of A . We show that each weakly equivariant Hopf algebra can be replaced by a Morita equivalent algebra A^{str} with a strict action of G and with a coalgebra structure that leads to a tensor equivalent representation category. However, the coproduct of this strictification cannot, in general, be chosen to be unital, so that a strictification of the G -action can only be found on a *weak* Hopf algebra A^{str} .

1 Introduction

This paper is a supplement to our paper [MNS11]. In that paper we constructed a 3-dimensional equivariant topological field theory which is a generalization of the well-known Dijkgraaf-Witten theory [DW90, FQ93]. Our generalization is equivariant with respect to a finite group G (which was called J there). Our motivation comes from orbifold constructions in conformal field theory.

It is well known that one can extract a modular category \mathcal{C} from a 3-dimensional topological field theory, at least up to some technical subtleties [BK01, Chapter 4 & 5], involving properties of the dualities. A modular category is, in particular, a tensor category. If the initial topological field theory is moreover G -equivariant, the category \mathcal{C} carries additionally a G -grading and an action of G that is compatible with the tensor product. Such a structure is called a G -equivariant tensor category resp. G -modular category [Kir04, Tur10].

In general the action of the group G on a G -modular category \mathcal{C} is given by tensor functors $\phi_g : \mathcal{C} \rightarrow \mathcal{C}$ together with compositors $\phi_g \circ \phi_h \xrightarrow{\sim} \phi_{gh}$, subject to coherence laws for threefold products. It has been demonstrated by Müger [Tur10, Appendix 5] that one can replace \mathcal{C} by an equivalent category \mathcal{C}^{str} with a *strict* action of G , i.e. there the compositors are given by the identity: $\phi_g \circ \phi_h = \phi_{gh}$.

Now consider the G -modular category \mathcal{C} which belongs to our equivariant Dijkgraaf-Witten theory mentioned at the beginning. Although the category \mathcal{C} can relatively easily be described abstractly, it is very hard to work with it explicitly when it comes to orbifolding and

showing modularity. Therefore in [MNS11, section 4] we realized \mathcal{C} as the representation category of a certain algebra A , which we called the equivariant Drinfel'd double. The fact that \mathcal{C} is a tensor category is reflected by the fact that A is a Hopf algebra. Furthermore there is also an algebraic structure on A belonging to the G -action on the representation category. This structure is not just a G -action on A , as one might naively expect, but a *weak* G -action, which is an action by Hopf algebra automorphisms $\varphi_g : A \rightarrow A$ such that $\varphi_g \circ \varphi_h$ equals φ_{gh} only up to an inner automorphism of A . This weakening of the G -action reflects the fact that the action on the category is only weak in the sense that we have isomorphisms $\phi_g \circ \phi_h \xrightarrow{\sim} \phi_{gh}$ of functors rather than equalities. In order to accommodate the example of the algebra A , we had to introduce the notion of Hopf algebra with weak G -action ([MNS11, definition 4.13]), generalizing the notion of Hopf algebra with strict G -action considered before [Tur10, Vir02].

In the light of Müger's observation that one can replace a G -equivariant tensor category \mathcal{C} by an equivalent category \mathcal{C}^{str} with strict G -action it is a natural question to ask whether one can replace a Hopf algebra A with weak G -action by a Hopf algebra A^{str} with strict G -action such that the representation categories are equivalent as tensor categories. A first result of this paper asserts that this is not possible in general, see Theorem 3.2. The reason is that the Hopf algebra axioms are too rigid: the tensor product of the representation category is, in the case of Hopf algebras, directly inherited from the underlying tensor product of vector spaces. Weak Hopf algebras [BNS99, BS00, NV02] have been introduced to provide a more flexible notion for the tensor product. Note that the qualifier weak here refers to a weakening of the bialgebra axioms (i.e. a weakening of the unitality of the coproduct or, equivalently, of the counitality of the product) and should not be confused with 'weak G -action'. We refer to the appendix for a table summarizing the situation.

Thus, a refined version of the question posed above would be whether one can replace a Hopf algebra A with weak G -action by a *weak* Hopf algebra A^{str} with *strict* G -action such that the representation categories are equivalent. The second main result of the present paper is to show that this is indeed possible, see Theorem 4.1. The given concrete construction of A^{str} is inspired by Müger's strictification procedure [Tur10, Appendix 5] on the level of categories. Nevertheless we present it in an independent and elementary manner which requires no knowledge about orbifold categories and other constructions that enter in the categorical strictification.

Acknowledgements. We thank Alexander Barvels and Jürgen Fuchs for helpful discussions. TN and CS are partially supported by the Collaborative Research Centre 676 "Particles, Strings and the Early Universe - the Structure of Matter and Space-Time" and the cluster of excellence "Connecting particles with the cosmos". JM and CS are partially supported by the Research priority program SPP 1388 "Representation theory".

2 Equivariant Hopf algebras and their representation categories

In the following, let G be a finite group.

Definition 2.1. *Let A be an (associative, unital) algebra over a field \mathbb{K} . A weak G -action on A consists of algebra automorphisms $\varphi_g \in \text{Aut}(A)$, one for every element $g \in G$, and*

invertible elements $c_{g,h} \in A^\times$, one for every pair of elements $g, h \in G$, such that for all $g, h, k \in G$ the following conditions are satisfied:

$$\varphi_g \circ \varphi_h = \text{Inn}_{c_{g,h}} \circ \varphi_{g,h} \quad \varphi_g(c_{h,k}) \cdot c_{g,hk} = c_{g,h} \cdot c_{gh,k} \quad \text{and} \quad c_{1,1} = 1. \quad (1)$$

Here Inn_x with x an invertible element of A denotes the algebra automorphism $a \mapsto xax^{-1}$. A weak action of a group G is called *strict* if $c_{g,h} = 1$ for all pairs $g, h \in G$.

We first demonstrate how a weak G -action on an algebra A induces a categorical action (see [MNS11] for the definition) on the representation category $A\text{-mod}$. Here by $A\text{-mod}$ we denote the category of right modules over A ; using left modules would lead to slightly more complicated formulas in the rest of the paper. We define for each element $g \in G$ a functor ϕ_g on objects by

$${}^g(M, \rho) := (M, \rho \circ (\text{id}_M \otimes \varphi_{g^{-1}}))$$

and on morphisms by the identity, ${}^g f = f$, and take, for the functorial isomorphisms, $\alpha_{g,h}(M, \rho) := \rho(\text{id}_M \otimes c_{h^{-1},g^{-1}})$. One can check, that the cocycle condition in (1) implies the equality

$$\alpha_{gh,k} \circ \alpha_{g,h} = \alpha_{g,hk} \circ {}^g \alpha_{h,k}.$$

We summarize this in the following lemma.

Lemma 2.2. *Given a weak action of G on a \mathbb{K} -algebra A , the functors ϕ_g and the natural transformations $\alpha_{g,h}$ define a categorical action on the abelian category $A\text{-mod}$ of right A -modules.*

In the following we will mostly be interested in Hopf algebras. We therefore adapt the definition of a G -action to Hopf algebras.

Definition 2.3. *A weak G -action on a Hopf algebra A is a weak G -action $((\varphi_g)_{g \in G}, (c_{g,h})_{g,h \in G})$ on the underlying algebra which in addition satisfies the following properties:*

- G acts by automorphisms of Hopf algebras.
- The elements $(c_{g,h})_{g,h \in G}$ are group-like, i.e. $\Delta(c_{g,h}) = c_{g,h} \otimes c_{g,h}$.

Remark 2.4. *Analogously, one can give the definition of a weak G -action on a weak Hopf algebra, where one should keep in mind that a group-like element in a weak Hopf algebra is an element c such that $\Delta(c) = (c \otimes c)\Delta(1)$.*

Lemma 2.5. *Given a weak action of G on a Hopf algebra A , the induced action on the tensor category $A\text{-mod}$ of right A -modules is by strict tensor functors and tensor transformations.*

We next turn to an algebraic structure that yields tensor categories with G -action and compatible G -grading, called G -equivariant tensor categories [Kir04].

Definition 2.6. *A G -Hopf algebra over \mathbb{K} is a Hopf algebra A with a weak G -action $((\varphi_g)_{g \in G}, (c_{g,h})_{g,h \in G})$ and a G -grading $A = \bigoplus_{g \in G} A_g$ such that:*

- The algebra structure of A restricts to the structure of an associative algebra on each homogeneous component so that A is the direct sum of the components A_g as an algebra.
- The action of G is compatible with the grading, i.e. $\varphi_g(A_h) \subset A_{ghg^{-1}}$.

- The coproduct $\Delta : A \rightarrow A \otimes A$ respects the grading, i.e.

$$\Delta(A_g) \subset \bigoplus_{p,q \in G, pq=g} A_p \otimes A_q .$$

Remark 2.7. 1. For the counit ϵ and the antipode S of a G -Hopf algebra, the compatibility relations with the grading $\epsilon(A_g) = 0$ for $g \neq 1$ and $S(A_g) \subset A_{g^{-1}}$ are immediate consequences of the definitions.

2. The restrictions of the structure maps endow the homogeneous component A_1 of A with the structure of a Hopf algebra with a weak G -action.
3. G -Hopf algebras with strict G -action have been considered under the name “ G -crossed Hopf coalgebra” in [Tur10, Chapter VII.1.2].
4. Hopf algebras with weak G -action give a special case of G -Hopf algebra, where the grading is concentrated in degree 1. Thus all results of this paper imply analogous results where the term G -Hopf algebra is replaced by Hopf algebra with weak G -action.

The category $A\text{-mod}$ of finite-dimensional modules over a G -Hopf algebra inherits a natural (left and right) duality from the duality of the underlying category of \mathbb{K} -vector spaces. The weak action described in Lemma 2.2 is even a monoidal action, since G is required to act by Hopf algebra morphisms. A grading on $A\text{-mod}$ can be given by taking $(A\text{-mod})_g = A_g\text{-mod}$ as the g -homogeneous component. From the properties of a G -Hopf algebra one can finally deduce that the tensor product, duality and grading are compatible with the G -action. We have thus arrived at the following statement:

Lemma 2.8. [MNS11, Lemma 4.15] *The category of representations of a G -Hopf algebra inherits the natural structure of a \mathbb{K} -linear, abelian G -equivariant tensor category with dualities.*

A similar result holds for G -weak Hopf algebras.

3 Strictification of the group action

The action of the group G on a G -equivariant tensor category \mathcal{C} can always be strictified (see [Tur10, Appendix 5]), i.e. there is an equivalent G -equivariant tensor category \mathcal{C}^{str} with strict G -action (all compositors are identities). If one starts with the representation category of a G -Hopf algebra A , it is natural to ask whether this strictification leads to the representation category of another G -Hopf algebra with strict G -action. We will make this precise in the next definition. A G -equivariant functor between G -equivariant tensor categories is a tensor functor F together with natural isomorphisms

$$\psi_g : F({}^g M) \xrightarrow{\sim} {}^g F(M)$$

such that for every pair $g, h \in G$ the obvious coherence diagrams of morphisms from $F({}^{gh} M)$ to ${}^{gh} F(M)$ commute. See also [Tur10, Appendix 5, Def. 2.5].

Definition 3.1. 1. Let A be a Hopf algebra with weak G -action. A strictification of A is a weak Hopf algebra B with strict G -action and an equivalence

$$A\text{-mod} \xrightarrow{\sim} B\text{-mod}$$

of tensor categories with G -action.

2. Let A be a G -Hopf algebra. A strictification of A is a G -weak Hopf algebra B with strict G -action and an equivalence

$$A\text{-mod} \xrightarrow{\sim} B\text{-mod}$$

of G -equivariant tensor categories.

We will now show that it is in general not possible to find a strictification that is a Hopf algebra, rather than a weak Hopf algebra. This shows that we really have to allow for weak Hopf algebras as strictifications. In the next chapter we then show that a strictification as a weak Hopf algebra always exists.

Consider the weak action of $\mathbb{Z}/2 = \{1, -1\}$ on the group algebra $\mathbb{C}[\mathbb{Z}/2]$ given by

$$\varphi_1 = \varphi_{-1} = \text{id}$$

and non-trivial compositors given by the grouplike elements

$$c_{1,1} = c_{1,-1} = c_{-1,1} = 1 \quad c_{-1,-1} = -1 \in \mathbb{C}[\mathbb{Z}/2].$$

In [MNS11, Section 3.1] we showed how weak actions correspond to extensions of groups. In this case, the relevant extension is given by the exact sequence of abelian groups

$$\mathbb{Z}/2 \rightarrow \mathbb{Z}/4 \rightarrow \mathbb{Z}/2.$$

Theorem 3.2. *There is no strictification as a Hopf algebra of $\mathbb{C}[\mathbb{Z}/2]$ with the $\mathbb{Z}/2$ -action given above.*

Remark 3.3. *Note that the algebra $\mathbb{C}[\mathbb{Z}/2]$ is not a priori endowed with a grading by $\mathbb{Z}/2$. We can consider it as being trivially graded. Alternatively we can consider the non-trivially graded Hopf algebra $H := \mathcal{D}^{\mathbb{Z}/2}(\mathbb{Z}/2)$ introduced in [MNS11, section 4.3]. The same proof as below goes through and shows that H can not be strictified as a Hopf algebra either.*

For the proof of proposition 3.2 we need the following elementary facts:

Lemma 3.4. *Let $A = \mathbb{C}[G]$ be the complex group algebra of a finite abelian group G .*

1. *Let A' be an arbitrary Hopf algebra. If $A\text{-mod} \cong A'\text{-mod}$ as tensor categories, then $A \cong A'$ as algebras (not necessarily as Hopf algebras).*
2. *The natural endomorphisms of the identity functor $\text{Id} : A\text{-mod} \rightarrow A\text{-mod}$ are given by the action of elements in A . More precisely there is an isomorphism of algebras*

$$A \xrightarrow{\sim} \text{End}(\text{Id})_{A\text{-mod}} .$$

3. Let $\varphi : A \rightarrow A$ be an algebra automorphism such that the restriction functor $res_\varphi : A\text{-mod} \rightarrow A\text{-mod}$ is naturally isomorphic to the identity functor. Then $\varphi = \text{id}$.

Proof. 1.) By the reconstruction theorem we know that we can recover the Hopf algebra A as endomorphisms of the fibre functor $F : A\text{-mod} \rightarrow \mathbb{C}\text{-mod}$ and A' as endomorphisms of the fibre functor $G : A\text{-mod} \xrightarrow{\sim} A'\text{-mod} \rightarrow \mathbb{C}\text{-mod}$. Now we claim that the underlying functors of F and G are naturally isomorphic. To this end note that for each simple representation V_i of A we have $V_i^n \cong 1$ where n is the order of the group. Thus we have $F(V_i) \cong \mathbb{C} \cong G(V_i)$ by the fact that F and G are tensor-functors. But it is easy to see that the \mathbb{C} -linearity and the fact that $A\text{-mod}$ is semisimple then already show that F and G are isomorphic as functors between abelian categories. This implies that $A \cong \text{End}(F) \cong \text{End}(G) \cong A'$. Note that the functors F and G still might have different tensor functor structures, leading to different Hopf algebra structures on A and A' .

2.) This follows from the fact that A is abelian and from the fact that the center of an algebra is isomorphic to the endomorphisms of the identity functor on its representation category.

3) The functor res_φ is an equivalence of categories. Hence it sends simple objects to simple objects. That means it acts on characters $\chi : G \rightarrow \mathbb{C}^*$. By the fact that this functor is naturally isomorphic to the identity this action has to be trivial. Hence we know $\chi \circ \varphi = \chi$ for each character χ . Because G is abelian, the characters form a basis of the dual space A^* . Thus $\varphi^* = \text{id}$ which implies $\varphi = \text{id}$. □

Proof of proposition 3.2. Assume that there is a Hopf algebra H with a strict action of $\mathbb{Z}/2$ by Hopf algebra automorphisms ψ_g together with an equivalence of categories $A\text{-mod} \rightarrow H\text{-mod}$. By lemma 3.4(1) we know that the underlying algebra of H is isomorphic to $\mathbb{C}[\mathbb{Z}/2]$. We choose an isomorphism and transport the action ψ_g on H to an action ψ'_g on $\mathbb{C}[\mathbb{Z}/2]$ (which is now only an action by algebra automorphisms and not necessarily by Hopf algebra automorphisms). By assumption there are now natural isomorphisms $res_{\psi'_g} \xrightarrow{\sim} res_{\psi_g} = \text{Id}$ hence by lemma 3.4(3) we have $\psi'_g = \text{id}$.

Now we have both times the trivial action on the algebra $\mathbb{C}[\mathbb{Z}/2]$, once with a nontrivial compositor $c_{-1,-1}$ and once with the trivial compositors. By Lemma 3.4(2), an isomorphism between the two induced actions on the representation categories is induced by elements $(a_g \in \mathbb{C}[\mathbb{Z}/2])_{g \in \mathbb{Z}/2}$ such that $a_{gh} \cdot c_{g,h} = a_g \cdot a_h$. It is easy to see that such elements cannot exist since the sequence of algebras

$$\mathbb{C}[\mathbb{Z}/2] \rightarrow \mathbb{C}[\mathbb{Z}/4] \rightarrow \mathbb{C}[\mathbb{Z}/2]$$

is not split. This contradicts the existence of the strictification A' of a Hopf algebra. □

4 Existence of a strictification

In this section, we will successively prove the following theorem which holds for Hopf algebras over an arbitrary field \mathbb{K} .

Theorem 4.1. 1. For any Hopf algebra with weak G -action there exists a strictification in the sense of definition 3.1(1).

2. For any G -Hopf algebra there exists a strictification in the sense of definition 3.1(2).

Note that the first part of Theorem 4.1 follows from the second part if we consider a Hopf algebra with weak G -action as a G -Hopf algebra with grading concentrated in degree 1, see also Remark 2.7(4). Therefore we will only prove the second part.

In the following let A be a G -Hopf algebra with unit 1_A , counit ϵ_A , coproduct Δ_A and a weak G -action $((\varphi_g)_{g \in G}, (c_{g,h})_{g,h \in G})$. The plan of this section is to construct step by step a strictification A^{str} .

In section 4.1 we construct the algebra A^{str} , in section 4.2 we endow it with a weak Hopf algebra structure and finally in section 4.3 we turn it into a G -weak Hopf algebra with strict G -action. Along the way, we also provide the necessary equivalence of representation categories

$$F : A\text{-mod} \xrightarrow{\sim} A^{str}\text{-mod}$$

and show that it preserves all the structure involved. This implies that A^{str} is a strictification, which proves Theorem 4.1.

4.1 The algebra

In the following we use the notation $\mathbb{K}(G)$ for the \mathbb{K} -vector space of functions on the finite group G , with basis $(\delta_g)_{g \in G}$. By $\mathbb{K}[G]$ we denote the \mathbb{K} -vector space underlying the group algebra with basis $(g)_{g \in G}$.

Definition 4.2. Set $A^{str} = \mathbb{K}(G) \otimes_{\mathbb{K}} A \otimes_{\mathbb{K}} \mathbb{K}[G]$ as a vector space and define a multiplication on A^{str} on the generators of A^{str} by

$$(\delta_g \otimes a \otimes h)(\delta_{g'} \otimes a' \otimes h') = \delta(gh, g')(\delta_g \otimes a\varphi_h(a')c_{h,h'} \otimes hh') \quad (2)$$

where $\delta(gh, g')$ is the Kronecker delta, i.e. $\delta(gh, g') = 1$ if $gh = g'$ and $\delta(gh, g') = 0$ otherwise. Furthermore we define a unit by

$$1 = \sum_{g \in G} \delta_g \otimes 1_A \otimes 1. \quad (3)$$

It can easily be checked that the product and the unit defined in (2) and (3) endow A^{str} with the structure of an associative unital algebra.

We next define a functor $F : A\text{-mod} \rightarrow A^{str}\text{-mod}$: Let M be an object in $A\text{-mod}$. Define an object in $A^{str}\text{-mod}$ which is $M \otimes_{\mathbb{K}} \mathbb{K}[G]$ as a vector space and has the following right action of the algebra A^{str} : For an element of the form $(m \otimes k)$ with $m \in M, k \in G$, it reads:

$$(m \otimes k) \cdot (\delta_g \otimes a \otimes h) := \delta(k, g)(m \cdot \varphi_k(a)c_{k,h} \otimes kh) \quad (4)$$

One checks that this really defines a right action of A . For a morphism $f \in \text{Hom}_A(M, N)$ we consider the morphism $f \otimes \text{id}_{\mathbb{K}[G]} \in \text{Hom}_{A^{str}}(M \otimes \mathbb{K}[G], N \otimes \mathbb{K}[G])$. Together this defines a functor:

$$F : A\text{-mod} \rightarrow A^{str}\text{-mod} \quad (5)$$

Proposition 4.3. The functor F is an equivalence of abelian categories.

Proof. We show that F is essentially surjective and fully faithful.

For the essential surjectivity, we take an object N in A^{str} -mod and first show that

$$N \cong N_1 \otimes \mathbb{K}[G]$$

as A^{str} -modules, where $N_1 := N.(\delta_1 \otimes 1_A \otimes 1)$ and we take the action of A^{str} on an element $(n \otimes k) \in N_1 \otimes \mathbb{K}[G]$ to be $(n \otimes k).(\delta_g \otimes a \otimes h) = (n.(\delta_g \otimes a \otimes 1) \otimes kh)$. Indeed, we have

$$N = N.1_{A^{str}} = \bigoplus_{g \in G} N.(\delta_g \otimes 1_A \otimes 1) = \bigoplus_{g \in G} N_1.(\delta_1 \otimes 1_A \otimes g)$$

and there is an obvious isomorphism to $N_1 \otimes \mathbb{K}[G]$ that commutes with the action of A^{str} . It is clear that F is faithful. In order to see that F is also full, consider a morphism $f \in \text{Hom}_{A^{str}}(M \otimes \mathbb{K}[G], N \otimes \mathbb{K}[G])$. We have

$$f(m \otimes k).(\delta_g \otimes 1_A \otimes 1) = f((m \otimes k).(\delta_g \otimes 1_A \otimes 1)) = \delta(g, k)f(m \otimes k),$$

so $f(m \otimes k) \in N \otimes \mathbb{K}k$. Since f is linear, we deduce $f = g \otimes \text{id}_{\mathbb{K}[G]}$, where $g \in \text{Hom}_A(M, N)$. \square

4.2 The weak Hopf algebra structure

We need the strictification algebra A^{str} to have more structure in order for its representation category to be a tensor category. In fact, we want it to be a weak bialgebra. For the definition and properties of weak bialgebras, see e.g. [NTV03].

Proposition 4.4. *The linear maps $\Delta : A^{str} \rightarrow A^{str} \otimes A^{str}$ and $\epsilon : A^{str} \rightarrow \mathbb{K}$ defined on the generators of A^{str} by*

$$\begin{aligned} \Delta(\delta_g \otimes a \otimes h) &= \sum_{(a)} (\delta_g \otimes a_{(1)} \otimes h) \otimes (\delta_g \otimes a_{(2)} \otimes h) , \\ \epsilon(\delta_g \otimes a \otimes h) &= \epsilon_A(a) \end{aligned}$$

endow A^{str} with the structure of a weak bialgebra. Furthermore, the linear map $S : A \rightarrow A$ given by

$$S(\delta_g \otimes a \otimes h) = (\delta_{h^{-1}g} \otimes c_{h^{-1},h} \cdot \varphi_{h^{-1}}(S_A(a)) \otimes h^{-1})$$

is an antipode for A^{str} , where S_A is the antipode of A .

Proof. The maps Δ and ϵ are a coassociative coproduct and a counit on A^{str} , as they are just the structural maps of the tensor product coalgebra of $\mathbb{K}(G)$, A and $\mathbb{K}[G]$ (where we consider the diagonal coproduct on both $\mathbb{K}(G)$ and $\mathbb{K}[G]$). We show that Δ is also a morphism of algebras, i.e. that

$$(m \otimes m) \circ (\text{id} \otimes \tau \otimes \text{id})(\Delta \otimes \Delta) = \Delta \circ m. \tag{6}$$

If we plug in two elements $(\delta_g \otimes a \otimes h), (\delta_{g'} \otimes a' \otimes h')$, we get for the left hand side of (6)

$$\begin{aligned} & \sum_{(a)} (\delta_g \otimes a_{(1)} \otimes h) \cdot (\delta_{g'} \otimes a'_{(1)} \otimes h') \otimes (\delta_g \otimes a_{(2)} \otimes h) \cdot (\delta_{g'} \otimes a'_{(2)} \otimes h') \\ &= \sum_{(a)} \delta(h, hg') (\delta_g \otimes a_{(1)} \varphi_h(a'_{(1)}) c_{h,h'} \otimes hh') \otimes (\delta_g \otimes a_{(2)} \varphi_h(a'_{(2)}) c_{h,h'} \otimes hh') \ , \end{aligned}$$

and for the right hand side

$$\begin{aligned} & \delta(g, hg') \Delta(\delta_g \otimes a \varphi_h(a') c_{h,h'} \otimes hh') \\ &= \delta(g, hg') \sum_{(a)} (\delta_g \otimes (a \varphi_h(a') c_{h,h'})_{(1)} \otimes hh') \otimes (\delta_g \otimes (a \varphi_h(a') c_{h,h'})_{(2)} \otimes hh') \\ &= \sum_{(a)} \delta(h, hg') (\delta_g \otimes a_{(1)} \varphi_h(a'_{(1)}) c_{h,h'} \otimes hh') \otimes (\delta_g \otimes a_{(2)} \varphi_h(a'_{(2)}) c_{h,h'} \otimes hh') \ . \end{aligned}$$

All equations follow just by definition, except for the last one, where we used that the co-product in A is a morphism of algebras, that the elements $c_{g,h}$ are group-like and that the action of G on A is a coalgebra-morphism.

Further equations concerning the compatibilities of the product with the counit, the coproduct with the unit and the antipode can be checked directly. \square

The representation category of a weak Hopf algebra H has the structure of a tensor category, where the tensor product of two modules M, N is given by the following truncation of the tensor product of the underlying vector spaces,

$$M \bar{\otimes} N := (M \otimes_{\mathbb{K}} N) \Delta(1) \ .$$

The tensor unit is the counital subalgebra

$$H_t := \epsilon_t(H)$$

with H -action given by $z.h := \epsilon_t(zh)$ for $h \in H, z \in H_t$.

Lemma 4.5. *For the algebra A^{str} , the target and source counital maps are given by*

$$\epsilon_t(\delta_g \otimes a \otimes h) = \epsilon_s(\delta_g \otimes a \otimes h) = \epsilon_A(a)(\delta_g \otimes 1_A \otimes 1) \quad (7)$$

and the target and source counital subalgebras are

$$A_t^{str} \cong A_s^{str} \cong \mathbb{K}(G) \ .$$

Proof. We calculate ϵ_t on an element $(\delta_g \otimes a \otimes h) \in A^{str}$:

$$\begin{aligned} \epsilon_t(\delta_g \otimes a \otimes h) &= (\epsilon \otimes \text{id})(\Delta(1)((\delta_g \otimes a \otimes h) \otimes 1)) \\ &= (\epsilon \otimes \text{id})((\delta_g \otimes a \otimes h) \otimes (\delta_g \otimes 1_A \otimes 1)) \\ &= \epsilon_A(a)(\delta_g \otimes 1_A \otimes 1) \ . \end{aligned}$$

Choose a basis $(a_i)_{i \in I}$ of the algebra A with $a_j = 1_A$ for a fixed $j \in I$, then a general element $b \in A^{str}$ is of the form

$$b = \sum_{g,h \in G, i \in I} \lambda(g, h, i) (\delta_g \otimes a_i \otimes h)$$

with $\lambda(g, h, i) \in \mathbb{K}$. We have:

$$\begin{aligned}\Delta(b) &= \sum_{g, h \in G, i \in I} \sum_{(a_i)} \lambda(g, h, i) (\delta_g \otimes (a_i)_{(1)} \otimes h) \otimes (\delta_g \otimes (a_i)_{(2)} \otimes h) \\ \Delta(1)(b \otimes 1) &= \sum_{g, h \in G, i \in I} \lambda(g, h, i) (\delta_g \otimes a_i \otimes h) \otimes (\delta_g \otimes 1_A \otimes h)\end{aligned}$$

By equating coefficients, we get $\lambda(g, h, i) = 0$ for $h \neq 1, i \neq j$ and therefore:

$$A_t^{str} = \langle \delta_g \otimes 1_A \otimes 1, g \in G \rangle \cong \mathbb{K}(G) .$$

□

Proposition 4.6. *The equivalence $F : A\text{-mod} \xrightarrow{\sim} A^{str}\text{-mod}$ can be promoted to an equivalence of tensor categories.*

Proof. The tensor unit in the representation category of the weak Hopf algebra A^{str} is given by the target counital subalgebra, which is by lemma 4.5 isomorphic to $\mathbb{K}(G)$. The algebra acts on the target counital subalgebra by the target counital map ϵ_t after multiplication (see [NTV03]), i.e. for an element $(\delta_k \otimes 1_A \otimes 1) \in A_t^{str}$ and $(\delta_g \otimes a \otimes h) \in A^{str}$, we have

$$(\delta_k \otimes 1_A \otimes 1) \cdot (\delta_g \otimes a \otimes h) = \epsilon_t((\delta_k \otimes 1_A \otimes 1)(\delta_g \otimes a \otimes h)) = \delta(k, g) \epsilon_A(a) (\delta_k \otimes 1_A \otimes 1) .$$

The action of an element $a \in A$ on the tensor unit \mathbb{K} in $A\text{-mod}$ is by multiplication with $\epsilon_A(a)$. So we get for the image of the tensor unit under F the vector space $\mathbb{K} \otimes \mathbb{K}[G]$ with A^{str} -action

$$(\lambda \otimes k)(\delta_g \otimes a \otimes h) = \delta(k, g) \epsilon_A(a) \lambda \otimes kh .$$

We clearly have an isomorphism $F(\mathbf{1}) \rightarrow \mathbf{1}$ in $A^{str}\text{-mod}$ given by

$$\eta_0 : (\lambda \otimes k) \mapsto \lambda \delta_k .$$

Let $M, N \in A\text{-mod}$. We have

$$F(M) \bar{\otimes} F(N) = \langle (m \otimes g \otimes n \otimes g), m \in M, n \in N, g \in G \rangle .$$

Thus the linear map

$$\eta_2(M, N) : (m \otimes g \otimes n \otimes g) \mapsto (m \otimes n \otimes g)$$

is an isomorphism $F(M) \bar{\otimes} F(N) \xrightarrow{\sim} F(M \otimes N)$. It can be seen to commute with the action of A^{str} and is natural in M, N . Moreover the isomorphisms η_2 clearly satisfy the coherence axioms for three objects. We have therefore established that (F, η_0, η_2) is a tensor functor. □

4.3 G -action and G -grading

We will now define a G -equivariant structure on A^{str} that induces a G -equivariant structure on the category $A^{str}\text{-mod}$. The last step of proving theorem 4.1 is then to show that the categories $A\text{-mod}$ and $A^{str}\text{-mod}$ are even equivalent as G -equivariant categories.

Definition 4.7. On the weak Hopf algebra A^{str} we have a strict left action φ^{str} of the group G given by translation in the first factor. Explicitly, an element $g' \in G$ acts on an element $(\delta_g \otimes a \otimes h) \in A^{str}$ by

$$\varphi_{g'}^{str}(\delta_g \otimes a \otimes h) = (\delta_{g'g} \otimes a \otimes h) .$$

The strict G -action on A^{str} gives us a strict left G -action on the category $A^{str}\text{-mod}$ by setting $\phi_g(M, \rho) = (M, \rho \circ (\text{id}_M \otimes \varphi_{g^{-1}}^{str}))$. We will now establish, that the equivalence $A\text{-mod} \cong A^{str}$ is compatible with the G -actions.

Proposition 4.8. The equivalence $F : A\text{-mod} \xrightarrow{\sim} A^{str}\text{-mod}$ given in (5) respects the G -action of the two categories, i.e. for every element $g \in G$ there are natural isomorphisms

$$\psi_g : F({}^g M) \xrightarrow{\sim} {}^g F(M)$$

such that for every pair $g, h \in G$ the obvious coherence diagrams of morphisms from $F({}^{gh} M)$ to ${}^{gh} F(M)$ commute.

Proof. For $M \in A\text{-mod}$ consider the linear map $\psi_g : M \otimes \mathbb{K}[G] \rightarrow M \otimes \mathbb{K}[G]$ defined by

$$\psi_g : (m \otimes k) \mapsto (m.c_{g^{-1},k} \otimes g^{-1}k).$$

We first show that ψ_g is a morphism of A^{str} -modules. To distinguish the actions on the different modules we use the notation “ \star ” for the action on $F({}^g M)$ and ${}^g F(M)$, “ \odot ” for the action on $F(M)$ and “.” for the action on M .

$$\begin{aligned} & \psi_g((m \otimes k) \star (\delta_x \otimes a \otimes h)) \\ &= \psi_g(\delta(k, x)(m.c_{g^{-1}}(\varphi_k(a)c_{k,h}) \otimes kh)) \\ &= \delta(k, x) m.c_{g^{-1}}(\varphi_k(a)c_{k,h})c_{g^{-1},kh} \otimes g^{-1}kh \\ &= \delta(k, x) m.c_{g^{-1},k}\varphi_{g^{-1}k}(a)(c_{g^{-1},k})^{-1}\varphi_{g^{-1}}(c_{k,h})c_{g^{-1},kh} \otimes g^{-1}kh \\ &= \delta(g^{-1}k, g^{-1}x) m.c_{g^{-1},k}\varphi_{g^{-1}k}(a)c_{g^{-1}k,h} \otimes g^{-1}kh \\ &= (m.c_{g^{-1},k} \otimes g^{-1}k) \odot (\delta_{g^{-1}x} \otimes a \otimes h) \\ &= \psi_g(m \otimes k) \star (\delta_x \otimes a \otimes h) . \end{aligned}$$

Moreover we have to verify that the ψ_g satisfy a coherence condition for two indices g and h . This condition can be checked similarly to the above computation, using the cocycle condition for the $c_{g,h}$. \square

Definition 4.9. We define a G -grading in the sense of 2.6 on the algebra A^{str} by:

$$(A^{str})_h = \bigoplus_{g \in G} (\mathbb{K}(\delta_g) \otimes A_{g^{-1}hg}) \otimes \mathbb{K}[G] . \quad (8)$$

Lemma 4.10. The algebra A^{str} is a G -weak Hopf algebra with strict G -action, i.e a weak Hopf algebra with strict G -action and compatible G -grading.

Note that the grading on A resp. A^{str} gives a grading on the representation category by $(A\text{-mod})_h := A_h\text{-mod}$ resp. $(A^{str}\text{-mod})_h = (A^{str})_h\text{-mod}$.

Proposition 4.11. The equivalence $F : A\text{-mod} \xrightarrow{\sim} A^{str}\text{-mod}$ given in (5) respects the G -grading of the two categories, i.e. for every element $h \in G$, we have

$$F((A\text{-mod})_h) \subset (A^{str}\text{-mod})_h.$$

Proof. Let $M \in A_h\text{-mod}$. We know that the action by the unit $(\mathbf{1}_A)_h$ of A_h is the identity on M . We need to show, that the h -component of the unit in A^{str} , which is $\mathbf{1}_h = \sum_{g \in G} (\delta_g \otimes (\mathbf{1}_A)_{g^{-1}hg} \otimes 1)$, acts as a projector on $F(M) = M \otimes \mathbb{K}[G]$. For any element of the form $(m \otimes k)$, we have:

$$\begin{aligned} (m \otimes k) \cdot \left(\sum_{g \in G} \delta_g \otimes (\mathbf{1}_A)_{g^{-1}hg} \otimes 1 \right) &= \sum_{g \in G} (\delta(k, g) m \cdot \varphi_k((\mathbf{1}_A)_{g^{-1}hg}) c_{k,1} \otimes k) \\ &= (m \cdot \varphi_k((\mathbf{1}_A)_{k^{-1}hk}) \otimes k) \\ &= (m \cdot (\mathbf{1}_A)_h \otimes k) \\ &= (m \otimes k) \end{aligned}$$

where in the first equality we used the definition of the action of A^{str} on $M \otimes \mathbb{K}[G]$ given in (4), in the third equality the fact that G acts by unital algebra morphisms and in the last equality that M is in the h -component of $A\text{-mod}$.

So we have $F(M) \cdot \mathbf{1}_h = F(M)$; therefore $F(M)$ lies in the component $(A^{str}\text{-mod})_h$. \square

5 Equivariant R-Matrix and ribbon-element

In [MNS11] we considered G -equivariant categories with a G -braiding and a G -twist as additional data (G -ribbon categories). For the definition see [Tur10, Kir04]. Since those categories were our main motivation to study the strictification in terms of algebras, we want to say a few words about the G -ribbon structure.

The definition of a G -equivariant R -matrix is rather involved even in the strict Hopf algebra case. We will refrain here from stating the axioms for it explicitly, but we will instead make an equivalent definition:

Definition 5.1. *Let A be a G -(weak) Hopf algebra.*

1. *A G -equivariant R -matrix is an element $R = R_1 \otimes R_2 \in \Delta^{op}(1)(A \otimes A)\Delta(1)$ such that for $V \in (A\text{-mod})_g$, $W \in A\text{-mod}$, the map*

$$\begin{aligned} c_{VW} : V \otimes W &\rightarrow {}^gW \otimes V \\ v \otimes w &\mapsto w \cdot R_2 \otimes v \cdot R_1 \end{aligned}$$

is a G -braiding, in particular a morphism of A -modules.

2. *A G -twist is an invertible element $\theta \in A$ such that for every object $V \in (A\text{-mod})_g$ the induced map*

$$\begin{aligned} \theta_V : V &\rightarrow {}^gV \\ v &\mapsto v \cdot \theta^{-1} \end{aligned}$$

is a G -twist in $A\text{-mod}$.

If A has a G -equivariant R -matrix and a G -twist, we call it G -(weak) ribbon-algebra.

Lemma 5.2. *A G -weak Hopf algebra A is a G -weak ribbon algebra if and only if $A\text{-mod}$ is a G -ribbon category.*

Proof. If A is a G -ribbon algebra, it follows from the definition that $A\text{-mod}$ is a G -ribbon category. If on the other hand, $A\text{-mod}$ is a G -ribbon category with G -braiding R and G -twist θ , define an R -matrix and a twist of A by

$$R = \tau \circ c_{A,A}(1_A \otimes 1_A) \quad \text{and} \quad \theta = \theta_A(1)^{-1} . \quad (9)$$

For $v \in V$, $w \in W$ let $\bar{v} : A \rightarrow V$, $\bar{w} : A \rightarrow W$ be the A -linear maps with $\bar{v}(1_A) = v$, $\bar{w}(1_A) = w$. We then have

$$\begin{aligned} \tau((v \otimes w).R) &= \tau(\bar{v} \otimes \bar{w}(R)) = (\bar{w} \otimes \bar{v})c_{A,A}(1_A \otimes 1_A) = c_{V,W}(v \otimes w) , \\ v.\theta^{-1} &= v.(\theta_A(1_A)) = \bar{v}(\theta_A(1_A)) = \theta_V \bar{v}(1_A) = \theta_V(v). \end{aligned}$$

Thus R and θ satisfy the conditions of definition 5.1 by construction. \square

As an immediate consequence of lemma 5.2, we have:

Corollary 5.3. *If A is a G -ribbon algebra, the strictification algebra A^{str} inherits the structure of a G -weak ribbon algebra such that the equivalence $F : A\text{-mod} \rightarrow A^{\text{str}}\text{-mod}$ is an equivalence of G -ribbon categories.*

A Table summarizing terminology

The following table summarizes the terminology for Hopf algebras with an action of a finite group G and their weakenings. We consider two types of weakenings: a weakening of the G -action corresponding to the two rows of the table, and a weakening of the unitality of the coproduct, corresponding to the two columns of the table.

Each square contains three different entries, depending on additional structure on the Hopf algebra. The objects in 1. only have the G -action and no additional structure (see Definition 2.3). The objects in 2. are equipped with a G -grading with the compatibilities introduced in Definition 2.6. The objects in 3. have, in addition to the G -equivariant structure, a G -equivariant R -matrix and a G -twist as introduced in Definition 5.1.

	Hopf algebra	weak Hopf algebra
strict G -action	1. Hopf algebra with strict G -action	1. weak Hopf algebra with strict G -action
	2. G -Hopf algebra with strict G -action	2. G -weak Hopf algebra with strict G -action
	3. G -ribbon algebra with strict G -action	3. G -weak ribbon algebra with strict G -action
weak G -action	1. Hopf algebra with weak G -action	1. weak Hopf algebra with weak G -action
	2. G -Hopf algebra	2. G -weak Hopf algebra
	3. G -ribbon algebra	3. G -weak ribbon algebra

References

- [BK01] B. Bakalov and A.A. Kirillov. *Lectures on tensor categories and modular functors*. American Mathematical Society, 2001.
- [BNS99] G. Böhm, F. Nill, and K. Szlachányi. Weak Hopf algebras. I. Integral theory and C^* -structure. *J. Algebra*, 221(2):385–438, 1999.
- [BS00] G. Böhm and K. Szlachányi. Weak Hopf algebras. II. Representation theory, dimensions, and the Markov trace. *J. Algebra*, 233(1):156–212, 2000.
- [DW90] R. Dijkgraaf and E. Witten. Topological gauge theories and group cohomology. *Comm. Math. Phys.*, 129(2):393–429, 1990.
- [FQ93] D.S. Freed and F. Quinn. Chern-Simons theory with finite gauge group. *Comm. Math. Phys.*, 156(3):435–472, 1993.
- [Kir04] A. Kirillov. On G -equivariant modular categories. *Arxiv preprint: math/0401119v1*, 2004.
- [MNS11] J. Maier, T. Nikolaus, and C. Schweigert. Equivariant modular categories via Dijkgraaf-Witten theory. *Arxiv preprint:1103.2963*, 2011.
- [NTV03] D. Nikshych, V. Turaev, and L. Vainerman. Invariants of knots and 3-manifolds from quantum groupoids. *Topology Appl.*, 127(1-2):91–123, 2003.
- [NV02] D. Nikshych and L. Vainerman. Finite quantum groupoids and their applications. In *New directions in Hopf algebras*, volume 43 of *Math. Sci. Res. Inst. Publ.*, pages 211–262. Cambridge Univ. Press, Cambridge, 2002.
- [Tur10] V. Turaev. *Homotopy Quantum Field Theory*. European Mathematical Society, 2010.
- [Vir02] A. Virelizier. Hopf group-coalgebras. *J. Pure Appl. Algebra*, 171(1):75–122, 2002.