

81-3-39
DEUTSCHE ATOMSIEBEN

DEUTSCHES ELEKTRONEN-SYNCHROTRON DESY

DESY 81-004
January 1981

SCHRÖDINGER REPRESENTATION AND CASIMIR EFFECT
IN RENORMALIZABLE QUANTUM FIELD THEORY

by

K. Symanzik

NOTKESTRASSE 85 · 2 HAMBURG 52

DESY behält sich alle Rechte für den Fall der Schutzrechtserteilung und für die wirtschaftliche Verwertung der in diesem Bericht enthaltenen Informationen vor.

DESY reserves all rights for commercial use of information included in this report, especially in case of apply for or grant of patents.

To be sure that your preprints are promptly included in the
HIGH ENERGY PHYSICS INDEX,
send them to the following address (if possible by air mail) :





DESY 81-004
January 1981

Erratum to "Schrödinger representation and Casimir effect in renormalizable quantum field theory" :

The conditions (E. 9), (E. 10), (E. 11b) have more solutions than those given in Appendix E. If $S = 0$, every A with $A^2 = 1$ (which excludes real A) and such that $A + Q$ is invertible for all momenta leads to a solution $X = -2(A + Q)^{-1}$. This excludes all rotationally noninvariant A , and excludes among the rotationally invariant ones

$$A = \alpha \gamma^0 + b \gamma_5 + c i \gamma^0 \gamma_5, \quad \alpha^2 + b^2 + c^2 = 1$$

if $m > 0$ those with $(1 - b^2)^{-1}(-\alpha \pm i b c) \in [1, \infty)$

and if $m = 0$ those with $b = \pm 1$. G becomes simple, however, only if $c = \pm 1$, $a = b = 0$, which yields the solution (E. 16).

If, instead of $S = 0$, (E. 14) is required, then necessarily $S = 1/2$, and

$X = \frac{1}{2}(1+A)$ with A the ones of the $S = 0$ solutions as described, which yields the same G as in those solutions.

The argument that, for the spin 1/2 case, the "Dirichlet" and the "Neumann" boundary conditions can both be upheld under Yukawa interaction and in QED will be given in the published version of the paper.

DESY 81-004
January 1981

Abstract

We show that the Schrödinger representation exists in renormalizable quantum field theory to all orders in the perturbation expansion. In this sense, completeness of the Schrödinger states also holds. However, the field operator that is being diagonalized on a smooth three-dimensional hypersurface differs from the usual renormalized one by a factor that diverges logarithmically if the distance from the hypersurface goes to zero. This requires a limit procedure to be employed if expectation values of the renormalized field operator are to be computed in this representation. The Schrödinger functional differential operator involves point splitting Δ and has coefficients depending logarithmically on Δ , and also some by factors Δ^{-1} , Δ^{-2} , Δ^{-4} . Details are given for the massless ϕ^4 theory, but the extension to other models, in particular with spin 1/2 fermions, is outlined. The Casimir potential for disjoint surfaces is shown to be finite to all orders in the perturbation expansion, and computed for a pair of parallel plates to first order in massless ϕ^4 .

Schrödinger representation and Casimir effect
in renormalizable quantum field theory

K. Symanzik

Deutsches Elektronen-Synchrotron DESY, Hamburg

Introduction

Soon after the invention of quantum field theory, also its Schrödinger representation was known, and it has been mentioned since in some text books [1]. Calculations, however, were done first in the interaction representation, which is formally related to the Schrödinger representation by a change of basis. Later, covariant four-dimensional formalisms (S-matrix calculus, Green's functions, and in particular functional integrals) were used almost exclusively. Even more than the interaction representation, which preserves a certain conceptual role in scattering theory, the Schrödinger representation fell into disrepute, the more so since it seems to have been considered to be nonrenormalizable, as the interaction representation indeed is [2,3].

More recently, however, the search for nonperturbative methods in strong-interaction theory led to the discussion of highly non-pointlike objects (dual strings, Wilson loops). The dynamics of such objects was formulated in terms of QFTs with boundaries [4,5], an attractive setting due to the high flexibility and perfection of the QFT formalism. Prescribing the value of the quantum field on a boundary, however, means using the Schrödinger representation, slightly extended from flat to curved boundaries. This led us to consider renormalizable QFTs with boundaries, since nonrenormalizable theories pose unsolved problems already in infinite space-time. Superrenormalizable theories, on the other hand, offer no particular difficulty in this respect and, above all, do not appear^{*} to describe interesting physics here.

* Recently, however, Migdal [5] has proposed a free-fermion theory with boundaries for model use in QCD.

We show here that in renormalizable models, the Schrödinger wave functional exists to all orders in perturbation theory, and give, what we believe to be strong arguments, that so also does the Schrödinger functional differential operator that appears in the Schrödinger equation. For simplicity, we treat in detail only the ϕ_4^4 theory, and mostly choose it massless since masses are inessential here, but describe the principles of the extension to other models. Dimensional regularization is used and mostly the Euclidean frame, since the transition to the Minkowski one is obvious. Renormalization group equations are given at every stage.

In sect. 1, we discuss how boundary conditions (in particular, homogeneous and inhomogeneous Dirichlet ones) are implemented by surface interaction theory. That the latter can be introduced in the renormalized infinite-space theory and hereby only cause divergences absorbable in logarithmic factors is crucial for the renormalizability, as we show in sect. 2. An immediate consequence is the finiteness of the Casimir effect (for a pair of parallel plates, for instance) to all orders, and we compute it to first order in massless ϕ_4^4 theory in sect. 3. In sect. 4 we show that if the renormalized field operator, or its derivative in the normal direction, approaches the boundary plane, a factor, different in the two cases, with logarithmic dependence on the distance from the boundary must be applied to keep matrix elements finite. In sect. 5 we construct the Schrödinger functional differential operator, which requires point splitting as it does already in the free field theory. Completeness and unitarity are discussed in sect. 6. That the field operator that is being diagonalized is not the renormalized one forces us to use a limit procedure if expectations of the field operator are to be computed. The extension of our method to other models we discuss in sect. 7, and give some

details for the spin 1/2 Majorana field in an Appendix. We also point out the divergences that arise if the transition to the interaction representation is attempted. We note that, unfortunately, the present methods are not applicable to the string Lagrangeans [4] as long as these are not shown to be renormalizable in infinite space. Conclusions are stated in sect. 8. Some technical material is relegated to Appendices.

1. Boundary conditions by surface interactions

Consider the theory of a free one-component scalar field with (Euclidean) action density in ν dimensions

$$(1.1) \quad \mathcal{L}_0 = -\frac{1}{2} \partial_\mu \phi \partial^\mu \phi - \frac{1}{2} m^2 \phi^2$$

Let \mathcal{T} be a simply connected (not necessarily bounded) region of space with $\nu-1$ dimensional smooth boundary $\partial\mathcal{T}$, described by $f(x) = 0$, $x \in R^\nu$, with $f(x) > 0$ in \mathcal{T} and $f(x) < 0$ in the complementary region \mathcal{T}' . Consider now the augmented action density

$$(1.2) \quad \mathcal{L}_n = \mathcal{L}_0 - \delta(f(x)) \phi(x) \partial_\mu \phi(x) \partial^\mu \phi(x)$$

where $\sim \partial f$ is an infinitesimal vector pointing from $x \in \partial\mathcal{T}$ outwards.

We show in Appendix A that the functional integral with source is

$$(1.3) \quad \begin{aligned} & \int \partial\phi \exp \left[\int_{\mathcal{T}} (\mathcal{L}_0 + \delta(f(x)) \phi(x) \partial_\mu \phi(x) \partial^\mu \phi(x)) \right] = \\ & = \text{const}(r) \exp \left[\frac{1}{2} \left(\iint_{\mathcal{T} \times \mathcal{T}'} f(x) G_D^{T'}(x, x') f(x') + \right. \right. \\ & \left. \left. + \frac{r}{2} \iint_{\mathcal{T} \times \mathcal{T}'} f(x) G_N^{T'}(x, x') f(x') \right) \right]. \end{aligned}$$

Here G_D^T is the Dirichlet Green's function in \mathcal{T}

$$(1.4a) \quad (m^2 - \partial_x^2) G_D^T(x, x') = \delta(x-x') \quad x, x' \in \mathcal{T}$$

$$(1.4b) \quad G_D^T(x, x') = 0 \quad x \in \partial\mathcal{T}, \quad x' \in \mathcal{T}'$$

and $G_N(x, x')$ the Neumann Green's function * in \mathcal{T}'

* If \mathcal{T}' is infinite and $m^2 = 0$, G_N vanishes in infinity strongly enough to render this function unique. (See also Appendix A.)

$$(1.5a) \quad (m^2 - \partial_x^2) G_N^{T'}(x, x') = \delta(x-x') \quad x, x' \in \mathcal{T}'$$

$$(1.5b) \quad \partial_n G_N^{T'}(x, x') = 0 \quad x \in \partial\mathcal{T}, \quad x' \in \mathcal{T}'$$

with ∂_n the (to \mathcal{T}' , interior) normal derivative at x .

We can say for short that in (1.3)

$$(1.6a) \quad \phi(x) \rightarrow 0, \quad x \rightarrow \partial\mathcal{T} \text{ from } \mathcal{T}'$$

$$(1.6b) \quad \partial_n \phi(x) \rightarrow 0, \quad x \rightarrow \partial\mathcal{T} \text{ from } \mathcal{T}'$$

meant in the sense of arguments of correlation functions. Note that in (1.3) there are no correlations between points in \mathcal{T}' and in \mathcal{T} , that is, \mathcal{T}' and \mathcal{T} have been decoupled from each other by the surface interaction.

We recall the familiar relations

$$(1.7a) \quad G_o^T(x, x') = G_o^T(x, x'), \quad G_v^T(x, x') = G_v^T(x, x')$$

$$(1.7b) \quad \lim_{\substack{x \rightarrow \partial T \\ x' \in \partial T}} G_o^T(x, x') \delta_{\nu}^{\mu} = \delta(x, x')$$

$$(1.7c) \quad \lim_{\substack{x \rightarrow \partial T \\ x' \in \partial T}} \partial_{\nu} G_o^T(x, x') \frac{\delta_{\nu}^{\mu}}{A(\partial T)} = \delta(x, x')$$

where $\delta(x, x')$ is the δ -function on ∂T . The function

$$\lim_{\substack{x \rightarrow \partial T \\ x' \in \partial T}} \partial_{\nu} G_o^T(x, x') \delta_{\nu}^{\mu} =$$

$$= \lim_{\substack{x \rightarrow \partial T \\ x' \in \partial T}} \partial_{\nu} G_o^T(x, x') \delta_{\nu}^{\mu} = \partial_{\nu} G_o^T \delta^{\mu}_{\nu}$$

is a negative definite kernel on ∂T which will appear often later.

Inhomogeneous Dirichlet boundary conditions are implemented by replacing

L_T of (1.2) by

$$(1.9) \quad L_{T,A} = L_T + \delta(f(x)) A(x) \partial_{\nu} G_o^T \delta_{\nu}^{\mu} \partial_{\mu} f(x).$$

In simplified notation,

$$(1.10) \quad L_{T,A} = -\frac{1}{2} \partial_{\nu} \phi \partial_{\mu} \phi - \delta(\sigma) A \partial_{\nu} \phi$$

+ $\delta(\sigma) \phi \partial_{\nu} \phi - \delta(\sigma) A \partial_{\nu} \phi$
where the operator ordering in the third term, specified in (1.2), must be kept in mind. From (1.3) follows by substitution for \check{J}

$$(1.11) \quad \begin{aligned} & \int \partial_{\nu} \phi \exp \left[\int L_T A + \int J \phi \right] = \\ & = \text{const}(T) \exp \left[\frac{1}{2} \int \int G_o^T + \frac{1}{2} \int \int G_v^T + \right. \\ & \quad \left. + \frac{1}{2} \int \int A \partial_{\nu} G_o^T \delta_{\nu}^{\mu} A - \int \int A \partial_{\nu} G_o^T J \right] \end{aligned}$$

where, however, we have discarded a singular linearly divergent term $\text{prop. } \int \partial_{\nu} A^2$ in the exponent which, to allow explicit subtraction in (1.10), requires smearing the last $\partial_{\nu} \phi$ in (1.10) with an L^2 function within a layer adjoining ∂T inside T and letting that layer shrink to zero width. (We shall put this in formulae in sect. 4.2.) From (1.11) and (1.7b) follows that now (1.6a) is replaced by

$$(1.12) \quad \phi(x) \rightarrow A(x \in \partial T), \quad x \rightarrow \partial T \text{ from } T$$

while (1.6b) remains unchanged.

If ∂T is a plane, there is no difficulty in taking space parallel to the plane $3-\mathcal{E}$ dimensional ($\mathcal{E} > 0$), such that $\nu = 4 - \mathcal{E}$. We give formulae, to be used later, for this case. We set in (1.2) $f(x) = y$, with x the $3 - \mathcal{E}$

* All underlined arguments are $3-\mathcal{E}$ -dimensional.

dimensional coordinate along the plane and the y -axis pointing into \mathcal{T}^* orthogonal to the plane, such that $y, y' > 0$ is the Dirichlet and $y, y' < 0$ the Neumann region. To simplify, we set $m = 0$ and then have

$$(1.13) \quad G_D(x, x') = \frac{1}{4} \pi^{-2 + \epsilon/2} T^*(1 - \frac{\epsilon}{2}\epsilon) \cdot$$

$$\cdot \left\{ (x-x')^2 + (y-y')^2 \right\}^{-1 + \frac{\epsilon}{2}} \cdot \overline{F((x-x')^2 + (y+y')^2)^{-\eta + \frac{\epsilon}{2}}} \epsilon \}$$

with Fourier transforms

$$\int d\mathbf{x} G_D(x, x') e^{ik(x-x')} =$$

$$(1.14)$$

$$= (2k)^{-1} \left[e^{-k/y-y'} \right] \overline{e^{-k/y+y'}} \equiv \tilde{G}_D(k, y, y')$$

where $k = |k|$ (or $= (k^2 + m^2)^{1/2}$ if $m > 0$). One notes the singularity in

$$(1.15) \quad \partial_{y'} \partial_{y'} \tilde{G}_D(k, y, y') = \delta(y-y') - k^2 \tilde{G}_N(k, -y, -y')$$

(1.8) becomes

$$(1.16) \quad F.T. \partial_n G_D^{xx'} = -k.$$

For one argument on the boundary with inner normal derivative,

$$(1.17) \quad \partial_{y'} G_D(x, y, y') \Big|_{y'=0} =$$

$$= -\pi^{-2 + \frac{\epsilon}{2}} \epsilon T^*(2 - \frac{\epsilon}{2}\epsilon) \gamma [(x-x')^2 + y^2]^{-2 + \frac{\epsilon}{2}}$$

with Fourier transform e^{-ky} .

The Fourier transform with respect to all variables is (again, for $m \geq 0$)

$$(1.18)$$

$$\begin{aligned} & \int dy e^{-iy} \int dy' e^{-iy'} \left[\theta(y) \theta(y') \tilde{G}_D(k, y, y') + \right. \\ & \quad \left. + \theta(-y) \theta(-y') \tilde{G}_N(k, y, y') \right] = \\ & = 2\pi \delta(y+y') (k^2 + y^2)^{-\eta} \left[k^2 + ik(y+y') + yy' \right]. \end{aligned}$$

Here the first term is the free-space function, and the second term, being generated by the surface interaction in (1.2), is a sum of factorizing parts. (1.18) is less singular than are the Fourier transforms of the Dirichlet or Neumann function alone.

2. Renormalization

To be able to add to (1.10) the interaction term, we must choose a regularization. While the renormalized theory is fixed by the renormalization conditions for the superficially divergent (s.d.) vertex functions (the amputated one-particle-irreducible (1PI) parts of the connected Green's functions) alone, the precise form of the counter terms depends on the regularization employed.

For most of our considerations, a plane $\mathcal{D}\mathcal{T}^*$ is sufficient, and then, for calculations, dimensional regularization is the most convenient one: $3-\epsilon$ space dimensions ($\text{Re } \epsilon > 0$), one (Euclidean or Minkowskian) time dimension,

This is as effective as introducing a lattice in space while keeping time continuous, but by itself does not break Lorentz invariance such that renormalization of the speed of light is not needed. Leaving the time direction unregularized means, however, that the ordering prescriptions in (1.2) and (1.9) must be kept in mind and that the form of the counter terms to these "bare" terms is in general not the one expected on the basis of naive canonical reasoning. (Pauli-Villars regularization is stronger and more generally applicable, but more cumbersome to calculate with.)

Therefore, with plane $\partial\mathcal{T}$, we add to (1.10) the interaction term

$$(2.1) \quad L_{int} = -\frac{1}{24} g \mu \epsilon \phi^4$$

and counter terms to cancel the singularities prop. \mathcal{E}^{-n} developing in Green's functions when $\epsilon \rightarrow 0$. Such singularities stem from coalescing vertices in (here, Euclidean) s.d. 1PI Feynman graphs. Counter terms need be associated only with final subtractions, since divergences due to coalescence of a subset of the vertices are compensated by the counter terms for final subtractions of the corresponding subgraph [3,6]. In the final subtraction, all vertices of the graph coalesce.

It is now convenient to separate the Dirichlet and Neumann propagator (1.13) or (1.14) or (1.18) into the free-space part and the remainder, which involves the surface vertex in (1.2) at least once and which we call surface propagator. (This separation is manifest also for nonplanar $\partial\mathcal{T}$, cp. (A.11), (A.14).) If a graph has at least one surface propagator, it can coalesce only on $\partial\mathcal{T}$. Thus, the 1PI graphs requiring final subtraction are either free-space ones,

if they involve free-space propagators only, or surface ones. The free-space ones require in L the usual covariant counter terms

$$(2.2) \quad \Delta L = -(Z_3 - 1) \frac{1}{2} \partial_\mu \phi \partial_\nu \phi -$$

$$- (Z_1 - 1) \frac{1}{24} g \mu \epsilon \phi^4 - (Z_2 - 1) \frac{1}{2} m^2 \phi^2$$

since the s.d. of these graphs is the usual one:

$$(2.3) \quad \mathcal{D}_E = 4 - E$$

where E is the (even) number of external lines.

2.2 Surface graphs and surface counter terms

It here suffices to consider $\partial\mathcal{T}$ as effectively flat. The singularity of a surface propagator, with both endpoints close to $\partial\mathcal{T}$ and also to each other, is the same as for the free-space propagator, see (1.13). Therefore, if one vertex in a coalescing group of vertices is fixed near $\partial\mathcal{T}$, the s.d. therefrom is the same as for the corresponding free-space graph. There is in addition, however, the integration of the fixed vertex over a small distance across $\partial\mathcal{T}$. This reduces the s.d. relative to the free-space graph by one, such that

$$(2.4) \quad \mathcal{D}_{\partial\mathcal{T}} = 3 - E$$

Thus, for $E = 0$, a "vacuum" graph, the divergence is cubic (we return to this

in sect. 3), and for $E = 2$ linear. The latter requires to add to the action density

$$(2.5) \quad \Delta L_{\partial\mathcal{T}} = (Z_4 - 1) \delta(\sigma) \phi \partial_n \phi + \\ + (c_1 A + c_2 R^{-1} \ln A) \delta(\sigma) \delta(\sigma) A \phi + \\ + (c_5 A + c_6 R^{-1} \ln A) \delta(\sigma) \delta(\sigma) A^2.$$

In dimensional regularization, $Z_4 - 1$ can be chosen as a power series in ϵ^{-1} .

The linear divergence also requires the term proportional to the cutoff (up to logarithms), absent if dimensional regularization were sufficient, and the term proportional to a typical curvature R^{-1} of $\partial\mathcal{T}$, with logarithmically divergent coefficient (here indicated symbolically), vanishing for flat $\partial\mathcal{T}$.

Lastly we consider the effect of the A-vertex in (1.10). It binds one external leg of the graph with a normal derivative to the surface. The s.d. of a surface graph is then the same as if that line were amputated, since a gain of three powers (2+1) from the line and derivative is compensated by the same loss due to the three-dimensional integration over the surface, supposing that the endpoint on the surface is smeared with a smooth function A. (It actually suffices that the endpoint does not coincide with some other endpoint of the graph on $\partial\mathcal{T}$. Thus, altogether

$$(2.6) \quad \partial_{\partial\mathcal{T}} A = \beta - E - E_A$$

and $\beta = 1$ for $E = E_A = 1$ and for $E = 0$, $E_A = 2$. This demands in addition to (2.5) the counter terms

$$(2.7) \quad \Delta L_{\partial\mathcal{T}} A = -(Z_5 - 1) \delta(\sigma) A \partial_n \phi +$$

$$+ (c_3 A + c_4 R^{-1} \ln A) \delta(\sigma) \delta(\sigma) A \phi + \\ + (c_7 A + c_8 R^{-1} \ln A) \delta(\sigma) \delta(\sigma) A^2.$$

Again, $Z_5 - 1$ can be chosen in dimensional regularization to be a power series in ϵ^{-1} , and to the c-terms what was said after (2.5) applies.

Collecting the action and adding here, for completeness, also the source term for ϕ^2 we have

$$(2.8) \quad L = -\frac{1}{2} Z_3 \partial_n \phi \partial_n \phi - \frac{1}{24} Z_7 g \mu \epsilon \phi^4 - \\ - \frac{1}{2} Z_2 m^2 \phi^2 + J \phi + \frac{1}{2} K Z_2 \phi^2 + \\ + Z_4 \delta(\sigma) \phi \partial_n \phi - Z_5 \delta(\sigma) A \partial_n \phi + \\ + Z_6 \mu^{-\epsilon} m^2 K - Z_6 \mu^{-\epsilon} m^{-\epsilon} K^2 + \\ + Z_7 \delta(\sigma) \mu^{-\epsilon} \partial_n K + \text{c-terms}$$

where the last ones now also encompass

$$(c_7 A + c_8 R^{-1} \ln A) \delta(\sigma) \mu^{-\epsilon} K.$$

The perturbation expansion with (2.8) (setting $K \equiv 0$) we discuss in Appendix B, and here only summarize the results:

The calculations on the Dirichlet and on the Neumann side can be done separately, due to the decoupling of the two regions mentioned after (1.3). The (unamputated!) Green's functions on the Dirichlet side obey the Dirichlet condition where $A = 0$, and are, as a consequence of this, independent of the renormalization condition that fixes the choice of Z_4 which, in dimensional renormalization, is

$$(2.9) \quad Z_4 = 1 + \frac{1}{6} \pi^{-2} \varepsilon^{-1} g + O(g^2).$$

(That it differs from the usual

$$(2.10) \quad Z_3 = 1 - (3 \cdot 2^{10} \pi^4 \varepsilon)^{-1} g^2 + O(g^3)$$

Here as usual

is due to the incompleteness of the regularization as emphasized in sect. 2.1.)

Also the c -terms in (2.5) and (2.7) are ineffective on the Dirichlet side.

On the Neumann side, the Neumann boundary condition (1.6b) cannot be upheld already to first order in perturbation theory, due to the necessity, for finiteness, of the c_1 -term in (2.5). However, since the Neumann part factors off and is Λ -independent (for this the Neumann property (1.5b) of the bare propagators suffices), we need not discuss it further.

2.3 Renormalization_group_properties

The functional integral with action density (2.8) (setting $K \equiv 0$) we denote by $\mathcal{W}(A/J)$. It factorizes into the Dirichlet part, depending on A and on J in \mathcal{T} , and the Neumann part, depending on J in \mathcal{T}' , and we disregard the second factor until sect. 3.

From (2.8) one derives $[J]$ by differentiation with respect to J the renormalization group equation

$$(2.11) \quad \left[\mu \frac{\partial}{\partial \mu} + \beta(g, \varepsilon) \frac{\partial}{\partial g} + \sigma(g) \int_{\mathcal{T}} \frac{\partial}{\partial J} - \sigma(g) \int_{\mathcal{T}'} A \frac{\partial}{\partial A} + \eta(g) m^2 \frac{\partial}{\partial m^2} \right] \mathcal{W}(A/J) = 0.$$

$$(2.12a) \quad \beta(g, \varepsilon) = -\varepsilon g + \beta(g) = -\varepsilon g \left[1 + g \frac{\partial^2}{\partial g^2} \ln(Z, Z^{-2}) \right]^{-1} =$$

$$= -\varepsilon g + (16\pi^2)^{-1} 3g^2 + O(g^3)$$

$$(2.12b) \quad \sigma(g) = \frac{1}{2} \beta(g, \varepsilon) \frac{\partial}{\partial g} \ln Z_3 = (3 \cdot 2^{10} \pi^4)^{-1} g^2 + O(g^3)$$

$$(2.12c) \quad \eta(g) = \beta(g, \varepsilon) \frac{\partial}{\partial g} \ln(Z_2 Z_3^{-1}) = (16\pi^2)^{-1} g + O(g^3)$$

while

$$(2.12d) \quad \sigma(y) = \beta(y, \epsilon) \frac{\partial}{\partial y} \ln (Z_5 z_3^{-1}) =$$

$$= (32\pi^2)^{-1}y + O(y^2)$$

is a new parametric function, ϵ -free under minimal choice of

$$(2.13) \quad Z_5 = 1 - (32\pi^2\epsilon)^{-1}y + O(y^2).$$

Note that the differentiation leading to (2.9) produces also the insertion

$$(2.14) \quad \Delta L = \beta(y, \epsilon) \frac{\partial}{\partial y} \ln (Z_5 z_3^{-1}) \cdot Z_4 \delta(t) \phi \partial_n \phi$$

in the functional integral. However, this insertion gives zero due to the Dirichlet condition emphasized at the end of sect. 2.2, and we may consider the $-A \partial_n \phi$ vertex in (2.8) as going to $\partial \mathcal{T}$ from \mathcal{T} latest as we shall make explicit in sect. 4.4.

To keep the notation simple, we shall now take as $\partial \mathcal{T}$ the (Euclidean) time plane $y = 0$, as we did at the end of sect. 1. We write the sources $J(xy)$ and $A(z)$. The Green's functions

$$(2.15) \quad G(z_1, \dots, z_L/x, y_1, \dots, y_n) =$$

$$= \prod_{j=1}^L \int dA(x_j) \int dT [D/\partial x_j(x_j; x_i)] \ln \psi(A(x_i)) / \int dA = 0$$

are connected, and zero unless $n + L = \text{even}$. (2.9) becomes

$$(2.16) \quad \left[n \frac{\partial^2}{\partial \epsilon^2} + \beta(y, \epsilon) \frac{\partial^2}{\partial y^2} + n \sigma(y) - L(\sigma, y) + \right. \\ \left. + 2(y/n)^2 \frac{\partial^2}{\partial \epsilon^2} \right] G(z_1, \dots, z_L/x, y_1, \dots, y_n) = 0.$$

3.1 General discussion

Renormalization renders all Green's functions finite. It leaves untouched the quartically UV-divergent vacuum graphs, i.e. those without external lines, for which final subtraction (which would simply remove the whole graph) is not prescribed. It can be meaningful, however, to compare the vacuum energy, obtained (for the scalar field) by

$$(3.1) \quad E_{\bar{\mathcal{T}}} = - \lim_{T \rightarrow \infty} T^{-1} \ln \int d\mathbf{x} \int d\mathbf{x}' \exp \left[\int d\mathbf{x} \int d\mathbf{x}' \mathcal{L}(\phi, \dot{\phi}) \right]$$

for different $\nu-1$ -dimensional regions $\bar{\mathcal{T}}$ and with different boundary conditions on the $\nu-2$ -dimensional boundary $\partial \bar{\mathcal{T}}$. (The relation to the notation so far is: $T = \bar{\mathcal{T}} X[\phi, T]$, $\partial T = \partial \bar{\mathcal{T}} X[\phi, T]$, plus the irrelevant $t = 0$ and $t = T$ closures.)

One easily sees, however, that the quantity that is simple to compute is not the vacuum (i.e. ground state) energy in $\bar{\mathcal{T}}$, but the total energy, which is

the ground state energy in $\bar{\mathcal{T}}$ with e.g. Dirichlet boundary conditions plus the one in the complementary region $\bar{\mathcal{T}}'$ with Neumann boundary conditions. Then the boundary-independent free-space energy can be omitted, and the remainder is given by surface graphs only. In particular, in the simplest setting of Dirichlet conditions on the inner sides of two parallel plates, in distance L , the Neumann part is independent of L such that the L -dependence of the total energy is the same as if the two Neumann regions were absent, disregarding the free-space part.

As we shall see, the Casimir potential between disjoint surfaces is always well defined. The one for a single surface, e.g. a sphere, in general is not, at least for the family of boundary conditions discussed in Appendix A, due to divergence already of the free-field part if taken absolutely and not relative to e.g. some other shape. In such a case, the physical problem requires a more complete formulation, which will then imply some other boundary condition [8] than the (idealizing) Dirichlet (or Neumann) one.

3.2 Free field

Here, the surface graphs have merely the $\phi \partial_n \phi$ -vertices on $\partial\mathcal{T}$ shown in (1.2), and it is easy to derive the simple graphical expansion *

* (3.2) is a special case of a formula of Balian and Duplantier [9].

$$(3.2) \quad E_{Dir} + E_{Neu\bar{\mathcal{T}}} - const = - \lim_{r \rightarrow \infty} \frac{1}{2} r^{-1} \left[2 \text{Tr} \overline{\partial_n G} + \frac{1}{2} 2^2 \text{Tr} \overline{\partial_n G} \cdot \overline{\partial_n G} + \frac{1}{3} 2^3 r \overline{\partial_n G} \cdot \overline{\partial_n G} \cdot \overline{\partial_n G} + \dots \right]$$

where Tr is the trace on the surface $\partial\mathcal{T} = \partial\bar{\mathcal{T}} \times \{0, \tau\}$ and $\overline{\partial_n G}$ is defined in (A.5b). The first term in the square bracket is zero under appropriate (e.g., Pauli-Villars) regularization for symmetry reasons. (This reflects the well-known fact that the strongest, cubically divergent part of the surface energy has the opposite sign in the Dirichlet and Neumann case.) The higher terms in (3.2) all vanish for flat $\partial\mathcal{T}$ (i.e., $\overline{\partial_n G}$) since then $\overline{\partial_n G}$ is zero (see Appendix A).

In the arrangement of two parallel plates in distance L , however, $\overline{\partial_n G}$ is not zero if its two arguments are on different plates. In the massless theory in ν dimensions, i.e. two $\nu-2$ -dimensional parallel plates in $\nu-1$ -dimensional space,

$$(3.3) \quad \overline{\partial_n G} = -\frac{1}{2} r^{-\nu/2} \text{Tr} \left(\frac{1}{2} \omega_x \right) \left(\frac{1}{2} \omega_{x'} \right)^2 \angle^{2\nu-2} \angle^{-\nu/2} \angle$$

where x and x' are the $(\nu-2+1)$ -dimensional arguments on the two plates extended in Euclidean time. While on the r.h.s. of (3.2) the odd terms all vanish, the even ones are easily summed and lead to the well-known result

$$(3.4) \quad \left(E_{Dir} + E_{Neu\bar{\mathcal{T}}} - const \right) / \text{area} = -2^{-\nu} r^{-\nu/2} \text{Tr} \left(\frac{1}{2} \omega_x \right) \zeta(\nu) L^{-\nu+1} = -\frac{1}{1440} \pi r^2 L^{-3} e^{\nu\pi/4}.$$

Hereby in $\zeta(\nu) = \sum_{n=1}^{\infty} n^{-\nu}$ the n th term is obtained from the "one-loop" polygon with n vertices on each plate. - (3.2) also shows that the

Casimir potential decays exponentially in the massive case.

3.3 Interacting field

The higher-order vacuum surface graphs are (for $\nu = 4$) cubically divergent ($E = 0$ in (2.3)) provided they can shrink to a point on $\partial\mathcal{T}$. If, however, $\partial\mathcal{T}$ consists of two disjoint pieces $\partial\mathcal{T}_1$ and $\partial\mathcal{T}_2$, the surface graphs on $\partial\mathcal{T}_1$ are independent of the location of $\partial\mathcal{T}_2$, and vice versa. Only the surface graphs with vertices on both $\partial\mathcal{T}_1$ and $\partial\mathcal{T}_2$ depend on the relative location, and since they cannot be shrunk to a point they are finite provided all subdivergences have been subtracted by counter terms. These are the usual free-space counter terms (2.2) and the ones on $\partial\mathcal{T}_1$ and $\partial\mathcal{T}_2$, as prescribed by (2.5), that render the surface graphs for these sub-surfaces finite. Thus, the Casimir effect is finite and computable for any configuration of disjoint surfaces at each of which the theory has been made finite by counter terms as if the other surfaces did not exist.

Since we know (see Appendix 2) that the homogeneous Dirichlet condition can be implemented for the plane without need of a new renormalization parameter, the Casimir energy of a pair of parallel plates with these conditions on the two insides obeys, in massless ϕ_4^4 theory, according to (3.1) and (2.16),

$$(3.5a) \quad \int \mu \frac{\partial}{\partial \mu} + \beta(g, \epsilon) \frac{\partial}{\partial g} \int E(g, L, \mu, \epsilon) = 0.$$

Thus

$$(3.5b) \quad E(g, L, \mu, \epsilon)/\text{area} = \int^{-3+\epsilon} e^{(g/\mu)(\gamma_{\mu}-\gamma_{\epsilon})} \epsilon^2 = \int^{-3+\epsilon} \sum_{n=0}^{\infty} C_n(\epsilon) \left[\bar{g}(g, (\gamma_{\mu}-\gamma_{\epsilon})\mu) \right]^n + O(g^2).$$

where \bar{g} is the usual sliding coupling constant and the $c_n(\epsilon)^*$ are computable,

* we do not discuss here the IR problems [10] of the expansion in (3.5b) if $\epsilon > 0$.

with $c_0(\epsilon)$ from (3.4).

Since the massless ϕ_4^4 theory is "asymptotically free in the infrared", \bar{g} in (3.5b) vanishes, if $\epsilon = 0$, proportional to $(\ln L)^{-1}$ as $L \rightarrow \infty$, such that then only the $c_0(0)$ -term survives. This universality (i.e., independence of g) is a counter part of the one observed by Lüscher [11] in a two-dimensional problem, and is, since insensitive to UV cutoffs, not affected by the fairly well established [12] nonexistence of the continuum ϕ_4^4 theory in the UV.

3.4 First order calculation

In terms of zeroth-order Dirichlet Green's functions, we have, from (3.1, 5b)

$$(3.6) \quad \begin{aligned} & \int^{-3+\epsilon} g(\epsilon) \bar{g}(g, (\gamma_{\mu}-\gamma_{\epsilon}), \epsilon) = \\ & = \frac{1}{g} \gamma_{\mu} \epsilon \int \int \int \bar{g}_0^4(\partial_Y \bar{g}_0(\partial_Y \partial_Y) - G(00))^2 - \\ & - \int G_0(\partial_{\mu} \partial_{\mu} - G(00))^2 - \\ & - \int G_0(\partial_{\mu} \partial_{\mu} - G(00))^2 - \\ & - 2 \int \int \bar{g}_0^2 [G_0(\partial_Y \partial_Y) - G(00)]^2 + O(g^2). \end{aligned}$$

Here,

$$(3.7) \quad G_d^L(x, y, x', y') = \\ = \frac{1}{4} \pi^{-2 + \varepsilon/2} T(1 - \frac{\varepsilon}{2}) \cdot \sum_{n=-\infty}^{\infty} \left\{ \left[(\gamma - x')^2 + (\gamma - y' + 2Ln)^2 \right]^{-1 + \varepsilon/2} \right. \\ \left. - \left[(\gamma - x')^2 + (\gamma + y' + 2Ln)^2 \right]^{-1 + \varepsilon/2} \right\}$$

is the Dirichlet Green's function for a pair of $3-\varepsilon$ -dimensional plates at $y = 0$ and $y = L$, while G_D is given in (1.13) and $G(\dots)$ is the free-space function. The cross term from the first square bracket in the curly bracket in (3.6) removes the free-space mass renormalization to the one-loop (i.e., zeroth order) graph discussed in sect. 3.2. The second and third term in the curly bracket in (3.6) remove the two single-plate surface graphs to first order. The last integral in (3.6) extends these two last graphs to the whole half-space relevant for the single-plate problem. (Namely, otherwise the integration over $0 \dots L$ only would have given an unallowed L -dependence to the single-plate subtractions.) Apart from this, in (3.6) the integration over $0 \dots L$ only is permitted since the energy contributions from the two outer regions are, after the free-space parts are subtracted, L -independent.

We now insert into (3.6)

$$(3.8) \quad G_d^L(\varrho_y, \varrho_y) - G(\varrho_0) = \frac{1}{4} \pi^{-2 + \varepsilon/2} (2L)^{-\varepsilon/2} T(1 - \frac{\varepsilon}{2}) \cdot \\ \cdot \left[2 \Im(\varrho - \varepsilon) - \Im(\varrho - \varepsilon, y/L) - \Im(\varrho - \varepsilon, 1 - y/L) \right], \\ = (4\pi L^2)^{-1} \left[1 - 3(\sin(\pi y/L))^{-2} \right] \text{ if } \varepsilon = 0,$$

where $\Im(z, \alpha)$ is the generalized \Im -function, and find

$$(3.9) \quad C_1(\varepsilon) = 2^{-9 + 2\varepsilon} \pi^{-4 + \varepsilon} T(1 - \frac{\varepsilon}{2})^2 \cdot \\ \cdot \left[\Im(\varrho - \varepsilon)^2 + (1 - \cos \pi \varepsilon) B(-1 + \varepsilon, 3 - 2\varepsilon) \Im(3 - 2\varepsilon) \right]$$

$$= 2^{-7\pi} \cdot 3^{-2} \quad \text{if } \varepsilon = 0,$$

* This result was obtained by Toms [13] using dimensional regularization.

$$= - \frac{3}{3} 2^{-9} (1 + 48\pi^{-2}) \quad \text{if } \varepsilon = 1 \\ \text{where } \partial_x = \frac{1}{2} \partial^2 / \partial s^2 \left[(s - 1) \Im(s) \right] \Big|_{s=1}.$$

4. Behaviour at the boundary

4.1 Calculations to first order

For the Green's functions (2.13), we define Fourier transforms by

$$\int dz_1 \dots dz_n \int d\chi_1 \dots d\chi_n \exp \left[i \sum K_j z_j + i \sum \rho_i \chi_i \right].$$

$$\cdot G(z_1 \dots z_n, \chi_1 \dots \chi_n) = \\ = (2\pi)^{4-\varepsilon} \delta(\sum \rho_i + \sum K_i) \tilde{G}(K_1, \dots, K_n / \rho_1, \chi_1 \dots \chi_n / \rho_n).$$

Using the propagator (1.14), we find to first order in g

$$(4.1) \quad \tilde{G}(\rho\gamma(-\rho)\gamma') = \theta(\gamma-\gamma') \left\{ \rho e^{-\rho t} \operatorname{sh}(\rho\gamma') + \right.$$

$$+ 2^{-5+\varepsilon} \pi^{-2+\varepsilon/2} \rho^2 g \rho \varepsilon \left[\int dt t^{-2+\varepsilon} (\operatorname{sh}(\rho t))^\varepsilon + \right.$$

$$+ e^{-\rho t} \operatorname{sh}(\rho\gamma') \left(\int dt t^{-2+\varepsilon} \operatorname{sh}(gt) e^{-\rho t} \right) e^{-\rho t} +$$

$$+ \operatorname{sh}(\rho\gamma') \operatorname{sh}(\rho\gamma') \left(\int dt t^{-2+\varepsilon} e^{-2\rho t} \right) e^{-\rho t} +$$

$$\left. + (\gamma \rightarrow \gamma') \right\}$$

where $\rho = \frac{1}{2}t$. While the Dirichlet condition $\tilde{G}((\rho\gamma(-\rho))0) = 0$ is satisfied, $\partial_{\gamma'} \tilde{G}((\rho\gamma(-\rho))\gamma')$ has, if $\varepsilon > 0$ but $\gamma' = 0$, an ε^{-1} singularity as $\varepsilon \rightarrow 0$, whereas, if $\varepsilon = 0$, it blows up logarithmically as $\gamma' \rightarrow 0$.

In the first case, the remedy is the factor Z_5 , given in (2.13) to this order, while at $\varepsilon = 0$,

$$(4.2) \quad \lim_{\gamma' \rightarrow 0} \left[1 + (32\pi^2)^{-1} g^2 \ln(g\gamma') \right] \partial_{\gamma'} \tilde{G}((\rho\gamma(-\rho))\gamma') =$$

$$= \tilde{G}(-\rho\gamma(-\rho)) = e^{-\rho\gamma} \left[1 + (32\pi^2)^{-1} g^2 (4\gamma^2 - \ln(g\gamma')) \right]$$

$$- (32\pi^2)^{-1} g^2 e^{\rho\gamma} \int dt t^{-1} e^{-2\rho t} \gamma'$$

The r.h.s. differs insignificantly from the corresponding minimal-subtraction function computed directly from (2.8) using the prescriptions of Appendix B.

Letting $\gamma \rightarrow 0$ in the $\varepsilon > 0$ formula for $\tilde{G}(-\rho/\rho\gamma)$ again produces a $1/\varepsilon$ singularity, remedied by the factor $1 + (32\pi^2)^{-1} g^2$, which is Z_5^{-1} to this order. At $\varepsilon = 0$, we must again introduce a logarithmic factor,

$$(4.3) \quad \lim_{\gamma \rightarrow 0} \left\{ \left[1 - (32\pi^2)^{-2} (1 + \ln(g\gamma)) \right] \tilde{G}(-\rho/\rho\gamma) \right\} = 1$$

If we form $\partial_{\gamma'} \tilde{G}(-\rho/\rho\gamma)$, however, we must, to let $\gamma \rightarrow 0$, make a subtraction, which we choose at zero momentum. Thereupon, at $\varepsilon = 0$, the same factor as in (4.2) is required,

$$(4.4) \quad \lim_{\gamma \rightarrow 0} \left\{ \left[1 + (32\pi^2)^{-1} g^2 \ln(g\gamma) \right] \partial_{\gamma'} \tilde{G}(-\rho/\rho\gamma) - \partial_{\gamma'} \tilde{G}((\rho(-\rho))/) \right\} =$$

$$= -\rho \left[1 - (16\pi^2)^{-1} (\ln(2\rho\mu^{-1}) - \varphi(2)) \right].$$

whereas at $\varepsilon > 0$ again the factor Z_5 from (2.13) is appropriate. As in (4.2), the r.h.s. in (4.4) differs insignificantly from the corresponding minimal-subtraction result. The functions in (4.1), (4.2), and (4.4) obey (2.16) with (2.12) to first order in g .

For computations in Appendix D it is useful to observe that the r.h.sides of

(4.2) and (4.4) are also given by

$$(4.5a) \quad G(-\rho/\rho y) = e^{-\rho y} + \\ + (64\pi^2\rho)^{-1} \int_0^\infty dt e^{-\rho t} (e^{-\rho/y-t} - e^{-\rho(y+t)}) \mu^2 \rho (\frac{t}{y})^{-2}$$

4.2_Generating_surface_arguments

(2.8) shows that for $\epsilon > 0$,

$$(4.7) \quad Z_S \partial_y \phi(x,y) \xrightarrow[y \rightarrow 0]{} \partial/\partial A(x).$$

and

(4.2) indicates that, for $\epsilon = 0$, this needs to be replaced by

$$(4.5b) \quad G(-\rho/\rho) = -\rho + (32\pi^2)^{-1} \int_0^\infty dt e^{-\rho/y-t} \mu^2 \rho (\frac{t}{y})^{-2}$$

where

$$(4.6a) \quad \rho_x x^{-2} = \lim_{\Delta \rightarrow 0} [x^{-2} \theta(x-\Delta) - x^{-2} \theta(x)] - \delta'(x) \ln \Delta]$$

is the one-sided principal value related by

$$(4.6b) \quad \rho_x x^{-2} = \int x^{-2} - (x-2)^{-1} \delta'(x) / 2^{-2}$$

to the function defined by Gelfand and Shilow [4].

The other first-order graph is the lowest-order contribution to the four-point function $G(k_1 k_2 y_2 k_3 y_3 (-k_1 -k_2 -k_3) y_4)$. One easily sees that it, and all first y -derivatives, are ordinary functions of the k and the y , and vanish if one or more y is set zero without derivative.

$$(4.8b) \quad \lim_{y \rightarrow 0} \{ c_{xy} [\partial_y G(z_1 \dots z_n / x, y_1 \dots y_n) - \\ - \partial_{n+1} \partial_{n+2} \partial_x G(z_1 \dots z_n - x, y_1 \dots y_n)] \} =$$

$$= (\partial/\partial A(x)) \psi(A) \cup.$$

Here we recognize the subtraction corresponding to a particular choice of c_5 in (2.7), while the c_3 -term does not contribute due to the Dirichlet property. In terms of Green's functions (2.15), (4.8b) is

$$(4.8c) \quad \lim_{y \rightarrow 0} \{ c_{xy} [\partial_y G(z_1 \dots z_n / x, y_1 \dots y_n) - \\ - \partial_{n+1} \partial_{n+2} \partial_x G(z_1 \dots z_n - x, y_1 \dots y_n)] \} = \\ = G(z_1 \dots z_n / x, y_1 \dots y_n)$$

valid, however, only in the sense of distributions with smooth test functions A and J. We can choose

$$(4.9) \quad C(\gamma) = [\partial_y \tilde{G}(-\mu/\rho\gamma) - \partial_y \tilde{G}(0/\rho\gamma)]^{-1} \tilde{G}(\rho(-\mu)/\rho) / \rho_0$$

which defines $c(y)$ up to a merely g (and, in the massive case, $\ln \mu/\rho$) dependent factor that depends on the convention used in computing the r.h.s. of (4.9).

$c(y)$ as defined by (4.9) satisfies

$$(4.10) \quad \int \mu \frac{\partial^2}{\partial y^2} + \rho(y) \frac{\partial}{\partial y} - \sigma(y) - \sigma'(y) + \sigma(y) m^2 \frac{\partial}{\partial m} \tilde{G}(y) = 0$$

which renders (4.8c) consistent with (2.16). In the massless case, $c(y)$ is a power series in g with coefficients polynomials in $\ln \mu/\gamma$. In the massive case (4.9) also yields a dependence on m which can, however, be factored away under neglect of also $O(y)$ terms, since the UV effect to be achieved by $c(y)$ is m -independent. - Note that, while $z_5 z_3^{-1}$ in (2.10d) is, in minimal subtraction convention, uniquely determined by $\sigma(g)$, this is not so, except in the leading and next-to-leading logs, for $c(y)$ from (4.10) even if $m = 0$.

(4.8c) is easily interpreted: An external leg of G is upon normal-differentiation bent to the boundary. Therefore, we need discuss only the cases of superficial divergence, $n = 1, l = 0$ and $n = 0, l = 1$, since the other cases are then covered by skeleton expansions (Appendix B). In the first case,

$$(4.11) \quad \lim_{y \rightarrow 0} \left\{ c(y) \partial_y G(1/x, y, y_n) \right\} = G(x/x_n y_n)$$

we must remember (see Appendix B) that the functions here are singular if any

y goes to zero; $G(\underline{x}, y, y_1)$ is comparable to a four-point function of the covariant theory, with x_0 and \underline{x}_1^0 the two suppressed arguments, and $G(\underline{x}, y)$ is comparable to a three-point (i.e. unamputated mass vertex) function of the covariant theory, with \underline{x}_1^0 the suppressed argument. In this sense, (4.11) means that the renormalized vertex function is gotten from the four-point function by binding two legs together with a point-split bare vertex, multiplying by a factor that depends logarithmically on the splitting distance (and on g and μ) only, and letting that distance go to zero.

In the covariant case, the reason for this factorization is a Wilson short-distance expansion [15], which to the order required here can be derived elementarily by manipulating algebraically the Bethe-Salpeter equation (see, e.g., [16]). In analogy, the reason for validity of (4.11) is a small- y expansion of the "four-point function" on the l.h.s., the leading term being y times powers of $\ln(\mu/y)$. The proof, on the basis of the Bethe-Salpeter equation, will not be given here. We merely note that the reason for $c(y)$ being merely logarithmic, inspite of superficial linear divergence, is the appearance of the factor y due to the Dirichlet property.

The other case,

$$(4.12) \quad \lim_{y \rightarrow 0} \left\{ c(y) \left[\partial_y G(z/x, y) - \partial_z G(x-z, y) \right] \right\} = G(z/x)$$

is analogous to forming a two-mass-vertices correlation function in the covariant theory. Also there, an additive renormalization is required before

the remaining divergence can be removed multiplicatively. Overlapping divergences are disentangled by the subtraction (or momentum differentiation), which corresponds to multiplication by $\underline{x}_i - \underline{z}_i$. Again, the proof, analogous to the one in the covariant case, will not be given here. - In QED, the analogous procedure is applied when computing a quadratically divergent current correlation function or photon self energy part or linearly divergent electron self energy part.

4.3 Boundary value of ϕ

Going in (2.8) with ϕ to the boundary where it vanishes requires crossing the $\sim Z_5 A_{\partial x}$ term, and the canonical commutation relation gives

$$(4.13) \quad Z_5^{-1} Z_3 \phi(\underline{x}\underline{y}) \xrightarrow[\underline{y} \rightarrow 0]{} A(\underline{x})$$

for $\epsilon > 0$, while, as we saw in sect. 4.1, for $\epsilon = 0$ this needs to be replaced by

$$(4.14a) \quad \alpha(\underline{y}) \phi(\underline{x}\underline{y}) \xrightarrow[\underline{y} \rightarrow 0]{} A(\underline{x})$$

or, explicitly,

$$(4.14b) \quad \lim_{\underline{y} \rightarrow 0} \left\{ \alpha(\underline{y}) [\delta(\underline{x}/\underline{x}\underline{y})] \psi(A/\underline{y}) \right\} = A(\underline{x}) \psi(A/\underline{y})$$

or

$$(4.14c) \quad \lim_{\underline{y} \rightarrow 0} \left\{ \alpha(\underline{y}) G(z_1 \dots z_c / \underline{x}\underline{y} \underline{x}_1 \underline{y}_1 \dots \underline{x}_n \underline{y}_n) \right\} =$$

$$= \delta_{i1} \delta_{n0} \delta(\underline{x} - \underline{z}_1),$$

valid, again, only in the sense of distributions with smooth test functions A and J . We can choose

$$(4.15) \quad \alpha(\underline{y}) = \tilde{G}(\underline{0}/\underline{0}\underline{y})^{-1}$$

which obeys, according to (2.16),

$$(4.16) \quad \left[\mu \frac{\partial^2}{\partial \underline{y}^2} + \beta(\underline{y}) \frac{\partial}{\partial \underline{y}} - \delta(\underline{y}) + \sigma(\underline{y}) + e(\underline{y}) i m^2 \frac{\partial}{\partial \underline{m}_2} \right] \alpha(\underline{y}) = 0$$

and what was said after (4.10) on $c(y)$ applies also to $a(y)$.

(4.14c) is easily interpreted: If $l = 1$, $n = 0$, the l.h.s. vanishes due to the Dirichlet property if $\underline{x} \neq \underline{z}_1$. Thus, the r.h.s. if finite, and it is finite by the choice (4.15), must be a distribution with support on $\underline{x} = \underline{z}$, and by power counting, this must be a \mathcal{D} -function. In all other cases (i.e. unless $l = 1$, $n = 0$) power counting does not allow a singularity as strong as a \mathcal{D} -function on the r.h.s. of (4.1 c) if \underline{x} coalesces with any \underline{z} , the remaining arguments being understood not to coalesce with these or as being integrated over with smooth test functions.

4.4 A formula for $\psi_{(A/J)}$

By repeating the operation (4.8b) indefinitely, we can build up $\psi_{(A/J)}$ from $\psi_{(0/J)}$. However, due to the \mathcal{D} -singularity (1.15) of $\partial_{\underline{y}} \partial_{\underline{y}'} G(\underline{M}\underline{Y}\underline{x}'\underline{y}')$ at $\underline{x} = \underline{x}'$, $\underline{y} = \underline{y}'$, we must in the repetition employ a square-integrable smearing function $c_A(y)$ that, as $A \rightarrow \infty$, approaches $\delta(y)$, e.g.

$$(4.17) \quad C_{12}(y) = A^{-1} (n!)^{-1} (\gamma/1)^n \exp(-y/1), \quad n \geq 1.$$

Then

$$(4.18) \quad \begin{aligned} \psi(A/J) &= \\ &= \lim_{A \rightarrow \infty} e^{\chi p} \left[\int dz A(z) \int_0^\infty dy c_A(y) c(y) \partial_y (\partial_z c_A(y)) - \right. \\ &\quad \left. - \frac{1}{2} \int dy c_A(y) c(y) \int dz' c_A(y') c(y') \partial_y \partial_{y'} \tilde{G}(10y \partial_{y'}) \right]. \end{aligned}$$

$$\cdot \int dz A(z)^2 \] \psi(0/J)$$

where the A -smearing and limit is understood. The subtraction removes the singular part if the first term in the exponent is expanded in A , and effects $\tilde{G}(001) = 0$. Only the J, A -two point function is operative in yielding (4.14b) from (4.20). The compact formula (4.20) will be useful in the following section.

5. Schrödinger equation

5.1. Generalities

which is, in the limit $A \rightarrow \infty$, using (4.11) easily checked to be consistent with (4.8b). The subtraction in the exponent in (4.18) corresponds to the c_5 term in (2.7), and is needed even in the free-field case as remarked after (1.11). If $\psi(0/J)$ satisfies (2.11) with $A = 0$, then, due to (4.10), $\psi(A/J)$ does with $A \neq 0$.

In view of (2.15), we write

$$(4.19) \quad \psi(0/J) = \exp G(J)$$

Then (4.18) becomes, in symbolic notation

$$(5.1) \quad \int dx \partial_y J(x) [\partial_z \partial_y \psi(x)] \psi(A/J) = H(A, \delta/\delta A) \psi(A/J)$$

provided J has support at $y > 0$ only. $H(A, \delta/\delta A)$ is the Hamiltonian as a

$$(4.20) \quad \psi(A/J) = \exp \{ G(J + c A \partial_y) -$$

$$- \frac{1}{2} \int A^2 [c \partial_y G \partial_y, c] \}$$

The Schrödinger functional $\psi(A/J)$ is the scalar product $\langle A / e^{J\phi} \rangle$ of a state specified by the function A at (Euclidean) time zero with the state obtained from the vacuum by operating on it with sources at various $y > 0$. (More details will be given in sect. 6.) The dependence on the time of specification of A by shifting this time by $\Delta \tau$ in the negative y -direction is the same as when shifting the sources by $\Delta \tau$ in the positive y -direction, which means replacing $J(xy)$ by $J(x-y-\Delta \tau)$, since the vacuum state is translation invariant. The Schrödinger equation expresses this dependence by an operation on the functional dependence on A alone. Infinitesimally,

functional differential operator. In particular, (5.1) implies

$$(5.2) \quad H(A, \partial A) \psi(A|J) = 0$$

i.e. $H(A, \partial A)$ has the vacuum energy subtracted. We shall construct $H(A, \partial A)$ by analyzing the l.h.s. of (5.1) with the help of (4.20).

5.2 Free field and combinatorics

Disregarding renormalization problems, we can write, according to (1.3)

$$(5.3) \quad \psi(A|J) = \exp[-P(\phi|A)] \exp\left[\frac{1}{2} J G_0 J\right]$$

where $P(\phi)$ is the interaction part of the action. (Not to have derivatives in $P(\phi)$, ϕ should be the unrenormalized field.) Then, from (1.11)

$$(5.4a) \quad \psi(A|J) = \exp[-P(\phi|A)] \psi^0(A|J)$$

where

$$(5.4b) \quad \psi^0(A|J) = \exp\left[\frac{1}{2} J G_0 J - A \partial_n G_0 J + \frac{1}{2} A \partial_n G_0 \partial_n A\right]$$

with ∂_n the outer normal derivative $- \partial_J|_{J=0}$. Straightforward calculation gives

$$\begin{aligned} (5.5) \quad & \partial_J \left[\partial_J \psi^0 \right] \psi^0(A|J) = \\ & = \exp[-P(\phi|A)] \left\{ -\frac{1}{2} J (\partial_J G_0 + G_0 \partial_J) J + \right. \\ & \quad \left. + \int \partial_J G_0 \partial_n A - P'(\phi|A) \partial_J G_0 \partial_n \right\} \psi^0(A|J). \end{aligned}$$

The following equations hold (the dot means integration over the $y = 0$ plane)

$$(5.6a) \quad \partial_J G_0 + G_0 \partial_J = G_0 \partial_n \cdot \partial_n G_0$$

$$(5.6b) \quad \partial_J G_0 \partial_n' = G_0 \partial_n \cdot \partial_n G_0 \partial_n'$$

$$(5.6c) \quad \partial_n G_0 \partial_n' \cdot \partial_n' G_0 \partial_n'' = \partial_n' G_0 \partial_n' + m^2 \partial_n'.$$

Namely, (5.6a) holds if (5.6b) does since, as a function of the right argument, both sides of (5.6a) are solutions of the Poisson equation with the same boundary value, due to (1.4), of the normal derivative at $y = 0$, and vanish at infinity. (5.6b) holds since, as a function of the left argument both sides are solutions of the Poisson equation with common boundary value, due to (1.7b), and vanish at infinity. (5.6c) follows from (5.6b) due to (1.4b) and (1.7b). Inversely, from (5.6a) (5.6b) follows. (5.6a) is actually a special case of the familiar formula for variation of G_0 by boundary variation

$$(5.6d) \quad \int \partial/\partial x(x) G_0(x', x'') = G_0(x', x) \delta/\delta x(x) G_0(x, x'')$$

with x' and x'' away from x , for smooth ∂T with $x \in \partial T$.

Using (5.6a,b) in (5.5) we have

$$\begin{aligned} (5.7) \quad & \partial_J \left[\partial_J \psi^0 \right] \psi^0(A|J) = \\ & = \exp[-P(\phi|A)] \left[-\frac{1}{2} J G_0 \partial_n \cdot \partial_n G_0 J + \right. \\ & \quad \left. + \int G_0 \partial_n \cdot \partial_n G_0 A - P'(\phi|A) \partial_J G_0 \partial_n \right\} \psi^0(A|J). \end{aligned}$$

However, from (5.4)

$$[\partial/\partial A] \Psi(A/J) = \exp[-P(\partial/\partial A)].$$

$$\cdot \left\{ -\partial_n G_0 J - \partial_n G_0 \partial_n^2 A \right\} \Psi^0(A/J).$$

and similarly for $[\partial^2/\partial A \partial A]$. Using this and (5.6c) in (5.7) and observing that the last term in the curly bracket in (5.7) contains a complete differential allows to obtain from (5.1)

$$(5.8) \quad H(A, \partial A) \Psi(A/J) =$$

$$= \int \partial z \left[-\frac{1}{2} [\partial^2/\partial A \partial A] + \frac{1}{2} \partial A \partial A + \frac{1}{2} m^2 A^2 + \right. \\ \left. + P(A) + \frac{1}{2} (\partial_n G_0 \partial_n^2 A) - \text{const} \right] \Psi(A/J)$$

$$\text{where const} = \lim_{J \rightarrow \infty} P(\partial/\partial A)(\Sigma_J) \Psi(A/J).$$

(5.8) is the expected result. We have given this combinatorial (i.e. "graphical") derivation since the actual calculations in renormalized perturbation theory, the results of which we shall present below, follow the same combinatorial pattern.

Even in the free unregularized theory, (5.8) is not usable as it stands, as seen from the meaningless subtraction term. One needs to introduce point splitting, i.e. with the first and (second to) last term as

$$\lim_{\Delta \rightarrow 0} \int dz \left[-\frac{1}{2} [\partial^2/\partial A \partial A](z, z + \Delta) \right] + \\ + \frac{1}{2} (\partial_n G_0 \partial_n^2 A)(z, z + \Delta) \right\}.$$

A local Schrödinger equation, with local Hamiltonian density $\mathcal{H}(A, \partial A)$ instead of the Hamiltonian $H(A, \partial A)$, holds for local deformation of a generally nonflat surface (cp. (5.6d)). Calculations hereto for the free theory in two-dimensional flat space with smooth ∂T are given in Appendix E of [17].

5.3. Ansatz for the interacting theory

In the regularized interacting theory, in view of (2.8), (4.7), and (4.15), (5.8) becomes

$$(5.9) \quad H(A, \partial/\partial A) = \int dz \left[-\frac{1}{2} Z_3 Z_5^{-2} (\partial^2/\partial A \partial A) + \right. \\ \left. + \frac{1}{2} Z_3^{-1} Z_5^2 \partial A \partial A + \frac{1}{2} m^2 Z_3^{-2} Z_2 Z_5^2 A^2 + \right. \\ \left. + \frac{1}{24} g_{\mu\nu} \epsilon Z_1 Z_3^{-4} Z_5^4 A^4 + \text{const} \right].$$

To render this expression meaningful for the transition $\epsilon \rightarrow 0$, we again must use point splitting and, with hindsight, add some terms that have no analog in (5.9) since they involve $\Delta = / \Delta$ / also nonlogarithmically:

$$(5.10) \quad H(A, \partial/\partial A) = \lim_{\Delta \rightarrow 0} \int dA^2 \left[-\frac{1}{2} \bar{K}_2(\Delta) (\partial^2/\partial A^2) \bar{K}_2(\Delta) + \frac{1}{2} K_2'''(\Delta) A^4 + \frac{1}{24} g K_4(\Delta) A^4 + \bar{K}(\Delta) \bar{\sigma}^2 K(\Delta) A^4 + \right. \\ \left. + \frac{1}{2} K_2'''(\Delta) \partial^2 A^2 + \frac{1}{2} K_2''(\Delta) \Delta^{-2} (\partial^2 A^2)^2 + \right. \\ \left. + \Delta^{-4} K(\Delta) + m^2 \Delta^{-2} K'(\Delta) + m^4 K''(\Delta) \right].$$

Here all K-functions are logarithmic functions of $\Delta \mu$, and the form of (5.10) is guessed in parallel to the work on renormalized field equations [18]. Comparison of (5.10) with (5.9) suggests, in parallel to the relations between (2.8) on one hand and (4.10), (4.18) on the other, the renormalization group equations, with

$$m \frac{\partial^2}{\partial g^2} + \beta(g) \frac{\partial^2}{\partial g^2} = \Delta \frac{\partial^2}{\partial A^2} + \beta(g) \frac{\partial^2}{\partial g^2} = \partial \rho$$

$$(5.11a) \quad [\partial \rho + 2\sigma(g)] (K_2, K_2') K_2''' = 0$$

$$(5.11b) \quad [\partial \rho - 2\sigma(g)] (K_2, K_2') K_2''' = 0$$

$$(5.11c) \quad [\partial \rho - 4\sigma(g)] K_4 = 0$$

$$(5.11d) \quad \partial \rho \bar{K} = 0$$

$$(5.11e) \quad \partial \rho K = 0$$

$$(5.11f) \quad \partial \rho K = 0$$

$$(5.11g) \quad [\partial \rho + \sigma(g)] K' = 0$$

$$(5.11h) \quad [\partial \rho + 2\sigma(g)] K'' = 0.$$

The $m = 0$ equations(i.e., except c,g,h) we shall verify to first order in g.

We now insert (4.20) and (5.10) into (5.1). With integrals suppressed in the notation, and setting $m = 0$ to simplify computations later on, the result ^{*} is

* Functional differentiation w.r.t. J at $x = \underline{x}y$ is indicated by a subscript x to the functional.

$$(5.12) \quad \frac{1}{2} \bar{K}_2 \partial_x \partial_y G_{x \rightarrow g} (J + A \partial_x) - \\ - \bar{K}_2 A \partial_x G_x (J + A \partial_x) [\partial_x G_2 \partial_x' C] + \\ + \frac{1}{2} \bar{K}_2 A^2 [\partial_x G_2 \partial_x' C] + \frac{1}{2} [\partial_x G_x (J + A \partial_x) \partial_x C]^2 - \\ - \bar{K} \Delta^{-1} A \partial_x G_x (J + A \partial_x) - \bar{K} A^2 \Delta^{-1} [\partial_x G_2 \partial_x C] - \\ - \Delta^{-4} K + \partial_x J G_x (J + A \partial_x) = \\ = \frac{1}{2} K_2 \partial_x A \partial_x A + \frac{1}{4} g K_4 A^4 + \frac{1}{2} K_2' \Delta^{-2} A^2 + \\ + \frac{1}{2} K_2'' \Delta^{-2} (\Delta^{-2} A)^2 + O(\Delta \ln \Delta)$$

where the last term means equality up to terms that disappear linearly (up to logarithms) as $\Delta \rightarrow 0$. Recalling that in the free-field case, from (5.6a) follows (5.6b) and from that (5.6c), we shall here first show that (5.12) holds with $A = 0$, and then consider (5.12) successively in increasing powers of A . This corresponds to putting in the $A = 0$ equation J -arguments with a normal derivative successively on the boundary in the sense of the $\nearrow \nearrow \nearrow$ limit (4.17,18). We shall see that the r.h.s. of (5.12) hereby accounts for the superficial divergences that arise in this process.

In passing only, we note that if the Schrödinger equation is generalized to curved boundary as mentioned at the end of sect. 5.2, in the Hamiltonian density corresponding to (5.10), the terms with factors Δ^{-n} , $n \geq 1$, get further contributions involving the curvature of $\partial\mathcal{T}$ explicitly, cp. (2.5) and (2.9).

5.4 $A = 0$ - equation and \bar{K}_2 determination

We rewrite (5.12) for $A = 0$

$$(5.13) \quad - \int d\omega \partial_J G_J(\omega) = \\ = \int d\omega \left[\frac{1}{2} \bar{K}_2(\omega) c_{\omega} \partial_{\omega} c_{\omega} G_2(\omega) + \frac{1}{2} \bar{K}_2(\omega) [c_{\omega}^2 G_2(\omega)]^2 - \Delta^{-4} K(\omega) \right]$$

which is the extension of (5.6a) to the interacting theory. We denote the terms in (5.13) from left to right L , R_1 , R_2 , and R_3 , R_3 is given by R_1 for $J = 0$, whereupon R_1 diverges quartically as $\Delta \rightarrow 0$. This yields K from

\bar{K}_2 . Next we decompose $G_x(J)$ in L into free-space and surface part, all J -arguments being at $y > 0$. Since the free-space part is translation invariant, to L only the surface part contributes.

For two J -arguments, (5.13) becomes, with Fourier transform taken w.r.t. the space variables,

$$(5.14) \quad (\partial_y + \partial_{y'}) \tilde{G}(1/\rho y(-\rho)) y' =$$

$$= \bar{K}_2(\omega) \left[\frac{1}{2} (2\pi)^3 \int dk e^{ikx} \tilde{G}(k(-\rho)) / \rho y(-\rho) y' \right] + \\ + \tilde{G}(-\rho/\rho y) \tilde{G}(\rho y(-\rho) y')] + O(\Delta \ln \Delta)$$

The square bracket on the r.h.s. has the form of an unamputated unrenormalized vertex, defined by point splitting. The corresponding bare vertex is $\partial_y \partial_{y'}$ or, more precisely,

$$c(y) c_{y'}(y) \partial_y c(y') c_{y'}(y') \partial_{y'} / 1 + \infty$$

according to sects. 4.2 and 4.4. The l.h.s. of (5.14) is the actual renormalized unamputated vertex. Although the superficial divergence of the vertex function is 2 (see (2.4)), $\bar{K}_2(\omega)$ is only logarithmically divergent as $\omega \rightarrow 0$: The amputated vertex requires the counter terms

$$A(\omega) \partial_y \partial_{y'} + B(\omega) (\partial_y^2 + \partial_{y'}^2) + C(\omega) \rho^2 + \\ + D(\omega) (\partial_y + \partial_{y'}) \rho + \Delta^{-1} E(\omega) (\partial_y + \partial_{y'}) + \\ + \Delta^{-1} F(\omega) \rho + \Delta^{-2} G(\omega),$$

however, of these due to the Dirichlet condition only the first term contributes, $\tilde{K}_2(\Delta)$ may, to logarithmic accuracy in Δ , be defined by the l.h.s. of (5.14) divided by the square bracket on the r.h.s., at some values of μ , y , y' or, rather, in Fourier variables w.r.t. y and y' (e.g., $\mu = 0$, $q = q' = \mu$). That the $\tilde{K}_2(\Delta)$ so defined is (to logarithmic accuracy in Δ) independent of that choice follows by an argument analogous to the one used in sect. 4.2 to ascertain the independence of $c(y)$ of the other variables in (4.11) and (4.12): One forms superficially convergent differences and then uses skeleton expansions, which will be the same on both sides of (the Fourier-transformed) (5.14), upon use of (5.13) and (5.14) to the appropriate lower order.

By Legendre transformation, one shows that in (5.13) he may restrict himself to the amputated 1PI equations in the $j^4, j^6 \dots$ case; the ambiguity in inverting the full Dirichlet propagator does not affect (in perturbation theory at least) the higher-point functions (cp. Appendix B). The 1PI j^4 equation has a logarithmic superficial divergence, which means that the amputated function requires a counter term proportional to a product of Δ -functions on the boundary. Such counter terms, however, annihilated when undoing the amputation due to the Dirichlet condition. Therefore, the j^4 equation of (5.13) is, recursively, a consequence of the j^2 equation due to skeleton expansion (see Appendix B), and so are the $j^6, j^8 \dots$ equations.

The actual computation to order g is easily done using the representation (4.5a) of sect. 4.1. One finds from (5.13)

$$(5.15) \quad \tilde{K}_2(\Delta) = 1 - (16\pi^2)^{-1} g \ln(g\Delta/2) + O(\Delta^2)$$

and verifies the μ, y, y' -independence. The logarithm in (5.15) stems from the Δ' term in

$$(5.16) \quad [4y^2 + \Delta^2]^{-1} = \frac{1}{4} \mu^2 \delta_{yy}^{-1} + \frac{1}{4} \Delta^{-2} \pi \delta(\gamma) + \frac{1}{4} \delta^{1/2}(\gamma) \ln(\mu\Delta/2) + O(\Delta/\ln\Delta)$$

with the principle value from (4.6b). The 1PI j^4 equation to (5.13) is to order g trivial, requires only (5.6a) to order g^2 , and to order g^3 requires (5.14) only to order g . Finally,

$$(5.17) \quad \langle \Delta \rangle = (2\pi^2)^{-1} + (32\pi^4)^{-1} g_1 - \frac{1}{2} + \ln 2 + O(g^2).$$

5.5 A δ -0-equations

The equations obtained from (5.13) are identities in their arguments (as $\Delta \rightarrow 0$) if all $y > 0$ and, of course, also upon differentiating w.r.t. some y . Putting undifferentiated arguments on the boundary gives (unless there is already a $c(y) \delta_{yy}|_{y=0}$ argument at the same point) zero on both sides in accordance with (4.14c). Putting a $s(y) \delta_{yy}$ -argument on the boundary gives, in general position, an A -argument as set free by differentiating (5.12) w.r.t. A .

However, that (4.14c) only holds in the sense of distributions has the effect that at coinciding such A -arguments, in general δ -type singularities and

their derivatives appear, as the only possible singularities with point-like support. While the coincidence of two (or more) λ -arguments is taken care of by the distribution character of (5.12) in A , the coincidence of the extra surface arguments on the r.h.s. with A -arguments leads to precisely those point-like singularities as arise from the polynomial terms on the r.h.s. as power counting and invariance considerations show, with K_2 , K_4 , K_2^* and K_2^{**} being logarithmic in Δ . (Note that J has support only at $y > 0$.) The \bar{K} terms on the l.h.s., stemming from the $A[\delta/\partial A]$ term in (5.10), subtract

$\Delta \rightarrow 0$ - divergences arising upon application of $[\delta/\partial A(z)]\langle K_2(\omega) \delta/\partial A(z+\epsilon)\rangle$ to expressions $1/(3!) \int \text{AAAG}(\dots, l_{xy})$ by acting on a corresponding $\int A G(\cdot/x) y$ in $\psi(A/J)$. The $A \delta/\partial A J$ term does not spoil the symmetry (as a functional differential operator) of $\text{H}(A, \delta/\partial A)$ since this operator is defined with the limit $\Delta \rightarrow 0$ performed first.

Again, with the help of (5.16) the order-g-calculations are straightforward.

The JA-equation from (5.12) yields again (5.15) and in addition

$$(5.18) \quad \bar{K}(\omega) = -(32\pi)^{-1} g + O(g^2)$$

and the AA-equation yields

$$(5.19) \quad \bar{K}_2(\omega) = 1 + (16\pi^2)^{-1} \ln(\Delta \omega^2) + O(g^2).$$

To order g , K_2^* and K_2^{**} are zero. All these results verify the pDEs (5.11) to order g .

6. Completeness and unitarity

6.1 Free field

The free-field Schrödinger functional for a Dirichlet region \mathcal{T} is, according to (1.11)

$$\begin{aligned} \phi^\circ(A/J) = & \text{const } \exp \left[\frac{i}{2} A \partial_\mu g_0 \partial_\mu A - A \partial_\mu g_0 J + \right. \\ & \left. + \frac{i}{2} J \bar{g}_0 J \right]. \end{aligned}$$

Let \mathcal{T} be divided by a surface $\partial\mathcal{T}$ into two subregions \mathcal{T}_1 and \mathcal{T}_2 , such that $\partial\mathcal{T}_{1,2} = \partial\mathcal{T}_{1,2} + \partial\mathcal{T}$. Let \bar{A} denote the common boundary value of ϕ on $\partial\mathcal{T}$. Denote the restrictions of A to $\partial\mathcal{T}_{1,2}$ by $\bar{A}_{1,2}$, and the restrictions of J to $\mathcal{T}_{1,2}$ by $J_{1,2}$. Then from composition formulae for Green's functions, which are easy to derive, in obvious notation

$$\begin{aligned} (6.1) \text{ const } \partial\bar{A} \psi_{\mathcal{T}_1}^*(\bar{A}, \bar{A}/J_{1,2}) \psi_{\mathcal{T}_2}^*(\bar{A}_2 \bar{A}'_1 J_{1,2}) = \\ = \psi_{\mathcal{T}}^*(A/J) \end{aligned}$$

follows, the \bar{A} -integral being the obvious Gaussian one. This formula is a consequence of the Markov property of the Gaussian random field involved. Upon letting \mathcal{T} be the infinite space, and choosing $\partial\mathcal{T}$ flat, a Wick rotation from the Euclidean to the Minkowskian frame can be performed, and (6.1) becomes equivalent to the ordinary completeness relation for the free-particle Fock space.

Note that if $\partial\mathcal{T}$ is infinite, the integral (6.1) will have a volume divergence which reflects Haag's theorem [19]. This can be handled by introducing a space

cutoff in such case, and extracting from the integral a, in the limit divergent, $J_{1,2}$ independent factor absorbed in the const. on the l.h.s.
We will in the following tacitly understand this device being employed where necessary.

6.2 Interacting_Fields

If Pauli-Villars regularization is introduced, (6.1) can be extended
* The functional-integration concept used in the proof [20] is the one of Friedrichs and Shapiro [21].

to fields in local nonderivative polynomial interaction, whereby if N regulator fields are used or, more directly, derivatives up to the $N + 1^{\text{st}}$ occur in the kinetic part, the $\bar{D}A$ integration must be replaced by one over \tilde{A} and the first N normal derivatives [20]. We are interested in the renormalized theory, however, where cutoffs are removed. Since hereby A undergoes only multiplicative renormalization, we expect that (6.1) can be upheld for the interacting theory. For T again the infinite space and $\bar{\mathcal{D}}T$ flat, (6.1) then becomes

$$(6.2) \text{const} \int \partial A \Psi(A/J) \Psi(A/J') =$$

$$= \langle (\exp \{ \int J \phi \})_+ \rangle$$

where the r.h.s. is the generating functional of the ordinary covariant

Schwinger functions, where

$$\mathcal{J}(x/y) = \begin{cases} J'(x(-y)) & y < 0 \\ J(x/y) & y > 0, \end{cases}$$

$$\phi(x/y) = e^{yH} \phi(x/0) e^{-yH},$$

$(\cdot)_+$ denotes y -ordering (increasing y from right to left), and H is the Hamilton operator of the Schrödinger equation. The verification of (6.2) to first order in g proceeds in parallel to the computation in Appendix D.

By letting $y \rightarrow z$ and analytic continuation from real z to $z = i$, one obtains from $\Psi^{(A/J)}$ the Minkowskian Schrödinger wave functional, in terms of the Volterra expansion in A and J if (2.15) is used. Then (6.2) becomes

$$(6.3) \text{const} \int \partial A \Psi(A/J)_{y \rightarrow it} \Psi(A/J')_{y \rightarrow it} = \\ = \langle (\exp[i \int J \phi])_+ \rangle,$$

with \int as before (writing t in place of y), and $\phi(x/t) = e^{itH} \phi(x/0) e^{-itH}$, $(\cdot)_+$ being now the usual time ordering. (6.3) expresses the completeness of the states with diagonal A . Similarly,

$$(6.4) \text{const} \int \partial A \Psi(A/J)_{y \rightarrow it} \Psi(A/J')_{y \rightarrow -it} = \\ = \langle (\exp[i \int J \phi])_+ (\exp[-i \int \phi J' J])_- \rangle$$

where $(\cdot)_-$ denotes anti-time ordering and

$$\tilde{J}(\underline{x}_t) = J(\underline{xy})|_{y=t}, \quad \tilde{j}(\underline{x}_t) = j^*(\underline{xy})|_{y=t}$$

If $J = J^*$ the r.h.s. of (6.4) is independent of J , and (6.4) expresses that

$$\left(\exp(-i\phi J) \right)_-$$

has unit norm. (6.3) and (6.4) can be verified to first order in ϵ in the same way as (6.2).

With the help of the usual asymptotic condition, one can from $\Psi(A|J)$ obtain matrix elements $\Psi(A|\text{in-state})$ and $\Psi(A|\text{out-state})$, and (6.4) becomes

the completeness relation of the diagonal states in Minkowski space.

6.3 Computation_of_expectation_values

(4.13) has the consequence that, in regularized form, (6.2) leads to

$$(6.5) \text{ const } \int dA \Psi(A|J) F(Z_5 Z_3^{-1} A) \Psi(A|J') =$$

$$= \langle (F(\phi) \exp(i\phi J))_+ \rangle$$

where $F(\phi)$ is e.g. a polynomial in time-zero smeared fields, such that the r.h.s. is finite. Since $Z_5 Z_3^{-1}$ diverges upon regularization removal (already to first order in ϵ) this factor must be absorbed, similarly as such factors were absorbed by point-splitting and split-dependent factors in the transition from (5.9) to (5.10). Here, the necessary splitting is easily seen to be one in time:

$$(6.6) \quad Z_5 Z_3^{-1} A(\underline{z}) \Psi(A|J) \rightarrow \lim_{y \rightarrow 0} [S/\partial y(z-y)] \Psi(A|J) = \\ = \Psi(A|J) \lim_{y \rightarrow 0} \sum_{n=0}^{\infty} \sum_{l=0}^{\infty} (y^n)^{-1} (y^l)^{-1}.$$

$$\int dz_1 \dots dz_n \int dx_1 dy_1 \dots dx_n dy_n J(x_1) \dots J(x_n) A(z_1) \dots A(z_n).$$

using (2.15), where this relation is meant in the sense of a replacement on the l.h.s. of (6.5). Higher powers of $Z_5 Z_3^{-1} A$ must be replaced by the expression obtained by repeated application of the functional differentiation in (6.6) in the obvious way, whereby some $Z_5 Z_3^{-1} A$ may be generated by acting on $\Psi(A|J)$ and the others by acting on $\Psi(A|J')$. We verify (6.5) with the replacement (6.6) to first order in ϵ in Appendix D.

Note that the limit $y \rightarrow 0$ in (6.6) cannot be taken under the integral sign in (6.5), since this would formally yield $\Psi(A|J) \lim_{y \rightarrow 0} y^{-1} A(y) = \infty$ according to (4.14c). (4.14c) implies, however, that the replacement (6.6) in (6.5) introduces an infinite series in A and J , the coefficients of which become the smaller the smaller y is, apart from the term linear in A which approaches a δ -function with diverging coefficient. In a "non-asymptotically-free" theory, where ultraviolet coefficients are not under control, they must, if they appear, be cancelled identically (and in perturbation theory, they always must), which makes it unlikely that then the fairly complicated limit prescription given here can be replaced by a simpler one at this stage. In an "asymptotically free" theory, however, under the usual assumptions one can "sum the logarithms up", and then the factors $Z_5 Z_3^{-1}$ in (6.5) could presumably

be removed in a simpler manner than by the replacement (6.6).

7. Discussion

7.1 Other models

- 1) The techniques of this paper are applicable to any renormalizable bosonic theory, according to the following prescription (in the free or dimensionally regularized theory): 1) Choose a first-order formulation of the field equation (i.e., in the scalar case, the Kemmer [22] representation). 2) Find that local linear transformation of the field components that corresponds to time reversal. (It is not necessary that the interaction part allows time reversal as an invariance, provided it involves no time-derivatives. This will always be so in a renormalizable theory in four dimensions.) The field components that change sign under time reversal are called Neumann ones, the others Dirichlet ones.
- 3) Construct that integral over (three) space that, if commuted with the field components, causes them to be transformed into the time reversed ones. Hereto, place the Dirichlet components on one layer and the Neumann components on an infinitesimally (in time) neighbouring one. 4) This integral, inserted in the Lagrangean at some time, implements Dirichlet/Neumann boundary conditions at that time. (The steps 1) - 4) we shall demonstrate in Appendix E in the spin 1/2 case.) 5) In order to implement inhomogeneous Dirichlet conditions, distribute the Neumann components over space, multiplied by the Dirichlet-component valued source function (analogous to A in (2.5)) and let the space integral approach the time-reversal one from the Dirichlet side as in sects. 2.2 and 4.4. 6) Go from the Minkowskian to the Euclidean frame and, if desired, replace

the plane surface by a curved one in the obvious way. Since the operator density on the surface has mass dimension three, power counting of superficial divergences is essentially the same as described in sect. 2.2. While, under interaction, the terms already present on the surface will need logarithmically divergent factors, there may also arise new ones as counter terms. (The possible counter terms are restricted by the invariance not broken by the surface interaction.)

In gauge theory, the natural gauge to choose is the obvious (time) axial one. In the Schrödinger equation, in order to preserve invariance under time-independent gauge transformations, it may be useful to introduce ordered exponentials similar to those considered by Brandt [23] in the construction of point-split renormalized field equations in QED. However, the effect of such exponentials can also be obtained [23] from polynomial terms, in the limit $\Delta \rightarrow 0$.

Computable large-momenta behaviour ("asymptotic freedom") allows to obtain the "precise" small-y behaviour of the analogs of the functions $a(y)$ and $c(y)$ of sect. 4. The leading factor is a, in general broken, power of $\ln \mu y$. Knowledge of this kind may allow in these theories to replace the complicated limit procedure of sect. 6.3 by a simpler one as remarked there.

In a fermion theory, the only new feature is that the "Dirichlet-component valued" source function is an anticommuting function, as is the ordinary space (time)-source function. Therefore, the Green's functions analogous to the ones in (2.15) are antisymmetric rather than symmetric in the two groups of arguments, whereby the components located on the boundary are Neumann ones. That the Dirichlet components of the fermion field take the anticommuting

source function as boundary "value" is expressed by the validity of the analog of (4.14c). The point-split Schrödinger operator involves, of course, the fermionic (Dirichlet) source field and functional derivatives w.r.t. it, but the corresponding c-number equations are analogous to (5.6) in the free-field case, and to (5.14) etc. in the interacting one. – For clarity, we verify the construction of the Dirichlet–Neumann surface interaction for the spin 1/2 case in Appendix E. There is an arbitrariness in which components are called Dirichlet and which Neumann ones. The implications for the spin 1/2 case of our finding that in the scalar case, in the presence of interaction we were unable to implement Neumann conditions in the strict sense (cp. end of sect. 2.2), we have not investigated.

There are models that are not renormalizable in perturbation theory but in other expansions, notably the $O(N)$ nonlinear σ -model in less than four dimensions is in $1/N$ expansion [24]. We expect that, following the prescription outlined before, homogeneous and inhomogeneous Dirichlet conditions (for the fundamental field) can be obtained, and renormalized in $1/N$ expansion.

7.2 Interaction representation

Expanding $\Psi_{(A|J)}$, as a functional of A , in terms of Hermite functionals, which are the $\Psi^0_{(A|J)}$ of (5.4b) (e.g. for one-time sources J , and which are well known [1] to form a complete orthonormalizable system in Fock space) yields the Euclidean analog of the interaction representation. This is seen simplest from (6.1) or its special case, for flat $\overline{\partial T}$, (6.2). If $\Psi_{(A|J')}$ is replaced by $\Psi^0_{(A|J')}$, in the resulting Euclidean functional integral on the r.h.s., the coupling constant is set zero in that part of four-space, such that there "free propagation" takes place.

The divergences that arise are the infrared ones (if $\overline{\partial T}$ is infinite) already mentioned in sect. 6.1, and the ultraviolet ones pointed out by Stückelberg [27] (see also [25]). Their nature was clarified by Bogoliubov and Shirkov [3]: If the coupling constant is space-time dependent, renormalization requires beyond the counter terms in (2.2), where g is to be replaced by $g(x)$, also the terms $U(g(x)) \phi(x) \partial_\mu g(x) \partial_\nu \phi(x)$ and $V(g(x)) \phi(x)^2 \partial_\mu g(x) \partial_\nu g(x)$, with U and V logarithmically divergent (e.g., in dimensional regularization, power series in ϵ^{-1}). The first counter term becomes ambiguous and the second one meaningless for $g(x)$ a step function, and neither counter term is provided for in the original definition of the interaction representation.

A similar conclusion is obtained from a discussion of the UV divergences arising in the integration in (6.2) with $\Psi_{(A|J')}$ replaced by $\Psi^0_{(A|J')}$: The only (linearly) superficially divergent surface diagrams not yet subtracted have two J , or two J' , or one J and one J' argument. The original definition of the interaction representation has no compensating terms for these, and these divergences are not related to the ones removed by Z_5 in the Schrödinger representation. – That in the Dirichlet case, the Dirichlet property suppresses divergences brought about by the sharp boundary we showed in Appendix C.

7.3 Applications

We do not expect the results of this paper to have interesting applications in the conventional renormalizable theories, e.g. QED or QCD. Rather, our starting point was the Lagrangean formulation of string theory [4] as an approximate model of Wilson loop behaviour [26]. The expectation of the Wilson

loop is approximated by the Schrödinger functional of a four-component field, in nonrenormalizable self interaction, with continuous boundary values $X_\mu(s)$ prescribed on the circumference of a two-dimensional domain. In this approximation, the QCD string tension is obtained [17, 11] from the Casimir potential *

* The connection with the Casimir effect was pointed out to the authors of [17] by Y. Nambu.

for two parallel lines,

We found in sect. 3 that the Casimir effect is finite (at least in perturbation expansion) provided the theory in the half-space is made finite (in that expansion) by appropriately chosen counter terms. The difficulty here is that, although one can make any polynomial theory "finite" in perturbation expansion by counter terms [3, 27], this expansion is most likely meaningless for a non-renormalizable theory since strong arguments have been given [28] that a non-renormalizable theory has, if it makes sense at all, in its correct perturbation expansion also terms involving the logarithm of the coupling constant.

A systematic construction of such "improved" perturbation expansion succeeds so far only in cases where it can be derived from an alternative expansion with stable power counting, e.g. from the 1/N expansion in the nonlinear σ -model in less than four dimensions [24]. Of course, with such well-behaved expansion at hand, one attempt to apply the considerations of this paper directly, since e.g. the mentioned 1/N expansion is again one in terms of graphs.

Therefore, the prerequisite for an application of the methods of this paper to the string Lagrangeans and similar models for extended structures is to find expansions in which these models become renormalizable in infinite space. So far, attempts by the author to find such expansions for the Nambu and Eguchi Lagrangeans [4, 7] have failed.

8. Conclusions

We have shown that, in every renormalizable theory, a) the Schrödinger representation exists, b) in this representation, a Schrödinger equation (with point splitting as already needed in the free theory) holds, c) the field operator that is being diagonalized is not the renormalized (nor the unrenormalized) one, but differs from it by a factor that diverges logarithmically if the distance from the boundary (which, if nonzero, acts like a cutoff) goes to zero, d) this last feature requires a limit process to be employed in the calculation of expectation values, e) the Casimir effect, for disjoint surfaces, is computable to all orders in renormalized perturbation theory.

For simplicity, we gave details only for the Φ_4^4 theory, but we explained the principles of the extension to other models, in particular, also those with fermions. Our reasoning was heuristic at times, and explicit calculations were given only for the first (already nontrivial), and in one case second order in the perturbation expansion. The author is convinced, however, that the conclusions hold to all orders.

Our motivation was the intended application to string-Lagrangean models of the Wilson loop. Unfortunately, this application requires first to find a

Appendices

renormalizable expansion for such models in infinite space, which has not yet been found. Meanwhile, we recall that Dirac regretted [29] the lack of a Schrödinger equation in quantum electrodynamics.

Acknowledgment

The author is indebted to M. Lüscher and T.T. Wu for discussions.

The Gaussian integral (1.3) is evaluated using the ϕ field equation, which becomes a linear integral equation for the correlation function. Herby we must distinguish whether, seen from \mathcal{T} , an argument goes to $\partial\mathcal{T}$ first, like ϕ , or second, like $\partial_{\mathcal{T}}\phi$, and we denote this order by a subscript to a vertical bar which means putting that argument on $\partial\mathcal{T}$: $|_1$ means approach from \mathcal{T}' and $|_2$ approach from \mathcal{T} . It is instructive to give to the boundary interaction in (1.2) the general coefficient c rather than 1. Denoting by G the free-space Green's function

$$(A.1) \quad G(x-x') = (2\pi)^{-\nu} \int d^{\nu}p e^{ip(x-x')} (p^2 + m^2)^{-1},$$
$$= \frac{1}{4} \pi^{-\nu/2} \mathcal{T}(\frac{1}{2}\nu - 1) / |x-x'|^{-\nu+2} \text{ if } m=0$$

and by G_c the correlation function with $c \neq 0$, we find

$$(A.2) \quad G_c = G + c G \delta_n |_1 \cdot |_1 / G_c + c G \cdot |_2 / \partial_n G_c$$

where the dot means integration over $\partial\mathcal{T}$. Herefrom

$$(A.3a) \quad g/G_c = |G + c G \delta_n |_2 \cdot |_1 / G_c + c G \cdot |_1 / \partial_n G_c$$

and

$$(A.3b) \quad g/G_c = |G + c G \delta_n |_2 \cdot |_1 / G_c + c G \cdot |_1 / \partial_n G_c$$

and similarly

$$(A.4a) \quad \frac{1}{\partial_n} G_c = \frac{1}{\partial_n} G + c \frac{1}{\partial_n} G \delta_{n1} \cdot_1 \frac{1}{\partial_n} G_c + \\ + c \frac{1}{\partial_n} G /_1 \cdot_2 \frac{1}{\partial_n} G_c$$

and

$$(A.4b) \quad \frac{1}{\partial_n} G_c = \frac{1}{\partial_n} G + c \frac{1}{\partial_n} G \delta_{n1} \cdot_1 \frac{1}{\partial_n} G_c + \\ + c \frac{1}{\partial_n} G /_2 \cdot_2 \frac{1}{\partial_n} G_c$$

due to

$$(A.5a) \quad \frac{1}{\partial_n} G /_1 = \frac{1}{\partial_n} G /_2, \quad \frac{1}{\partial_n} G \delta_{n1} /_1 = \frac{1}{\partial_n} G \delta_{n1} /_2.$$

We now note the discontinuity relation

$$(A.5b) \quad \frac{1}{\partial_n} G /_2 = \frac{1}{\partial_n} I + \overline{\frac{1}{\partial_n} G}$$

where I is the δ -function on ∂T , and $\overline{\frac{1}{\partial_n} G}$, defined by this equation, is an integral kernel on ∂T well known in potential theory [30], of form

$$(A.6) \quad \overline{\frac{1}{\partial_n} G} = -\frac{1}{4} \pi^{-\nu/2} T(\frac{1}{2}\nu) / R^{-1} / x - x' / -\nu + 2 +$$

+ less singular terms

where R^{-1} is the signed curvature of ∂T along $x-x'$. $\frac{1}{\partial_n} G$ vanishes on flat portions of ∂T , and for smooth compact ∂T , $\frac{1}{\partial_n} G$ is a Fredholm kernel if $\nu \neq 2k$. Using (A.5b) in (A.3a,b) gives

$$(A.7a) \quad \frac{1}{\partial_n} G_c = (1-c) \frac{1}{\partial_n} G_c$$

and from (A.4a,b)

$$(A.7b) \quad (1-c) \frac{1}{\partial_n} G_c = \frac{1}{\partial_n} G_c.$$

These homogeneous jump relations, where the right argument may be in T or T' , completely characterize the effect of the surface interaction. With the help of Green's formula, (A.7a,b) lead back to (A.2).

To solve (A.2), we insert (A.7a) and obtain

$$(A.8) \quad G_c = G + (1-c)^{-1} c G \delta_{n1} /_1 \cdot_2 \frac{1}{\partial_n} G_c + c G /_2 / \partial_n G_c.$$

If both arguments are in T , Green's formula gives

$$(A.9) \quad G_c = G - G \delta_{n1} /_1 \cdot_2 \frac{1}{\partial_n} G_c + G /_2 / \partial_n G_c$$

Herefrom and from (A.8)

$$(A.10a) \quad G_c = G + (2c-c^2) G /_2 / \partial_n G_c$$

and

$$(A.10b) \quad G_c = G + (1-c)^{-2} c (2-c) G \overset{\leftrightarrow}{\partial}_n / \cdot \cdot_2 / G_c$$

The solution of (A.10a), using (A.5b), is

$$(A.11) \quad (\mathcal{T}'\mathcal{T}): \quad G_c = G + 2f(c) G / \cdot [1-2f(c)] \overset{\leftrightarrow}{\partial}_n G J^{-1} \cdot / \partial_n G$$

where

$$(A.12) \quad f(c) = [1 + (1-c)^2]^{-1} [1 - (1-c)^2].$$

Similarly, (A.10b) yields the transposed of (A.11), which verifies the symmetry of G_c (which also follows from known properties of the functions on the r.h.s. of (A.11)).

In potential theory, the convergence of the iteration solution of the inverse in (A.11) is proven for $|f(c)| < 1$. G_1 is in ($\mathcal{T}'\mathcal{T}$) the Dirichlet function and $G \neq \infty$ the Neumann function, as follows already from (A.7).

For the left argument in \mathcal{T}' , the right argument in \mathcal{T} , (A.9) is replaced by

$$0 = G - G \overset{\leftrightarrow}{\partial}_n / \cdot \cdot_2 / G_c + G / \cdot \cdot_2 / \partial_n G_c = \\ = G_c - G \overset{\leftrightarrow}{\partial}_n / \cdot \cdot_1 / G_c + G / \cdot \cdot_1 / \partial_n G_c.$$

Herefrom, and from (A.7), we obtain the two forms

$$(A.16) \quad (\mathcal{T}'\mathcal{T}): \quad (2/\partial_n) G_c = 2(1-c)^{-1} G_c / \cdot_2 \cdot \cdot_2 / \partial_n G_c = \\ = (1-c)^{-1} [G_c / \cdot_2 \cdot \cdot_2 / \partial_n G_c + G \overset{\leftrightarrow}{\partial}_n / \cdot_2 \cdot \cdot_2 / G_c]$$

$$(A.13) \quad (\mathcal{T}'\mathcal{T}'): \quad G_c = \\ = (1-c)^{-1} \{ G + 2f(c) G \overset{\leftrightarrow}{\partial}_n / \cdot [1-2f(c)] \overset{\leftrightarrow}{\partial}_n G J^{-1} \cdot / \partial_n G \} \\ = (1-c)^{-1} \{ G + 2f(c) G / \cdot [1-2f(c)] \overset{\leftrightarrow}{\partial}_n G J^{-1} \cdot / \partial_n G \}$$

which show that in this case, $G_c \geq 0$ if and only if $c = 1$ or $c = \pm\infty$.

Finally, with both arguments in \mathcal{T}' , one derives in a similar way

$$(A.14) \quad (\mathcal{T}'\mathcal{T}'): \quad G_c = G + 2f(c) G / \cdot [1-2f(c)] \overset{\leftrightarrow}{\partial}_n G J^{-1} \cdot / \partial_n G$$

which, upon comparison with (A.11) and noting the direction of the normal, shows that effectively a change of sign of $f(c)$ occurred, i.e. $c \rightarrow c' = \frac{c}{c-1}$ as also obtainable directly from (A.2) and (A.7). Thus, G_1 is in ($\mathcal{T}'\mathcal{T}'$) the Neumann and $G \neq \infty$ the Dirichlet function. This proves (1.3).

Interchanging in (1.2) the two layers, i.e. posing the ϕ -layer inside the $\partial_n \phi_{\text{one}}$, yields the correlation function G_c' which obeys

$$(A.15) \quad G_c' = G_c / (1+c).$$

Thus, the order of the two layers does matter.

From (A.11) one easily derives by differentiation

which due to (A.7a) agrees with the general formula obtained from (1.3)

$$(\partial/\partial c) G_c = G_c \partial_n /_2 + 1/G_c + G_{c1} 1 \cdot \partial_n G_c.$$

It is obvious that in all formulae, c could be taken to be a function of the point on $\partial\mathcal{T}$. All these boundary conditions have in common that they can be implemented by a bilinear interaction local on $\partial\mathcal{T}$, with dimensionless coefficient.

B. Perturbation expansion

The derivation of the action density (2.8), which provides all the possibly needed counter terms, also prescribes how to compute,

First, we set $A = 0$. For a graph, one notes the 1PI parts as usual. Keeping full Dirichlet, Neumann, or zero lines (cp. (1.3)) for the connecting links and external legs, one decomposes the other propagators into free-space parts and surface parts as described in sect. 2.1. This yields free-space and surface 1PI parts. One must now revoke the one-particle-reducibility in any chain of

$E = 2$ subgraphs if the two subgraphs at the ends are surface ones. For interpretation of this prescription one visualizes in these cases $\partial\mathcal{T}$ (i.e. the plane $y = 0$) as analogous to one line of a covariant ϕ^4 graph: If two $E = 2$ surface graphs are connected by one (possibly covariantly corrected) line "and by the surface", they are connected one-particle-irreducibly. Thus, a chain of $E = 2$ surface graphs, each one 1PI in the usual sense, must be computed like a covariant four-point vertex function that is the same number of times two-particle-reducible in some channel. This means that subtractions for each

possible sub-four-point vertex must be made to treat the overlapping divergences correctly. One possible way to do that is by suitably arranged subtractions on a Bethe-Salpeter equation, see e.g. [16]. In the present case, in each step the divergence is linear rather than logarithmic which requires to make two subtractions rather than one.

Consider, e.g. an unsubtracted two-point surface vertex $\mathcal{T}_{(xyx'y')}$ 1PI in the usual sense. Its Fourier transform $\tilde{\mathcal{T}}(\underline{k}_y(-\underline{k})y')$ is in general, if $\epsilon = 0$, not a distribution in y or y' , however,

$$(B.1) \lim_{\epsilon \rightarrow 0} [\tilde{\mathcal{T}}_{\epsilon,0}(\underline{k}_y(-\underline{k})y') - \\ - A(\epsilon) (\delta(y)\delta'(y') + \delta(y)\delta(y'))] \equiv \tilde{\mathcal{T}}_{\text{ren}}(\underline{k}_y(-\underline{k})y')$$

is, with the subtraction provided by a part of $Z_4 - 1$ in (2.8). If to the expression in the square bracket two Dirichlet or two Neumann lines are attached, with the other argument of that line strictly positive, the subtraction term is annihilated. This means that in this case, the renormalized unamputated two-point function is insensitive to the subtraction convention for the amputated function, which determines the finite part of $A(\epsilon)$.

Let $\tilde{\mathcal{T}}_{\text{sub}}(\underline{k}_q(-\underline{k})q')$ be the Fourier transform of the square bracket in (B.1) w.r.t. y and y' . If, with (1.18)

$$2\pi i^{-2} \int dq'' \delta(q'' - \underline{k}_q(-\underline{k})q'') G(k, q'' q'').$$

$$\cdot \tilde{\mathcal{T}}_{\text{sub}}(\underline{k}_q''(-\underline{k})q')$$

is formed, as $\epsilon \rightarrow 0$ its singular part has the form $-ia'(\epsilon)(q+q')$, and if

$\underline{k} = 0$, needs the ordinary free-space subtraction. The general two-point loop, with factor $1/2$ included, is

$$\begin{aligned}
 (C.1) \quad & \frac{1}{32} \pi^{-4} \epsilon \mu \epsilon \int d^4x e^{ik \cdot x} \left\{ \left[x^2 + (z_1 - z_2)^2 \right]^{-1+\epsilon/2} \right. \\
 & - \left. \left[x^2 + (z_1 + z_2)^2 \right]^{-1+\epsilon/2} \right\}^2 = \\
 & = (32\pi^2)^{-1} \left\{ e^{-K/z_1 - z_2} / P_\mu/z_1 - z_2 /^{-1} - \right. \\
 & - (K z_1 z_2)^{-1} \left(e^{-K/z_1 - z_2} - e^{-K(z_1 + z_2)} \right) + \\
 & + (z_1 + z_2)^{-1} e^{-K(z_1 + z_2)} + \\
 & \left. + \delta(z_1 - z_2) \left[\epsilon^{-1} + \ln 2 + 1 + \frac{1}{2} \ln \pi + \frac{1}{2} \psi(1) \right] \right\} + O(\epsilon)
 \end{aligned}$$

where P_μ denotes the two-sided principal value,

$$(C.2) \quad P_\mu/z/^{-1} = \int \mu \left(\mu/z/ \right)^{-1} + 2(z-1)^{-1} \delta(z) /_{z=1}$$

denoted by $\mu(z)$ by Gelfand and Shilow [14]. The part prop. ϵ^{-1} is absorbed by the usual covariant coupling constant renormalization. The integration in B, with $\underline{k} = 0$, yields a term prop. z_1^{-2} and one prop. $z_1^{-2} \ln(z/\mu)$ due to the principal value in (C.1). The remaining z_1 integration is possible since, for $y_1 > z_1$, $y_2 > z_1$, the external-leg Dirichlet functions give rise to a factor z_1^2 .

Graph C yields a sum of nine double integrals corresponding to the possible time orderings, all of them unproblematic.

A contains the free-space-propagator part

$$\begin{aligned}
 & \frac{1}{6} 2^{-6} \pi^{-6+3\epsilon/2} \mu^{2\epsilon} \int d^4x e^{ik \cdot x} \cdot \\
 & \cdot \left[x^2 + (z_1 - z_2)^2 \right]^{-3+3\epsilon/2} = \\
 & = \frac{1}{3} \mu^{2\epsilon} T^{-1/(3-3\epsilon/2)} - 1 / \left(\frac{3}{2} - \epsilon \right) 2^{-\epsilon} \pi^{\epsilon-9/2} \\
 & \int (z_1 - z_2)^{-3+2\epsilon} - \frac{1}{2} K^2 (1-2\epsilon)^{-1} / z_1 - z_2 /^{-7+2\epsilon} + \\
 & + \text{non sing.}
 \end{aligned}$$

Use herein of (C.2) and [14]

$$(C.3) \quad P_\mu/z/^{-3} = \int \mu^3 \left(\mu/z/ \right)^{-2} + (z-3)^{-1} \delta''(z) /_{z=3}$$

gives the $\epsilon \rightarrow 0$ singular part

$$(C.4) \quad (3/2)^{10} \pi^{4\epsilon} T^{-1} \left[\delta''(z_1 - z_2) / -K^2 \delta(z_1 - z_2) \right]$$

which is absorbed by the covariant amplitude renormalization using (2.10).

(The remaining finite part is Euclidean invariant, which verifies explicitly to this order the absence of a renormalization of the speed of light, cp. sect. 2.1.) The term in A with two free-space propagators and one surface propagator involves the loop (C.1), the $1/\epsilon$ term of which is again absorbed

it is removed the limit $\epsilon \rightarrow 0$ exists. More generally, one subtracts from integrals of this kind the Taylor expansion to first order in \underline{k} , \underline{q} , \underline{q}' at $\underline{k} = \underline{q} = \underline{q}' = 0$, and adds $i\epsilon(\underline{q}+\underline{q}')$ with some c to satisfy the renormalization condition. The unamputated two-point function on the Dirichlet side (cp. below) is independent of the choice of that condition.

Skeleton expansions of 1PI $E \geq 4$ graphs are obtained as follows: One first identifies 1PI $E \leq 4$ subgraphs as usual. The one-particle links between these are again left as Dirichlet, Neumann, or zero propagators. The other propagators are decomposed into free-space and surface parts, and 1PI $E = 2$ surface subgraphs are identified as described before. $E = 4$ surface subgraphs are then further decomposed until the original graph is obtained as a sum of graphs with 1PI $\mathcal{G} \geq 0$ subgraphs connected by Dirichlet, Neumann, or zero propagators. For these subgraphs the appropriate subtractions must be made.

A consequence of this construction is: While the convolution inverse of the full two-point function is not unique since the Dirichlet propagator vanishes on the boundary, the amputated 4-point, 6-point etc. functions can be defined (in perturbation theory) directly in terms of graphs contributing to them. This we shall make use of in sect. 5.

Consider now $A \neq 0$ in (2.8), which gives rise to one-leg surface vertices. If one leg of a two-point function is attached to such A-vertex, the graph is a surface one apart from the other-end propagator and a possible free-space propagator-correction there. The surface graph must be computed as in the covariant theory an ordinary (e.g. mass) vertex is, with all possible subtractions (cp. (2.6)) to linearly divergent subgraphs, of which the sub-vertex is a

prominent one. The subtraction convention for this subgraph does matter, and possible choices would be $\frac{\partial}{\partial q^2} \mathcal{T}(k^2)/_{q=0}, \frac{\partial}{\partial q^2} \omega_{\mu\nu} = \epsilon$ or minimal Z_5 as in (2.13). The relation between Green's functions with and without A-arguments is discussed in sect. 4.2.

In the description so far, both the Dirichlet and the Neumann regions were used. If the graphs from all splittings of the lines into free-space and surface ones are added up, the two regions again decouple. The line separation was needed only to show the sufficiency for finiteness of the counter terms in (2.8). The actual computation can, conveniently in $\underline{k}\text{-space}$, be carried out using either the Dirichlet or the Neumann side alone, with the latter possessing no A-vertices. On the Dirichlet side, power counting shows that the Dirichlet condition is satisfied for each unamputated two-point function, and this renders all unamputated Green's functions (and also all vacuum graphs, see sect. 3.3) insensitive to the subtraction prescription that fixes $Z_4 = 1$. In fact, in computing these functions, the subtraction terms in (2.5) are inactive due to preservation of the Dirichlet condition in every step. To illustrate the role of the Dirichlet condition we discuss in Appendix C the computation to second order of the two-point function in some detail.

The circumstances on the Neumann side are described at the end of sect. 2.2.

C. Two-point function and Dirichlet condition

The graphs for the two-point function to second order are shown in Fig. 1. In the one-vertex loops in B and C, only the surface propagator, with result prop. $z^{2+\epsilon}$, is to be used since the free-space one is absorbed in the covariant mass renormalization. The two-point graph in B, which transmits

by covariant coupling constant renormalization.

Collecting the finite parts we find, after some calculation,

$$\begin{aligned}
 (C.5) \text{ F.P. of } A &= (3.2^2 \pi^4)^{-1} \rho_\mu [e^{-\kappa s} / (\Delta^{-3} / (\tau + \kappa \Delta)) - \\
 &\quad - 12 \Delta^{-1} (s^2 \Delta^2)^{-1} + 48 \kappa^{-1} (s^2 \Delta^2)^{-2} - (\Delta \leftrightarrow s)] = \\
 &= \rho_\mu (3.2^2 \pi^4)^{-1} (s - \Delta)^3 \left\{ \Delta^{-3} s^{-3} (\Delta + s)^{-2} (s^2 + \kappa s \Delta + \Delta^2) - \right. \\
 &\quad \left. - \frac{1}{2} \kappa^2 \Delta^{-1} s^{-1} (\Delta + s)^{-2} + O(\kappa^4) \right\}
 \end{aligned}$$

$$\begin{aligned}
 &- 12 \Delta^{-1} (s^2 \Delta^2)^{-1} + 48 \kappa^{-1} (s^2 \Delta^2)^{-2} - (\Delta \leftrightarrow s)] = \\
 &= \rho_\mu (3.2^2 \pi^4)^{-1} (s - \Delta)^3 \left\{ \Delta^{-3} s^{-3} (\Delta + s)^{-2} (s^2 + \kappa s \Delta + \Delta^2) - \right.
 \end{aligned}$$

$$- \frac{1}{2} \kappa^2 \Delta^{-1} s^{-1} (\Delta + s)^{-2} + O(\kappa^4) \Big\}$$

where $\Delta = i(z_1 - z_2)$ and $s = z_1 + z_2$. ρ_μ prescribes the two-sided principal value w.r.t. Δ . of (C.2,3). (C.5) gives in A for $y_1 > 0$, $y_2 > 0$ a well defined integral, due to the factor $z_1 z_2$ from the two Dirichlet lines if $y_1 > z_1$, $y_2 > z_2$:

$$(C.6) \quad \frac{1}{4} \kappa^{-2} \int_{z_1}^{\infty} dz_2 \left[e^{-\kappa/y_2} - e^{-\kappa(y_2 + z_2)} \right] \rho_\mu (z_1 - z_2) / \tau^{-3} =$$

$$\begin{aligned}
 &\left[e^{-\kappa/y_2} - e^{-\kappa(y_2 + z_2)} \right] \rho^2 / (\kappa (\mu/\mu^{-1}) - \psi(3)) \Big]. \\
 &\cdot (e^{i\theta y_1} - e^{-\kappa y_1}) (e^{-i\theta y_2} - e^{-\kappa y_2})
 \end{aligned}$$

where we used the Fourier representation of the principal value [14] and

$$\begin{aligned}
 \frac{1}{2} \kappa^{-1} \int_0^\infty dz [e^{-\kappa/y_1 - z_1} - e^{-\kappa(y_1 + z)}] e^{i\theta z} = \\
 = (\kappa^2 + \mu^2)^{-1} (e^{i\theta y_1} - e^{-\kappa y_1}).
 \end{aligned}$$

The p-integral in (C.6) is absolutely convergent, and vanishes if $y_1 > 0$ or $y_2 > 0$. In higher orders in perturbation theory, there appears in (C.6) $\rho_\mu (ln/z_1 - z_2) / \tau^{-3}$ which also gives a well-defined integral. - Also the formula

$$\begin{aligned}
 (C.7) \quad &\int_0^\infty dz_1 z_1 e^{iy_1 z_1} \int_0^\infty dz_2 z_2 e^{-iy_2 z_2} \rho_\mu (z_1 - z_2) / \tau^{-3} \\
 &= \int -2 \psi(3) + \ln[\mu^{-2} (y_1 + i\theta)(y_2 + i\theta)] \Big].
 \end{aligned}$$

is instructive; for convergence again the factor $z_1 z_2$ is needed.

In the interaction representation, as discussed in sect. 7.2, the integral (C.6) appears but with the second terms in the square brackets missing since only the interaction, not also the kinetic part is switched off at the boundary (see also [25]). Then, that the principal value is integrable only with a C^2 test function leads to a divergence.

D. Completeness check to first order

The Schrödinger functional (4.20) we write as

$$(D.1) \quad \psi(A|J) = \exp \left[-\frac{1}{2} A K A - A \partial_n G J + \frac{1}{2} J G_D J + P(A) + Q(A|J) + R(J) \right]$$

where the terms in the first line are the zero-order ones (cp. (5.4b)) and the (connected) $Q(A|J)$ shall have no term depending only on A or on J . For brevity, however, we will set $J = 0$, i.e. consider in (6.5) the vacuum expectation values, since the calculation with $J = 0$ offers no new feature. To first order,

$$(D.2) \quad P(A) = -\frac{1}{24} g^2 \int dx [A \partial_n G_D(x)]^4 + + \frac{1}{2} g (32\pi^2)^{-1} \int dx \int dy' [C A \partial_n G_D(x) C(y')]^2 \mu^2 \rho(\mu) r^{-2}$$

according to (4.5b).

By Gaussian integration, we find

$$(D.3) \quad \begin{aligned} & \text{const} \int \partial A F(A) \psi(A|0) \psi(A|0) = \\ & \approx \text{const}' F(C/\delta B) \exp [2P(C/\delta B)] \exp [\frac{1}{4} B K^{-1} B] \Big|_{B=0} \\ & \cdot \exp [S(\frac{1}{4} L \partial A) K^{-1} (\partial A) - S(0)]. \\ & \cdot \exp \left(\frac{1}{4} L \partial A \right) K^{-1} (\partial A) F(A) \Big|_{A=0} \end{aligned}$$

where we introduced

$$(D.4) \quad \begin{aligned} & \exp \left(\frac{1}{4} L \partial A \right) K^{-1} (\partial A) \exp \vartheta P(A+B) \Big|_{A=0} = \\ & =: e^{\chi \vartheta} S(CB). \end{aligned}$$

Like P , Q , and R , S is connected due to the linked-cluster theorem. In (D.3), $\exp(S(0))$ is the infrared (by Haag's theorem) and ultraviolet divergent factor absorbed (see sect. 6.1) in the const. in (6.2) and (D.3).

We choose $F(A) = A(z_1)A(z_2)$, which suffices to display all difficulties. The correct two-point vacuum expectation value to first order is the free one, $\frac{1}{2} K_{12}^{-1} = \frac{1}{2} \text{F.T.} \int k f^{-1}$. From (D.3), however, we find a divergent result, formally (or, under regularization) proportional to K_{12}^{-1} , since it involves the last term in (4.5b) minus the same integral with the principal-value sign omitted. This second term stems from the first term on the r.h.s. of (D.2) and the evaluation

$$(D.5) \quad \begin{aligned} & \frac{1}{4} \left[\partial A \partial A \right] K^{-1} (\partial A) \frac{1}{2} \int (A \partial_n G_D(x) r)^2 \Big|_{A=0} = \\ & = \frac{1}{4} (2\pi)^{-3} \int dk K^{-1} e^{-2kr} = (32\pi^2 r^2)^{-1}. \end{aligned}$$

This remaining divergence is in accord with (6.5), where Z_5 diverges to first order, see (2.13).

$$\text{The prescription of sect. 6.3 gives, for approach of both operators in } F(A)$$

to $y = 0$ from the same side

$$(D.6) \quad F(z_s z_s^{-1} A) \rightarrow F(A, \gamma_1 \gamma_2) = \\ = \left\{ - (A \partial_n G_0)(z_1 \gamma_1) - \right. \\ \left. - g (32\pi^2)^{-1} \int dx \int dy' (A \partial_n G_0)(x y') G_0(x y', z_1 \gamma_1) \right. \\ \left. + \frac{g^2}{6} \rho(y y')^{-2} + \frac{1}{6} g \int dx \int dy' [(A \partial_n G_0)(x y')]^3 G_0(x y', z_1 \gamma_1) \right\}.$$

of same expression with $z_1 \rightarrow z_2, \gamma_1 \rightarrow \gamma_2$

$$+ G_0(z_1 \gamma_1, z_2 \gamma_2) + g \int dx \int dy' \left\{ (32\pi^2 y'^2)^{-1} - \right. \\ \left. - \frac{1}{2} [(A \partial_n G_0)(x y')]^2 G_0(x y', z_2 \gamma_2) G_0(x y', z_2 \gamma_2) \right\}$$

where only the terms up to first order in g are to be kept, and (4.5a) has been used. At fixed function A , the limit $y_1 \rightarrow 0, y_2 \rightarrow 0$ does not exist on the r.h.s. of (D.6), the factor $z_5^{-2} z_3^2$ missing. The A -independent terms, however, vanish in this limit.

Working out (D.3), with insertion of (D.6) and (D.2), term by term is trivial, the characteristic integration being again (D.5). Divergences do appear, however, in the y' integrations in separate terms. They are made unambiguous by dimensional regularization, whereby it is advantageous to replace the principal values on the r.h.s. of (D.6) by (4.6b). Then, all divergences are found to cancel, and

the sum of finite terms to order g vanishes linearly, as $y_1 \rightarrow 0, y_2 \rightarrow 0$, as it should.

The same result is obtained if in (6.5), $A(z_1)$ and $A(z_2)$ are shifted into different factors $\psi(A|0)$ in the sense (6.6). Some fewer terms then appear, but otherwise the calculations are identical to the ones just described.

E. Dirichlet conditions for the Majorana field

The prescription given in sect. 7.1 is simplest to illustrate for the spin $1/2$ Majorana field, wherefrom the transition to the Dirac field is obvious. The Majorana Lagrangian with surface term is

$$(E.1a) \quad L = \frac{1}{2} i \bar{\psi} \alpha^\mu \partial_\mu \psi - \frac{1}{2} m \bar{\psi} \beta \psi + i \bar{\psi} \gamma^\mu \psi \gamma^\nu \partial_\nu \psi$$

where, with two independent sets σ_i and τ_i of Pauli matrices, we may choose

$$(E.1b) \quad \alpha^0 = 1, \quad \alpha^1 = \sigma_3 \tau_1, \quad \alpha^2 = \sigma_3 \tau_3, \quad \alpha^3 = \sigma_1, \\ \beta = \sigma_2, \quad \psi = \psi^+$$

In (E.1a), χ is an as yet unspecified matrix, and subscripts 1 and 2 indicate approach to the time-zero plane from positive and negative times, respectively.

From the field equations the integral equations for the Feynman Green's functions follow,

$$(E.2) \quad G = G^0 + G^0 \cdot x_2 G + G^0 \cdot x^\tau G = \\ = G^0 - G_1 \cdot x G^0 + G_2 \cdot x^\tau G^0$$

where

$$(E.3) \quad G^0 = -i \int [(\omega - i0)/\beta - \cos \alpha d\omega]^{-1}$$

and the dot indicates integration on the time-zero plane. The integral operators on the boundary

$$(E.4a) \quad {}_1 G_2^0 = :P_+ , \quad - {}_2 G_1^0 = :P_-$$

obey

$$(E.4b) \quad P_+ \cdot P_+ = P_+ , \quad P_- \cdot P_- = P_- \\ P_+ \cdot P_- = P_- \cdot P_+ = 0 , \quad P_+ + P_- = 1.$$

Going in (E.3) to the boundary and using (E.4) yields

$$(E.5) \quad (1-x^\tau)_1 G = (1-x)_2 G , \quad G_1(1-x) = G_2(1-x^\tau).$$

This allows to rewrite (E.2) as

$$(E.6) \quad G = G^0 \cdot {}_2 G + G^0 \cdot {}_1 G = G - G_1 \cdot G^0 + G_2 \cdot G^0$$

and herefrom

$$(E.7) \quad G = G^0 + G^0 \cdot X \cdot G^0 , \quad X = {}_{11} G_2 - {}_{11} G_1 - {}_{22} G_1 + {}_{22} G_2$$

where easily follows. The subscript 11 indicates that the argument goes to the boundary from the 1 direction but later than the ones denoted by 1 or 2. The consequences of (E.2) or (E.7)

$${}_{11} G_1 - {}_{11} G_2 = 1 = - {}_{22} G_2 + {}_{22} G_2$$

are also consequences of the canonical anticommutation relations.

We set

$$(E.8) \quad \chi = S + A , \quad S = S^T , \quad A = -A^T .$$

With the notation $P_+ - P_- = Q$, (E.5) is, using (E.7), equivalent to

$$(E.9) \quad (1 - S A + A Q) \cdot \chi = -2 A \chi = \chi \cdot (1 - S A + Q A) .$$

Consistency requires

$$(E.10) \quad [S, A] = 0 = [\chi, \chi^\tau]$$

which, apparently, is the condition that the symmetry property of the Green's function, expressed by the simultaneous validity of the two integral equations in (E.2), is not in contradiction to the Lagrangean (E.1a) which, in general, is not hermitean. The solution of (E.10) $A = 0$ yields $X = 0$ for generic S ,

and by continuity also if S has eigenvalue 1. This corresponds to the surface term in (E.1a) being ineffective.

For the Green's function to vanish if the two arguments are on opposite sides of the boundary,

$$(E.11a) \quad G_2 = 0 = \frac{G}{2} \Big|_1$$

is necessary, which with (E.7) becomes

$$(E.11b) \quad P_+ \cdot X \cdot P_+ = -P_+, \quad P_- \cdot X \cdot P_- = P_-.$$

That (E.11b) is also sufficient follows from the frequency property

$$G_1^0 \cdot P_- = \langle_0 G_1 \cdot P_+ = P_+ \cdot G_0^0 = P_- \cdot G_0^0 = 0.$$

With (E.9), if $m > 0$ and $S = 0$, the only solutions of (E.11b) are

$$(E.12) \quad A = \pm i\sigma_3 \tau_2 = \pm i\mathcal{J}^0 \mathcal{J}^0, \quad X = -A\mathbf{I} - Q$$

as one finds from special choices of momenta. The time reflection matrix here obeys

$$AP_+ = P_- A, \quad P_+ A = AP_-.$$

(If $m = 0$, there is the family of solutions

$$\begin{aligned} S &= 0, \quad A = e^{i\omega\sigma_3\tau_2} \sigma_3 \tau_2 e^{-i\omega\sigma_3\tau_2} = \\ &= e^{i\mathcal{J}^0 \mathcal{J}^0} e^{-i\mathcal{J}^0 \mathcal{J}^0}, \quad \text{arbitrary.} \end{aligned}$$

The boundary conditions (E.5) become

$$\begin{aligned} (E.13) \quad (1 \pm \sigma_3 \tau_2) \Big|_1 G_{>0} = 0 &= \rangle_0 G_1 (1 \mp \sigma_3 \tau_2) \\ (1 \mp \sigma_3 \tau_2) \Big|_2 G_{<0} = 0 &= \langle_0 G_2 (1 \pm \sigma_3 \tau_2) \end{aligned}$$

for the two choices in (E.12).

For the solutions given by (E.12), in (E.1a)^{the "1"} and the "2"^{the "2"} components in the surface term do not anticommute among themselves. Comparison with (1.2) suggests to choose χ such that the two groups of components do anticommute among themselves. This requires

$$(E.14) \quad \chi^\tau \chi = 0 = \chi \chi^\tau$$

The only solution of (E.11b) with (E.9), (E.14) is, if $m > 0$,

$$(E.15) \quad S = \frac{1}{2}, \quad A = \pm \frac{1}{2} \sigma_3 \tau_2.$$

it leads to the same X as (E.12) does and thus the same boundary conditions (E.13). (If $m = 0$, there are again the additional solutions corresponding to the ones mentioned before.) The surface term in (E.1a) has now the properties described in sect. 7.1. The decoupling of positive from negative times is possible since L from (E.1a), with χ from (E.12) or (E.15), is not hermitean.

For a simple characterization of Dirichlet and Neumann components one would,

in view of (E.13), have to go to a representation where $\sigma_3 \bar{\sigma}_5$ is diagonal. This is not possible in a Majorana representation since the imaginary matrix $\sigma_3 \bar{\sigma}_2$ cannot be made real by a real orthogonal transformation.

However, if the computation of this Appendix is performed with a surface interaction on the timelike plane $\sum_{i=1}^3 n_i x^i = 0$, with the condition of decoupling of the two sides for all times, the matrix $\sigma_3 \bar{\sigma}_2 \bar{\sigma}_3 \sigma_5$ in (E.13) becomes replaced by *

* That at a timelike plane the boundary conditions $(1 \pm n_i \bar{\sigma}_i \bar{\sigma}_5) \psi = 0$ are meaningful has been noted by T.T. Wu [30] on the basis of different considerations.

$$n_i \bar{\sigma}_i \bar{\sigma}_5 = -n_1 \tau_3 + n_2 \tau_1 - n_3 \sigma_2 \tau_2$$

which can be made diagonal in a Majorana representation. The Lagrangean is hermitean in this case.

For $x^0 > 0, x^{0'} > 0$, the decoupling solutions (E.12) or (E.15) are

$$\begin{aligned} (E.16a) \quad G = & \frac{1}{2} (1 \pm \sigma_3 \tau_2) [(m\beta - i\alpha \bar{\sigma}_2) G_0 + \\ & + i\alpha \bar{\sigma}_2 \partial_0 G_0] + \\ & + \frac{1}{2} (1 \mp \sigma_3 \tau_2) [(m\beta - i\alpha \bar{\sigma}_2) G_N + i\alpha \bar{\sigma}_2 \partial_0 G_0] = \\ & = \frac{1}{2} (1 \pm i\alpha \bar{\sigma}_5) [(m\beta - i\alpha \bar{\sigma}_2) G_0 + i\alpha \bar{\sigma}_2 \partial_0 G_N] \end{aligned}$$

Here,

$$\begin{aligned} G_P = & (2\pi)^{-3} \int d^3 k (2k_0)^{-1} e^{i\vec{k} \cdot \vec{x}} (x^0 - x^{0'}) \\ & \cdot [e^{-ik_0 / (2\pi x^{0'})} \mp e^{-ik_0 / (2\pi x^{0'})}] \end{aligned}$$

with $k_0 = (\underline{k}^2 + \underline{m}^2)^{1/2}$ are the Minkowski-space versions of (1.14). Note that the first line of (E.16a) does not equal the first line of (E.16b).

If $m = 0$, the upper-sign solution for G is transformed into the lower-sign

one by $\psi \rightarrow i\sigma_1 \tau_2 \psi = i\bar{\psi} \tau_2$. This means that there is, concerning UV behaviour, no qualitative distinction between the Dirichlet-region and the Neumann one since the sign of the Fermion mass term is then not essential. To call that region the "Dirichlet" one in which, by a boundary source that has no effect on the other region (step 5) in sect. 7.1), inhomogeneous "Dirichlet" boundary conditions are imposed, is therefore, for the spin 1/2 field, merely a convention.

References

- 1) E.g. K.O. Friedrichs, "Mathematical Aspects of the Quantum Theory of Fields", New York, Interscience Publ. 1953
- 2) F.C.G. Stueckelberg, Phys. Rev. 81, 130 (1951)
- 3) N.M. Bogoliubov, D.V. Shirkov, "Introduction to the Theory of Quantized Fields", New York, Interscience Publ. 1959
- 4) C. Rebbi, Phys. Rep. 12C, 1 (1974) and refs. given there;
T. Eguchi, Phys. Rev. Lett. 44, 126 (1980)
- 5) A.A. Migdal, "QCD = Fermi String Theory" (Landau Inst. prepr.)
- 6) K. Hepp, Comm. Math. Phys. 2, 301 (1966), "Théorie de la renormalisation"
Lecture notes in Physics 2, Eds. J. Ehlers et al., Berlin, Springer 1969
- 7) G. 't Hooft, Nucl. Phys. B61, 455 (1973)
- 8) E.g., J. Schwinger, L.L. De Raad Jr., K.A. Milton, Ann. Phys. (N.Y.) 115, 1 (1978); K.A. Milton, Phys. Rev. 22, 1441, 1444 (1980); T. Kunimasa, IC/80/107 Trieste
- 9) R. Balian, B. Duplantier, Ann. Phys. (N.Y.) 112, 165 (1978), formula (5.27)
- 10) Cp. K. Symanzik, Lett. Nuovo Cimento 8, 771 (1973), Cargèse Lectures 1973 (DESY 73/58, unpublished); G. Parisi, Nucl. Phys. B150, 163 (1979);
R. Jackiw, S. Templeton, MIT-CTP 895 (Nov. 1980)
- 11) M. Lüscher, DESY 80/87
- 12) K.G. Wilson, J. Kogut, Phys. Rep. 12C, 75 (1974) Sect. 13;
G.A. Baker, J.M. Kincaid, Phys. Rev. Lett. 42, 1431 (1979)

- 13) D.J. Toms, Phys. Rev. D21, 2805 (1980)
- 14) I.M. Gelfand, G.E. Schilow, Verallgemeinerte Funktionen (Distributionen) I, Berlin, VEB Dtsch. Verl. d. Wiss. 1960
- 15) W. Zimmermann, Ann. Phys. (N.Y.) 77, 536, 570 (1973)
- 16) K. Symanzik, in "Particles, Fields, and Statistical Mechanics", Eds. M. Alexanian, A. Zepeda, Lecture Notes in Physics 32, p. 20, Berlin, Springer 1975
- 17) M. Lüscher, K. Symanzik, P. Weisz, Nucl. Phys. B173, 365 (1980)
- 18) W. Zimmermann, Comm. Math. Phys. 6, 161 (1967), ibid. 10, 325 (1968) and refs. given there
- 19) R. Haag, Kgl. Danske Videnskab. Selskab. Mat.-Fys. Medd. 29, No. 12 (1955)
- 20) K. Symanzik, IMM-NYU 327 (1964) (unpublished)
- 21) K.O. Friedrichs, H.N. Shapiro, Proc. Natl. Acad. Sci., U.S.A., 43, 336 (1951)
- 22) e.g. I. Akhiezer, V.B. Berestetskii, Quantum Electrodynamics, New York, Intersc. Publ. 1965
- 23) R.A. Brandt, Fortschr. d. Phys. 19, 249 (1970)
- 24) e.g. K.G. Wilson, Phys. Rev. D7, 2912 (1973);
I. Ya. Aref'eva, E.R. Nissimov, S.J. Pacheva, Comm. Math. Phys. 71, 213 (1980); K. Symanzik, DESY 71/05 (1977) (unpublished)
- 25) H. Lehmann, Z. Naturforsch. 8a, 579 (1953)
- 26) Y. Nambu, Phys. Lett. 80B, 372 (1979); J.-L. Gervais, A. Neveu, Nucl. Phys. B163, 189 (1980); A.M. Polyakov, Nucl. Phys. B164, 171 (1980);
I. Ya. Aref'eva, Lectures at the 1980 Bulgarian School, Primorsko
(to be published)
- 27) S. Epstein, V. Glaser, Ann. Inst. Henri Poincaré 19, 211 (1973)
- 28) G. Parisi, Lett. Nuovo Cimento 6, 450 (1973), in "New Developments in Quantum Field Theory and Statistical Mechanics" Cargèse 1976, p. 281, Eds. M. Lévy, P.K. Mitter, New York, Plenum 1977; K. Symanzik, Comm. Math. Phys. 45, 79 (1975), DESY 75/24 (unpublished)
- 29) P.A.M. Dirac, in "Fundamental Interactions at High Energy" I, 1 (1969), II, 144 (1970), New York, Gordon and Breach
- 30) E.g., O.D. Kellogg, "Foundations of Potential Theory", Berlin, Springer 1929
- 31) T.T. Wu (unpublished)




Fig. 1

Figure Caption

Fig. 1 Second-order contributions to the two-point function.

Lines are Dirichlet Green's functions.

