

81-9-219

DEUTSCHES ELEKTRONEN-SYNCHROTRON **DESY**

DESY 81-048
August 1981

OBSERVATION OF HIGH P_T JETS IN TWO-PHOTON INTERACTIONS

by

JADE Collaboration

NOTKESTRASSE 85 · 2 HAMBURG 52

DESY behält sich alle Rechte für den Fall der Schutzrechtserteilung und für die wirtschaftliche Verwertung der in diesem Bericht enthaltenen Informationen vor.

DESY reserves all rights for commercial use of information included in this report, especially in case of apply for or grant of patents.

To be sure that your preprints are promptly included in the
HIGH ENERGY PHYSICS INDEX ,
send them to the following address (if possible by air mail) :

DESY
Bibliothek
Notkestrasse 85
2 Hamburg 52
Germany

OBSERVATION OF HIGH P_T JETS IN TWO-PHOTON INTERACTIONS

JADE Collaboration

W. Bartel, D. Cords, P. Dittmann, R. Eichler, R. Felst, D. Haidt,
H. Krehbiel, K. Meier, B. Naroska, L.H. O'Neill 1), P. Steffen,
H. Wenninger 2), Y. Zhang 3)

Deutsches Elektronen-Synchrotron DESY, Hamburg, Germany

E. Elsen, M. Helm 4), A. Petersen, P. Warming, G. Weber

II. Institut f. Experimentalphysik der Univers. Hamburg, Germany

S. Bethke, H. Drumm, J. Heintze, G. Heinzelmann, K.H. Hellenbrand,
R.D. Heuer, J. von Krogh, P. Lenner, S. Kawabata, H. Matsumura,
T. Nozaki, J. Olsson, H. Rieseberg, A. Wagner

Physikalisches Institut der Univers. Heidelberg, Germany

A. Bell, F. Foster, G. Hughes, H. Wriedt

University of Lancaster, England

J. Allison, A.H. Ball, G. Bamford, R. Barlow, C. Bowdery,
I.P. Duerdoth, J.F. Hassard, B.T. King, F.K. Loebinger,
A.A. Macbeth, H. McCann, H.E. Mills, P.G. Murphy, K. Stephens

University of Manchester, England

D. Clarke, M.C. Goddard, R. Marshall, G.F. Pearce

Rutherford Laboratory, Chilton, England

T. Kobayashi, S. Komamiya, M. Koshiha, M. Minowa, M. Nosaki,
S. Orito, A. Sato, T. Suda 5), H. Takeda, Y. Totsuka, Y. Watanabe,
S. Yamada, C. Yanagisawa

Lab. of Int. Coll. on Elementary Particle Physics and Department of
Physics, University of Tokyo, Japan

1) Present address: Bell Laboratories, Whippany, N.J., USA.

2) On leave from CERN, Geneva, Switzerland.

3) Visitor from Institute of High Energy Physics, Chinese Academy
of Science, Peking, Peoples' Republic of China.

4) Present address: Texaco AG., Hamburg, Germany.

5) Present address: Cosmic Ray Laboratory, University Tokyo, Japan.

Abstract:

Events with a characteristic 2-jet topology have been observed in two-photon interactions. The production cross section is found to be higher than the point-like $\gamma\gamma \rightarrow q\bar{q}$ cross section, which is approached only at transverse momenta larger than 3 GeV/c.

Two-photon quark exchange scattering $\gamma\gamma \rightarrow q\bar{q}$ is of as fundamental importance as quark pair production in e^+e^- annihilation. This process [1] is expected to dominate at high four-momentum transfers which imply high two-photon invariant masses. Therefore, one has to look for a hard scattering process which is characterized in general by two non-back-to-back jets of high transverse momenta. This letter reports on a jet analysis carried out on the data from the JADE detector [2] and discusses briefly the question of the quark charges which enter to the fourth power in the cross section.

In order to avoid a high background from e^+e^- annihilation, we restrict our analysis in this first stage to single tagging triggers having an electron shower recorded in one of two sets of lead glass counters, each of which has 92 elements and surrounds the beam pipe about 5 m away from the interaction point covering an approximate angular range from 34 to 75 mrad with respect to the beam line. In addition the trigger required that an energy of at least 1 GeV was deposited in the cylindrical lead glass array or at least two charged tracks registered in the central drift chambers [2].

The recorded events were further selected by demanding a tagging cluster of at least 4 GeV, a well defined vertex (of at least 2 charged tracks) within 8 cm of the nominal interaction point, and at least four particles (charged or neutral) in total. Charged tracks were accepted down to 20° and photon clusters down to 12° with respect to the beam line. Since the bulk of low mass two-photon events is of no interest for this type of analysis, it was required further that the total visible energy E_{vis} (assuming all particles to be pions and photons) exceeded the momentum component along the beam line $P_{vis,z}$ by 3 GeV. Events with a visible energy greater than 20 GeV were excluded. With these selection criteria and a total integrated luminosity of 9730 nb^{-1} , ($29.9 \leq \sqrt{s} \leq 36.7 \text{ GeV}$) 538 events were found. The visible energy spectrum of these events is shown in Fig. 1a. Even for tagged events a contamination from e^+e^- annihilation must be considered and in Fig. 1b we show the energy spectrum of simulated annihilation events [3] with a radiated photon heading towards one of the tagging counters. Since electrons and photons cannot be distinguished in the tagging counter all "two-photon" events between 15 and 20 GeV can be associated with annihilation processes. One then obtains for the region from 3 to 10 GeV an upper limit of 2% for the annihilation background and from 10 to 15 GeV an upper limit of 20%.

Beam gas background was subtracted in the standard way by choosing control regions along the beam line away from the interaction point, and its contribution was found to be of the order of 1%. The background from tau pair production is calculated to be about 1%.

In order to select events with a characteristic 2-jet structure, a computer scan of the data was performed which resembles the visual grouping of tracks into narrow cones. Advantage was taken of the fact that a general jet search method [4] has been proposed and an efficient Monte-Carlo events at 13 and checked against e^+e^- annihilation Monte-Carlo events at 13 and 30 GeV. This search method allows for an arbitrary number of jets and proceeds via the following steps:

- Particles within 30° of each other are combined to form preclusters. Single isolated particles are also regarded as preclusters.
- Preclusters within 45° of each other are combined to form clusters. Single isolated preclusters are also regarded as clusters.
- Clusters with an energy of at least 2 GeV and with at least 2 particles (charged or neutral) are defined to be jets.

The search for jets was performed in the laboratory frame and also in the centre of mass system of the measured charged particles and photons, and it was found that our final results did not depend on the reference system. Here we shall only present the jet analysis performed in the centre of mass system. Out of the original data sample 119, 104, and 9 events are assigned as 1-jet, 2-jet, and more than 2-jet topologies, respectively. The 2-jet events, being of prime interest for our discussion, contain on the average 85% of the visible energy in the jets.

The inclusive particle spectra are shown in Fig. 2 as a function of the transverse particle momenta P_T with respect to the centre of mass direction of motion: (a) for all selected events, (b) for 2-jet events selected as described above, and

and the Monte-Carlo simulation leads to an expected number of 20 2-jet events with at least one $P_T(\text{JET})$ exceeding 2 GeV/c. Since $R_{\gamma\gamma}$ is close to one, we expect roughly the same number of muon pairs. In fact, 18 $\gamma\gamma \rightarrow \mu^+\mu^-$ events were observed which were selected under identical kinematic conditions. However, the actually observed number of high P_T 2-jet events was 42 and therefore twice as high as expected. The transverse momentum dependence of the 2-jet event cross section is shown in Fig. 3 as a function of $x_T(\text{JET}) = P_T(\text{JET})/E_{\text{beam}}$ and compared to the curve expected for $\gamma\gamma \rightarrow q\bar{q}$. It appears that the point-like $\gamma\gamma \rightarrow q\bar{q}$ cross section is approached from above [9]. Whereas the measurement and the prediction differ by a factor of nearly 4 at $P_T(\text{JET})$ of 2 GeV/c, this difference drops down to a factor of about 1.5 for $P_T(\text{JET})$ values above 3 GeV/c. The excess of events may possibly be accounted for by the high P_T tail of diffractive events which dominate at low P_T and also by processes [1,6] like $\gamma q \rightarrow qg$ or $\gamma g \rightarrow q\bar{q}$ which may have a steeper P_T dependence than $\gamma\gamma \rightarrow q\bar{q}$.

The measured invariant mass of the observed high P_T 2-jet events ranges from 5 to 15 GeV. Because of this high mass the quark was included in the charge counting for the quark coupling which also determines the normalization in Fig. 3. The above calculations assumed standard fractional charged quarks. If, however, one follows the conjecture [10] of integer charged quarks, then the curve in Fig. 3 has to move up by a factor of 2.65, and the data points beyond $P_T(\text{JET})$ of 3 GeV/c would fall below this curve by nearly a factor of two. Even with our limited number of events this conjecture seems to be ruled out. Another point of interest is that within the scheme of integer charged quarks the prediction should increase by an additional

(c) for only those 42 2-jet events with at least one of the jets having a $P_T(\text{JET})$ which exceeds 2 GeV/c. One observes that the P_T distribution becomes flatter as more selections are applied to the data. The high P_T 2-jet class of events agrees best in shape with what is expected [6] for $\gamma\gamma \rightarrow q\bar{q}$ when in addition to the QED type process, a standard fragmentation of the quarks is taken into account. The curve [6] in Fig. 2 is normalized to the high P_T 2-jet events and depicts the expected approximate x_T^{-4} dependence where $x_T = P_T/E_{\text{beam}}$.

This definition of jets is purely experimental and it was the purpose of a detailed Monte-Carlo simulation to see how well the experimental findings resembled the leading order quark scattering mentioned above. The Monte-Carlo calculation employed a two-photon event generator [7] to produce in a QED-like fashion quark pairs (with a mass of 300 MeV for u,d,s and of 1.5 GeV for c) which then turn into hadrons via the standard Field-Feynman fragmentation functions [8]. It turned out that this model described the shape of various distributions of the original data sample of 538 events, for example the charged and neutral multiplicity, the two-photon invariant mass spectrum, and the thrust in the centre of mass system. An exception is the inclusive particle P_T distribution which came out too flat and only fitted the high P_T 2-jet events, as we have already discussed above.

The experimental 2-jet cross section may now be compared with the absolute prediction for the $\gamma\gamma \rightarrow q\bar{q}$ process.

Including the first four quark flavours u,d,s,c one obtains:

$$R_{\gamma\gamma} = \frac{\sigma(e^+e^- \rightarrow \gamma\gamma \rightarrow q\bar{q})}{\sigma(e^+e^- \rightarrow \gamma\gamma \rightarrow \mu^+\mu^-)} = 3 \sum_{i=u,d,s,c} Q_i^4 = \frac{34}{27}$$

factor of 1.8 when the colour threshold is crossed. This threshold effect is not expected for fractional charged quarks, and for integer charged quarks it is certainly excluded in the energy range of our two-photon 2-jet events.

In conclusion, we observe that the high P_T 2-jet cross section in two-photon interactions approaches the theoretical point-like $\chi\chi \rightarrow q\bar{q}$ cross section from above and that the scheme of integer charged quarks is excluded from this process by four standard deviations. More data would be needed to decide at which P_T (JET) value above 3 GeV/c the point-like $\chi\chi \rightarrow q\bar{q}$ limit is reached.


Acknowledgement:

We acknowledge the excellent support of the PETRA machine group and the efforts of all the engineers and technicians who have participated in the construction and maintenance of our detector. This experiment was supported by the Bundesministerium für Forschung und Technologie, by the Education Ministry of Japan and by the U.K. Science Research Council through the Rutherford Laboratory. The visiting groups wish to thank the DESY directorate for their hospitality.

REFERENCES

[1] H. Suura, T.F. Walsh, B.L. Young, Lett. Nuovo Cimento 4 (1972) 505.
 C. Llewellyn Smith, Phys. Lett. 79B (1978) 83.
 S. Brodsky, T. DeGrand, J. Gunion, J. Weis, Phys. Rev. Lett. 41 (1978) 672 and Phys. Rev. D19 (1979) 1418.
 K. Kajantie, Physica Scripta 29 (1979) 230 and Acta Physica Austriaca, Suppl. XXI (1979) 663.
 [2] JADE-Collab., W. Bartel et al., Phys. Lett. 88B (1979) 171.
 [3] Events were generated with the Lund Monte-Carlo program:

B. Anderson, G. Gustafson, T. Sjöstrand, Phys. Lett. 94B (1980) 211,
 and radiative corrections were taken into account according to:
 G. Bonneau and P. Martin, Nucl. Phys. B27 (1971) 381.
 [4] K. Lanius, preprint DESY 80/36 (1980).
 [5] H.J. Daum, H. Meyer, J. Buerger, Zeitschr. f. Phys. C8 (1981) 167.
 [6] K. Kajantie, Proceed. of 4th Intern. Coll. on Photon-Photon Interactions, Paris, France (1981), to be published.
 [7] P. Bhattacharya, J. Smith, G. Grammer, Phys. Rev. D15 (1977) 3267.
 J. Smith, J.A.M. Vermaseren, G. Grammer, Phys. Rev. D15 (1977) 3280.
 J.A.M. Vermaseren, Program Write-Up (1978), unpublished.
 [8] R.D. Field and R.P. Feynman, Nucl. Phys. B136 (1978) 1:
 The constant in the fragmentation function has been chosen $a=0.77$ for u,d,s quarks and $a=0$ for the c quark. The transverse momentum within a jet is limited by $\sqrt{2} G(p_T) = 350 \text{ MeV}$. Sea quark contributions are u:d:s = 2:2:1 and the vector to pseudo-vector part is 1:1.
 [9] A similar analysis is in progress for the PLUTO and TASSO experiments:
 W. Wagner, Proceed. of XXth Intern. Conf. on High Energy Physics, Madison, Wisconsin, USA, p.576 (1980).
 D. Cords, Proceed. of 4th Intern. Coll. on Photon-Photon Interactions, Paris, France (1981), to be published.
 [10] Y. Nambu and M.-Y. Han, Phys. Rev. D10 (1974) 674.
 J.C. Pati, Proceed. of Intern. Summer Inst. on Theor. Particle Phys., Hamburg, Germany, P. 384 (1975).
 H.K. Lee, Preprint Intern. Centre of Theor. Phys., Trieste, Italy, IC/78/95 (1978).



32923

Fig. 2: Inclusive particle distribution for multihadronic two-photon events compared with the expected slope of $d\sigma/dx_T$ where $x_T = P_T/E_{beam}$ (see ref. 6, Fig. 4).




Fig. 1: Visible energy of (a) single tag multihadronic two-photon events (see text) and (b) Monte-Carlo simulated annihilation events with a radiated photon heading towards one of the tagging counters.




Fig. 3: Transverse momentum distribution of jets for single tag data compared with absolute predictions for fractional charged quarks (curve). $x_T(JET) = P_T(JET)/E_{beam}$ is taken with respect to the centre of mass direction of motion. The cross section is given for the JADE tagging condition on the lefthand scale and integrated over all electron angles and energies on the righthand scale.