

82-3-336

DESY 82-008  
Erratum



Determination of the Radiative Widths of the  $\eta'$  and  $A_2$  from Two Photon

Exchange Production

Physics Letters 114B (1982) 378

The joint angular distribution quoted for the cascade decay

$A_2 \rightarrow \rho^\pm \pi^\mp$ ,  $\rho^\pm \rightarrow \pi^\pm \pi^0$  is incorrect. It should be replaced by the expression


$$\frac{d^3\sigma}{dcos\theta_1 dcos\theta_2 d\phi_2} \sim \sin^2\theta_1 \sin^2\theta_2 (1 - \sin^2\theta_1 \sin^2\phi_2)$$

where  $\theta_1$  and  $\theta_2$  are defined as before and  $\phi_2$  is the angle between the production and decay planes of the  $\rho$  in the  $\rho$  rest frame. We are grateful to H. Kolanoski for pointing out this error to us. Recalculating the acceptance for the  $A_2$  within our cuts we find a 19% reduction.

We have also noticed an error in our calculation of the two photon luminosity function which results in 14%, 15% increases respectively for the radiative widths of the  $\eta'$ ,  $A_2$ . Taking into account both the above corrections we find :

$$\Gamma_{\gamma\gamma}(\eta') = 6.2 \pm 1.1 \text{ (stat.)} \pm 0.8 \text{ (syst.) KeV}$$

$$\Gamma_{\gamma\gamma}(A_2) = 0.81 \pm 0.19 \text{ (stat.)} \begin{array}{l} + 0.42 \\ - 0.11 \end{array} \text{ (syst.) KeV}$$



DESY 82-008  
February 1982

DETERMINATION OF THE RADIATIVE WIDTHS OF THE  $n'$  AND  $A_2$   
FROM TWO PHOTON EXCHANGE PRODUCTION

by

*CELLO Collaboration*

NOTKESTRASSE 85 · 2 HAMBURG 52

**DESY behält sich alle Rechte für den Fall der Schutzrechtserteilung und für die wirtschaftliche  
Verwertung der in diesem Bericht enthaltenen Informationen vor.**

**DESY reserves all rights for commercial use of information included in this report, especially in  
case of filing application for or grant of patents.**

**To be sure that your preprints are promptly included in the  
HIGH ENERGY PHYSICS INDEX ,  
send them to the following address ( if possible by air mail ) :**

**DESY  
Bibliothek  
Notkestrasse 85  
2 Hamburg 52  
Germany**

DETERMINATION OF THE RADIATIVE WIDTHS OF THE  $\eta'$  AND  $A_2$   
FROM TWO PHOTON EXCHANGE PRODUCTION

ABSTRACT: From the two photon exchange processes

$$e^+ e^- \rightarrow e^+ e^- \eta' \quad (958) \rightarrow e^+ e^- \rho^0 \gamma$$

and

$$e^+ e^- \rightarrow e^+ e^- A_2 \quad (1310) \rightarrow e^+ e^- \pi^+ \pi^- \pi^0$$

observed using the CELLO detector at PETRA the radiative widths of the  $\eta'$  and  $A_2$  have been determined with the results:

$$\Gamma_{\gamma\gamma}(\eta') = 5.4 \pm 1.0 [\text{stat.}] \pm 0.7 [\text{syst.}] \text{ keV}$$

$$\Gamma_{\gamma\gamma}(A_2) = 0.59 \pm 0.14 [\text{stat.}] \pm 0.31 [\text{syst.}] \text{ keV}$$

CELLO Collaboration

H.J. Behrend, Ch. Chen<sup>1)</sup>, H. Fenner, J.H. Field<sup>2)</sup>, U. Gimpel, V. Schröder,

H. Sirdt  
Deutsches Elektronen Synchrotron DESY, Hamburg, Germany

G. D'Agostini, W.-D. Apel, S. Banerjee, J. Bodenkamp, D. Chrobaczek, J. Engler,  
G. Flügge, D.C. Fries, W. Fues, K. Gamerding, G. Hopp, H. Küster, H. Müller,  
H. Randoll, G. Schmidt, H. Schneider  
Kernforschungszentrum Karlsruhe and Universität Karlsruhe, Karlsruhe, Germany

W. Boer, G. Buschhorn, G. Grindhammer, P. Grossmann, B. Gunderson,  
C. Kiesling, R. Kotthaus, U. Kruse<sup>3)</sup>, H. Lierl, D. Lüters, T. Meyer<sup>4)</sup>,  
H. Oberlack, P. Schacht, M.J. Schachter<sup>5)</sup>, A. Snyder<sup>6)</sup>, H. Steiner<sup>7)</sup>  
Max-Planck-Institut für Physik und Astrophysik, München, Germany

G. Carnesecchi, P. Colas, A. Cordier, M. Davier, D. Fourrier, J.F. Grivaz,  
J. Haissinski, V. Journe, A. Klarsfeld, F. Laplanche, F. Le Diberder, U. Mallik,  
J.J. Veillet  
Laboratoire de l'Accélérateur Linéaire, Orsay, France

R. George<sup>8)</sup>, M. Goldberg, B. Grosssetete, O. Hamon, F. Kovacs,  
G. London<sup>9)</sup>, L. Poggiali, M. Rivocal  
Laboratoire de la Physique Nucléaire et Hautes Energies, University of Paris,  
Paris, France

R. Aleksan, J. Bouchez, G. Cozika, Y. Ductros, A. Gaidot, S. Jadach<sup>9)</sup>,  
Y. Lavagne, J. Pamela, J.P. Pansart, F. Pierre  
Centre d'Etudes Nucléaires, Saclay, France

1) Visitor from the Institute of High Energy Physics, Chinese Academy of Science, Peking, People's Republic of China

2) On leave of absence, presently at University of Paris

3) Visitor from the University of Illinois, Urbana, USA

4) Now at the University of Wisconsin, Madison, USA

5) Now at DESY, Hamburg, Germany

6) Now at Rutgers University, New Brunswick, USA

7) Alexander von Humboldt Foundation, Senior American Scientist, University of California, Berkeley, USA

8) On leave of absence, presently at University of California, Berkeley, USA

9) Visitor from the University of Cracow, Poland

**INTRODUCTION.**  $\eta'$  and  $\Lambda_2$  production have been observed in two photon interactions in the CELLO detector at PETRA. For this analysis data with two visible charged tracks in the central detector and one photon in the liquid argon calorimeter were used. The scattered electron and positron in the final state were unobserved. The average CMS energy was 34.6 GeV and the total integrated luminosity  $11.15 \text{ pb}^{-1}$ . A detailed description of the CELLO detector is given in ref. [1]. Charged tracks are measured in a cylindrical wire chamber detector consisting of 7 drift and 5 proportional chamber layers in a magnetic field of  $1.3 \text{ T}$ . The liquid argon calorimeter samples electromagnetic showers 17 times in depth leading to 6 spatial measurements of the shower, up to a maximum depth of 20 radiation lengths. The fast trigger [2] used required two tracks in a plane perpendicular to the beam axis (R0) and one in the plane containing the beam axis (RZ). The trigger accepts tracks with transverse momentum greater than 200 MeV/c and coming from within  $\pm 10 \text{ cm}$  of the interaction point along the beam axis. The trigger condition was verified using more detailed track information read directly from the central wire chambers by an on line filter program.

**EVENT SELECTION.** Candidates for low multiplicity events produced in two photon interactions were first selected using track information recorded by the fast trigger. Events were required to have  $\leq 4$  charged tracks. Bhhabha events satisfying the neutral trigger condition ( $E > 6 \text{ GeV}$ ) were rejected. The selected events were processed through the reconstruction programs for charged tracks in the central detector and for showers in the liquid argon calorimeter. The charged track reconstruction efficiency was 99%. Events with two oppositely charged tracks having the vertex within  $\pm 2 \text{ cm}$  around the interaction point were kept for further analysis. Cosmic  $\mu$  background was rejected by a cut on the acollinearity angle  $\alpha$  in the RZ-projection:  $|\cos\alpha| < 0.995$ .

After this cut the residual background from cosmic  $\mu$ 's and beam-gas events was  $\sim 2\%$ . Background from the QED reaction

$$e^+ e^- \rightarrow (e^+ e^-) e^+ e^-$$

was removed by rejecting all events where one of the two detected tracks has been identified in the liquid argon calorimeter as an electron. For the electron identification the total energy of the shower linked to the charged track, and the shower development in depth in the liquid argon were used. Electrons accompanied by bremsstrahlung photons were removed by rejecting events with photons within a cone of 100 mrad around a track. Photon candidates in the liquid argon calorimeter were selected using the following cuts:

- energy  $> 100 \text{ MeV}$
  - more than 20% of the total charge is deposited between 3 and 8 radiation lengths in depth
  - no charge is found beyond 16 radiation lengths
  - the shower has to be reconstructed in depth by at least two clusters.
- In addition all photon candidates possibly originating in the interaction of charged particles in the liquid argon were removed. The background from exclusive  $\mu^+ \mu^-$ ,  $\pi^+ \pi^-$  and residual  $e^+ e^-$  production in two photon interactions has been reduced using the following cuts:
- the momentum transverse to the beam direction of the pair has to be larger than 50 MeV/c
  - The acollinearity angle between the two charged tracks in the plane perpendicular to the beam axis has to be at least 200 mrad.

Finally, the background from photons simulated by noise in the liquid argon calorimeter was reduced by the following cuts:

- one and only one photon candidate is found within  $\pm 250 \text{ mrad}$  of the direction of the missing transverse momentum vector of the pion pair
- the total transverse momentum of the two pions and the photon has to be  $< 200 \text{ MeV}/c$ .

To improve the energy measurement of the photons we corrected the measured photon energy by multiplying it with the ratio of transverse momentum of the two pions and the transverse momentum of the photon [3]. After this correction a more stringent cut for the transverse momentum of the  $\pi^+ \pi^- \gamma$  system was applied:

$P_{\perp}(\pi^+ \pi^- \gamma) < 100 \text{ MeV}/c$ .

**RESULTS FOR  $\eta'$ .** The  $\pi^+ \pi^- \gamma$  mass distribution after the cuts detailed above is shown in Fig. 1. A clear signal at the  $\eta'$  mass region can be seen. Fitting this distribution in the region of the  $\eta'$ -peak with a gaussian and a linear background gives for  $m(\eta') = 957 \pm 3 \text{ MeV}$ , in good agreement with ref. [4].

Fig. 2 shows the  $\pi^+ \pi^- \gamma$  mass distribution for events with  $600 < m_{\pi^+ \pi^-} < 900 \text{ MeV}/c^2$ . Almost all events in the  $\eta'$  mass peak are compatible with the  $\eta'$  decay  $\eta' \rightarrow \rho \gamma$ . The subtraction of a handdrawn background gives a signal of 43 events above a background of 32 events. To correct the data for all losses due to acceptance, data selection and reconstruction inefficiency, we generated Monte Carlo (MC) events using a detailed detector simulation. In the MC generator we used exact transverse photon luminosity functions [5] and used particle data group values for the  $\eta'$ ,  $\rho$  masses, the  $\rho$  width and the  $\eta' \rightarrow \rho \gamma$  decay branching ratio [4].

The MC events were passed through the same reconstruction programs as the real data. Good agreement between the data and the MC events is obtained for the  $\pi^+\pi^-$  mass distribution and the transverse momentum distributions of the  $\pi^+\pi^-$  and the  $\pi^+\pi^-Y$  systems. With all cuts mentioned above and the  $\eta'$  mass cut  $850 < m_{\pi^+\pi^-} < 1050 \text{ MeV}/c^2$ , the overall acceptance was found to be 0.78%. This yields a radiative width of the  $\eta'$  of  $\Gamma_{YY}(\eta') = 5.4 \pm 1.0 \text{ keV}$ , where the error includes the statistical error and the error due to the background subtraction only. We have checked the dependence of this value on the minimum photon energy required and found it to stay constant within errors. To estimate the systematic errors detailed MC studies were made varying the noise level in the liquid argon detector. To this error (12%) we have added in quadrature the errors due to the uncertainty in the overall normalization (3%), the trigger efficiency (2%) and data selection efficiency (4%), yielding a total systematic error of 0.7 keV.

RESULTS FOR  $A_2$ . Fig. 1 shows in addition to the  $\eta'$  signal a peak at  $m_{\pi^+\pi^-} = 1300 \text{ MeV}/c^2$ . We interpret this as evidence for two photon production of the  $A_2$  in the reaction

$$e^+e^- \rightarrow (e^+e^-) A_2 \rightarrow (e^+e^-) \pi^+\pi^-\pi^0$$

where one low energy photon from the  $\pi^0$  decay is not detected. To confirm this interpretation we generated MC events and passed them through the standard reconstruction programs. The virtual photons were generated as for the  $\eta'$ . The  $A_2$  was assumed to be in a pure helicity 2 state [6]. Neglecting relativistic corrections, the joint decay distribution of the  $A_2$  is

$$\frac{d^2\sigma}{dcos\theta_1 dcos\theta_2} \sim 4 \sin^4 \theta_1 \cdot \cos^2 \theta_2 + \sin^2 \theta_1 (\sin^2 \theta_1 + 2 \cos^4 \theta_1) \cdot \sin^2 \theta_2$$

where  $\theta_1$  is the angle between the virtual photon direction and the  $p$  direction of flight in the  $YY$  c.m. system, and  $\theta_2$  is the angle between the  $p$  direction of flight and a  $\rho$  decay pion in the  $p$  c.m. system. Relativistic Breit-Wigner shapes were used for the  $A_2$  and  $\rho$  with standard values for the masses and widths and for the  $A_2 \rightarrow \rho^+\pi^+ \rightarrow \pi^+\pi^-\pi^0$  decay branching ratio [4]. We calculated the acceptance for this reaction detecting both photons from the  $\pi^0$  decay and demanding only a minimum transverse momentum of 50 MeV/c of the two charged pions. With the full reconstructed  $\pi^0$  we found the acceptance to be down by a factor of 5.1 in comparison to the detection of only one photon including all the cuts as described for the  $\eta'$  selection. The latter acceptance was found to be 1.96%. Thus our statistics is not large enough to see the  $A_2$  signal in the full reconstructed  $\pi^+\pi^-\pi^0$  mode. However demanding only one

detected photon, we get good agreement for the  $\pi^+\pi^-Y$  mass distribution between the data and the MC generated  $A_2$  events. Subtracting a handrawn background we obtain a signal of 35 events above a background of 40 events. This yields a radiative width of

$$\Gamma_{YY}(A_2) = 0.59 \pm 0.14[\text{stat.}] \pm 0.31[\text{syst.}] \text{ keV.}$$

Again this value was checked to be independent within errors of the minimum photon energy required. The systematic error comes mainly from changes in acceptance resulting from uncertainties in the liquid argon noise level. The detection efficiency for the second low energy photon of the  $\pi^0$  decay depends strongly on this noise level. Because we demand for the acceptance calculation that one and only one photon be detected within the angular cone of 250 mrad around the missing transverse momentum vector of the charged pion pair, the acceptance decreases when the noise level in the liquid argon calorimeter is increased. Therefore the resulting systematic error shows a strong asymmetry.

#### CONCLUSIONS. Our value of the radiative width of the $\eta'$ of

$$\Gamma_{YY}(\eta') = 5.4 \pm 1.0[\text{stat.}] \pm 0.7[\text{syst.}] \text{ keV}$$

confirms with improved accuracy the values obtained by the MARK II detector at SPEAR of  $\Gamma_{YY}(\eta') = 5.9 \pm 1.6[\text{stat.}] \pm 1.2[\text{syst.}] \text{ keV}$  [3] and the preliminary result from JADE at PETRA of  $\Gamma_{YY}(\eta') = 7.5 \pm 1.7[\text{stat.}] \pm 3.5[\text{syst.}] \text{ keV}$  [7].

It also agrees with the prediction (6 keV) of the quark model with fractionally charged quarks and rules out the integrally charge quark model which predicts a value of 25.6 keV [8]. For the radiative width of the  $A_2$  we obtained, using the decay mode  $A_2 \rightarrow \rho^+\pi^+ \rightarrow \pi^+\pi^0$

$$\Gamma_{YY}(A_2) = 0.59 \pm 0.14[\text{stat.}] \pm 0.31[\text{syst.}] \text{ keV}$$

which may be compared with the preliminary value of JADE of

$$\Gamma_{YY}(A_2) = 1.2 \pm 0.4[\text{stat.}] \pm 0.5[\text{syst.}] \text{ keV} [9].$$

Our result is in agreement with and of comparable accuracy to that of the Crystal Ball detector, using the decay mode  $A_2 \rightarrow \eta\pi^0 \rightarrow 4\gamma$ , of

$$\Gamma_{YY}(A_2) = 0.77 \pm 0.18[\text{stat.}] \pm 0.27[\text{syst.}] \text{ keV} [7,10].$$

Assuming ideal mixing in the  $\pi^+$  nonet, the ratio  $\Gamma_{YY}(A_2)/\Gamma_{YY}(f_0)$  is predicted to be 9:25 [11]. Taking the weighted average of the three measurements of  $\Gamma_{YY}(A_2)$  quoted above (statistical and systematic errors have been added in quadrature) one obtains  $\Gamma_{YY}(A_2) = 0.75 \pm 0.22 \text{ keV}$ . Taking the corresponding average of the

five values quoted in ref. [9] for  $\Gamma_{\gamma\gamma}(f^0)$  leads to the value of  $\Gamma_{\gamma\gamma}(f^0) = 3.1 \pm 0.3$  kev. Thus the measured ratio  $\Gamma_{\gamma\gamma}(A_2)/\Gamma_{\gamma\gamma}(f^0)$  is  $0.24 \pm 0.08$ , to be compared with the above prediction of 0.36. A recent detailed analysis of our data for the two photon production of  $f^0$  in the  $\pi^+\pi^-$  decay mode indicates a significantly smaller value for  $\Gamma_{\gamma\gamma}(f^0)$  than quoted in ref. [9]. This is due to an important contribution of a dipion continuum production under the  $f^0$  peak. Taking this into account will improve the agreement of the above ratio with the ideal mixing prediction.

**ACKNOWLEDGMENT.** We are indebted to the PETRA machine group and the DESY computer center for their excellent support during the experiments. We acknowledge the efforts of our engineers and technicians and in particular the support of Dr. Horlitz and G. Mayaux and their groups in the operation of the magnet system. The visiting groups wish to thank the DESY directorate for the support and kind hospitality extended to them. This work was partly supported by the Bundesministerium für Forschung und Technologie.

#### REFERENCES

- [1] CELLO - A new detector at PETRA. H.-J. Behrend et al., Phys. Scripta 23 (1981) 610.
- [2] H.-J. Behrend, Computer Physics Communications 22 (1981) 365; V. Schroeder, CERN 81-07 (1981) 22.
- [3] G.S. Abrams et al., Phys. Rev. Lett. 43 (1979) 477.
- [4] Particle Data Group, Rev. Mod. Phys. 52 (1980) 51.
- [5] J.H. Field, Nucl. Phys. B168 (1980) 477 and erratum.
- [6] This assumption is in agreement with theoretical expectations:  
B. Schrempp-Otto, F. Schrempp, T.F. Walsh, Phys. Lett. 36B (1971) 463;  
J.L. Rosner, Phys. Reports C11 (1974) 189;  
P. Grassberger, R. Kögerler, Nucl. Phys. B106 (1976) 451;  
Also a direct measurement of the Crystal Ball detector (see ref. [7,10])  
of  $f^0 \rightarrow \pi^0 \pi^0$  production indicates the dominance of the helicity 2 amplitude.
- [7] J.H. Field, Rapporteur talk at the International Colloquium on Photon Photon Interactions (Paris, 1981).
- [8] H. Suma, T.F. Walsh and B.-L. Young, Nuovo Cimento Lett. 4 (1972) 505.
- [9] Ch. Berger, Rapporteur talk at the EPS Int. Conf. on High Energy Physics, Lisbon, 1981, CERN-EP/81-154;
- R. J. Wedemeyer, invited talk given at the "International Symposium on Lepton and Photon Interactions at High Energies", Bonn, 1981, BONN-HE-81-25;
- [10] D. L. Burke, invited talk at the IVth International Colloquium on Photon Photon Interactions, (Paris 1981), SLAC-PUB-2745.
- [11] F.J. Gilman, Proceedings of the 1979 Conference on Two Photon Interactions, (Lake Tahoe) 215.




FIGURE CAPTIONS

Fig. 1: Distribution of  $\pi^+\pi^-\gamma$  effective mass.

Fig. 2: Distribution of  $\pi^+\pi^-\gamma$  effective mass with the additional cut

$600 < m_{\pi^+\pi^-} < 900 \text{ MeV}/c^2$ .




Fig.2