

82-12-310

DEUTSCHES ELEKTRONEN-SYNCHROTRON **DESY**

DESY 82-075
November 1982

COMPARISON OF IMPACT-PICTURE PREDICTIONS WITH DATA FROM CERN $p\bar{p}$ COLLIDER

by

C. Bourrely

*Centre de Physique Théorique,
Centre National de la Recherche Scientifique, Marseille*

and

Physics Department, Westfield College, London

J. Soffer

*Centre de Physique Théorique,
Centre National de la Recherche Scientifique, Marseille*

Tai Tsun Wu

Deutsches Elektronen-Synchrotron DESY, Hamburg

NOTKESTRASSE 85 · 2 HAMBURG 52

DESY behält sich alle Rechte für den Fall der Schutzrechtserteilung und für die wirtschaftliche Verwertung der in diesem Bericht enthaltenen Informationen vor.

DESY reserves all rights for commercial use of information included in this report, especially in case of filing application for or grant of patents.

**To be sure that your preprints are promptly included in the
HIGH ENERGY PHYSICS INDEX ,
send them to the following address (if possible by air mail) :**

**DESY
Bibliothek
Notkestrasse 85
2 Hamburg 52
Germany**

COMPARISON OF IMPACT-PICTURE PREDICTIONS WITH DATA FROM CERN $p\bar{p}$ COLLIDER

C. BOURRELY J. SOFFER Tai Tsun WU *

Centre de Physique Théorique, Centre National de la Recherche Scientifique,
Marseille, France, Deutsches Elektronen-Synchrotron DESY, Hamburg, Germany
and Physics Department, Westfield College, London, England

ABSTRACT :

Recent data on the total cross section and the elastic slope parameters at small momentum transfers from the UA1 and UA4 Collaborations at the CERN $p\bar{p}$ Collider are compared with the impact-picture predictions.

* On leave from Harvard University, Cambridge, Massachusetts.
Work supported in part by the U.S. Department of Energy Contract
no DE-AS02-76ER03227.

In 1970 Cheng and Wu predicted the rising total cross section [1] on the basis of a systematic study of the high-energy behavior of quantum field theory [2]. The first phenomenological analysis incorporating this prediction was carried out [3] using Serpukhov data and on the assumption of the absence of exotic exchange. This analysis was greatly improved [4] a year later when the ISR data [5] showed that the increase in total cross section was only about 50% higher than that from the first analysis [3]. Numerous predictions were presented at the 1973 Stony Brook Conference [6]. Other results were summarized at the 18th International Conference on High-Energy Physics [7].

Recently, data became available from the CERN $p\bar{p}$ Collider at a high energy of $\sqrt{s} = 540$ GeV [8,9]. It is the purpose of this note to compare the 1973 predictions with the 1982 data.

In reference 6 two sets of predictions were given, labelled as $c' = 0$ and $c' = 1$ respectively in the notation there. The case of $c' = 0$ is the same as that given in reference 4. In Fig.1 here we show the total pp and $p\bar{p}$ cross sections of reference 6 (Fig.8 there) together with the UA4 data [8]. In Fig.2, we show the slope parameters of reference 6 (Fig.14 there) with the UA1 and UA4 data [8,9]. From these comparisons, which are also given in Table I, the case $c' = 1$ is favored.

Both the total cross section from UA4 [8] and the slope parameters from UA1 [9] are in good agreement with the early impact-picture predictions. However, the slope parameter from UA4 [8] for smaller values of momentum transfer disagrees with the predictions. The UA4 slope parameter is much larger, as discussed in the last paragraph of reference 8. In order to get a better feeling of this discrepancy, the unnormalized differential cross section is compared directly in Fig.3. It is seen that there is no serious discrepancy between the data and the predictions. The cause of the large slope parameter of UA4 is due mainly to two data points at $|t| = 0.18$ and 0.19 , which are in disagreement with the UA1 data. In other words, the discrepancy is entirely an experimental problem. We expect the UA4 slope parameter to be an overestimate.

The phenomenology of the impact-picture was improved [10] several years ago so far as polarization and Regge backgrounds are concerned. These improvements have no significant effect for small momentum transfers at these very high energies. The curve of Fig.3 is essentially unchanged.

It is clear that the impact-picture predictions of nearly a decade ago are very adequate to describe the data from the CERN $p\bar{p}$ collider for small momentum transfers. We look forward to many further comparisons in the near future.

CB acknowledges Westfield College for hospitality and the Royal Society for a fellowship. TTW thanks Professor Hans Joos, Professor Erich Lohrmann, Professor Paul Söding, Professor Volker Soergal, and Professor Thomas Walsh for their kind hospitality at DESY.

REFERENCES

-0-

- [1] H. Cheng and T.T. Wu, Phys. Rev. Letters 24 (1970), 1456.
- [2] H. Cheng and T.T. Wu, Phys. Rev. Letters 22 (1969) 666, 23 (1969) 1311, and 24 (1970) 759; Phys. Rev. 182 (1969) 1852 and D1 (1970) 2775.
- [3] H. Cheng, J.K. Walker, and T.T. Wu, "Impact-Picture of Very-High-Energy Hadron Interactions (With Numerical Results)," submitted to the 16th. International Conference on High Energy Physics at Batavia, Illinois (1972).
- [4] H. Cheng, J.K. Walker, and T.T. Wu, Phys. Letters 44B (1973) 97.
- [5] U. Amaldi et al., Phys. Letters 44B (1973) 112; S.R. Amendolia et al., Phys. Letters 44B (1973) 119.
- [6] T.T. Wu and H. Cheng, Fifth International Conference on High-Energy Collisions, Stony Brook, N.Y. 1973 (A.I.P. Conference Proceedings n° 15), p.54.
- [7] T.T. Wu, Proceedings of the 18th. International Conference on High-Energy Physics, Tbilisi, U.S.S.R. (1976), Vol.I, p.A5-23.
- [8] UA4 Collaboration, R. Battiston et al., Phys. Letters 115B (1982) 333.
- [9] UA1 Collaboration, G. Arnison et al., "Small Angle Elastic Scattering at the CERN Proton-Antiproton Collider", preprint CERN-EP/82-123, Proceedings of the 21st International Conference on High-Energy Physics, Paris (1982).
- [10] C. Bourrely, J. Soffer, and T.T. Wu, Phys. Rev. D19 (1979) 3249.

FIGURE CAPTIONS

-0-

- FIG.1 Comparison of the prediction of rising total cross-sections from ref 6 with the recent UA4 data [8]. The solid lines correspond to $c' = 0$ and the dashed lines to $c' = 1$.
- Fig.2 Comparison of the prediction of the slope parameters from ref 6 with the recent UA1 data [8] and UA4 data [9].
- Fig.3 $|t|$ -distribution of elastic scattering events in the $|t|$ -range from 0.05 to 0.26 $(\text{GeV}/c)^2$ taken from [9] and compared to the $c' = 1$ prediction from ref 6.

TABLE 1 : COMPARISON OF PREDICTIONS WITH DATA

	Theory		Experiment	
	$c' = 0[4,6]$	$c' = 1[6]$	UA1[9]	UA4[8]
Total cross section (mb)	58	62	-	66 ± 7
Slope parameter $(\text{GeV}/c)^{-2}$				
$0.05 < t < 0.19$	13.8	14.2	-	17.2 ± 1.0
$0.14 < t < 0.26$	13.1	13.6	13.3 ± 1.5	-




FIG. 1




FIG. 2




FIG. 3

