

DESY 83-053
June 1983



SEARCH FOR MAGNETIC MONOPOLES IN ELECTRON-POSITRON
COLLISIONS AT 34 GeV C.M. ENERGY

by

P. Musset and M. Price
CERN, Genève, Switzerland

and

E. Lohrmann
II. Institut für Experimentalphysik der Universität Hamburg, Germany

ISSN 0418-9833

NOTKESTRASSE 85 · 2 HAMBURG 52

DESY behält sich alle Rechte für den Fall der Schutzrechtserteilung und für die wirtschaftliche Verwertung der in diesem Bericht enthaltenen Informationen vor.

DESY reserves all rights for commercial use of information included in this report, especially in case of filing application for or grant of patents.

**To be sure that your preprints are promptly included in the
HIGH ENERGY PHYSICS INDEX ,
send them to the following address (if possible by air mail) :**

**DESY
Bibliothek
Notkestrasse 85
2 Hamburg 52
Germany**

DESY 83-053
June 1983

Search for Magnetic Monopoles in Electron-Positron
Collisions at 34 GeV c.m. Energy

P.Musset and M.Price

CERN, Genève, Switzerland, and

E.Lohrmann

II.Institut für Experimentalphysik, Universität Hamburg, Germany.

Abstract:

A search for heavily ionizing particles was carried out at the electron-positron storage ring PETRA with the help of Kapton detectors placed inside the vacuum pipe. The total integrated luminosity was 90 pb^{-1} at c.m. energies above 34 GeV. The search was sensitive to particles with magnetic charge from $1 \cdot (137/2)e$ to $5 \cdot (137/2)e$ and with masses up to $10 - 16 \text{ GeV}/c^2$. No such particle was found, leading to an upper limit of the production cross section of monopole-antimonopole pairs of $4 \cdot 10^{-38} \text{ cm}^2$ (95% c.l.).

Since Dirac's work on magnetic monopoles¹⁾ the question of the actual existence of such particles has remained an important experimental issue^{2,7)}. The magnetic charge g of a monopole is given by

$$e \cdot g = n \cdot \frac{1}{2} hc \quad (1)$$

e = elementary electric charge = charge of positron, $n = 1, 2, 3, \dots$

New arguments for the existence of monopoles come from modern gauge theories³⁾. These theories place the mass of monopoles well beyond the reach of present accelerators or storage rings. Nevertheless, considering the general difficulty of predicting masses of elementary particles, it is important to continue the search for monopoles at accelerators or storage rings with the largest available energies. The present search was also undertaken with the aim to be sensitive to rather large magnetic charges. If for example the "true" elementary electric charge was $1/3e$ instead of e , the smallest monopole charge according to equ.(1) could be as large as $g = 3 \cdot (\frac{137}{2})e$, and it was our aim to include this possibility in our search.

The experiment was carried out at the PETRA e^+e^- -storage ring of the DESY Laboratory. The detector consisted of 7 cylindrical layers of Kapton foils, 40 cm long, 18.4 cm diameter, 75 μ thick, placed at the SW interaction region inside the vacuum chamber of the storage ring. The solid angle covered was 85% of 4π , using a length of 30 cm of the detector.

The importance of having no material between the interaction point and the detector can be seen by considering the ionization rate of monopoles in matter, which for not too small values of the velocity β is approximately constant, given by

$$\frac{dE}{dx} \approx g^2 \cdot \left. \frac{dE}{dx} \right|_m = n^2 \cdot (137/2)^2 \cdot \left. \frac{dE}{dx} \right|_m \quad (2)$$

where $\left. \frac{dE}{dx} \right|_m$ = ionisation rate of a minimum ionizing particle.

For $n = 1$ one has $dE/dx \approx 9 \text{ GeV/g cm}^{-2}$, but if the "true" elementary electric charge were $e/3$, and this corresponds to $n = 3$, then $dE/dx \approx 80 \text{ GeV/g cm}^{-2}$. A monopole of this kind would be stopped by the PETRA vacuum chambers and therefore escape detection if the detector was placed outside.

Kapton was used as detector, because the ionization threshold above which tracks can be detected in this material is rather high, $dE/dx \geq 4 \text{ GeV/g cm}^{-2}$, and therefore it is insensitive to background. Also, because of its small degassing rate it can be exposed inside the vacuum chamber of the storage ring.

Monopoles would be detected in these foils by their high ionization rate. After exposure the foils are developed in 15% NaCl solution for 4 hours and any tracks would appear as holes in the plastic since the etchant preferentially attacks the region of the tracks. The foils are placed between pads soaked with electrolyte and, if a hole were present, electrical contact would be established between the two sides. A survey of this technique can be found in Ref. 4 Ad.7.

In order to check that kapton does not lose its detection properties at the conditions inside the storage ring a small piece of kapton, which had previously been exposed to heavy ions, was kept inside the PETRA vacuum chamber for 9 months. No change in its sensitivity was observed.

The foils were exposed in the period from February 1981 to January 1983. The total integrated luminosity was 1.9 pb^{-1} , 3.4 pb^{-1} , 89 pb^{-1} , 1.5 pb^{-1} at c.m. energies of 14 GeV, 22 GeV, 34.5 GeV and 38 GeV respectively.

After development no hole was found which could be interpreted as being produced by a singly ionizing particle.

In order to compute a limit for the production cross section, we assume the following mechanism for the production of a monopole antimonopole pair:

$$e^+e^- \rightarrow M^{(+)} M^{(-)}$$

The energy of the monopole is then equal to the beam energy of the storage ring, and therefore the velocity β of the monopole as function of its mass m is known.

The kapton foils are only sensitive for monopoles above a threshold velocity β_{th} , because the ionisation rate is a decreasing function of β . Using the work of Ref. (5) on the ionisation rate of monopoles, and a safe value of 6 GeV/g cm^{-2} for the sensitivity limit of kapton, one obtains $\beta_{th} = 0.6$ for $n = 1$ or $g = (137/2)e$. For larger values of the magnetic charge, the limit on β is given by the requirement that the monopole should penetrate at least one foil at the largest angle inside the acceptance of the detector. These values of β_{th} were again calculated using Ref. (5). The values of β_{th} , which depend on the magnetic charge $g = n \cdot (137/2)e$, imply upper limits for the mass of a monopole which can be detected. These mass limits are shown in Fig. 1 as a function of the magnetic charge. Also included in Fig. 1 are the limits obtained in a similar experiment with PEP at SLAC⁶⁾.

In order to compute an upper limit for the pair production cross section σ , we assumed isotropic production of the monopoles. Using only the luminosity above 34 GeV c.m. energy, one obtains

$$\sigma < 4 \cdot 10^{-38} \text{ cm}^{-2} \quad (95\% \text{ c.l.})$$

This value can be compared with the limit of $0.9 \cdot 10^{-36} \text{ cm}^{-2}$ obtained in a similar experiment at SLAC⁶⁾. It can also be compared with the QED point cross section σ_0 of producing a fermion antifermion pair with charge e

$$\sigma_0 = \frac{2\pi}{3} \frac{\alpha^2}{E_{c.m.}^2} \beta (3 - \beta^2)$$

which is $0.7 \cdot 10^{-34}$ for 34.6 GeV c.m. energy and $\beta = 1$.

Comparison with a recent experiment at the $p\bar{p}$ collider at CERN⁷⁾ is more difficult, because the production mechanism is different. A cross section limit about 10^{-32} cm^{-2} was given there, with mass limits of course larger than those obtained here.

Acknowledgements:


We greatly appreciate the support given to us by Prof. G.A.Voss and Prof. S.C.C.Ting, and the MARK J Group, who allowed us to expose our foils inside the PETRA vacuum and at the position of the MARK J detector. We thank Dr. J.Kouptsidis and the PETRA vacuum group for their help, and the PETRA machine group for the efficient operation of the machine. Furthermore we thank Dr. T.Walsh for his useful theoretical comments.

References:

- 1) P.A.M. Dirac, Proc. Royal Soc. of London, A-133 (1931) 60
- 2) For a recent review, see e.g. Review of Particle Properties, Phys.Lett. 111 B (1982) 111;
see also J.Bartelt et al. Phys.Rev.Lett.50 (1983) 655,
D.E.Groom et al., Phys.Rev.Lett. 50 (1983) 573 and references contained therein
- 3) G. 't Hooft, Nucl.Phys. 79B (1974) 276; 105B (1976) 538
A.M.Polyakov JETP Lett. 20 (1974) 194
For a Review see P.Langacker, Phys.Rep. 72 C (1981) 181
- 4) R.L.Fleischer, P.B.Price and R.M.Walker,
Nuclear Tracks in Solids (Univ. of Calif. Press, Berkeley 1975)
- 5) S.P.Ahlen, Phys.Rev. D 17 (1978) 229
S.P.Ahlen and K.Kinoshita, Phys.Rev. D 26 (1982) 2342
For small values of β , see S.Drell et al., Phys.Rev.Lett. 50 (1983) 644
- 6) K.Kinoshita, P.B. Price and D.Fryberger, Phys.Rev.Lett. 48 (1982) 77
- 7) B.Aubert et al., Phys.Lett. 120B (1983) 465.

Figure Caption:

Fig. 1 - The monopole mass m , for which this experiment was sensitive, is shown (shaded), g = magnetic charge g of the monopole, n = magnetic charge of monopole in multiples of the Dirac charge $(137/2)e$; also shown for comparison are the limits obtained by Ref. (6).



35790

Fig. 1