



DEUTSCHES ELEKTRONEN – SYNCHROTRON

DESY 91-037

May 1991



Quasi Hopf Quantum Symmetry in Quantum Theory

G. Mack, V. Schomerus

II. Institut für Theoretische Physik, Universität Hamburg

ISSN 0418-9833

NOTKESTRASSE 85 · D - 2000 HAMBURG 52

DESY behält sich alle Rechte für den Fall der Schutzrechtserteilung und für die wirtschaftliche Verwertung der in diesem Bericht enthaltenen Informationen vor.

DESY reserves all rights for commercial use of information included in this report, especially in case of filing application for or grant of patents.

**To be sure that your preprints are promptly included in the
HIGH ENERGY PHYSICS INDEX,
send them to the following (if possible by air mail):**

**DESY
Bibliothek
Notkestrasse 85
D-2000 Hamburg 52
Germany**

Quasi Hopf Quantum Symmetry in Quantum Theory

GERHARD MACK and VOLKER SCHOMERUS
II. Institut für Theoretische Physik, Universität Hamburg *

May 1, 1991

Abstract

In quantum theory, internal symmetries more general than groups are possible. We show that quasitriangular quasi Hopf algebras \mathcal{G}^* as introduced by Drinfeld [1] permit a consistent formulation of a transformation law of states in the physical Hilbert space \mathcal{H} , of invariance of the ground state, and of a transformation law of field operators which is consistent with local braid relations of field operators as proposed by Fröhlich [2]. All this remains true when Drinfeld's axioms are suitably weakened in order to build in truncated tensor products. Conversely, all the axioms of a weak quasitriangular quasi Hopf algebra are motivated from what physics demands of a symmetry. Unitarity requires in addition that \mathcal{G}^* admits a $*$ -operation with certain properties. Invariance properties of Greens functions follow from invariance of the ground state and covariance of field operators as usual. Covariant adjoints and covariant products of field operators can be defined. The R-matrix elements in the local braid relations are in general operators in \mathcal{H} . They are determined by the symmetry up to a phase factor. Quantum group algebras like $U_q(sl_2)$ with $|q| = 1$ are examples of symmetries with special properties. We show that a weak quasitriangular quasi Hopf algebra \mathcal{G}^* is canonically associated with $U_q(sl_2)$ if $q^p = -1$. We argue that these weak quasi Hopf algebras are the true symmetries of minimal conformal models. Their dual algebras \mathcal{G} ("functions on the group") are neither commutative nor associative.

1 Introduction

In quantum physics, states $|\psi\rangle$ are elements of a Hilbert space \mathcal{H} , and

- Elements of a symmetry act as transformations of physical states (i.e. as operators in \mathcal{H}) $|\psi\rangle \mapsto |\psi'\rangle$
- The transformation law is consistent with statistics
- There is a relation between spin and statistics. Bosons have spin $s = 0, 1, 2, \dots$ fermions have spin $s = \frac{1}{2}, \frac{3}{2}, \dots$, but other values for s are possible in 2 space dimensions, and are associated with braid group statistics.

*E-mail: 102MAC@DHHEDESYS3.BITNET

In this paper we assume that braid group statistics is implemented through local braid relations among the field operators which create and annihilate particles or excitations, as was proposed by Fröhlich [2]. We examine what symmetries are consistent with this assumption. They are called quantum symmetries and are not groups in general.

To find signs of (nonabelian) quantum symmetry in nature, experimental solid state physicists should look for degenerate multiplets of excitations with non-half-integer fractional spin in 2-dimensional quantum systems.

It appears that the possibility of quantum symmetries is special to quantum theory. In classical mechanics, symmetries are groups of transformations which act on phase space. The observables form a commutative algebra \mathcal{A} which consists of functions f on phase space; the Hamiltonian is one of these. In quantum mechanics, the points of phase space have gone away, but a (noncommutative) algebra of observables or field operators persists. It is surprising that it took 60 years to discover that quantum theory admits a larger class of symmetries than groups. Groups are topological spaces G whose points are the group elements a . When the points of phase space on which the group elements act have gone away, one does not need the group elements a any more either. It is said that "quantum theory may be regarded as mechanics on noncommutative phase space" [9] and that "quantum groups are the symmetries of noncommutative spaces" [10,11]. But actually they are only examples with special properties as we shall see.

The reader is invited to think of (second quantized) nonrelativistic many particle systems or of relativistic quantum field theories as examples of quantum mechanical systems.

When there is a symmetry group G , the Hilbert space of physical states carries a (unitary) representation of an algebra \mathcal{G}^* (the group algebra) which contains the group elements as distinguished elements that generate it. \mathcal{G}^* is the dual of the commutative algebra \mathcal{G} of functions on the group G , i.e. its elements $\phi \in \mathcal{G}^*$ are linear maps

$$\phi : \mathcal{G} \rightarrow \mathbb{C}.$$

Actually \mathcal{G} is a "bialgebra" $\mathcal{G} = \mathcal{G}^{-1}$ and this yields a product in \mathcal{G}^* known as the convolution product. The notion of invariance of the ground state or vacuum can be formulated in this language, without reference to group elements, and also the transformation law of field operators which create particles or excitations. This is all that is needed to formulate the notion of a symmetry and to extract from it the usual consequences.

It is therefore possible to admit as symmetry algebra \mathcal{G}^* the dual of a more general bialgebra \mathcal{G} than the commutative algebra of functions on a group. Quantum groups [7] are among these more general symmetries. They are associated with a special type of braid relations which involve solutions of the Yang-Baxter equations [2]. But this is not the most general possibility as will be seen. Nonassociative algebras \mathcal{G} are also admissible when they are the dual of a quasitriangular quasi Hopf algebra \mathcal{G}^* as introduced by Drinfeld [1]. We will exhibit the appropriate generalization of Fröhlich's local braid relations of field operators which is quasi-Hopf invariant. (It

¹A bialgebra is an algebra with a product and a coproduct, a unit element and a counit - see later

involves solutions of quasi-Yang-Baxter equations). But this is still not the most general possibility. We show that all the physical requirements, including in particular validity of local braid relations of field operators, remain satisfied when Drinfeld axioms are weakened in such a way that the truncated tensor products which have appeared in conformal field theory are accommodated. In this case we speak of a weak quasitriangular quasi Hopf algebra. Conversely, all the axioms of a weak quasitriangular quasi Hopf algebra will be motivated by properties which one wants quantum fields to have, and we believe therefore that this is the general algebraic structure of a symmetry in low dimensions.

In all cases a $*$ -operation with certain properties is needed in order to formulate unitarity. This is a somewhat subtle issue. We will explain in some detail why quantum group algebras $U_q(sl_2)$ with $|q| = 1$ are consistent with this requirement.

In this paper, \mathcal{G}^* will be regarded as the fundamental symmetry algebra. Emphasis is on its representations in the Hilbert space \mathcal{H} of physical states, as opposed to symmetries of Green's functions. The transformation law of field operators is the same as proposed by Buchholz, Mack and Todorov for quantum group algebras [4]. There is an alternative picture, proposed by Reshetikhin and Smirnov [15] where \mathcal{H} is regarded as a comodule for \mathcal{G} . The relation between both interpretations will be explained in section 8.

Models with quantum groups as symmetries were studied by many authors, [4,5, 13,14,15,16,17,19,20,21,22] and others. Integrable models, conformal field theory, and the massive sine Gordon model provide examples. Greens functions of an orbifold model with a quasi Hopf algebra as symmetry were examined by Dijkgraaf, Pasquier and Roche [3].

Some pedagogical material that is well known to experts is included in this paper for the benefit of the reader. But the construction of quasi-Hopf covariant products (5. 4) and adjoints (7. 2) of field operators, and the resulting quasi-Hopf invariant form of local braid relations of field operators (5. 7)ff are believed to be new, and so is the weaker form of the quasi Hopf axioms which permits to incorporate truncated tensor products, and the identification of what we believe to be the general algebraic structure of a symmetry in low dimensions.

Our results on the incorporation of truncated tensor products are in sections 6, 9. We show in section 9 that a weak quasitriangular quasi Hopf algebra \mathcal{G}^* is associated in a canonical way to every quantum group algebra $U_q(sl_2)$ with $q^p = -1$. The physical representations of $U_q(sl_2)$ are representations of \mathcal{G}^* , and the coproduct of \mathcal{G}^* yields the truncated tensor product of physical representations of $U_q(sl_2)$. Since this truncated tensor product is the "physical" tensor product in conformal models [19], it appears appropriate to regard \mathcal{G}^* rather than $U_q(sl_2)$ as the true symmetry of the conformal models. Our results on local braid relations imply for instance that an appropriate form of local braid relations for field operators holds in the *whole* physical Hilbert space of the conformal Ising model [5].² There is no need to introduce unphysical states to achieve this. This is in contrast with what happens when one interprets the full algebra $U_q(sl_2)$ as a symmetry, instead of the appropriate algebra $\mathcal{G}^* = U_q(sl_2)/\mathcal{J}$ with a non-coassociative coproduct as proposed here. The $U_q(sl_2)$ braid relations are

²The fields which satisfy these braid relations are precisely those constructed in [5]. The proof will be detailed in [6].

either only valid on a subspace of \mathcal{H} , or else one has to introduce unphysical states [5,30].³

The weak quasi-Hopf character of the symmetry affects not only the braid relations, but also the operator product expansions. It is the (nonassociative) covariant product of field operators which involves numerical Clebsch Gordan coefficients. Upon transition to (associative) ordinary products, these numerical Clebsch Gordan coefficients get multiplied with representation operators of the symmetry algebra in the physical Hilbert space.

2 Abstraction from symmetry groups

We begin with well known facts about group representations in quantum mechanics. The idea will be to reformulate them so that the group elements can be discarded in the end.

First we remind the reader that the Hilbert space \mathcal{H} carries a unitary representation of the group algebra \mathcal{G}^* . Then we will deduce the transformation law of field operators under \mathcal{G}^* . This will involve the commultiplication. The statement of invariance of the ground state will involve the counit. The transformation law of adjoints of field operators will involve the antipode. And the transformation law of states can be recovered from the transformation law of field operators and invariance of the ground state.

To state all this in more detail, we begin with the definition and representations of \mathcal{G}^* . Let G be a compact symmetry group of a quantum mechanical system. Then there will be a unitary operator $U(a)$ associated with every group element a . These operators act in the Hilbert space of physical states,

$$U(a) : \mathcal{H} \rightarrow \mathcal{H} .$$

Let \mathcal{G} be the commutative algebra of ("good") functions f on G ,

$$f : G \mapsto \mathbf{C} . \tag{2.1}$$

The product \cdot in \mathcal{G} is defined by pointwise multiplication, $f \cdot g(a) = f(a)g(a)$.

The dual \mathcal{G}^* of \mathcal{G} consists of linear maps

$$\phi : \mathcal{G} \mapsto \mathbf{C} . \tag{2.2}$$

The group elements $a \in G$ can be regarded as *special elements* of \mathcal{G}^* by virtue of the assignment

$$a(f) = f(a) \quad \text{for} \quad f \in \mathcal{G} . \tag{2.3}$$

Compact symmetry groups admit a volume element da known as Haar measure, which is uniquely determined by the requirement of invariance and normalization. Elements ϕ of \mathcal{G}^* are determined by generalized functions [35] Φ on G . We use the shorthand notation

$$\phi = \int_G da \Phi(a) a .$$

³The unphysical states may transform nontrivially under the ideal \mathcal{J} which we propose to factor out.

It follows that

$$\phi(f) = \int da \Phi(a) a(f) = \int da \Phi(a) f(a)$$

by eq.(2. 3). A group element a , regarded as an element of \mathcal{G}^* , is identified with the δ -function on G supported at a .

The product in \mathcal{G}^* is given by convolution of generalized functions, since

$$\phi_1 \phi_2 = \int da \int db \Phi_1(a) \Phi_2(b) ab = \int dc (\Phi_1 * \Phi_2)(c) \quad (2.4)$$

with

$$(\Phi_1 * \Phi_2)(c) = \int da \Phi_1(a) \Phi_2(a^{-1}c). \quad (2.5)$$

For group elements, regarded as elements of \mathcal{G}^* in the manner just explained, this definition agrees with the product definition in G .

The algebra \mathcal{G} admits a unit element 1, the constant function 1 on G . One defines a counit on \mathcal{G}^* , $\epsilon : \mathcal{G}^* \rightarrow \mathbb{C}$ which is a homomorphism of algebras and is defined by

$$\epsilon(\phi) = \phi(1) = \int da \Phi(a). \quad (2.6)$$

Later on we shall need also the coproduct Δ on \mathcal{G}^* . It is a linear map $\Delta : \mathcal{G}^* \rightarrow \mathcal{G}^* \otimes \mathcal{G}^*$ and is explicitly defined by reference to the product in \mathcal{G} , viz.

$$\Delta(\phi)(f, g) = \phi(f \cdot g) \quad (2.7)$$

When \mathcal{F} is a space of functions of one variable, then $\mathcal{F} \otimes \mathcal{F}$ consists of functions of two variables. Therefore elements of $\mathcal{G}^* \otimes \mathcal{G}^*$ may be regarded as maps of pairs of functions (f, g) into complex numbers. $\mathcal{G}^* \otimes \mathcal{G}^*$ is an algebra with product $(\xi_1 \otimes \xi_2)(\eta_1 \otimes \eta_2) = \xi_1 \eta_1 \otimes \xi_2 \eta_2$, and Δ is in fact a homomorphism of algebras.

The representation of G induces a representation of the algebra \mathcal{G}^* by operators

$$U(\phi) : \mathcal{H} \rightarrow \mathcal{H} \quad (2.8)$$

in the Hilbert space of physical states, viz.

$$U(\phi) = \int_G da \Phi(a) U(a). \quad (2.9)$$

The representation property holds,

$$U(\phi_1 \phi_2) = U(\phi_1) U(\phi_2). \quad (2.10)$$

Unitarity of the representation translates into

$$U(\phi^*) = U(\phi)^*. \quad (2.11)$$

The action of the $*$ -operations on elements $\phi \in \mathcal{G}^*$ is defined by $a^* = a^{-1}$, and more generally by $\phi^* = \int da \overline{\Phi(a)} a^{-1}$. Finally, G -invariance of the ground state $|0\rangle$ translates into

$$U(\phi)|0\rangle = |0\rangle \epsilon(\phi). \quad (2.12)$$

We ended up with statements which involve only the algebra \mathcal{G}^* and the $*$ -operation and counit on it, but make no reference to group elements. Summing

up, the Hilbert space carries a unitary representation of the algebra \mathcal{G}^* . Unitarity is expressed through eq.(2. 11) and invariance of the ground state through eq.(2. 12).

Let us now suppose that the Hilbert space of physical states is created from the ground state $|0\rangle$ by application of field operators $\Psi_i^{\dagger}(\mathbf{r}, t)$ which create particles or excitations. I, i are some quantum numbers which distinguish these. Later on, I will specify the transformation law (isospin, for instance) while i distinguishes members of a multiplet. Let us suppose that these field operators transform covariantly under the symmetry group G . This means that there exist finite dimensional representations of G by matrices $\tau^I(a)$ such that

$$U(a)\Psi_i^{\dagger} = \Psi_k^{\dagger} \tau_{ki}^I(a) U(a). \quad (2.13)$$

Summation over repeated indices will be understood throughout. We will neglect to write arguments (\mathbf{r}, t) of field operators when their values do not matter. Eq.(2. 13) says that Ψ^{\dagger} is a tensor operator.

We will generalize this formula to arbitrary elements $\phi \in \mathcal{G}^*$. The resulting formula will involve the coproduct Δ . By definition $\Delta(\phi)$ is an element of $\mathcal{G}^* \otimes \mathcal{G}^*$ and is therefore of the following form. To every ϕ there exist sequences of elements ϕ_p^1 and ϕ_p^2 of elements of \mathcal{G}^* such that

$$\Delta(\phi) = \sum_p \phi_p^1 \otimes \phi_p^2. \quad (2.14)$$

If ϕ is a group element, then there exists only one term in the sum. We may admit integrals in place of sums \sum_p and use our shorthand notation to write an explicit general formula,

$$\Delta(a) = a \otimes a \text{ for } a \in G, \quad \Delta(\phi) = \int da \Phi(a) a \otimes a. \quad (2.15)$$

The covariance property generalizes to

$$U(\phi)\Psi_i^{\dagger} = \sum_p \Psi_m^{\dagger} \tau_{mi}^I(\phi_p^1) U(\phi_p^2). \quad (2.16)$$

$U(\phi)$ was defined in eq.(2. 9) and $\tau(\phi)$ is defined in the same way.

Let us check eq.(2. 16) with the use of the explicit formula (2. 15). The right hand side of eq.(2. 16) will read

$$\int da \Phi(a) \Psi_m^{\dagger} \tau_{mi}^I(a) U(a).$$

By virtue of eq.(2. 13) this is indeed equal to

$$U(\phi)\Psi_i^{\dagger} = \int da \Phi(a) U(a) \Psi_i^{\dagger}.$$

Let us now determine the transformation law of the special states

$$|I, i\rangle = |\Psi_i^{\dagger}|0\rangle$$

that follow from this. Denoting the identity map by id , coproduct and counit have the property that

$$(id \otimes \epsilon)\Delta = id. \quad (2.17)$$

This means that validity of eq.(2. 14) implies

$$\sum_p \phi_p^1 \otimes \epsilon(\phi_p^2) = \phi. \quad (2.18)$$

Since $\epsilon(\phi)$ is a complex number, the symbol \otimes is redundant in this formula. It stands for ordinary multiplication with a complex number.

Using eqs.(2. 16),(2. 12), and (2. 18) in turn, we find

$$U(\phi)\Psi_i^I|0\rangle = \sum_p \Psi_m^I \tau_{mi}^I(\phi_p^1)U(\phi_p^2)|0\rangle \quad (2.19)$$

$$= \sum_p \Psi_m^I \tau_{mi}^I(\phi_p^1)|0\rangle \epsilon(\phi_p^2) \quad (2.20)$$

$$= \Psi_m^I|0\rangle > \tau_{mi}^I(\phi). \quad (2.21)$$

This shows that the "1-excitation states" $|I, i\rangle = \Psi_i^I|0\rangle >$ transform according to the representation τ^I . The derivation of this formula used covariance of field operators under \mathcal{G}^* , invariance (2. 12) of the ground state $|0\rangle >$, and property (2. 17) of the counit. All of these will be retained in the case of more general symmetries later on. Similarly, the transformation law of states

$$|I_1, i_1, \dots, I_n, i_n\rangle = \Psi_{i_1 \dots i_n}^{I_1 \dots I_n}|0\rangle >$$

is determined. Applying $U(\phi)$ to such a state, one shifts the representation operator $U(\cdot)$ through the fields one by one from left to right, using covariance property (2. 16). Finally, an operator $U(\cdot)$ stands next to the ground state, so that one can convert it into a c-number using invariance (2. 12) of the ground state.

As a result one finds the well known result that the states $|I_1, i_1, \dots, I_n, i_n\rangle >$ transform according to the tensor product

$$(\tau^{I_1} \otimes \tau^{I_2} \otimes \dots \otimes \tau^{I_{n-1}} \otimes \tau^{I_n} \dots) \quad (2.22)$$

of representations under the group G . The brackets in this formula are actually unnecessary because the tensor product is associative. But if one does not make use of this fact in the derivation, eq.(2. 22) is what one gets. To generalize this formula, one observes that the tensor product of group representations can be defined in an abstract way by use of the coproduct Δ , as follows.

We distinguish in notation between the standard tensor product \otimes of matrices, and more generally of algebras and of their elements, and the tensor product \otimes of representations. If τ^I and τ^J are representations of \mathcal{G}^* by matrices $\tau^I(\xi)$ etc., then the standard tensor product $\tau^I \otimes \tau^J(A)$ is defined for elements $A \in \mathcal{G}^* \otimes \mathcal{G}^*$ such as $A = \xi \otimes \eta$ in the obvious way, $(\tau^I \otimes \tau^J)(\xi \otimes \eta) = \tau^I(\xi) \otimes \tau^J(\eta)$, where the right hand side involves a tensor product of matrices. By contrast, the tensor product of representations $(\tau^I \otimes \tau^J)(\xi)$ is going to be defined for arguments $\phi \in \mathcal{G}^*$ through the formula

$$(\tau^I \otimes \tau^J)(\phi) = (\tau^I \otimes \tau^J)(\Delta(\phi)) = \sum_p \tau^I(\phi_p^1) \otimes \tau^J(\phi_p^2). \quad (2.23)$$

Indeed, for group elements $\phi = a$,

$$(\tau^I \otimes \tau^J)(\Delta(a)) = (\tau^I \otimes \tau^J)(a \otimes a) = \tau^I(a) \otimes \tau^J(a)$$

by eq.(2. 15). Retaining definition (2. 23) of the tensor product of representations for the group algebra \mathcal{G}^* , the tensor product transformation law of states continues to hold for arbitrary elements $\phi \in \mathcal{G}^*$.

By taking the adjoint of the transformation law (2. 13) of field operators under group elements, one finds that the adjoint of a field operator Ψ^I transforms according to the contragredient representation $\bar{\tau}^I$ of τ^I . This generalizes to \mathcal{G}^* in the same way as before. The contragredient representation is defined by

$$\bar{\tau}(\phi) = {}^t\tau(\mathcal{S}(\phi))$$

where \mathcal{S} is the antipodal map. Its action on group elements is given by $\mathcal{S}(a) = a^{-1}$. This extends to an antiautomorphism of the group algebra in the obvious way, viz. $\mathcal{S}(\phi) = \int_G d\alpha \bar{\Phi}(\alpha)a^{-1}$. The transformation law of the adjoint field operator can be written as

$$U(\phi)\bar{\Psi}_i^I = \sum_p \Psi_m^I \bar{\tau}_{mi}(\phi_p^2)U(\phi_p^1). \quad (2.24)$$

The interchange of ϕ_p^1 and ϕ_p^2 compared to the transformation law (2. 16) of field operators is immaterial here. This is true because the coproduct is cocommutative in the case of group algebras. Indeed, if P interchanges factors, $P(a \otimes b) = b \otimes a$, then $\Delta(a) = a \otimes a = P\Delta(a)$ for $a \in G$, and this extends to \mathcal{G}^* .

However, if one wants to derive the transformation law of adjoints from the transformation law (2. 16) of fields using abstract properties of the antipode but not cocommutativity of the coproduct, eq.(2. 24) is what one gets, see Appendix A. This formula was first obtained for the quantum group algebra $U_q(sl_2)$ with $|q| = 1$ in [5].

3 Quantum symmetry

Abstracting from the example of a symmetry group as discussed in the last section, we find conditions on a symmetry algebra \mathcal{G}^* which permit to spell out features that are commonly associated with symmetries in quantum theory,

- A transformation law of states in the Hilbert space \mathcal{H} which determines a unitary representation of \mathcal{G}^* in \mathcal{H} ,
- Covariance properties of field operators,
- Invariance of the ground state.

Locality and statistics will be discussed in the next section. The conditions are as follows:

1. \mathcal{G}^* is a $*$ -algebra with unit element ϵ .
2. A comultiplication $\Delta : \mathcal{G}^* \rightarrow \mathcal{G}^* \otimes \mathcal{G}^*$ is defined which is a $*$ -homomorphism of algebras, i.e. a linear map with properties as follows:

$$\Delta(\xi\eta) = \Delta(\xi)\Delta(\eta), \quad (3.1)$$

$$\Delta(\xi^*) = \Delta(\xi)^*. \quad (3.2)$$

3. There exists a counit $\epsilon : \mathcal{G}^* \rightarrow \mathbf{C}$ which is a *-homomorphism of algebras subject to the extra condition

$$(id \otimes \epsilon)\Delta = id = (\epsilon \otimes id)\Delta. \quad (3.3)$$

The *-homomorphism property means that ϵ is a linear map with $\epsilon(e) = 1$ and

$$\epsilon(\xi\eta) = \epsilon(\xi)\epsilon(\eta), \quad (3.4)$$

$$\epsilon(\xi^*) = \overline{\epsilon(\xi)}. \quad (3.5)$$

4. There exists an antipodal map $S : \mathcal{G}^* \rightarrow \mathcal{G}^*$ which is a *-antiautomorphism, i.e. a linear map obeying

$$S(\xi\eta) = S(\eta)S(\xi), \quad (3.6)$$

$$S(\xi^*) = S(\xi)^*. \quad (3.7)$$

It should obey the further conditions

$$m_*((S \otimes id \otimes id)(id \otimes \Delta)\Delta(\phi)) = e \otimes \phi, \quad (3.8)$$

$$m'_*((id \otimes S \otimes id)(id \otimes \Delta)\Delta(\phi)) = \phi \otimes e, \quad (3.9)$$

with multiplication operators defined by

$$m_*(\xi \otimes \eta \otimes \zeta) = \eta\xi \otimes \zeta, \quad m'_*(\xi \otimes \eta \otimes \zeta) = \zeta \otimes \eta\xi.$$

Remark: Note that we do not require that $\Delta(e) = e \otimes e$. Instead, $\Delta(e)$ may be a nontrivial projector which commutes with all $\Delta(\xi)$, $\xi \in \mathcal{G}^*$. In this case representation operators $(\pi^I \otimes \pi^J)(\xi) = \pi^I \otimes \pi^J(\Delta(\xi))$ will annihilate a proper subspace of the representation space $V^I \otimes V^J$ for some I, J and all $\xi \in \mathcal{G}^*$. This is so because $\Delta(\xi) = \Delta(e\xi) = \Delta(e)\Delta(\xi)$. In this way Drinfelds remark [1], which suggests that minimal conformal models cannot have quasi-Hopf algebras as symmetries, will be amended - see later

The standard choice of the *-operation on $\mathcal{G}^* \otimes \mathcal{G}^*$ is

$$(\xi \otimes \eta)^* = \xi^* \otimes \eta^*, \quad (3.10)$$

but it is also permissible to adopt the alternative choice

$$(\xi \otimes \eta)^* = \eta^* \otimes \xi^*, \quad (3.11)$$

and to interpret condition (3. 2) accordingly. The interchange of factors matches with the formula for adjoints of operators, $(AB)^* = B^*A^*$.

For the quantum group algebras $U_q(sl_2)$ with q a root of unity which were proposed as symmetries of conformal field theory models, the above conditions will be fulfilled if we adopt the second choice (3. 11) of *-operation on $\mathcal{G}^* \otimes \mathcal{G}^*$. The consequences of this choice will be discussed later on.

Given an algebra \mathcal{G}^* with the structure described above, it may be called a symmetry if the Hilbert space of physical states carries a unitary representation U of \mathcal{G}^* ,

the ground state $|0\rangle$ is invariant, and field operators transform covariantly. This means that for all $\phi \in \mathcal{G}^*$ we should have

$$U(\phi^*) = U(\phi)^*, \quad (3.12)$$

$$U(\phi)|0\rangle = |0\rangle \epsilon(\phi), \quad (3.13)$$

$$U(\phi)\Psi_i^j = \sum_p \Psi_{m_i}^{j'}(\phi_p^1)U(\phi_p^2), \quad (3.14)$$

$$\text{if } \Delta(\phi) = \sum_p \phi_p^1 \otimes \phi_p^2. \quad (3.15)$$

From covariance properties of field operators and invariance of the ground state one obtains invariance properties of Greens functions in the standard way, see below.

Neither cocommutativity nor coassociativity of the coproduct were required.

The coproduct Δ is said to be coassociative if

$$(id \otimes \Delta)\Delta = (\Delta \otimes id)\Delta. \quad (3.16)$$

If the comultiplication is coassociative, then the properties (3. 8, 3. 9) are implied by the requirement

$$m'_*(S \otimes id)\Delta(\phi) = m'_*(id \otimes S)\Delta(\phi) = \epsilon(\phi)e \quad (3.17)$$

for all $\phi \in \mathcal{G}^*$. The multiplication operator m' is defined by $m'(\xi \otimes \eta) = \eta\xi$.

Given one comultiplication Δ ,

$$\Delta(\phi) = \sum_p \phi_p^1 \otimes \phi_p^2, \quad (3.18)$$

there exists another one

$$\Delta'(\phi) = \sum_p \phi_p^2 \otimes \phi_p^1, \quad (3.19)$$

The comultiplication is said to be cocommutative if $\Delta = \Delta'$.

Compact symmetry groups yield group algebras \mathcal{G}^* with a comultiplication that is coassociative and cocommutative. Hopf algebras \mathcal{G}^* are examples which fulfill all our conditions except possibly those involving the *-operation. In a Hopf algebra, the comultiplication is coassociative but not necessarily cocommutative. In a quasi Hopf algebra as introduced by Drinfeld the comultiplication is not coassociative but only "quasi-coassociative". This and further properties of quasitriangular quasi Hopf algebras which are motivated by locality will be discussed in the next section.

When comultiplication is not cocommutative, there exist actually two possible comultiplications Δ and Δ' , and therefore also two possible covariant field transformation laws, and two possible tensor products of representations \otimes and \otimes' .

If we adhere to the definition (3. 11) of the *-operation on $\mathcal{G}^* \otimes \mathcal{G}^*$, as is necessary for $U_q(sl_2)$ with $|q| = 1$ for instance, then validity of the one covariant field transformation law (3. 14) for the field Ψ^j will imply that its adjoint Ψ^{j*} will transform the other way,

$$U(\phi)\Psi_i^{j*} = \sum_p \Psi_{m_i}^{j'}(\phi_p^2)U(\phi_p^1). \quad (3.20)$$

This involves the contragredient representation,

$$\bar{\tau}(\phi) = \tau(S(\phi)) \quad (3.21)$$

where i stands for the transpose. In addition to that, the role of ϕ_i^j and ϕ_j^i got interchanged compared to (3. 14). This means that Δ got replaced by Δ' . The necessary computations to establish this formula will be presented in Appendix A. We shall see in the next section that the presence of an R -element in $\mathcal{G}^* \otimes \mathcal{G}^*$ will permit to define a covariant adjoint Ψ^j which transforms in the same way as Ψ^i , apart from the appearance of the contragredient representation $\bar{\tau}^j$ in place of τ^j .

Let us now turn to the transformation law of states

$$|I_1, i_1, \dots, I_n, i_n\rangle = \Psi_{i_1}^{j_1} \dots \Psi_{i_n}^{j_n} |0\rangle .$$

It is derived in the same way as described at the end of the last section. This derivation makes no reference to group elements. One uses the covariance property of field operators (3. 14) to shift representation operators $U(\cdot)$ through field operators from left to right until they act on the ground state $|0\rangle$. Then invariance of the ground eq.(2. 12) is used to convert the operators $U(\cdot)$ into numbers. As a result one finds that "1-excitation states" $|I, i\rangle = \Psi_i^j |0\rangle$ transform according to representation τ^j of \mathcal{G}^* ,

$$U(\phi)|I, i\rangle = \sum_k |I, k\rangle \tau_{ki}^j(\phi) . \quad (3.22)$$

More generally, the states transform according to tensor products of representations

$$U(\phi)|I_1, i_1, \dots, I_n, i_n\rangle = |I_1, k_1, \dots, I_n, k_n\rangle (\tau^{j_1})_{k_1 i_1} \dots (\tau^{j_n})_{k_n i_n} (\phi) \quad (3.23)$$

Summation over k_1, \dots, k_n is understood. The tensor product of representations \otimes is defined with the help of the coproduct as before,

$$(\tau^j \otimes \tau^j)(\phi) = (\tau^j \otimes \tau^j)(\Delta(\phi)) . \quad (3.24)$$

The standard tensor product \otimes of matrices is associative by definition. But the tensor product \otimes of representations is not associative unless the coproduct is coassociative. It is associative for Hopf algebras \mathcal{G}^* (quantum group algebras), though.

The representations τ^j furnish linear functions τ_{ik}^j on \mathcal{G}^* . Such linear functions may be regarded as elements of the dual \mathcal{G} of \mathcal{G}^* . The coproduct Δ in \mathcal{G}^* yields a product \cdot in \mathcal{G} such that

$$(\tau^j \otimes \tau^j)_{ij,kl}(\phi) = (\tau_{ik}^j \cdot \tau_{jl}^j)(\phi) . \quad (3.25)$$

The details of this construction shall be recalled in the section 8. In general this product is neither commutative nor associative. In this new notation, the transformation law (3. 23) can be written in the equivalent form

$$U(\phi)|I_1, i_1, \dots, I_n, i_n\rangle = |I_1, k_1, \dots, I_n, k_n\rangle (\tau_{k_1 i_1}^{j_1}) \dots (\tau_{k_n i_n}^{j_n}) (\phi) . \quad (3.26)$$

Let us finally turn to the invariance properties of Greens functions. Taking the adjoint of the eq.(2. 12) which expresses invariance of the ground state, and using the unitarity conditions on U and ϵ we find that

$$\langle 0|U(\phi) = \bar{\epsilon}(\phi) \langle 0|$$

It follows that

$$\bar{\epsilon}(\phi) \langle 0|\Psi_{i_1}^{j_1} \dots \Psi_{i_n}^{j_n}|0\rangle = \langle 0|U(\phi)\Psi_{i_1}^{j_1} \dots \Psi_{i_n}^{j_n}|0\rangle .$$

Inserting the transformation law of states derived before, one obtains the invariance property

$$\bar{\epsilon}(\phi) \langle 0|\Psi_{i_1}^{j_1} \dots \Psi_{i_n}^{j_n}|0\rangle = \langle 0|\Psi_{m_1}^{j_1} \dots \Psi_{m_n}^{j_n}|0\rangle (\tau_{m_1 i_1}^{j_1}) \dots (\tau_{m_n i_n}^{j_n}) (\phi) . \quad (3.27)$$

This involves a tensor product of representations, written in the compact notation which involves the noncommutative product \cdot in \mathcal{G} , see eqs.(3. 25) above.

In special cases it is known that the most general Green function which satisfies these invariance properties is a weighted sum of products of suitably defined Clebsch Gordan coefficients for tensor products of irreducible representations, much as in the case of symmetry groups. This is true for $U_q(sl_2)$ for instance when q is not a root of unity [24]. In general one will need information about the decomposition of the representation U into irreducibles to obtain such a representation. This is why one prefers semisimple symmetry algebras whose representations are all fully reducible.

4 Locality

Local Fermi or Bose fields obey anticommutation or commutation relations

$$\Psi_i^j(\mathbf{r}_1, t)\Psi_j^i(\mathbf{r}_2, t) = \pm \Psi_j^i(\mathbf{r}_2, t)\Psi_i^j(\mathbf{r}_1, t) \text{ if } \mathbf{r}_1 \neq \mathbf{r}_2 . \quad (4.1)$$

In the case of a Lorentz covariant theory this has to hold in an arbitrary reference frame and implies local (anti-)commutation relations at arbitrary spacelike distances.

It has been discussed in detail in the literature that there are more general possibilities in 1 and 2 space dimensions [2,26,27]. When one wants to interchange positions of two particles or arguments of wave functions \mathbf{r}_1 and $\mathbf{r}_2 \neq \mathbf{r}_1$ one may encounter two different situations which we distinguish symbolically as $\mathbf{r}_2 > \mathbf{r}_1$ and $\mathbf{r}_2 < \mathbf{r}_1$.⁴ These two different situations may require two different interchanging operations σ_1 and σ_1^{-1} . More generally, when there are N particles there may be two operations σ_i and σ_i^{-1} which interchange particle i and $i+1$. These operations must satisfy the Artin relations

$$\sigma_i \sigma_k = \sigma_k \sigma_i \text{ if } |k-i| \geq 2, \quad \sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1} . \quad (4.2)$$

As a result, $\sigma_1 \dots \sigma_{n-1}$ generate the braid group B_n on n threads. Using the graphical notation of figure 1a, a general element

$$b = \sigma_{i_1}^{\pm 1} \dots \sigma_{i_n}^{\pm 1}$$

can be represented by a braid, as in figure 1b, and multiplication of group elements corresponds with juxtaposition of braids. Braids which can be continuously deformed

⁴In 1 space dimensions $\mathbf{r}_2 > \mathbf{r}_1$ if the real coordinate r_2 is bigger than r_1 . In 2 space dimensions, wave functions may be multivalued, i.e. have cuts, and the two possibilities differ in the relative positions to the cut - see [27]

into each other (holding ends fixed) with the help of operations as in figure 2 are regarded to be the same. The possible deformations express the Artin relations and the further relation $\sigma_i^{-1} \sigma_i = \sigma_i \sigma_i^{-1} = 1$. As a result, the braid shown in figure 1b is the product of a braid b (above the dashed line) with its inverse, and is thus equal to the identity braid (parallel lines). Every braid determines a permutation in the obvious way, but σ_i and σ_i^{-1} determine the same permutation $i \rightarrow i+1$. Therefore the permutation group S_n is a factor group of the braid group B_n . Bose Fermi statistics can be regarded as a special case of braid group statistics with $\sigma_i^{-1} = \sigma_i$.

Fröhlich proposed that field operators should obey local braid relations of the following form in the case of braid group statistics,

$$\Psi_i^j(\mathbf{r}_1, t) \Psi_j^k(\mathbf{r}_2, t) = \Psi_m^j(\mathbf{r}_2, t) \Psi_n^k(\mathbf{r}_1, t) \mathcal{R}_{nm,ij}^{JK} \text{ if } \mathbf{r}_2 > \mathbf{r}_1, \quad (4.3)$$

and a similar relation involving \mathcal{R}^{JK} if $\mathbf{r}_2 < \mathbf{r}_1$. The two matrices are related as follows. Denoting by $\tilde{\mathcal{R}}$ the matrix obtained from \mathcal{R} by interchange of the first two indices n, m , the two matrices $\mathcal{R}^>$ and $\mathcal{R}^<$ are related by

$$\tilde{\mathcal{R}}^{JK} = (\mathcal{R}^{JK})^{-1}. \quad (4.4)$$

The local braid relations for $\mathbf{r}_2 < \mathbf{r}_1$ follow from those for $\mathbf{r}_2 > \mathbf{r}_1$ by inversion using (4.4) and will therefore not lead to independent consistency conditions.

Fröhlich proposed that local braid relations (4.3) should hold with a numerical matrix \mathcal{R} . In this case, consistency requires that matrices \mathcal{R} should satisfy the Yang-Baxter equations. This assures that \mathcal{R} yields a representation of the braid group. The Yang-Baxter equations are a translation of the Artin relations - see [2].

As we shall see, quasi-Hopf symmetry leads in a natural way to \mathcal{R} -matrices which are representation operators of certain elements of \mathcal{G}^* . Only in the special case of Hopf symmetry (quantum groups) can they be expected to be numerical matrices.

It is nevertheless instructive to examine first the conditions that come from the requirement that both sides of the local braid relations transform in the same way under the symmetry, assuming that \mathcal{R} is a numerical matrix. The calculation is presented in Appendix B. The result is as follows.

Suppose that there exists an element R in the algebra,

$$R = \sum_a r_a^1 \otimes r_a^2 \in \mathcal{G}^* \otimes \mathcal{G}^* \quad (4.5)$$

such that

$$\mathcal{R}_{tu,sv}^{JK} = c(\tau^J \otimes \tau^K)_{tu,sv}(R) = c \sum_a \tau_a^J(\tau_a^1) \tau_a^K(\tau_a^2), \quad (4.6)$$

where c is a numerical phase factor. Then the following condition is sufficient.

$$(R \otimes e) \cdot (id \otimes \Delta) \Delta(\phi) = (P_{12}(id \otimes \Delta) \Delta(\phi)) \cdot (R \otimes e), \quad (4.7)$$

where $P_{12}(a_1 \otimes a_2 \otimes a_3) = a_2 \otimes a_1 \otimes a_3$ and we wrote \cdot for the product in $\mathcal{G}^* \otimes \mathcal{G}^* \otimes \mathcal{G}^*$ for the sake of clarity.

In a Hopf algebra, comultiplication is coassociative, so that eq.(3.16) holds. Then eq.(4.7) is fulfilled if

$(R \otimes e) \cdot (\Delta \otimes id)(\xi \otimes \eta) = (P_{12}(\Delta \otimes id)(\xi \otimes \eta))(R \otimes e)$ for all $\xi \in \mathcal{G}^*$.
This is equivalent to

$$R \Delta(\xi) = \Delta'(e) R \text{ for all } \xi \in \mathcal{G}^*, \quad (4.8)$$

where Δ' differs from Δ by the interchange of factors cp. eq.(3.19).

Hopf algebras with an element $R \in \mathcal{G}^* \otimes \mathcal{G}^*$ satisfying eq.(4.8) are called quasi-triangular, or quantum group algebras. $U_q(sl_2)$ are examples.

In conclusion we see that quantum group symmetry is consistent with local braid relations with a numerical \mathcal{R} -matrix that satisfies the Yang-Baxter equations. It is a special case of quantum symmetry in which the comultiplication is coassociative.

By Einstein causality, observables have to satisfy ordinary Bose commutation relations. A nontrivial transformation law under a quantum symmetry which does not come from a group is consistent with an appropriate form of local braid relations but not in general with Bose commutation relations. It follows that observables should be invariant under quantum symmetries. (Global) symmetries which leave the observables invariant are known as *gauge symmetries* (of first kind). They occur in theories with superselection sectors [5,25,26,28]. Fields with a nontrivial transformation law under such a gauge symmetry make transitions between different superselection sectors.

5 Quasi Hopf symmetry

The comultiplication Δ is said to be quasi-coassociative if there exists an invertible element $\varphi \in \mathcal{G}^* \otimes \mathcal{G}^* \otimes \mathcal{G}^*$ such that

$$\varphi(\Delta \otimes id) \Delta(\xi) = (id \otimes \Delta) \Delta(\xi) \varphi \text{ for all } \xi \in \mathcal{G}^*. \quad (5.1)$$

If the comultiplication is not coassociative, then the tensor product of representations of \mathcal{G}^* is not associative. However, quasiassociativity of multiplication implies that representations

$$(\pi^1 \otimes \pi^2) \otimes \pi^3 \text{ and } \pi^1 \otimes (\pi^2 \otimes \pi^3) \quad (5.2)$$

are equivalent.

Quasitriangular quasi-Hopf algebras as introduced by Drinfeld [1] are bialgebras in which the multiplication is quasi-coassociative. They possess a counit, antipode and R-element $R \in \mathcal{G}^* \otimes \mathcal{G}^*$ which satisfy certain properties. We shall see that symmetry under a quasitriangular Hopf algebra is consistent with an appropriate form of local braid relations.

If comultiplication is not coassociative, then the covariant field transformation law (3.14) does not imply a covariant transformation law of products of fields. But coassociativity helps to define a new product of field operators which is covariant. Let

$$\varphi = \sum_{\sigma} \varphi_{\sigma}^1 \otimes \varphi_{\sigma}^2 \otimes \varphi_{\sigma}^3, \quad \varphi_{\sigma}^i \in \mathcal{G}^*. \quad (5.3)$$

We define

$$(\Psi^I \times \Psi^J)_{ij} = \sum_{\sigma} \Psi_m^I \Psi_n^J \tau_{mi}(\varphi_{\sigma}^1) \tau_{nj}(\varphi_{\sigma}^2) U(\varphi_{\sigma}^3). \quad (5.4)$$

Straightforward computation shows that $\Psi^I \times \Psi^J$ transforms according to the tensor product $\tau^I \otimes \tau^J$ of representations of \mathcal{G}^* in the sense that it obeys transformation law (3. 14).

The product \times is not associative in general. But if φ satisfies Drinfelds relation

$$(id \otimes id \otimes \Delta)(\varphi)(\Delta \otimes id \otimes id)(\varphi) = (\varepsilon \otimes \varphi)(id \otimes \Delta \otimes id)(\varphi \otimes \varepsilon). \quad (5.5)$$

then the product \times is quasi-associative in the sense that

$$((\Psi^I \times \Psi^J) \times \Psi^K)_{ijk} = (\Psi^I \times (\Psi^J \times \Psi^K))_{i'j'k'}(\tau^I \otimes \tau^J \otimes \tau^K)_{i''j''k''}(\varphi). \quad (5.6)$$

Explicitly

$$(\tau^I \otimes \tau^J \otimes \tau^K)_{i'j'k'}(\varphi) = \sum_{\sigma} \tau_{i'i}^I(\varphi^1) \tau_{j'j}^J(\varphi^2) \tau_{k'k}^K(\varphi^3).$$

Next we turn to the discussion of local braid relations for field operators. It is consistent to demand that

$$(\Psi^I \times \Psi^J)_{ij} = (\Psi^J \times \Psi^I)_{j'i'} \bar{\mathcal{R}}_{\tau^I, \tau^J, ij} \quad (5.7)$$

if the arguments of the field operators, which we neglect to write, satisfy conditions as in eq.(4. 3). Because of the covariance properties of field operators, both sides will transform in the same way if there exists an element $R \in \mathcal{G}^* \otimes \mathcal{G}^*$ satisfying the relation

$$\Delta^I(\eta)R = R\Delta(\eta) \text{ for all } \eta \in \mathcal{G}^* \quad (5.8)$$

and \bar{R} in eq.(5. 7) is a numerical matrix which is obtained from R , up to a phase factor c which may depend on I and J , as

$$\bar{R} = c(\tau^I \otimes \tau^J)(R). \quad (5.9)$$

If $R = \sum \tau_a^I \otimes \tau_a^J$ then this reads explicitly as follows

$$\bar{\mathcal{R}}_{\tau^I, \tau^J, ij} = c^{IJ} \sum_{\alpha} \tau_{i'i}^I(\tau_a^I) \tau_{j'j}^J(\tau_a^J). \quad (5.10)$$

We neglect to write superscripts $I, J >$ on $\bar{\mathcal{R}}$.

Later on we shall wish to relax invertibility of φ . But for now, we adopt the hypothesis that φ has an inverse which we write as

$$\varphi^{-1} = \sum_{\sigma} \xi_{\sigma}^I \otimes \xi_{\sigma}^J \otimes \xi_{\sigma}^K. \quad (5.11)$$

It follows that ordinary products of field operators can be recovered from covariant ones,

$$\Psi_i^I \Psi_j^J = \sum_{\sigma} (\Psi^I \times \Psi^J)_{i'j'} \tau_{i'i}^I(\xi_{\sigma}^I) \tau_{j'j}^J(\xi_{\sigma}^J) U(\xi_{\sigma}^K). \quad (5.12)$$

Inserting the local braid relations in the form (5. 7) and using the definition of the covariant product once again, one finds that Fröhlich's local braid relations (4. 3) hold. But the R -matrix elements $\bar{\mathcal{R}}_{\tau^I, \tau^J, ij}$ are not numbers any more but operators in Hilbert space,

$$\bar{\mathcal{R}}_{\tau^I, \tau^J, ij} = c^{IJ} \sum_{\tau, \sigma, \rho} \tau_{i'i}^I(\varphi_a^1 \xi_{\tau}^1) \tau_{j'j}^J(\varphi_a^2 \xi_{\sigma}^2) U(\varphi_a^3 \xi_{\rho}^3). \quad (5.13)$$

This is a linear combination of representation operators $U(\varphi_a^s \xi_{\tau}^s)$ and is therefore equal to a representation operator $U(\eta)$ of some element $\eta \in \mathcal{G}^*$. Expression (5. 13) involves the element

$$\sum_{\tau, \sigma, \rho} \varphi_a^1 \tau_{i'i}^1 \xi_{\tau}^1 \otimes \varphi_a^2 \tau_{j'j}^2 \xi_{\sigma}^2 \otimes \varphi_a^3 \xi_{\rho}^3 = \quad (5. 14)$$

$$\varphi_{213}(\mathcal{R} \otimes \varepsilon) \varphi^{-1} \in \mathcal{G}^* \otimes \mathcal{G}^* \otimes \mathcal{G}^*. \quad (5. 15)$$

Here and in the following we use the notation

$$\varphi_{s(1)4(2)3(3)} = \sum_{\sigma} \varphi_{\sigma}^{s-1(1)} \otimes \varphi_{\sigma}^{s-1(2)} \otimes \varphi_{\sigma}^{s-1(3)} \quad (5.16)$$

for any permutation s of 1, 2, 3 if φ is given by eq.(5. 3). The same notation will be used for $\xi = \varphi^{-1}$.

Consistency requires that the matrix $\bar{\mathcal{R}}$ satisfies constraints which come from the possibility of interchanging triples of fields in two different ways, leading to the same result. We write Ψ^i for $\Psi^i(\mathbf{r}_1, t)$ and τ^i for τ^i . The constraint is that the following diagram must be commutative.

$$\begin{array}{ccc} (\Psi^1 \times \Psi^2) \times \Psi^3 & \mapsto & (\Psi^2 \times \Psi^1) \times \Psi^3 \mapsto \Psi^2 \times (\Psi^1 \times \Psi^3) \mapsto (\Psi^3 \times \Psi^2) \times \Psi^1 \\ \downarrow & & \downarrow \\ \Psi^1 \times (\Psi^2 \times \Psi^3) & & \Psi^1 \times (\Psi^3 \times \Psi^2) \mapsto (\Psi^1 \times \Psi^2) \times \Psi^3 \mapsto \Psi^3 \times (\Psi^1 \times \Psi^2) \mapsto \Psi^3 \times (\Psi^1 \times \Psi^2) \end{array} \quad (5.17)$$

Let us suppose that $\mathbf{r}_1 < \mathbf{r}_2 < \mathbf{r}_3$. The operations represented by arrows may then be performed by use of the local braid relations (5. 7) and quasiassociativity (5. 6) of the product \times . Using eq.(5. 9) one sees that the resulting relation for $\bar{\mathcal{R}}$ is fulfilled if $R \in \mathcal{G}^* \otimes \mathcal{G}^*$ satisfies Drinfelds quasi Yang Baxter equations. We use the standard notation

$$R_{13} = \sum \tau_a^1 \otimes 1 \otimes \tau_a^2 \in \mathcal{G}^* \otimes \mathcal{G}^* \otimes \mathcal{G}^*$$

etc. (ε =unit element in \mathcal{G}^*). The quasi Yang Baxter equations read

$$R_{12} \varphi_{312} R_{13} \varphi_{132}^{-1} R_{23} \varphi = \varphi_{321} R_{23} \varphi_{231}^{-1} R_{13} \varphi_{213} R_{12}. \quad (5.18)$$

In the case that the coproduct is associative, φ is equal to the identity and the quasi Yang Baxter equation becomes the ordinary Yang Baxter equation

$$R_{12} R_{13} R_{23} = R_{23} R_{13} R_{12}. \quad (5.19)$$

It is known to follow from general principles that the Hilbert space of physical states carries a unitary representation of the braid group [26]. The quasi Yang Baxter equations are a reflection of this fact. Consider states which are created from the ground state $|0\rangle$ by application of a covariant product $\Psi^1 \times \dots \times \Psi^N$ of field operators $\Psi^1 \dots \Psi^N$ which depend on arguments $\mathbf{r}_1, \dots, \mathbf{r}_N$. Suppose that $\mathbf{r}_1 < \mathbf{r}_2 < \dots < \mathbf{r}_N$; for instance. Complete specification of the multiple covariant product requires putting brackets, but we may use quasiassociativity to rearrange the brackets. Suppose this is done in a suitable way so that $(\Psi^1 \times \Psi^{i+1})$ appears. Then we may define the action

of σ_i on the state by interchanging Ψ^i and Ψ^{i+1} . (This is true if $r_i < r_{i+1}$. In the other case this operation is σ_i^{-1} .) This action of σ_i 's must satisfy the second Artin relation (4. 2). Commutativity of the diagram (5. 17) assures this.

Finally one would like to have validity of local braid relations also for composite fields. Therefore it is natural to require that braid relations (5. 7) are valid when Ψ^i is a covariant product of $\Psi^{i_1} \times \Psi^{i_2}$ of field operators, with a braid matrix (5. 9) involving representation $\tau^i \equiv \tau^{i_1} \otimes \tau^{i_2}$. Thus we should have that

$$(\Psi^1 \times (\Psi^2 \times \Psi^3))_{ijk} = ((\Psi^2 \times \Psi^3) \times \Psi^1)_{i_1 j_1 k} \bar{R}_{i_2 j_2 i_3 j_3} \quad (5.20)$$

with

$$\bar{R} = c \sum_{\alpha} \tau^{\alpha}(r_{\alpha}^1) \otimes (\tau^{\alpha} \otimes \tau^{\alpha})(r_{\alpha}^2) \quad (5.21)$$

where c is a phase factor. Similarly, consideration of composite Ψ^j leads to

$$((\Psi^1 \times \Psi^2) \times \Psi^3)_{ijk} = (\Psi^3 \times (\Psi^1 \times \Psi^2))_{i_1 j_1 k} \bar{R}'_{i_2 j_2 i_3 j_3} \quad (5.22)$$

with

$$\bar{R}' = c' \sum_{\alpha} (\tau^{\alpha} \otimes \tau^{\alpha})(r_{\alpha}^1) \otimes \tau^{\alpha}(r_{\alpha}^2) \quad (5.23)$$

The result of the exchange (5. 20) can be determined from the simple braid relation (5. 7) and quasiasociativity as shown in the following diagram.

$$\begin{array}{ccc} (\Psi^1 \times \Psi^2) \times \Psi^3 & \mapsto & \Psi^3 \times (\Psi^1 \times \Psi^2) \\ \downarrow & & \uparrow \\ \Psi^1 \times (\Psi^2 \times \Psi^3) & \mapsto & \Psi^1 \times (\Psi^2 \times \Psi^3) \mapsto (\Psi^1 \times \Psi^3) \times \Psi^2 \mapsto (\Psi^3 \times \Psi^1) \times \Psi^2 \end{array} \quad (5.24)$$

This leads to the same result as (5. 20), (5. 21) if

$$(id \otimes \Delta)(R) = \varphi_{231}^{-1} R_{13} \varphi_{213} R_{12} \varphi^{-1} \quad (5.25)$$

Similarly, eqs.(5. 22), (5. 23) are valid if

$$(\Delta \otimes id)(R) = \varphi_{312} R_{13} \varphi_{132}^{-1} R_{23} \varphi \quad (5.26)$$

These two relations are true by definition in a quasitriangular quasi Hopf algebra [1].

Conversely we have now motivated all the relations (5. 5), (4. 8), (5. 18), (5. 25), (5. 26) between R , φ and Δ , which are valid in a quasitriangular quasi Hopf algebra, from natural postulates on properties of quantum fields. (The quasi Yang Baxter equations (5. 18) are actually consequences of eqs.(5. 25) and (5. 26) [1].) Invertibility of φ and R are actually consequences of eqs.(5. 25) and (5. 26) [1].

There is also a counit and an antipode in a quasi Hopf algebra. They serve to state invariance of the ground state and the transformation law of adjoints of field operators as was explained earlier.

6 Weak quasi Hopf algebras

We would like to admit the possibility that

$$\Delta(e) \neq e \otimes e \quad (6.1)$$

Since $e^2 = e = e^*$ and Δ is supposed to be a *-morphism, $\Delta(e)$ is a projector, i.e. selfadjoint and idempotent. If it is a nontrivial projector, then for some L, J the range and cokernel of the tensor product representation $\pi^i \otimes \pi^j$ will have dimension less than the product of the dimensions of the representations π^i and π^j . This is so because $e\xi = \xi$ implies

$$\pi^i \otimes \pi^j(\xi) = \pi^i \otimes \pi^j(\Delta(\xi)) \quad (6.2)$$

$$= P_{LJ} \pi^i \otimes \pi^j(\xi) = \pi^i \otimes \pi^j(\xi) P_{LJ} \quad (6.3)$$

$$P_{LJ} = (\pi^i \otimes \pi^j)(\Delta(e)) \quad (6.4)$$

In this way we can hope to accommodate the truncated tensor products of representations of $U_q(\mathfrak{sl}_2)$ which appear in conformal field theory - see the last section. To accommodate this possibility, we must weaken the requirement that R and the associator φ both have inverses. Otherwise, the right hand side of eq.(5. 25) has an inverse and the left hand side does not.

Instead we may require that there exist quasi-inverses, still denoted by φ^{-1} and R^{-1} , such that

$$\varphi \varphi^{-1} = (id \otimes \Delta)\Delta(e) \quad (6.5)$$

$$\varphi^{-1} \varphi = (\Delta \otimes id)\Delta(e) \quad (6.6)$$

$$R R^{-1} = \Delta^*(e) \quad (6.7)$$

$$R^{-1} R = \Delta(e) \quad (6.8)$$

The existence of such a quasiinverse permits to deduce from eq.(5. 1) that

$$\varphi(\Delta \otimes id)\Delta(\xi)\varphi^{-1} = (id \otimes \Delta)\Delta(\xi) \quad (6.9)$$

$$(\Delta \otimes id)\Delta(\xi) = \varphi^{-1}(id \otimes \Delta)\Delta(\xi)\varphi \quad (6.10)$$

for all $\xi \in \mathcal{G}^*$. This suffices to ensure that the tensor product representations (5. 2) are equivalent.

There are various consistency checks which one can make to get familiar with the setup. For instance, one may multiply eq.(5. 26) for $(\Delta \otimes id)(R)$ with $(\Delta \otimes id)\Delta(e)$ from the right. On both sides this amounts to doing nothing. One may also push the factor $(\Delta \otimes id)\Delta(e)$ through the operators on the right hand side from right to left. In each step it gets converted into another projector which matches with the range of the preceding operator and with the cokernel of the next one.

Retaining Drinfelds relations (3. 3), (5. 5), (4. 8), (5. 18), (5. 25), (5. 26) between counit ϵ , R , φ and Δ , but interpreting φ^{-1} as a quasi-inverse and weakening invertibility of R as stated in (6. 5)ff, we arrive at a class of algebras which contain quasitriangular Hopf algebras as special cases. We call them *weak quasitriangular quasi Hopf algebras*. An antipode should exist subject to the conditions stated in section 3. We shall also demand that

$$(id \otimes id \otimes \epsilon)(\varphi) = \Delta(e) \quad (6.11)$$

We will not make an attempt to derive this property from the others, but we explain why it should hold. The counit defines the trivial 1-dimensional representation, $\xi \mapsto$

$\epsilon(\xi)$. Identity (3. 3) assures that tensoring with this 1-dimensional representation amounts to doing nothing. Therefore the map which implements the equivalence of the two triple products of representations (5. 2) should be trivial when the representation π^3 is the trivial 1-dimensional one. Identity (6. 11) assures that.

We would like to argue that the conclusions of the last section remain unchanged when we weaken invertibility requirements in the manner described above, and that we should therefore regard weak quasitriangular quasi Hopf algebras as possible symmetries just as well.

There is only one point which needs discussion. We used the existence of an inverse of φ to recover local braid relations involving ordinary products of fields from the local braid relation (5. 7) for covariant products.

Let $\Delta(\epsilon) = \sum \epsilon_p^i \otimes \epsilon_p^j$. Given the fields Ψ^I , we may define new fields $\tilde{\Psi}^I$ by

$$\tilde{\Psi}_i^I = (\Psi^I(\tau^I \otimes U)(\Delta(\epsilon)))_i \equiv \sum_p \tilde{\Psi}_{i,p}^I(\epsilon_p^1)U(\epsilon_p^2). \quad (6.12)$$

Using the field transformation law, property (3. 3) of the counit, and $U(\epsilon) = 1$, it is straightforward to verify by induction that

$$\tilde{\Psi}_{i_1}^{I_1} \dots \tilde{\Psi}_{i_n}^{I_n} |0\rangle = U(\epsilon) \Psi_{i_1}^{I_1} \dots \Psi_{i_n}^{I_n} |0\rangle = \tilde{\Psi}_{i_1}^{I_1} \dots \Psi_{i_n}^{I_n} |0\rangle. \quad (6.13)$$

But the Hilbert space of physical states is spanned by vectors of the form $\Psi_{i_1}^{I_1} \dots \Psi_{i_n}^{I_n} |0\rangle$. As operators in this Hilbert space, $\tilde{\Psi}^I$ and Ψ^I are therefore just the same. We may therefore assume without loss of generality that

$$\Psi^I = \tilde{\Psi}^I(\tau^I \otimes U)(\Delta(\epsilon)). \quad (6.14)$$

Writing $(\Psi^I \otimes \Psi^J)_{ij}$ for $\Psi_i^I \Psi_j^J$, it follows from the field transformation law that

$$\Psi_i^I \Psi_j^J = \tilde{\Psi}_i^I \tilde{\Psi}_j^J = ((\Psi^I \otimes \Psi^J)(\tau^I \otimes \tau^J \otimes U)(id \otimes \Delta(\epsilon)))_{ij}. \quad (6.15)$$

Using this fact, one verifies that the ordinary product of field operators can be recovered from the covariant product through the same formula (5. 12) as before, but with φ^{-1} interpreted as a quasi-inverse. The local braid relations for ordinary products follow then from those for covariant products in the same manner as before.

Next we discuss consequences of the postulate (6. 11). Together with invariance of the ground state (2. 12) eq.(6. 11) implies that

$$(\Psi^I \times \Psi^J)_{ij} |0\rangle = \Psi_i^I \Psi_j^J |0\rangle, \quad (6.16)$$

and more generally

$$(\Psi^{I_1} \times (\Psi^{I_2} \times \dots (\Psi^{I_{n-1}} \times \Psi^{I_n} \dots)))_{i_1 \dots i_n} |0\rangle = \Psi_{i_1}^{I_1} \Psi_{i_2}^{I_2} \dots \Psi_{i_{n-1}}^{I_{n-1}} \Psi_{i_n}^{I_n} |0\rangle. \quad (6.17)$$

Thus, the ordinary product of field operators agrees with the covariant product with *multiplications performed in a definite order*, when it is applied to the invariant ground state. Using quasiassociativity (5. 6) of the product \times one can convert to a different order of performing the multiplications.

This makes it easy to read off the braid relations for states and for correlation functions. We give an example. Using quasiassociativity (5. 6) we have that

$$\Psi_j^J \Psi_k^K \Psi_l^L |0\rangle = (\Psi^J \times (\Psi^K \times \Psi^L))_{jkl} |0\rangle \quad (6. 18)$$

$$= ((\Psi^J \times \Psi^K) \times \Psi^L)_{rpq} |0\rangle = (\tau^J \otimes \tau^K \otimes \tau^L)_{rpq, jkl} (\varphi^{-1}) \quad (6. 19)$$

$$= (\Psi^J \times \Psi^K)_{rp} \Psi_q^L |0\rangle = (\tau^J \otimes \tau^K \otimes \tau^L)_{rpp, jkl} (\varphi^{-1}) \quad (6. 20)$$

Assuming the arguments of the fields Ψ^J and Ψ^K , which we neglected to write, are in the proper order, we can now apply the local braid relations (5. 7), (5. 10) to invert the order of the field operators Ψ^J and Ψ^K . Then one goes backwards, using quasiassociativity and relation (6. 11) again. The final result is that

$$\Psi_j^J \Psi_k^K \Psi_l^L |0\rangle = \Psi_{kl}^K \Psi_j^J \Psi_l^L |0\rangle > T_{j^k l^r, jkl}^{JKL}(\sigma_2), \quad (6.21)$$

where

$$T_{j^k l^r, jkl}^{JKL}(\sigma_2) = (\tau^J \otimes \tau^K \otimes \tau^L)_{j^k l^r, jkl} (\varphi_{213}(R \otimes \epsilon) \varphi^{-1}) c_{JK}, \quad (6.22)$$

where c_{JK} is the phase factor in the braid relations which is left undetermined by the symmetry; it may depend on J and K . The notation $T(\sigma_2)$ was chosen to take note of the fact that a representation of the braid group shows up. The braid relation for 4-point correlation functions is obtained from eq.(6. 21) by taking the scalar product with states $\langle 0 | \Psi_i^I$.

Let us finally add a mathematical note. The definition of the covariant product of field operators amounts to introducing a \mathcal{G}^* -submodule $V^I \times V^J$ of $V^I \otimes V^J \otimes \mathcal{G}^*$, and more generally \mathcal{G}^* -submodules $(\dots (V^{I_1} \times V^{I_2}) \dots \times V^{I_n})$ etc. of $V^{I_1} \otimes V^{I_2} \otimes \dots \otimes V^{I_n} \otimes \mathcal{G}^*$. These are the appropriate modules to consider in our context. It may be of interest to note that they are actually bimodules, since right and left action of \mathcal{G}^* on them are defined. Bimodules appear naturally [29] when one starts from the algebraic theory of superselection sectors [28].

7 Covariant adjoints of field operators

The R-element $R = \sum r_a^i \otimes r_a^j \in \mathcal{G}^* \otimes \mathcal{G}^*$ serves another purpose beyond the consistent formulation of local braid relation. It makes it possible to define a covariant adjoint of field operators. This is of interest also in the quantum group context.

The transformation law of adjoints of field operators is derived in Appendix A. If there exists an antipodal map S of the algebra \mathcal{G}^* with properties as described at the beginning of section 3., then the transformation law (3. 14) of field operators Ψ^I leads to a different transformation law for adjoints Ψ^{I*} of field operators, assuming the $*$ -operation on $\mathcal{G}^* \otimes \mathcal{G}^*$ is defined as in (3. 11).

$$U(\phi) \Psi_i^{I*} = \sum_p \Psi_m^I \bar{\tau}_{mi}(\phi_p^2) U(\phi_p^1). \quad (7.1)$$

This differs from the transformation law of fields by the interchange of ϕ_p^1 and ϕ_p^2 , apart from the expected appearance of the contragredient representation

$$\bar{\tau}(\phi) = {}^t \tau(S(\phi)).$$

This complication can be eliminated by defining a covariant adjoint

$$\bar{\Psi}_i^j = \sum_a \Psi_k^j \tau_{ki}^{-1}(\tau_a^{-1}) U(\tau_a^2) \quad (7.2)$$

The main property $\Delta'(\phi)R = R\Delta(\phi)$ of the R-element reads explicitly

$$\sum_{a,b} \phi_a^2 \tau_a^1 \otimes \phi_b^1 \tau_a^2 = \sum_{a,b} \tau_a^1 \phi_a^1 \otimes \tau_a^2 \phi_a^2.$$

Using this it is straightforward to deduce from the transformation law (7. 1) of adjoint field operators that $\bar{\Psi}_i^j$ transforms covariantly according to the representation $\bar{\tau}^j$ which is contragredient to τ^j ,

$$U(\phi) \bar{\Psi}_i^j = \sum_a \bar{\Psi}_k^j \tau_{ki}^{-1}(\phi_a^1) U(\phi_a^2). \quad (7.3)$$

This differs from the transformation law for Ψ^j only by the appearance of $\bar{\tau}^j$ in place of τ^j .

Our construction of the covariant adjoint resembles what one does in the theory of the Dirac equation when one introduces an adjoint Dirac spinor $\bar{\Psi} = \Psi^* \gamma_0$ which transforms covariantly under the Lorentz group.

8 The comodule picture

We regarded the Hilbert space \mathcal{H} of physical states as a representation space for \mathcal{G}^* because we wanted to generalize the well known notion of a symmetry in quantum theory where \mathcal{H} is a representation space for a group G and its group algebra \mathcal{G}^* . This point of view was proposed by Buchholz, Mack and Todorov [4].

There exists an alternative point of view, introduced by Reshetikhin and Smirnov [15] where the dual \mathcal{G} of \mathcal{G}^* plays the central role, and \mathcal{H} is regarded as a comodule for \mathcal{G} . In this section we will explain the connection between both pictures.

A representation U of \mathcal{G}^* in \mathcal{H} defines a linear map

$$U(\xi) : \mathcal{H} \rightarrow \mathcal{H} \quad (8.1)$$

for every $\xi \in \mathcal{G}^*$. Therefore, U may be regarded as a map

$$U : \mathcal{G}^* \otimes \mathcal{H} \rightarrow \mathcal{H}. \quad (8.2)$$

Given such a representation U , \mathcal{H} becomes a comodule for the dual \mathcal{G} of \mathcal{G}^* in a natural way. This means that there is a linear map

$$\bar{U} : \mathcal{H} \rightarrow \mathcal{G} \otimes \mathcal{H}. \quad (8.3)$$

It is determined by U as follows. Elements $f \in \mathcal{G}$ may be regarded as linear maps of \mathcal{G}^* into \mathbb{C} , $\xi \mapsto f(\xi)$. Therefore, elements $f \otimes |v\rangle \in \mathcal{G} \otimes \mathcal{H}$ may be regarded as linear maps $\mathcal{G}^* \rightarrow \mathcal{H}$, viz.

$$\xi \mapsto (f \otimes |v\rangle)(\xi) = |v\rangle f(\xi) \in \mathcal{H}. \quad (8.4)$$

\bar{U} is defined by

$$\bar{U}|v\rangle(\xi) = U(\xi)|v\rangle \text{ for } |v\rangle \in \mathcal{H}. \quad (8.5)$$

If \mathcal{G}^* is a bialgebra then \mathcal{G} is also a bialgebra. The coproduct Δ in \mathcal{G}^* gives the product \cdot in \mathcal{G} ,

$$(f \cdot g)(\xi) = f \otimes g(\Delta(\xi)). \quad (8.6)$$

And the product in \mathcal{G}^* yields a coproduct δ in \mathcal{G} ,

$$\delta f(\xi, \eta) = f(\xi \eta). \quad (8.7)$$

Unit and counit are similarly interchanged.

When the coproduct in \mathcal{G}^* is not coassociative then \mathcal{G} is a nonassociative algebra. So far the comodule picture has been used only for associative \mathcal{G} as far as we know, and we will restrict the discussion to this case to begin with.

We postulated a covariant transformation law of fields under \mathcal{G}^* . This translates into commutation relations of \bar{U} with fields. In careful notation

$$\bar{U} \Psi_i^j = (1 \otimes \Psi_j^i)(\tau_{ji}^j \otimes 1) \bar{U}. \quad (8.8)$$

Herein, the first 1 is the unit element of \mathcal{G} and $1 \otimes \Psi_i^j$ is a linear map on $\mathcal{G} \otimes \mathcal{H}$ which takes $f \otimes |v\rangle$ into $f \otimes \Psi_i^j |v\rangle$. τ_{ji}^i are the elements of \mathcal{G} which are defined as linear maps on \mathcal{G}^* , $\xi \mapsto \tau_{ji}^i(\xi)$ by the representation matrices encountered previously. A product of such elements of \mathcal{G} is defined through eq.(8. 6).

Let us demonstrate validity of commutation relation (8. 8). We introduce an orthonormal basis $|v_n\rangle$ in \mathcal{H} . We see from eq.(8. 5) that

$$\bar{U}|v_n\rangle = u_{mn} \otimes |v_m\rangle \quad (8.9)$$

with $u_{mn} \in \mathcal{G}$ given by

$$u_{mn}(\xi) = \langle v_m | U(\xi) | v_n \rangle. \quad (8.10)$$

Consider now $\bar{U} \Psi_i^j |v_n\rangle$. From the definition (8. 5) and the covariance property of fields under \mathcal{G}^* we deduce

$$\bar{U} \Psi_i^j |v_n\rangle(\xi) = U(\xi) \Psi_i^j |v_n\rangle = \sum_a \Psi_k^j \tau_{ki}^a(\xi_a^1) U(\xi_a^2) |v_n\rangle \quad (8.11)$$

$$= \sum_a \Psi_k^j |v_m\rangle \tau_{ki}^a(\xi_a^1) u_{mn}(\xi_a^2). \quad (8.12)$$

if $\Delta(\xi) = \sum_a \xi_a^1 \otimes \xi_a^2$. On the other hand

$$(1 \otimes \Psi_j^i)(\tau_{ji}^j \otimes 1) \bar{U} |v_n\rangle(\xi) = (1 \otimes \Psi_j^i)(\tau_{ji}^j \otimes 1)(u_{mn} \otimes |v_n\rangle)(\xi) \quad (8.13)$$

$$= \Psi_j^i |v_m\rangle \tau_{ji}^j(\xi) \quad (8.14)$$

$$= \sum_a \Psi_j^i |v_m\rangle \tau_{ji}^j(\xi_a^1) u_{mn}(\xi_a^2). \quad (8.15)$$

In the last equation, the multiplication law (8. 6) in \mathcal{G} was used. We see that both sides of eq.(8. 8) are indeed equal when applied to an arbitrary basis vector. This proves the commutation relations (8. 8).

The statement of invariance of the ground state translates into

$$\tilde{U}|0\rangle = 1 \otimes |0\rangle \quad (8.16)$$

where 1 is the unit element in \mathcal{G} . Regarded as a linear function on \mathcal{G}^* it is given by the counit on \mathcal{G}^* ,

$$1 : \xi \mapsto \epsilon(\xi) \quad (8.17)$$

Let us introduce a shorthand notation. Elements of $\mathcal{G} \otimes \mathcal{H}$ will be written $f|v\rangle$ or $|v\rangle f$ in place of $f \otimes |v\rangle$, and factors $1 \in \mathcal{G}$ will be dropped (but keep eq.(8.17) in mind!) Then, commutation relations with fields and invariance of the ground state reads

$$\tilde{U}\Psi_i^j = \Psi_j^i \tilde{U} \quad (8.18)$$

$$\tilde{U}|0\rangle = |0\rangle \quad (8.19)$$

$$\tilde{U}\Psi_{i_1}^{j_1} \dots \Psi_{i_n}^{j_n} |0\rangle = \Psi_{j_1}^{i_1} \dots \Psi_{j_n}^{i_n} |0\rangle \quad (8.20)$$

It follows that

When both sides of the state transformation law (8.20) are applied to an arbitrary element $\phi \in \mathcal{G}^*$ we recover the old transformation law (3.26) of states.

Elements $\tau \in \mathcal{G}$ commute with operators Ψ on \mathcal{H} . However, the multiplication in \mathcal{G} is not commutative if the comultiplication in \mathcal{G}^* is not commutative. Therefore the order of the factors τ_j^i matters.

From the definition (8.6) we see that the product in \mathcal{G} is in fact equal to the tensor product (of representations) introduced earlier in the sense that

$$\tau_{ik}^j \cdot \tau_{jl}^i = (\tau^j \otimes \tau^i)_{j,kl} \quad (8.21)$$

The transformation law (8.18) and ff are very much reminiscent of the formula (2.13) for the transformation law of fields under a group, the main difference being that the τ_{ik}^j don't commute. The similarity is still more deceptive when τ^j is the fundamental representation of a matrix group so that $\tau_{ik}^j(a) = a_{ik}$. One might then be tempted to write a_{ik} in place of τ_{ik}^j for the function which maps a into a_{ik} . But mathematical decency requires that one distinguish between functions and the values which they take. The τ_{ik}^j in eq.(8.20) are generalizations of functions on a group, not values of such functions.

In the case of a quasi-Hopf symmetry, the product \cdot in \mathcal{G} is not associative. The formulas obtained so far remain valid, except that it is necessary to put brackets to indicate the order in which multiplications in \mathcal{G} are to be performed. For instance, the state transformation law (8.20) is to be read as

$$\tilde{U}\Psi_{i_1}^{j_1} \dots \Psi_{i_n}^{j_n} |0\rangle = \Psi_{j_1}^{i_1} \dots \Psi_{j_n}^{i_n} |0\rangle \quad (8.22)$$

9 The quantum group algebra $U_q(sl_2)$

It is instructive to consider as an example the matrix pseudogroup \mathcal{G} whose dual is (or rather: contains) the quantum group algebra $U_q(sl_2)$ [9,8]. \mathcal{G} is an associative but

noncommutative one parameter deformation⁵ of the algebra $Fun(\mathcal{G})$ of functions on the group $G = SL(2, \mathbb{C})$ of complex unimodular 2×2 matrices $a = (a_{ij})_{i,j=1,2}$.

Let us first consider the commutative algebra $Fun(SL(2, \mathbb{C}))$. This algebra is generated by simple functions τ_{ij} , $i, j = 1, 2$. They are defined by

$$\tau_{ij}(a) = a_{ij} \quad (9.1)$$

In the old language, τ is the fundamental representation. These generators of \mathcal{G} obey the relations

$$\det \tau \equiv \tau_{11} \cdot \tau_{22} - \tau_{12} \cdot \tau_{21} = 1 \quad (9.2)$$

$$\tau_{ij} \cdot \tau_{kl} = \tau_{kl} \cdot \tau_{ij} \quad (9.3)$$

We remember that the multiplication \cdot of functions is pointwise defined, $f \cdot g(a) = f(a)g(a)$. Eq.(9.3) expresses commutativity of this product.

The deformations \mathcal{G} are parametrized by a complex number q . \mathcal{G} is again generated by elements τ_{ij} , but the relations (9.2) and (9.3) get replaced by

$$\det \tau \equiv \tau_{11} \cdot \tau_{22} - q\tau_{12} \cdot \tau_{21} = 1, \quad (9.4)$$

$$\mathcal{R}_{ik,rs} \tau_{rv} \cdot \tau_{sw} = \tau_{kl} \cdot \tau_{im} \mathcal{R}_{ml,sv}, \quad (9.5)$$

where \mathcal{R} is our old element $R \in \mathcal{G}^* \otimes \mathcal{G}^*$ in the fundamental representation τ of \mathcal{G}^* ,

$$\mathcal{R}_{ik,rs} = (\tau \otimes \tau)_{ik,rs}(R)$$

When rows and columns are labelled in lexicographic order, \mathcal{R} has the explicit form

$$\mathcal{R} = \begin{pmatrix} q & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & q - q^{-1} & 1 & 0 \\ 0 & 0 & 0 & q \end{pmatrix} \quad (9.6)$$

In the alternative notation

$$\tau = \begin{pmatrix} \tau_{11} & \tau_{12} \\ \tau_{21} & \tau_{22} \end{pmatrix} = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \quad (9.7)$$

the relations (9.5) are given by

$$ab = qba \quad (9.8)$$

$$ac = qca \quad (9.9)$$

$$ad - da = (q - q^{-1})bc \quad (9.10)$$

$$cd = qdc \quad (9.11)$$

$$bd = qdb \quad (9.12)$$

$$bc = cb \quad (9.13)$$

It remains to exhibit the generators of \mathcal{G} as complex linear functions on $U_q(sl_2)$. The formulas that do this have been found by Reshetikhin, Takhtajan and Faddeev [23].

⁵GL(2) possesses a two parameter deformation [31]

10 Weak quasi-Hopf algebras associated with $U_q(sl_2)$ for $q^p = -1$

Now we turn to the problem of how to truncate the algebra $U_q(sl_2)$ in such a way that its unphysical representations are eliminated but constructions of physical interest, such as the construction of tensor products of physical representations, remain possible.

The quantum group algebra $U_q(sl_2)$ with $q^p = -1$ has physical representations π^K , $2K = 0, 1, \dots, p-2$, which act in representation spaces V^K . The tensor product of two such physical representations decomposes in general into physical representations, plus unphysical subrepresentations with quantum dimension 0.

Let \mathcal{J} be the ideal in $U_q(sl_2)$ which is annihilated by all physical π^K . We set

$$\mathcal{G}^* = U_q(sl_2)/\mathcal{J}.$$

Equivalently,

$$\mathcal{G}^* = \pi(U_q(sl_2)), \text{ where } \pi = \bigoplus_K \pi^K. \quad (10.1)$$

Let \otimes_q be the tensor product of $U_q(sl_2)$ representations. It comes from a coproduct Δ_q for $U_q(sl_2)$. In application to conformal field theory one invents an extra rule which says that the extra terms represented by dots ... in the tensor product decomposition

$$\pi^I \otimes_q \pi^J = \bigoplus_{|I-J| \leq K \leq \min(I,J)} \pi^K + \dots, \quad (10.2)$$

$$u(IJ) \equiv \min\{|I+J|, p-2-I-J\} \quad (10.3)$$

are to be thrown away. The left out part has quantum dimension 0 [16]. Supplemented with this extra rule, the tensor product decomposition reproduces the fusion rules of minimal conformal models. We would like to define a coproduct Δ in the algebra \mathcal{G}^* in such a way that the resulting tensor product \otimes decomposes as in eq.(10.2) but without any extra terms ...

Let

$$C_q(IJ|K) : V^I \otimes V^J \mapsto V^K$$

be the Clebsch Gordan maps for $U_q(sl_2)$. By definition, they enjoy the intertwining property

$$C_q(IJ|K)(\pi^I \otimes_q \pi^J)(\xi) = \pi^K(\xi)C_q(IJ|K) \quad (10.4)$$

for all ξ . Using these Clebsch Gordan maps, one can define a nonassociative truncated tensor product \otimes of representations of $U_q(sl_2)$, and this tensor product is at the same time a representation of \mathcal{G}^* , i.e. it annihilates \mathcal{J} .

$$(\pi^I \otimes \pi^J)(\xi) = \sum_{|I-J| \leq K \leq \min(I,J)} C_q(IJ|K)^* \pi^K(\xi) C_q(IJ|K) \quad (10.5)$$

$$= (\pi^I \otimes \pi^J)(\Delta_q(\xi)) P_{IJ}, \quad (10.6)$$

with

$$P_{IJ} = \sum_{|I-J| \leq K \leq \min(I,J)} C_q(IJ|K)^* C_q(IJ|K) \quad (10.7)$$

The quantum group algebra $U_q(sl_2)$ is generated by elements $q^{\pm H/2}$ and S_{\pm} subject to the relations

$$q^{H/2} q^{-H/2} = q^{-H/2} q^{H/2} = 1, \quad (9.14)$$

$$q^{H/2} S_{\pm} = q^{\pm 1} S_{\pm} q^{H/2}, \quad (9.15)$$

$$[S_+, S_-] = \frac{q^H - q^{-H}}{q - q^{-1}}. \quad (9.16)$$

Here and in the following we write 1 in place of e for the unit element in $\mathcal{G}^* = U_q(sl_2)$. The comultiplication is given by

$$\Delta(q^{\pm H/2}) = q^{\pm H/2} \otimes q^{\pm H/2} \quad (9.17)$$

$$\Delta(S_{\pm}) = S_{\pm} \otimes q^{H/2} + q^{-H/2} \otimes S_{\pm}. \quad (9.18)$$

If $|q| = 1$, a $*$ -operation can be defined by

$$S_{\pm}^* = S_{\mp}, \quad (q^{\pm H/2})^* = q^{\mp H/2}$$

and this is an automorphism satisfying $\Delta(S^*) = \Delta(S)^*$ provided we adopt convention (3.11) for the adjoint on $\mathcal{G}^* \otimes \mathcal{G}^*$.

The counit is

$$\epsilon(q^{\pm H/2}) = 1, \quad \epsilon(S_{\pm}) = 0,$$

and the antipodal map S acts as

$$q^{\mp H/2} = S(q^{\pm H/2}), \quad -q^{\pm 1} S_{\pm} = S(S_{\pm}).$$

It is convenient to use matrix notation again. Following Reshetikhin, Takhtajan and Faddeev [23] one puts

$$L_+ = \begin{pmatrix} q^{H/2} & 0 \\ 0 & q^{-H/2} \end{pmatrix}, \quad L_- = \begin{pmatrix} q^{-H/2} & 0 \\ (q^{-1} - q)S_- & q^{H/2} \end{pmatrix}, \quad (9.19)$$

$$L_{\pm}^1 = L_{\pm} \otimes 1, \quad L_{\pm}^2 = 1 \otimes L_{\pm}, \quad (9.20)$$

$$T^1 = \tau \otimes 1, \quad T^2 = 1 \otimes \tau, \quad (9.21)$$

$$R^+ = PRP, \quad R^- = R^{-1}, \quad (9.22)$$

with $P(a \otimes b) = b \otimes a$.

In this notation the action of generators $q^{\pm H/2}$, S_{\pm} of $U_q(sl_2)$ on generators τ_{ij} of \mathcal{G} is given by

$$L_{\pm}^1 : T^2 \mapsto R^{\pm}. \quad (9.23)$$

Suppose now that there exists an element P of $U_q(sl_2) \otimes U_q(sl_2)$ such that

$$P_{IJ} = (\pi^I \otimes \pi^J)(P)$$

Then we may define a coproduct Δ in \mathcal{G}^* such that

$$(\pi^I \otimes \pi^J)(\xi) = (\pi^I \otimes \pi^J)(\Delta(\xi)). \quad (10.8)$$

This is achieved by setting

$$\Delta(\xi) = P\Delta_q(\xi).$$

Property (10. 8) is immediate. We show that $\Delta(\xi)$ is well defined for $\xi \in \mathcal{G}^*$, i.e. depends on $X \in U_q(sl_2)$ only through its equivalence class $\xi \text{ mod } \mathcal{J}$ in \mathcal{G}^* . In view of the definition (10. 1) it suffices to show that for arbitrary physical I, J we have $(\pi^I \otimes \pi^J)(\Delta(Z)) = 0$ if $Z \in \mathcal{J}$. But this follows right away from property (10. 8) and eq.(10. 5).

We will construct the desired projector from the Casimir invariant $C \in U_q(sl_2)$,

$$C = S^{-1}S^+ + \left(\frac{q^{(H+1)/2} - q^{-(H+1)/2}}{q - q^{-1}} \right)^2 - \left(\frac{q^{1/2} - q^{-1/2}}{q - q^{-1}} \right)^2. \quad (10.9)$$

It has eigenvalues

$$\lambda_J = \left[J + \frac{1}{2} \right]_q^2 - \left[\frac{1}{2} \right]_q^2, \quad J = 0, \frac{1}{2}, 1, \dots, \quad (10.10)$$

where we use the standard notation $[x]_q = (q^x - q^{-x})/(q - q^{-1})^{-1}$. The Casimir does not suffice to make a global distinction between physical representations and unphysical ones because of the symmetry of λ_K under $K \mapsto p - 1 - K$. But fortunately this is not needed, either. Given I, J we say that K is "permissible" if $|I - J| \leq K \leq |I + J|$ and $K \leq u(I, J)$ and "unwanted" if $|I - J| \leq K \leq |I + J|$ but $K > u(I, J)$. We see that it can never happen that K is unwanted but its brother $p - 1 - K$ is permissible, or the other way round.

Let θ_K be the real function, defined on the set E of distinct possible eigenvalues of C , which is equal to 1 for argument $x = \lambda_K$ and 0 otherwise,

$$\theta(x) = \prod_{\lambda_K \neq x} (x - \tau)(\lambda_K - \tau)^{-1}.$$

We consider the projectors

$$P^K = \theta_K(C) \in U_q(sl_2),$$

and set

$$P = \sum_{I, J \text{ phys}} (P^I \otimes P^J) \sum_{|I-J| \leq K \leq u(I, J)} \Delta_q(P^K) \in U_q(sl_2) \otimes U_q(sl_2). \quad (10.11)$$

P^I commutes with all elements of $U_q(sl_2)$. Therefore P is a sum of products of commuting projectors. It has the desired properties because of the intertwining property of Clebsch Gordan maps, and because there is no need to discriminate between representations which P^K fails to distinguish on account of the symmetry $K \mapsto p - 1 - K$.

The 1-dimensional representation π^0 gives the counit,

$$\epsilon(\xi) = \pi^0(\xi).$$

The counit will therefore remain the same as in $U_q(sl_2)$. When $I = 0$ or $J = 0$ then the restricted tensor product (10. 5) agrees with the old tensor product $\pi^I \otimes \pi^J$. Therefore

$$\pi^K(\xi) = (\pi^K \otimes \pi^0)(\xi) = (\pi^K \otimes \pi^0)(\Delta(\xi)) = \pi^K((id \otimes \epsilon)\Delta(\xi)),$$

and similarly for $\pi^0 \otimes \pi^K$. This is true for all physical representations π^K . Recalling definition (10. 1), it follows that the relation

$$(id \otimes \epsilon)\Delta = id = (\epsilon \otimes id)\Delta$$

holds true.

This completes the construction of the coproduct Δ for \mathcal{G}^* .

An R-element and its pseudoinverse exist such that $R\Delta(\xi) = \Delta'(\xi)R$ and relations (6. 7), (6. 8) hold. In terms of the R-element R_q of $U_q(sl_2)$ it is given by

$$R = R_q\Delta(\epsilon) = \Delta'(\epsilon)R_q, \quad (10. 12)$$

$$R^{-1} = R_q^{-1}\Delta'(\epsilon) = \Delta(\epsilon)R_q^{-1}. \quad (10. 13)$$

Next we will show that there exists an element $\varphi \in \mathcal{G}^* \otimes \mathcal{G}^* \otimes \mathcal{G}^*$ which possesses a quasi-inverse and which makes the coproduct quasi-coassociative in the sense that eq.(5. 1) holds, viz.

$$\varphi(\Delta \otimes id)\Delta(\xi) = (id \otimes \Delta)\Delta(\xi)\varphi \quad \text{for all } \xi \in \mathcal{G}^*. \quad (10.14)$$

One of the purposes of the associator φ is to ensure that the restricted tensor products of representations $(\pi^I \otimes \pi^J) \otimes \pi^K$ and $\pi^I \otimes (\pi^J \otimes \pi^K)$ are equivalent. By definition,

$$(\pi^I \otimes \pi^J) \otimes \pi^K(\xi) = \pi^I \otimes \pi^J \otimes \pi^K((\Delta \otimes id)\Delta(\xi)), \quad (10. 15)$$

$$\pi^I \otimes (\pi^J \otimes \pi^K)(\xi) = \pi^I \otimes \pi^J \otimes \pi^K((id \otimes \Delta)\Delta(\xi)). \quad (10. 16)$$

Both are representations of $U_q(sl_2)$ which are at the same time representations of \mathcal{G}^* . It is well known that equivalence of these two representations is indeed true [19,32]. There will therefore be maps

$$\varphi_{IJK} : V^I \otimes V^J \otimes V^K \mapsto V^I \otimes V^J \otimes V^K \quad (10.17)$$

such that

$$\varphi_{IJK}((\pi^I \otimes \pi^J) \otimes \pi^K)(\xi) = (\pi^I \otimes (\pi^J \otimes \pi^K))(\xi)\varphi_{IJK} \quad (10.18)$$

for all $\xi \in \mathcal{G}^*$, and φ_{IJK} possesses a quasiinverse φ_{IJK}^{-1} such that

$$\varphi_{IJK}\varphi_{IJK}^{-1} = (\pi^I \otimes (\pi^J \otimes \pi^K))(\epsilon) = (\pi^I \otimes \pi^J \otimes \pi^K)((id \otimes \Delta)\Delta(\epsilon)) \quad (10. 19)$$

$$\varphi_{IJK}^{-1}\varphi_{IJK} = ((\pi^I \otimes \pi^J) \otimes \pi^K)(\epsilon) = (\pi^I \otimes \pi^J \otimes \pi^K)((\Delta \otimes id)\Delta(\epsilon)) \quad (10. 20)$$

We seek $\varphi \in \mathcal{G}^* \otimes \mathcal{G}^* \otimes \mathcal{G}^*$, or a representative of it in $U_q(sl_2) \otimes U_q(sl_2) \otimes U_q(sl_2)$, such that

$$\varphi_{IJK} = (\pi^I \otimes \pi^J \otimes \pi^K)(\varphi). \quad (10.21)$$

Let \mathcal{M}^I be the full matrix algebra which consists of $2I + 1 \times 2I + 1$ matrices. By construction, \mathcal{G}^* is a finite dimensional semisimple algebra. Therefore it is a direct sum of full matrix algebras. Since the representations π^K are irreducible, it follows from the definition (10. 1) that $\pi^K(\mathcal{G}^*) = \mathcal{M}^K$. To every matrix $m \in \mathcal{M}^K$ there will therefore exist some element $\mu \in \mathcal{G}^*$ such that $m = \pi^K(\mu)$. By definition

$$\varphi_{IJK} \in \mathcal{M}^I \otimes \mathcal{M}^J \otimes \mathcal{M}^K.$$

It follows that there exists $\phi_{IJK} \in \mathcal{M}^I \otimes \mathcal{M}^J \otimes \mathcal{G}^*$ such that

$$\varphi_{IJK} = (id \otimes id \otimes \pi^K)(\phi_{IJK}).$$

Setting

$$\phi_{IJ} = \sum_K P^K \phi_{IJK},$$

we have that

$$\pi^K(\phi_{IJ}) = \varphi_{IJK}.$$

for all K . ϕ_{IJ} is a sum of terms $m^I \otimes m^J \otimes \xi_{IJ}$ with matrices m^I, m^J and $\xi_{IJ} \in \mathcal{G}^*$. Next we may use the same argument to write $m^J = \pi^J(\phi'_{IJ})$ and define

$$\phi'_I = \sum_J P^J \phi'_{IJ} \otimes \xi_{IJ} \in \mathcal{G}^* \otimes \mathcal{G}^*.$$

And so it goes. Repeating the same argument once more we arrive finally at an element $\varphi \in \mathcal{G}^* \otimes \mathcal{G}^* \otimes \mathcal{G}^*$ with the desired property (10. 21). This completes the construction of φ . The existence of its quasinverse φ^{-1} follows in the same way.

Given φ_{IJK} , φ is unique. We will now show that there exists an element $\varphi \in \mathcal{G}^* \otimes \mathcal{G}^* \otimes \mathcal{G}^*$, so that Drinfelds relations (5. 1), (5. 5), (5. 25), (5. 26) hold. To this effect we will give an explicit definition of φ_{IJK} by specifying its action on basis vectors. We will no longer distinguish in notation between φ and φ_{IJK} in the following. More generally we write ζ in place of $(\pi^I \otimes \pi^J \otimes \pi^K)(\zeta)$ etc.

A truncated tensor product \otimes is not only defined for representations, but also for basis vectors e_i^j of representation spaces V^I ,

$$e_i^j \otimes e_j^k = P_{IJ} e_i^k = \Delta(\epsilon_i^j) \otimes e_j^k.$$

The definition extends to reducible representations, so that multiple products are also defined. In particular

$$\epsilon_i^j \otimes (\epsilon_j^k \otimes \epsilon_k^l) = (id \otimes \Delta)(\epsilon_i^j \otimes \epsilon_j^k \otimes \epsilon_k^l), \quad (10. 22)$$

$$(\epsilon_i^j \otimes \epsilon_j^k) \otimes \epsilon_k^l = (\Delta \otimes id)(\epsilon_i^j \otimes \epsilon_j^k \otimes \epsilon_k^l). \quad (10. 23)$$

Vectors $\epsilon_i^j \otimes (\epsilon_j^k \otimes \epsilon_k^l)$ span a subspace $V_i^j \otimes (V_j^k \otimes V_k^l)$ of $V_i^j \otimes V_j^k \otimes V_k^l$, and vectors $(\epsilon_i^j \otimes \epsilon_j^k) \otimes \epsilon_k^l$ span another subspace $(V_i^j \otimes V_j^k) \otimes V_k^l$. The vectors $(\epsilon_i^j \otimes \epsilon_j^k) \otimes \epsilon_k^l$

will sometimes be called "physical states". They are not all linearly independent, but $(V_i^j \otimes V_j^k) \otimes V_k^l$ is spanned by the linearly independent vectors

$$\sum_{ijhp} C_{ij}^{(ijp)} C_{pq}^{(pjk)} C_{kl}^{(klh)} e_i^j \otimes e_j^k \otimes e_k^l$$

with $|I - J| \leq P \leq u(IJ)$ and $|P - K| \leq L \leq u(PK)$. The symbols $C_{ij}^{(ijp)}$ denote the usual Clebsch Gordon coefficients for $U_q(sl_2)$ which are determined by the above mentioned Clebsch Gordon maps. They will be the Clebsch Gordon coefficients for \mathcal{G}^* at the same time when $|I - J| \leq K \leq u(IJ)$. The other Clebsch Gordon coefficients for \mathcal{G}^* vanish.

We make an ansatz for the action of φ on these basis vectors

$$\varphi \sum_{ijhp} C_{ij}^{(ijp)} C_{pq}^{(pjk)} C_{kl}^{(klh)} e_i^j \otimes e_j^k \otimes e_k^l = \sum_{Q, ijhkq} F_{PQ} C_{ij}^{(ijp)} C_{kl}^{(klh)} C_{q}^{(qjk)} e_i^j \otimes e_j^k \otimes e_k^l. \quad (10.24)$$

The action of φ is extended to all of $V^I \otimes V^J \otimes V^K$ by imposing the requirement that $\varphi = \varphi(\Delta \otimes id)\Delta(\epsilon)$. By this ansatz, the intertwining property (5. 1) of φ is assured for arbitrary choice of the complex coefficients F_{PQ} .

This is verified by a short calculation as follows. Since

$$\varphi(\Delta \otimes id)\Delta(\epsilon) = \varphi(\Delta \otimes id)\Delta(\epsilon)(\Delta \otimes id)\Delta(\epsilon) = \varphi(\Delta \otimes id)\Delta(\epsilon)(\Delta \otimes id)\Delta(\epsilon)$$

it suffices to consider $\varphi(\Delta \otimes id)\Delta(\epsilon)$ on the physical states in the tensor product decomposition. Using the following transformation laws

$$(\Delta \otimes id)\Delta(\epsilon) \sum_{ijhp} C_{ij}^{(ijp)} C_{pq}^{(pjk)} C_{kl}^{(klh)} e_i^j \otimes e_j^k \otimes e_k^l = \sum_{ijhqp} \tau_{ij}^h(\epsilon) C_{ij}^{(ijp)} C_{pq}^{(pjk)} C_{kl}^{(klh)} e_i^j \otimes e_j^k \otimes e_k^l, \quad (10.25)$$

$$(id \otimes \Delta)\Delta(\epsilon) \sum_{ijhkq} C_{ij}^{(ijp)} C_{pq}^{(pjk)} C_{kl}^{(klh)} e_i^j \otimes e_j^k \otimes e_k^l = \sum_{ijhqp} \tau_{ij}^h(\epsilon) C_{ij}^{(ijp)} C_{pq}^{(pjk)} C_{kl}^{(klh)} e_i^j \otimes e_j^k \otimes e_k^l, \quad (10.26)$$

we calculate

$$\varphi(\Delta \otimes id)\Delta(\epsilon) \sum_{ijhkq} C_{ij}^{(ijp)} C_{pq}^{(pjk)} C_{kl}^{(klh)} e_i^j \otimes e_j^k \otimes e_k^l$$

$$= \varphi \sum_{ijhqp} \tau_{ij}^h(\epsilon) C_{ij}^{(ijp)} C_{pq}^{(pjk)} C_{kl}^{(klh)} e_i^j \otimes e_j^k \otimes e_k^l$$

$$= \sum_{Q, ijhkq} \tau_{ij}^h(\epsilon) F_{PQ} C_{ij}^{(ijp)} C_{pq}^{(pjk)} C_{kl}^{(klh)} e_i^j \otimes e_j^k \otimes e_k^l$$

$$= (id \otimes \Delta)\Delta(\epsilon) \sum_{Q, ijhkq} F_{PQ} C_{ij}^{(ijp)} C_{pq}^{(pjk)} C_{kl}^{(klh)} e_i^j \otimes e_j^k \otimes e_k^l$$

$$= (id \otimes \Delta)\Delta(\epsilon) \varphi \sum_{ijhp} C_{ij}^{(ijp)} C_{pq}^{(pjk)} C_{kl}^{(klh)} e_i^j \otimes e_j^k \otimes e_k^l. \quad (10. 27)$$

This completes the demonstration of the intertwining property (5.1). Now we turn to the other properties of φ in a weak quasi-Hopf algebra.

(5. 25), (5. 26) impose further restrictions on the coefficients $F_{PQ}[\dots]$. Acting with $(id \otimes id \otimes \Delta)(\varphi)(id \otimes id \otimes id)$ and $(e \otimes \varphi)(id \otimes \Delta \otimes id)(\varphi \otimes e)$ on physical states in the fourfold tensor product and comparing the coefficients, we may formulate (5. 5) as a relation for the $F_{PQ}[\dots]$.

$$F_{QQ'}[\begin{smallmatrix} K & L \\ P & M \end{smallmatrix}] F_{PP'}[\begin{smallmatrix} J & Q' \\ I & M \end{smallmatrix}] = \sum_{\mathbf{R}} F_{PR}[\begin{smallmatrix} J & K \\ I & Q \end{smallmatrix}] F_{RQ'}[\begin{smallmatrix} R & L \\ I & M \end{smallmatrix}] F_{RQ}[\begin{smallmatrix} K & L \\ J & Q' \end{smallmatrix}]. \quad (10.28)$$

Using the same procedure for the evaluation of $\varphi_{23}(id \otimes \Delta)(R)\varphi$ and $R_{13}\varphi_{213}R_{12}$ relation (5. 25) requires

$$\sum_{\mathbf{P}} (-1)^{P'} q^{\frac{P'}{2}} F_{P'Q}[\begin{smallmatrix} K & J \\ I & L \end{smallmatrix}] F_{PP'}[\begin{smallmatrix} K & I \\ J & L \end{smallmatrix}] = (-1)^{J'+K+L-Q-P} q^{\frac{Q+P'}{2}-\frac{J}{2}-\frac{Q}{2}-\frac{K}{2}-\frac{K'-C}{2}} F_{PQ}[\begin{smallmatrix} I & K \\ J & L \end{smallmatrix}]. \quad (10.29)$$

Here $c_J = (J+1)J$ and we used the relation

$$\sum_{\mathbf{I}, \mathbf{J}} (R^I)^J C[\begin{smallmatrix} J & K \\ I & J' & k, q \end{smallmatrix}] = (-1)^{J'+J-K} q^{\frac{K}{2}-\frac{J}{2}-\frac{J'}{2}} C[\begin{smallmatrix} I & K \\ J & I & k \end{smallmatrix}]. \quad (10.30)$$

A numerical solution of the restrictions (10. 28) and (10. 29) is explicitly known to be given by the fusion matrix,

$$F_{PQ}[\begin{smallmatrix} J & K \\ I & L \end{smallmatrix}] = \{ \begin{smallmatrix} I & J & P \\ K & L & Q \end{smallmatrix} \}_q. \quad (10.31)$$

Here $\{\dots\}_q$ denote the 6-j symbols of $U_q(sl_2)$. In the case where q is a root of unity it is well known that the corresponding identities for 6-j symbols (namely the q-analogs of Biedenharn-Elliott and Racah equations) restrict to summation over physical indices. The last identity (5. 26) is also satisfied by this choice of φ .

The definition of φ reduces to the identity when $\Delta(e) = \epsilon \otimes e$ and there is no truncation in the sum over representations. This situation prevails when q is not a root of unity. In our situation φ vanishes on a complement of the subspace $(V_i^I \otimes V_j^J) \otimes V_k^K$, and it maps this subspace onto $V_i^I \otimes (V_j^J \otimes V_k^K)$. This map is onto because it has an inverse φ^{-1} which is given by the inverse fusion matrices. This inverse extends to a quasinverse on all of $V_i^I \otimes V_j^J \otimes V_k^K$ by putting $\varphi^{-1} = \varphi^{-1}(id \otimes \Delta)\Delta(e)$. By construction, this quasi-inverse has the defining properties (6. 5), (6. 6) of a quasi-inverse. The property (6. 11) is also true. This completes the proof of the existence of φ with the properties that are required in a weak quasi-triangular quasi-Hopf algebra.

It may be of interest to note that the fusion matrix F which enters into the definition of φ may be computed from the algebra of observables and its morphisms in the algebraic field theory approach [26,33].

In conclusion we find that a weak quasitriangular quasi Hopf algebra is canonically associated with every quantum group algebra $U_q(sl_2)$ with $q^p = -1$, and its (non cocommutative) coproduct reproduces the truncated tensor product of physical representations of $U_q(sl_2)$.

11 Appendix A: Adjoint transformation law

We start from the transformation law (3. 14) of field operators under the algebra \mathcal{G}^* , viz.

$$U(\phi)\Psi_i^J = \sum_{\mathbf{P}} \Psi_m^I \tau_{m,(\phi_p^2)}^J(\phi_p^1) U(\phi_p^2) \quad (11.1)$$

if $\Delta(\phi) = \sum_p \phi_p^1 \otimes \phi_p^2$.

If we take the adjoint of this relation, the order of $U(\cdot)$ and Ψ gets interchanged, so that the resulting relation tells us how to move representation operators $U(\cdot)$ through adjoints of field operators from right to left, rather than from left to right. For the special case of the quantum groups $U_q(sl_2)$ with $|q|=1$, the inverse relation was found in [5],

$$U(\phi)\Psi_i^J = \sum_{\mathbf{P}} \Psi_m^I \tau_{m,(\phi_p^2)}^J(\phi_p^1) U(\phi_p^1), \quad (11.2)$$

where $\bar{\tau}$ is the contragredient representation to τ , defined with the help of the antipodal map \mathcal{S} ,

$$\bar{\tau}(\phi) = {}^t\tau(\mathcal{S}(\phi)). \quad (11.3)$$

We postulate general validity of the transformation law (11. 2) for adjoints of field operators and determine what properties of the antipodal map \mathcal{S} are required to assure this. It is immediately clear that \mathcal{S} must be an antihomomorphism of \mathcal{G}^* because otherwise $\bar{\tau}$ does not have the representation property.

We assume that Δ is a *-homomorphism of algebras with the choice (3. 11) of *-operation on $\mathcal{G}^* \otimes \mathcal{G}^*$, so that

$$\Delta(\phi^*) = \sum_p \phi_p^{2*} \otimes \phi_p^{1*}. \quad (11.4)$$

Substituting ϕ^* for ϕ in eq.(11. 2) and taking its adjoint we find with the help of eq.(11. 4) that transformation law (11. 2) is equivalent to

$$\Psi_i^J U(\phi) = \sum_{\mathbf{P}} U(\phi_p^2) \Psi_i^J \tau_{i,(\phi_p^1)}^J(\mathcal{S}(\phi_p^1)). \quad (11.5)$$

This relation tells us how one moves representation operators $U(\cdot)$ through field operators from right to left. Clearly, consistency of our postulate requires that one should obtain an identity if eq.(11. 5) is substituted into eq.(11. 1), or vice versa.

Let $\sum \Delta(\phi_p^2) = \sum_q \phi_{pq}^{21} \otimes \phi_{pq}^{22}$. Substituting eq.(11. 1) into eq.(11. 5) yields an identity if

$$\sum_{\mathbf{P}} \phi_{pq}^{21} \mathcal{S}(\phi_p^1) \otimes \phi_{pq}^{22} = \epsilon \otimes \phi,$$

where ϵ is the unit element of \mathcal{G}^* , so that $\tau_i^J(\epsilon) = \delta_{ik}$. Written more abstractly, this requirement reads

$$m_r((\mathcal{S} \otimes id \otimes id)(id \otimes \Delta)\Delta(\phi)) = \epsilon \otimes \phi, \quad (11.6)$$

with multiplication operator defined by

$$m_r(\xi \otimes \eta \otimes \zeta) = \eta \xi \otimes \zeta.$$

Similarly, substitution of eq.(11. 5) into eq.(11. 1) yields an identity if

$$m_r((id \otimes \mathcal{S} \otimes id)(id \otimes \Delta)\Delta(\phi)) = \phi \otimes e, \quad (11.7)$$

with multiplication operator defined by

$$m_r(\xi \otimes \eta \otimes \zeta) = \zeta \otimes \eta \xi.$$

It is instructive to verify validity of these relations in the case when \mathcal{G}^* is a group algebra. For group elements a we have $S(a) = a^{-1}$ and $\Delta(a) = a \otimes a$. Therefore

$$(id \otimes \Delta) \Delta(a) = (id \otimes id) a \otimes a = a \otimes a \otimes a,$$

and

$$\begin{aligned} m_r((S \otimes id) id) a \otimes a &= m_r(a^{-1} \otimes a \otimes a) = aa^{-1} \otimes a = e \otimes a, \quad (11. 8) \\ m'_r((id \otimes S) id) a \otimes a &= m'_r(a \otimes a^{-1} \otimes a) = a \otimes a^{-1} a = a \otimes e. \quad (11. 9) \end{aligned}$$

This illustrates that the required properties of the antipodal map S are natural generalizations of the property that it maps a group element into its inverse.

12 Appendix B: Consistency with a numerical R-matrix

We wish to examine what requirements are imposed by consistency of a quantum symmetry with local braid relations (4. 3) with a numerical R-matrix. Both sides of eq.(4. 3) will have to transform in the same way.

We multiply both sides of eq.(4. 3) with $U(\phi)$ from the left, push $U(\cdot)$ through the fields using covariance property (3. 14), and reinsert braid relations (4. 3) on the left hand side. Write

$$\Delta(\phi) = \sum_p \phi_p^1 \otimes \phi_p^2, \quad \Delta(\phi_p^2) = \sum_q \phi_{pq}^{21} \otimes \phi_{pq}^{22}.$$

Then we find that we should have

$$\sum_{p,q} \tau_{im}^j(\phi_p^1) \tau_{un}^l(\phi_{pq}^{21}) U(\phi_{pq}^{22}) R_{nm,ij} = \sum_{p,q} \tau_{ua}^i(\phi_p^1) \tau_{lj}^k(\phi_{pq}^{21}) U(\phi_{pq}^{22}) R_{al,kl}. \quad (12.1)$$

We dropped superscripts $l, j >$ on \mathcal{R} . Suppose that there exists an element R in the algebra,

$$R = \sum_a \tau_a^1 \otimes \tau_a^2 \in \mathcal{G}^* \otimes \mathcal{G}^*, \quad (12.2)$$

such that

$$\mathcal{R}_{lm,ar}^{lj} = (\tau^l \otimes \tau^r)_{lm,ar}(R) = \sum_a \tau_{ia}^l(\tau_a^1) \tau_{ar}^j(\tau_a^2). \quad (12.3)$$

Then requirement (12. 1) will be satisfied if

$$\sum_{p,q} \tau_a^1 \otimes \tau_a^2 \otimes \phi_{pq}^{21} \otimes \phi_{pq}^{22} = \sum_{pq} \phi_{pq}^{21} \otimes \phi_{pq}^{22} \otimes \tau_a^1 \otimes \tau_a^2.$$

Written in more abstract form, this requirement reads

$$(R \otimes \epsilon) \cdot (id \otimes \Delta) \Delta(\phi) = (P_{12}(id \otimes \Delta) \Delta(\phi)) \cdot (R \otimes \epsilon), \quad (12.4)$$

where $P_{12}(a_1 \otimes a_2 \otimes a_3) = a_2 \otimes a_1 \otimes a_3$ and \cdot is the product in $\mathcal{G}^* \otimes \mathcal{G}^* \otimes \mathcal{G}^*$.

Acknowledgements

We are grateful to K.H. Rehren for several discussions. One of us (G.M) would like to thank L. Faddeev and V. Pasquier for explaining their work. We were helped in our understanding of the comodule picture by a seminar of M. Karowski's in Hamburg [34].

V.S. wishes to thank R. Longo for the kind hospitality extended to him at the University of Rome.

Figure captions

Figure 1a. Segments of braids representing elementary operations σ_i and σ_i^{-1} .


Figure 1b. A braid multiplied with its inverse

Figure 2. Deformations of braids


References

- [1] V.G. Drinfel'd, *Quasi Hopf algebras and Knizhnik Zamolodchikov equations*, in: Problems of modern quantum field theory, Proceedings Alushta 1989, Research reports in physics, Springer Verlag Heidelberg 1989
- [2] J. Fröhlich, *Statistics of fields, the Yang-Baxter equation and the theory of knots and links*, in: Nonperturbative quantum field theory, G.'tHooft et al.(eds.), Plenum Press 1988
- [3] R. Dijkgraaf, V. Pasquier, P. Roche, *Quasi-quantum groups related to orbifold models*, in: Proc. Intl. Colloquium on modern quantum field theory, TATA Institute of Fundamental Research, Bombay, January 1990
- [4] D.Buchholz, G.Mack, I.T. Todorov, as reported in I. Todorov, in: Proceedings of Conf. on Quantum Groups, Clausthal Zellerfeld, July 1989 (reviewed in ref.[5]).
- [5] G. Mack, V. Schomerus, *Conformal field algebras with quantum symmetry from the theory of superselection sectors*, Commun. Math. Phys. **134**, 139 (1990)
- [6] G. Mack, V. Schomerus, Addendum to ref.[5], to be submitted to Commun. Matl. Phys.
- [7] V.G. Drinfel'd, *Quantum groups*, Proc. ICM (1987) 798
- [8] S.L. Woronowicz, *Compact matrix pseudogroups*, Commun. Math. Phys. **615** (1987)
- [9] S.L. Woronowicz, *Twisted $SU(2)$ group. An example of a Noncommutative differential calculus*, Publ. RIMS-D-Kyoto **23** 117 (1987)
- [9] L.D. Faddeev, *From integrable models to quantum groups*, Schladming lectures 1990

- [10] Yu.I. Manin, *Quantum groups and Non-commutative geometry*, Proc. Int. Congr. Math., Berkeley 1, 798 (1986)
- [11] J. Wess, B. Zumino, *Covariant differential calculus on the quantum hyperplane*, preprint CERN TH 5697/90 April 1990
- [12] L.D. Faddeev, N.Yu. Reshetikhin, L.A. Takhtajan, *Quantization of Lie groups and Lie algebras*, preprint LOMI-E-14-87 Leningrad 1987
- [13] L.A. Takhtajan, *Quantum groups and integrable models*, Advanced studies in pure mathematics 19 (1989)
- [14] G. Moore, N.Yu. Reshetikhin, *A comment on quantum symmetry in conformal field theory*, Nucl. Phys. **B328**, 557 1989
- [15] N.Yu Reshetikhin and F. Smirnov, *Hidden quantum symmetry and integrable perturbations of conformal field theory*, Commun. Math. Phys. **131**, 157 (1990)
- [16] V. Pasquier, H. Saleur, *Common structures between finite systems and conformal field theories through quantum groups*, Nucl.Phys. **B330**, 523 (1990)
- [17] H. Saleur, J.B. Zuber, *Integrable lattice models and quantum groups*, lectures at the 1990 spring school on string theory and quantum gravity, ICTP Trieste, April 1990
- [18] H. Saleur, *Lattice models and conformal field theories*, Phys. Rep. **184** 177 (1989)
- [19] L.Alvarez-Gaume, C.Gomez, G. Sierra, *Hidden quantum symmetry in rational conformal field theories*, Nucl.Phys. **B310** (1989)
- L.Alvarez-Gaumé, C.Gomez, G. Sierra, *Quantum group interpretation of some conformal field theories*, Phys. Lett. **220B** (1989) 142
- L.Alvarez-Gaumé, C.Gomez, G.Sierra, *Duality and quantum groups*, Nucl.Phys. **B330**, 347 (1990)
- [20] P.Furlan, A.Ch. Ganchev, V. Petkova, *Quantum groups and fusion rules multiplicities* preprint INFN/AE-89/15, Trieste 1989
- [21] K. Gawedzki *Classical origin of quantum group symmetries in Wess-Zumino-Witten conformal field theories*, IHES/P/90/92 Nov.1990
- [22] G. Felder, Ch. Wierczkowski, *Topological representations of the quantum group $U_q(sl_2)$* , ETH-TH 90-52 Zürich 1990
- [23] N.Yu. Reshetikhin, L.A. Takhtajan, L.D. Faddeev, *Algebra and Analysis* **1**, 178 (1989)
- [24] A.N.Kirillov, N. Reshetikhin, *Representations of the algebra $U_q(sl(2))$, q -orthogonal polynomials and invariants of links*, preprint LOMI-E-9-88, Leningrad 1988
- [25] G.G. Wick, E.P. Wigner, A.S. Wightman, *Intrinsic parity of elementary particles* Phys. Rev. **88** (1952) 101
- [26] K.Fredenhagen, K.H.Rehren, B.Schroer, *Superselection sectors with braid group statistics and exchange algebras, I, general theory*, Commun.Math.Phys. **125**, 201 (1989)
- [27] J. Fröhlich, F. Gabbiani, P.A. Marchetti *Braid statistics in 3-dimensional local quantum theory* preprint ETH-TH/89-36 Zürich 1989 / / - - *Superselection structure and statistics in 3-dimensional local quantum theory* preprint ETH-TH/89-22 Zürich 1989
- [28] D. Kastler (ed.), *The algebraic theory of superselection sectors. Introduction and recent results* World Scientific, Singapore 1990
- [29] B. Schroer, private communication
- K.H.Rehren, *Field operators for Plektons* preprint Hamburg 1991
- [30] L.Hadjivanov, R.Paunov, I.T.Todorov, *Quantum group extended chiral p -models* preprint 1990 (submitted to Nucl. Phys.)
- [31] A. Schirmacher, J. Wess, B. Zumino, *The two parameter deformation of $GL(2)$, its differential calculus and Lie algebra* Z.Phys.C (to appear)
- [32] G. Keller, *Fusion rules of $U_q(sl(2, C))$, $q^n = 1$* preprint MPI-PAE/PTh 47/90, München 1990
- [33] G. Mack, V. Schomerus, *Endomorphisms and quantum symmetry of the conformal Jacq model* in: D. Kastler (ed), *The algebraic theory of superselection sectors. Introduction and recent results*: World Scientific, Singapore 1990
- [34] M. Karowski, *Quantum groups and integrable models*, lecture presented at DESY Hamburg, 1990
- [35] I.M. Gelfand, G.E. Shilov, *Generalized functions*, vol 1, Academic press, New York 1964



σ_i



σ_i^{-1}

Figure 1a






Figure 1b



$\sigma_3 \sigma_1 = \sigma_1 \sigma_3$



$\sigma_2 \sigma_1 \sigma_2 = \sigma_1 \sigma_2 \sigma_1$

Figure 2