

DEUTSCHES ELEKTRONEN – SYNCHROTRON

DESY 91-048
Cavendish-HEP-90/26
May 1991



HERWIG - A Monte Carlo Event Generator for Simulating Hadron Emission Reactions with Interfering Gluons

G. Marchesini

Dipartimento di Fisica, Università di Parma

and

I.N.F.N., Gruppo di Parma

B.R. Webber, M.H. Seymour

Cavendish Laboratory, University of Cambridge

G. Abbiendi, L. Stanco

Dipartimento di Fisica, Università di Padova

and

I.N.F.N., Sezione di Padova

I.G. Knowles

Dept. of Physics and Astronomy, University of Glasgow


ISSN 0418-9833

NOTKESTRASSE 85 · D-2000 HAMBURG 52

DESY behält sich alle Rechte für den Fall der Schutzrechtserteilung und für die wirtschaftliche Verwertung der in diesem Bericht enthaltenen Informationen vor.

DESY reserves all rights for commercial use of information included in this report, especially in case of filing application for or grant of patents.

To be sure that your preprints are promptly included in the
HIGH ENERGY PHYSICS INDEX,
send them to the following (if possible by air mail):



Program Summary

HERWIG

a Monte Carlo event generator for simulating
Hadron Emission Reactions With Interfering Gluons*

Version 5.1—April 1991
by

G. Marchesini¹, B.R. Webber²,
G. Abbiendi³, I.G. Knowles⁴, M.H. Seymour² and L. Stanco³.

¹Dipartimento di Fisica, Università di Parma, and I.N.F.N., Gruppo di Parma.

²Cavendish Laboratory, University of Cambridge.

³Dipartimento di Fisica, Università di Padova, and I.N.F.N., Sezione di Padova.

⁴Department of Physics and Astronomy, University of Glasgow.

Abstract

HERWIG is a general-purpose particle physics event generator, which includes the simulation of hard lepton-lepton, lepton-hadron scattering and soft hadron-hadron collisions in one package. It uses the parton-shower approach for initial-state and final-state QCD radiation, including colour coherence effects and azimuthal correlations both within and between jets.

This article includes a brief review of the physics underlying HERWIG, followed by a description of the program itself. This includes details of the input and control parameters used by the program, and the output data provided by it. Sample output from a typical simulation is given and annotated.

Correspondence about this program should be sent by e-mail to one of the authors at the following addresses:

Decnet: vxpr:marchesini, cbhep:webber
cbhep:seymour, vxfpd:stanco
Bitnet: webber@phy.hep.cam.ac.uk
seymour@phy-hep.cam.ac.uk
gapas3@vms.gla.ac.uk

Title of the Program: HERWIG
Catalogue Number:
Program obtainable from: CPC Program Library, Queen's University of Belfast, N. Ireland
(see application form in this issue).

Computers on which the program is operable: VAX. Only minor modifications are needed to run on any machine running standard FORTRAN 77.

Operating system: VAX-VMS, and any other system which supports FORTRAN 77.

High speed storage required: 240 kwords.

No. of bits in a word: 32.

No. of lines in combined program and test deck: 10130.

Keywords: Monte Carlo, event generator, perturbative QCD, parton shower, coherent branching, fragmentation, hadronic final states.

Nature of the physical problem: Simulation of high energy particle collisions, especially those in which a high momentum transfer gives rise to the emission of jets of hadrons. The theoretical basis is the resummation of the perturbative QCD expansion including all asymptotic and some important subasymptotic contributions.

Method of solution: Monte Carlo simulation of a variety of primary collision processes, followed by parton shower generation using a coherent branching algorithm and hadron formation using a cluster fragmentation model.

Restriction on the complexity of the problem: To run at very high energy ($\sqrt{s} \gg 40$ TeV), the program would need recompiling with larger arrays to accommodate the high multiplicity expected.

Typical running time: Depends crucially on the process modelled. Can range from 1000 LEP $q\bar{q}$ events in 10 min, to 1000 SSC QCD $2 \rightarrow 2$ events in 3 hrs. Both times are for a 1 VUP (i.e. approximately 1 MIP) machine.

Reference to other published version of this program: CERN Program Library long write-up W5037.

*Submitted to Computer Physics Communications. Work supported in part by the U.K. Science and Engineering Council, and in part by the Italian Ministero della Pubblica Istruzione.

Contents

1 Introduction

1	Introduction	3
2	Physics Underlying HERWIG	4
2.1	Final state emission	6
2.2	Initial state emission	6
2.3	Elementary hard subprocess	7
2.4	Hadronization process	8
2.5	Cluster hadronization model	8
2.6	Underlying soft event	9
3	New Features of Recent Versions	10
4	Program Structure	11
5	Beams and Processes	14
6	Input Parameters	17
7	Common Block File	19
8	Form Factor File	22
9	Event Data	24
10	Status Codes	24
11	Event Weights	25
12	Heavy Flavour Decays	27
13	QCD Hard Subprocesses	28
14	QCD Direct Photon Subprocesses	29
15	Electroweak Subprocesses	29
16	Including New Subprocesses	31
17	Error Conditions	32
18	Sample Output	33
19	Guide to Sample Output	38
20	Subroutine Descriptions	45
	References	48

HERWIG is a general-purpose event generator for high energy hadronic processes, with particular emphasis on the detailed simulation of QCD parton showers. The program has the following special features:

- Simulation of hard lepton-lepton, lepton-hadron and hadron-hadron scattering and soft hadron-hadron collisions in one package.
 - Colour coherence of partons (initial and final) in hard subprocesses.
 - Heavy flavour hadron production and decay with QCD coherence effects.
 - QCD jet evolution with soft gluon interference via angular ordering.
 - Backward evolution of initial state partons including interference.
 - Azimuthal correlations within and between jets due to interference.
 - Azimuthal correlations within jets due to gluon polarization.
 - Cluster hadronization of jets via non-perturbative gluon splitting.
 - A similar cluster model for soft and underlying hadronic events.
- The program operates by setting up parameters in common blocks and then calling a sequence of subroutines to generate an event. Parameters not set in the main program HWIGPR are set to default values in the main initialisation routine HWIGIN. To generate events the user must first set up the beam particle names PART1, PART2 (type CHARACTER*4), in the common block /HWBEAM/, and the beam momenta PBEM1, PBEM2 (in GeV/c), a process code IPROC and the number of events required MAXEV in /HWPROC/. See Sect. 5 for beams and processes available.
- All analysis of generated events (histogramming, etc.) should be performed by the user-provided routines HWABEG (to initialise), HWANAL (to analyse an event) and HWAEND (to terminate). The default HWANAL subroutine writes event and jet information and stable particle data on unit LWEVT defined in HWIGIN (or simply returns if LWEVT = 0). See HWANAL for details of event information written.
- A detailed event summary is printed out for the first MAXPR events (default MAXPR = 1). Set IPRINT = 2 to list the particle identity codes and (simplified) particle decay schemes used in the program.
- The programming language is standard Fortran 77 as far as possible. However, the following may require modification for running on computers other than Vax's:
- Most common blocks are inserted by INCLUDE 'HERWIG51.INC' Vax Fortran statements (see Sect. 7 for contents of HERWIG51.INC).
 - Many common blocks are initialized by BLOCK DATA HWUDAT. Although BLOCK DATA is standard Fortran 77, it can cause linkage problems for some systems.
 - Subroutine HWUTIM (returning CPU time left) is machine dependent.

2 Physics Underlying HERWIG

The physics that underlies the original program was presented in detail in Ref. [1]. More recent improvements are discussed in Refs. [2–7]. Other relevant theoretical background may be found in Refs. [8–12]. We limit ourselves here to a review of the key components of the program and their theoretical basis.

The main theoretical justification for QCD Monte Carlo simulations lies in the factorization theorems for hard processes. This property is illustrated in Fig. 1 for the process $\bar{p}p \rightarrow W^+ + X$, with $W^+ \rightarrow t\bar{b}$.[†] Quarks and antiquarks are represented by single colour lines, gluons by double lines (the so-called planar approximation). Dotted lines represent colour-singlet particles (W bosons or leptons). Note that Fig. 1 is not a Feynman diagram: it represents the coherent sum of many real and virtual diagrams which are summed by the branching algorithm.

A process such as that in Fig. 1 can be factorized into the following subprocesses.

1. *Final state emission.* An outgoing virtual parton with large time-like mass generates a shower of partons with lower virtuality. The amount of emission depends on the upper limit on the virtual mass of the initiating parton, which is controlled by the momentum transfer scale Q of the hard subprocess, to be discussed below.
2. *Initial state emission.* A parton constituent of an incident hadron with low space-like virtuality radiates time-like partons. In the process it decreases its energy to a fraction x of that of the hadron, and increases its space-like virtual mass. This mass is bounded in absolute value by the scale Q of the hard subprocess. The initial state emission process leads to the evolution of the structure function $F(x, Q)$ of the incident hadron.
3. *Elementary hard subprocess.* This can be computed exactly to finite order in perturbation theory. For the process of Fig. 1 it is given by the $q\bar{q} \rightarrow W \rightarrow q\bar{q}$ amplitude squared. The hard subprocess momentum transfer scale Q , given here by the mass of the virtual W^+ , sets the boundary conditions for the initial and final state parton showers. A large variety of QCD and electroweak hard processes have the same basic structure as in Fig. 1, with different elementary subprocess matrix elements.
4. *Hadronization process.* In order to construct a realistic simulation one needs to convert the partons into hadrons. This process takes place at a low momentum transfer scale, for which the strong coupling α_s is large and perturbation theory is not applicable. Therefore we need to add to the above perturbative QCD processes a phenomenological hadronization model which fortunately (see later) does not conceal the perturbative structure.

An important consequence of the factorization theorem is that the distributions for any hard QCD process are obtained from the same four subprocesses described above. In lepton-lepton collisions we have only to consider the elementary and final state emission subprocesses, while in hadron-hadron collisions we need to consider also the emission from one incoming parton. In hadron-hadron collisions there is emission from two incoming partons: one from each hadron. In addition, in processes involving incoming hadrons there may be soft emission due to the presence of ‘spectator’ partons.

[†]This event was generated using an earlier version (2.2) of HERWIG. The decays of heavy quarks are handled slightly differently in version 5.1 – see Sect. 12. In addition we now know that the top quark is too heavy for the W to decay to $t\bar{b}$. Nevertheless, Fig. 1 illustrates the typical branching and colour structure of a simulated event.




Figure 1: Colour structure of a $\bar{p}p \rightarrow W^+ + X$, $W^+ \rightarrow t\bar{b}$ event.

As a consequence of factorization, one can construct a single Monte Carlo program, such as HERWIG, which can in principle simulate all hard processes. The main theoretical advantage of such a universal program is that the phenomenological parameters in the hadronization model can be tuned simultaneously by fitting data at different machines and energies. This enhances the predictive power of the program.

If one studies only totally inclusive quantities, the hard process of Fig. 1 is infrared finite. However we are interested also in exclusive distributions and for these we need to introduce a cutoff. In HERWIG this is done by requiring that the partons are emitted with a transverse momentum larger than some finite value Q_0 , which is selected in such a way that $\alpha_s(Q_0)$ is still a small number. (In fact the cutoff depends on the type of parton involved, as explained in Sect. 6, but we can take Q_0 here to represent the smallest cutoff.) As observed in some next-to-leading analyses [13], a transverse momentum cutoff corresponds to the $\overline{\text{MS}}$ scheme for the regularization of QCD.

In the Monte Carlo simulation one describes the entire process, within the resolution implied by the cutoff Q_0 , at the exclusive level. This can be done because the final and initial state emission processes also have a factorized structure, as we shall illustrate. Factorization in these cases holds to the leading order in collinear and infrared logarithms. However, as we shall recall, for some important distributions factorization can be extended beyond leading order.

We now discuss the above subprocesses separately in more detail.

2.1 Final state emission

Parton emission factorizes as a successive branching process which is characterized by the following properties:

1. The energy fractions are distributed according to the Altarelli-Parisi splitting functions.
2. The full available phase space is restricted to an angular-ordered region. Such a restriction is the result of interference and takes leading infrared singularities correctly into account. At each branching, the angle between the two emitted partons is smaller than that of the previous branching.
3. The emission angles are distributed according to the Sudakov form factors, which sum the virtual corrections. The Sudakov form factor normalizes the branching distributions to give the probabilistic interpretation needed for a Monte Carlo simulation. This fact is a consequence of field theory, in particular of unitarity and of the infrared finiteness of inclusive quantities.
4. The azimuthal angular distribution in each branching is determined by two effects: a) for a soft emitted gluon the azimuth is distributed according to the eikonal dipole distribution [1]; b) for non-soft emission one finds azimuthal correlations due to spin effects. See [2-4] for the method used to implement these correlations in full, to leading collinear logarithmic accuracy, in HERWIG.
5. In each branching the scale of α_S is the relative transverse momentum of the two emitted partons.
6. In the case of heavy flavour production the mass of the quark modifies the angular-ordered phase space. The most important effect is that the soft radiation in the direction of the heavy quark is depleted. One finds that the emission within an angle of order M/E vanishes (with M and E the mass and energy of the heavy quark emitting the soft gluon). This angular screening determines [5,14,15] the shape of the heavy flavour jet, which could prove to be an important signature. The angular screening can easily be taken into account in the coherent branching algorithm and is consistently included in HERWIG [5].

A branching algorithm characterized by the above properties will be called *coherent final state branching* and, in general, gives parton distributions which are correct to leading infrared order. In particular this algorithm correctly describes intra-jet distributions such as the string effects in three-jet events [16].

The coherent branching correctly describes [17-19] also the next-to-leading corrections to the distributions of soft partons, i.e. partons with momentum small compared with the hard scale Q but still large compared with the QCD scale Λ_{MS} . This fact is very important because the perturbative expansion for these distributions is singular. Therefore the next-to-leading corrections are large and for a reliable calculation at present energies they need to be taken into account. The importance of the next-to-leading contributions is clear in the LEP data

on the multiplicity and inclusive distributions in the soft region [20].

In the literature one can find different prescriptions for the branching subprocesses (with different phase space and/or argument of α_S , or without the angular screening due to the heavy quark mass). Some of these algorithms are based on old and partial results of perturbative QCD studies, and do not give the correct results even to leading infrared order.

2.2 Initial state emission

The theoretical analysis of this process is more complex than for the final-state case. Even to leading order, the structure function and associated radiation have only been analyzed quite recently for small x [21], x being the energy fraction of the incoming parton after the emission of initial state radiation. For lepton-hadron processes x corresponds to the Bjorken variable, while for hadron-hadron processes x is related to Q^2/s .

The main result is that for any value of x , even for $x \rightarrow 0$, the initial state emission process factorizes and can be described as a branching process suitable for Monte Carlo simulations, which we shall call *coherent initial state branching*. The properties which characterize this branching include all the properties discussed above for the final state emission. In the initial state emission the angular ordering restriction of the phase space (point 2 in the previous list) applies to the angles θ_i between the incoming hadron and the emitted partons i .

In the case of small values of x , the initial state branching process has the following additional properties, which are however not yet fully included in HERWIG:

7. For small x there are virtual corrections which are not included in the Sudakov form factors. These corrections are important in the case in which the energy fraction z_i of an exchanged space-like gluon is soft ($z_i \rightarrow 0$). In this case the Altarelli-Parisi splitting function contributes with a singular terms of the type C_A/z_i or C_F/z_i , according to whether the exchanged gluon is generated by a gluon or a quark. These virtual corrections exponentiate, giving a *non-Sudakov form factor* [21]. The important feature of this form factor is that it screens the $1/z_i$ singularity. This is the same effect as that of the Regge trajectory in the Lipatov equation [22].

8. A second important effect in the branching with a soft exchanged gluon is related to the angular ordering. For small z_i the angular ordering condition $\theta_{i+1} > \theta_i$ gives $q_{i+1} > z_i q_i$, where q_i is the transverse momentum of the i -th emission. Therefore for $z_i \rightarrow 0$ the lower bound on q_{i+1} vanishes, giving singular $\ln z_i$ contributions. However, these singular terms are cancelled by contributions from the non-Sudakov form factor.

A branching algorithm which includes these properties for small x leads to a structure function which satisfies the Lipatov equation for $x \rightarrow 0$ and the Altarelli-Parisi equation for finite x . Such an algorithm has been used to begin the construction of a new Monte Carlo simulation program [23], but much further work is needed before these developments can be included in a fully exclusive event generator.

For the moment, in HERWIG we take into account the fact that the most important singularities at small x , coming from the region $z_i q_i < q_{i+1} < q_i$ for $z_i \rightarrow 0$, are partially cancelled by contributions from the non-Sudakov form factor. Thus we ignore the non-Sudakov form factor and correspondingly reduce the phase space to the q -ordered region $q_i < q_{i+1}$ for $z_i \rightarrow 0$. This algorithm corresponds to neglecting non-leading singular contributions but generates an anomalous dimension which is correct in this region up to three loops.

For large x , the coherent branching correctly sums [6] not only the leading but also the

next-to-leading contributions. This accuracy allows us to identify the relation between the QCD scale used in the Monte Carlo program and the fundamental parameter $\Lambda_{\overline{\text{MS}}}$. This is achieved by using the one-loop Altarelli-Parisi splitting functions and the two-loop expression for α_S with the following universal relation between the scale parameter Λ_{MC} used in the simulation and $\Lambda_{\overline{\text{MS}}}$

$$\Lambda_{\text{MC}} = \exp \left(\frac{67 - 3\pi^2 - 10N_f/3}{2(33 - 2N_f)} \right) \Lambda_{\overline{\text{MS}}} \simeq 1.569 \Lambda_{\overline{\text{MS}}} \quad \text{for } N_f = 5. \quad (2.1)$$

Therefore a Monte Carlo simulation with next-to-leading accuracy can be used to determine $\Lambda_{\overline{\text{MS}}}$ from semi-inclusive data at large momentum fractions.

2.3 Elementary hard subprocess

In HERWIG version 5.1 there is a fairly large library of QCD and electroweak elementary subprocesses (see Sect. 5).

From the point of view of QCD coherence, the elementary subprocess plays an important rôle in defining the phase space of the initial and final state branching subprocesses. As we have seen, these branchings are ordered in angle from a maximum to a minimum value. The minimum values is fixed by the cutoff Q_0 , but the maximum value is determined by the elementary subprocess and is due to interference among soft gluons. The general result [1,17,24] is that the initial and final branchings are approximately confined within cones around the incoming and outgoing partons from the elementary subprocess. For the branching of parton i , the aperture of the cone is defined by the direction of the other parton j which is colour-connected to i .

For a general process there are various contributions with different colour connections. The HERWIG library of elementary subprocesses includes the separate colour connection contributions. See [1] for a discussion of the most complex case, namely the $2 \rightarrow 2$ QCD subprocesses generating two-jet events with high transverse energy. The relation between soft gluon interference and the colour connection structure of the elementary subprocess leads to detectable effects, such as the string effect in three-jet events [16].

Another important function of the elementary subprocess is to set up the polarizations of any electroweak bosons or gluon jets that may be involved. These polarizations give rise to angular asymmetries and correlations in boson decays and jet fragmentation. They are included in HERWIG for most of the subprocesses provided (but not yet for $W +$ jet production), using the approach of Refs. [2-4] to generate all correlations in jet fragmentation to leading-logarithmic accuracy.

2.4 Hadronization Process

For the most complicated hard processes such as that in Fig. 1, we have three type of non-perturbative contributions to consider: (a) the representation of the incoming partons as constituents of the incident hadrons; (b) the conversion of the emitted partons into outgoing hadrons; (c) the ‘underlying soft event’ associated with the presence of spectator partons.

We refer to all of these as aspects of ‘hadronization’. We first recall briefly the relevant features of perturbative field theory, then discuss the models used for (b) and (c) in the following Subsections.

The treatment of the *incoming parton* is related to the factorization theorem for collinear singularities. The distribution of the parton’s longitudinal momentum fraction at a low space-like scale Q , is described by a phenomenological input structure function. The transverse momentum at such a scale has a distribution characteristic of the size of the hadron.

Perturbative studies do not provide information on the confinement mechanism which converts the *emitted partons* into hadrons below the time-like cutoff scale Q_0 . However, provided this mechanism does actually exist, perturbative QCD predicts [25,26] that in hard processes confinement of partons is *local in colour* and *independent* of the hard scale Q . This ‘preconfinement’ property is due to the Sudakov form factor. In QED this form factor depletes the cross section for emission of a single charge without an accompanying cloud of photons, within a given resolution. Similarly, in QCD the Sudakov form factor inhibits the separation of the colour charges forming a singlet. In the emission of jets of partons, one finds [25] in perturbative QCD, as well as in the Monte Carlo simulations [9], that the mass distribution of two partons forming a colour singlet is concentrated around values of the order of Q_0 and is independent of the hard scale Q for large Q .

This property is confirmed by the phenomenological analysis of jet fragmentation by the Leningrad group [26]. In this analysis the fragmentation functions for π , K and p production in e^+e^- annihilation at various high energies Q were compared with the fragmentations function for partons, computed by resumming leading perturbative contributions. The result is that these fragmentation functions are proportional to the parton-fragmentation functions. The proportionality constants K_π , K_K and K_p are independent of Q and correspond to the hadronization conversion factors from partons to hadrons. Thus partons are converted into hadrons locally in phase space, a property that has been named Local Parton Hadron Duality (LPHD).

2.5 Cluster hadronization model

The preconfinement property is used in HERWIG by assuming a cluster hadronization model which is local in colour and independent of the hard process and the energy [1,10]. After the perturbative parton branching process, all outgoing gluons are split non-perturbatively, as shown in Fig. 1, into light (u or d) quark antiquark or diquark antiquark pairs (the default option is to suppress diquark splitting). At this point, each jet consists of a set of outgoing quarks and antiquarks (also possibly some diquarks and antidiquarks), and, in the case of spacelike jets, a single incoming valence quark or antiquark. The latter is replaced by an outgoing spectator carrying the opposite colour and the residual flavour and momentum of the corresponding beam hadron.

As may be seen in Fig. 1, a colour line can now be followed, in the planar approximation, from each quark to an antiquark or diquark with which it can form a colour-singlet cluster. These clusters satisfy the ‘pre-confinement’ discussed above — they have a distribution of mass and spatial size that peaks at low values, falls rapidly for large cluster masses and sizes, and is asymptotically independent of the hard subprocess scale.

The clusters thus formed are fragmented into hadrons. If a cluster is too light to decay into two hadrons, it is taken to represent the lightest single hadron of its flavour. Its mass is shifted to the appropriate value by an exchange of momentum with a neighbouring cluster in the jet. Those clusters massive enough to decay into two hadrons, but below a fission threshold to be specified below, decay isotropically into pairs of hadrons selected in the following way:

a flavour f is chosen at random from among u , d , s , the six corresponding diquark flavour combinations, and c . For a cluster of flavour $f_1\bar{f}_2$, this specifies the flavours $f_1\bar{f}$ and $f\bar{f}_2$ of the decay products, which are then selected at random from tables of hadrons of those flavours. The hadrons can be $J^P = 0^-$, 1^\pm or 2^+ mesons, or $\frac{1}{2}^+$ or $\frac{3}{2}^+$ baryons. For charmed hadrons, some of these states are just educated guesses. For b - and t -flavoured hadrons, only $J^P = 0^-$ mesons and $\frac{1}{2}^+$ baryons are included, and their binding energies are neglected. No diquark-antidiquark combinations are allowed. The selected choice of decay products is accepted in proportion to the density of states (phase space times spin degeneracy) for that channel. Otherwise, f is rejected and the procedure is repeated. In this way one obtains an unbiased selection of decay products that conserve flavour.

A small fraction of clusters have masses too high for isotropic two-body decay to be a reasonable ansatz, even though the cluster mass spectrum falls rapidly (faster than any power) at high masses. These are fragmented using an iterative fission model until the masses of the fission products fall below the fission threshold. In the fission model, the produced flavour f is limited to u , d or s and the product clusters $f_1\bar{f}$ and $f\bar{f}_2$ move in the directions of the original constituents f_1 and \bar{f}_2 in their c.m. frame. Thus the fission mechanism is not unlike string fragmentation [27]. There are two fission parameters, CLMAX and PSPLIT. The maximum cluster mass parameter CLMAX specifies the fission threshold M_f according to the formula

$$M_f^2 = \text{CLMAX}^2 + (Q_1 + Q_2)^2 \quad (2.2)$$

where Q_1 and Q_2 are the virtual mass cutoffs corresponding to flavours f_1 and f_2 . The parameter PSPLIT specifies the mass spectrum of the produced clusters, which is taken to be M_{PSPLIT}^2 within the allowed phase space. Provided the parameter CLMAX is not chosen too small (typically it is about 4 GeV), the gross features of events are insensitive to the details of the fission model, since only a small fraction of clusters undergo fission. However, the production rates of high- p_T or heavy particles (especially baryons) are affected, because they are sensitive to the tail of the cluster mass distribution.

2.6 Underlying soft event

In hadron-hadron and lepton-hadron collisions there are ‘beam clusters’ containing the spectators from the incoming hadrons. In the formation of beam clusters, the colour connection between the spectators and the initial-state parton showers is cut by the forced emission of a soft quark-antiquark pair. The underlying soft event in a hard hadron-hadron collision is then assumed to be a soft collision between the two beam clusters. In a lepton-hadron collision the corresponding ‘soft hadronic remnant’ is represented by a soft collision between the beam cluster and the adjacent cluster, i.e. the one produced by the forced emission mentioned above.

The necessity of adding an underlying soft event to the hard emission described in Subsect. 2.1 and 2.2 was analysed in Ref. [12], in which the “pedestal height” in hadronic jet production, i.e. the mean transverse energy per unit rapidity accompanying a high-transverse-energy jet, was studied. The observed pedestal height and its dependence on jet transverse energy [28] are accounted for by superposing on the hard emission an underlying event structure similar to that of a minimum-bias collision.

The model used for the underlying event is based on the minimum-bias $p\bar{p}$ event generator of the UA5 Collaboration [29], modified to make use of our cluster fragmentation algorithm.

combinations, and c . For a cluster of flavour $f_1\bar{f}_2$, this specifies the flavours $f_1\bar{f}$ and $f\bar{f}_2$ of the decay products, which are then selected at random from tables of hadrons of those flavours. The hadrons can be $J^P = 0^-$, 1^\pm or 2^+ mesons, or $\frac{1}{2}^+$ or $\frac{3}{2}^+$ baryons. For charmed hadrons, some of these states are just educated guesses. For b - and t -flavoured hadrons, only $J^P = 0^-$ mesons and $\frac{1}{2}^+$ baryons are included, and their binding energies are neglected. No diquark-antidiquark combinations are allowed. The selected choice of decay products is accepted in proportion to the density of states (phase space times spin degeneracy) for that channel. Otherwise, f is rejected and the procedure is repeated. In this way one obtains an unbiased selection of decay products that conserve flavour.

A small fraction of clusters have masses too high for isotropic two-body decay to be a reasonable ansatz, even though the cluster mass spectrum falls rapidly (faster than any power) at high masses. These are fragmented using an iterative fission model until the masses of the fission products fall below the fission threshold. In the fission model, the produced flavour f is limited to u , d or s and the product clusters $f_1\bar{f}$ and $f\bar{f}_2$ move in the directions of the original constituents f_1 and \bar{f}_2 in their c.m. frame. Thus the fission mechanism is not unlike string fragmentation [27]. There are two fission parameters, CLMAX and PSPLIT. The maximum cluster mass parameter CLMAX specifies the fission threshold M_f according to the formula

$$M_f^2 = \text{CLMAX}^2 + (Q_1 + Q_2)^2 \quad (2.2)$$

where Q_1 and Q_2 are the virtual mass cutoffs corresponding to flavours f_1 and f_2 . The parameter PSPLIT specifies the mass spectrum of the produced clusters, which is taken to be M_{PSPLIT}^2 within the allowed phase space. Provided the parameter CLMAX is not chosen too small (typically it is about 4 GeV), the gross features of events are insensitive to the details of the fission model, since only a small fraction of clusters undergo fission. However, the production rates of high- p_T or heavy particles (especially baryons) are affected, because they are sensitive to the tail of the cluster mass distribution.

2.6 Underlying soft event

In hadron-hadron and lepton-hadron collisions there are ‘beam clusters’ containing the spectators from the incoming hadrons. In the formation of beam clusters, the colour connection between the spectators and the initial-state parton showers is cut by the forced emission of a soft quark-antiquark pair. The underlying soft event in a hard hadron-hadron collision is then assumed to be a soft collision between the two beam clusters. In a lepton-hadron collision the corresponding ‘soft hadronic remnant’ is represented by a soft collision between the beam cluster and the adjacent cluster, i.e. the one produced by the forced emission mentioned above.

The necessity of adding an underlying soft event to the hard emission described in Subsect. 2.1 and 2.2 was analysed in Ref. [12], in which the “pedestal height” in hadronic jet production, i.e. the mean transverse energy per unit rapidity accompanying a high-transverse-energy jet, was studied. The observed pedestal height and its dependence on jet transverse energy [28] are accounted for by superposing on the hard emission an underlying event structure similar to that of a minimum-bias collision.

The model used for the underlying event is based on the minimum-bias $p\bar{p}$ event generator of the UA5 Collaboration [29], modified to make use of our cluster fragmentation algorithm.

The model starts from a parametrization of the $p\bar{p}$ inelastic charged multiplicity distribution as a negative binomial. As an option, for underlying events the value of \sqrt{s} used to choose the multiplicity n may be enhanced by a parameter ENSOF to allow for an enhanced underlying activity in hard events. The actual charged multiplicity is then taken to be n plus the sum of the moduli of the charges of the colliding hadrons or clusters. Next the clusters are hadronized using the model described above.

If the charged multiplicity from the beam clusters is equal to the selected value, no further clusters are created. If it is greater than the selected value, the beam cluster hadronization is redone. If it is less, ‘soft clusters’ $q_1\bar{q}_2$, $q_2\bar{q}_3$, ... are created and hadronized until the selected multiplicity is either reached, in which case cluster production stops, or else overshoot, in which case the whole procedure is repeated, starting from the beam cluster hadronization. The produced quarks q_i that define the flavour of the soft clusters are taken to be u or d only. The cluster masses are chosen from the distribution

$$P(M) \propto (M - M_0) \exp[-a(M - M_0)] \quad (2.3)$$

where $M_0 = 1$ GeV and $a = 2$ GeV $^{-1}$. Since we use the same hadronization model for soft clusters as for those that come from parton branching, our scheme differs from the original UA5 Monte Carlo in requiring no further parameters to specify the hadron distribution from a cluster of a given mass and flavour.

Once the preselected charged multiplicity has been achieved, the cluster momenta are generated with a simple longitudinal phase-space distribution and limited transverse momenta, as explained in Refs. [1, 29].

Unstable hadrons from clusters produced in both the hard and soft components of the event decay according to simplified decay schemes, which can be tabulated by specifying the print option IPRINT = 2. All decays are assumed to be quasi-two or -three body and modes are invented where necessary to make the branching ratios add up to 100%. Phase space distributions are assumed except for the decays of b and t -flavoured hadrons, for which a spectator model with charged-current decay of the heavy quark is assumed. Note that in the case of top decay the t quark is assumed to hadronize and depolarize before decaying, a model which could be unreliable for some distributions if the t mass is greater than about 130 GeV. After a b or t decay, secondary parton showers are produced by ongoing partons as discussed in Ref. [5]; these are hadronized in the same way as primary jets.

3 New Features of Recent Versions

We give here a summary of the main new features of HERWIG versions 4.1, 4.2, 4.3, 4.6, 5.0 and 5.1, all introduced since the preliminary program description [30].

For version 4.1, the program was largely rewritten to conform with the standards laid down by the Working Group on QCD Event Generators for LEP [31]. The standard event format, and any usages peculiar to HERWIG are discussed more fully later, especially in Sect. 9, 10, 16 and 19.

- All output event data appear in standard format in the LEP standard common block /HEPEVT/.

- Internal status codes and pointers in the program have been replaced by those in the standard common block. Status codes `ISTHEP` = 100–199 replace `HERWIG` codes `ISTK` = 0–99 of earlier versions, and the tasks of the `HERWIG` pointers `IPTRK` are now performed by the mother/daughter pointers `JMONEP` and `JDHEP`.
- All subroutine and non-standard common block names have been changed to the form `HW***`, to avoid conflicts with other programs to which `HERWIG` may be linked.
- The cluster formation and decay, and unstable particle and heavy flavour decay phases of event generation have been isolated in subroutines `HWCFOR`, `HWDCEG`, `HWDHAD` and `HWDENV` respectively, to enable any of these to be replaced by other packages if desired.

For version 4.2 the main new features were:

- Full azimuthal correlations due to gluon spin are now included in leading logarithmic approximation for both initial and final state parton showers, using the Knowles algorithms [2,4]. Azimuthal spin correlations between different jets are also included. In the case of heavy quark processes, correlations are treated in the massless approximation.
- Bugs in gluon splitting and Higgs cross section have been fixed. The $gg \rightarrow H^0$ cross section is now correctly included to lowest order for any combination of Higgs and top quark masses.
- Hard scattering and minimum bias soft events can now be generated for any combination of pion, nucleon and/or antinucleon beams.

- The identifier and colour connection vectors `IDN` and `ICU` for hard subprocesses have been extended to dimension 10 for users who wish to use them in their own multijet subprocess routines.
- All variable types are now declared explicitly so that the program can be compiled with `IMPLICIT NONE` where this option is available.

For version 4.3 the main new features were:

- Default values of the parameters `QCDLAM` and `CLMAX` (see Sect. 6) were changed to give better agreement with LEP data [32].
- Tau lepton decays are now generated from (simplified) Particle Data Tables, like strong hadronic decays, instead of by secondary parton showering (which is still used for heavy quark decays). The decay tables used can be seen by setting print option `IPRINT` = 2.
- Clusters too light to decay into two hadrons are now treated more covariantly; the mass is shifted to an appropriate single-hadron mass by 4-momentum transfer to another cluster via rescaling the two-cluster centre-of-mass momentum, rather than via a lab energy transfer.
- ‘Remnant’ clusters (ie. those containing spectator partons from incoming hadrons) are now split if possible into two clusters like those produced in minimum-bias hadronic events. This change resulted from a study of remnant fragmentation in deep inelastic lepton scattering.

For version 4.6 the main new features were:

- `QCD α_s` was changed to the full 2-loop formula throughout, with matching at flavour thresholds as specified by the Working Group on QCD at LEP [33]. This means that the parameter `QCDLAM` must be interpreted differently from earlier versions, and can now be related to $\Lambda_{\overline{\text{MS}}}$ in the semi-inclusive region (z or $z \rightarrow 1$) only. See Ref. [6] for proof of this relation. (Since default parameter values were not changed from those in version 4.3, agreement with LEP data became less good.)
- The program was converted to double precision and array sizes were increased to facilitate running at LHC/SSC energies. The protection against event overflows (`NMHEP > NMHEP`) was also improved.
 - Several hard processes were added:
 - New Higgs boson production and decay processes, including full matrix elements for Higgs production via WW fusion (`IPROC = 1900` etc),
 - Top production via W exchange, eg. $u + b \rightarrow d + t$ (`IPROC = 2000` etc),
 - $W +$ jet production, eg. $q + g \rightarrow W + q$ (`IPROC = 2100` etc).
 - `SAVE` statements were included to permit running in overlay mode.
 - Bugs in heavy quark matrix elements were fixed.
 - Default W and Z masses and widths were updated.
- `HERWIG` version 5.0 was the first version to attempt seriously to simulate lepton-nucleon scattering at HERA, by including the heavy flavour generator `HARHEA` and other program developments by Abbiendi and Stancic [7], who have primary responsibility for this area of the program. The main new features compared with version 4.6 are:
 - Neutral- and charged-current deep inelastic scattering of charged leptons were fully included with optionally polarized incident leptons.
 - Neutral- and charged-current heavy flavour electroproduction were included.
 - Several technical defects in the simulation of deep inelastic processes in earlier versions were corrected.
 - Bugs in heavy quark plus spectator cluster fragmentation were corrected.
- `REAL*8` was changed to `DOUBLE PRECISION` for machine independence, and common blocks re-ordered to put `DOUBLE PRECISION` variables first.
 - Parameter `QCDLAM` now equals the 5-flavour QCD scale $\Lambda_{\overline{\text{MS}}}^{(5)}$ in the high- x region. (Printed values of $\Lambda_{\overline{\text{MS}}}$ for $N_f = 3, 4, 5$ in version 4.6 were actually *not* $\Lambda_{\overline{\text{MS}}}$ so they are now suppressed).
 - Default parameter values were reset to those found to give a satisfactory fit to event shape data of the OPAL Collaboration at LEP [34].

The main further new features in the current version, 5.1, are:

- New e^+e^- processes: WW pair production ($\text{IPROC} = 200$, programmed by Kunszt [35]) and standard model Higgs boson production via the Bjorken process and boson-boson fusion ($\text{IPROC} = 300$ and 400 etc).
- All hard hadronic processes ($1000 < \text{IPROC} < 8000$) are available also for photon and lepton beams (and e^+e^-) via the hadronic component of the photon (Dress-Grassie structure functions [36], plus Weizsäcker-Williams approximation for photon emission by leptons).
- The soft hadron remnant fragmentation model outlined in Subsect. 2.6 is used in lepton-hadron processes ($\text{IPROC} = 9000$ etc). Set $\text{IPROC} = 19000$ etc to suppress soft fragmentation (see Sect. 5).
- Optional linking to the CERN nucleon structure function library PDFLIB [37]. For this option, remove the dummy subroutines `PDFSET` and `STRUCTF` and set the new HERWIG parameter `MOPPDF` to the value of the PDFLIB parameter `MODE` for the structure function set required.

- W and Z boson decay modes are now controlled by new parameters `MDBOS(i)`: see Sect. 15.
- The running coupling `HWDALF` is now matched continuously at the top quark mass as well as at the b and c masses (but the input parameter `QCDAFM` still represents the δ -flavour $\Lambda_{\text{MS}}^{(5)}$).
- The random number generator `HWGEN` is now one of those recommended in Ref. [38] (l'Ecuyer's algorithm).
- The reading and writing of the Sudakov form factor table also reads and writes the values of relevant parameters and checks for conflicts.
- The program prints more detailed information on cross sections, and on Higgs boson branching ratios if relevant.

- The reading and writing of the Sudakov form factor table also reads and writes the values of relevant parameters and checks for conflicts.
- The program prints more detailed information on cross sections, and on Higgs boson branching ratios if relevant.

Note that the following features are *not* yet included in the program: polarization of heavy quarks and leptons; weak decays of heavy quarks before hadronization; treatment of coherence in the small- x region of incoming jets [21,23]; multiple parton interactions and parton shadowing; diffractive processes; QED radiative corrections; $W/Z/\gamma$ gauge bosons as components of parton showers.

4 Program Structure

The main program `HWIGPR` has the following form:

```
PROGRAM HWIGPR
C--COMMON BLOCKS ARE INCLUDED AS FILE HERWIG51.INC
INCLUDE 'HERWIG51.INC'
INTEGER N
```

```
C--MAX NUMBER OF EVENTS THIS RUN
MAKEV=100
C--BEAM PARTICLES
PART1='P '
PART2='P '
C--BEAM MOMENTA
PBEM1=20000.
PBEM2=20000.
C--PROCESS
IPROC=1500
C--INITIALISE OTHER COMMON BLOCKS
CALL HWIGIN
C--USER CAN RESET PARAMETERS AT THIS POINT,
C OTHERWISE DEFAULT VALUES IN HWIGIN WILL BE USED.
PTMIN=1000.
C--COMPUTE PARAMETER-DEPENDENT CONSTANTS
CALL HWING
C--CALL HWUSTA TO MAKE ANY PARTICLE STABLE
CALL HWUSTA('IO ')
C--USER'S INITIAL CALCULATIONS
CALL HWABEG
C--INITIALISE ELEMENTARY PROCESS
CALL HWENINI
C--LOOP OVER EVENTS
DO 100 N=1,MAKEV
C--INITIALISE EVENT
CALL HWLINE
C--GENERATE HARD SUBPROCESS
CALL HWEPRO
C--GENERATE PARTON CASCADES
CALL HWBGEN
C--DO CLUSTER FORMATION
CALL HWCFOR
C--DO CLUSTER DECAYS
CALL HWDEC
C--DO UNSTABLE PARTICLE DECAYS
CALL HWDRAD
C--DO HEAVY FLAVOUR DECAYS
CALL HWHFLY
C--ADD SOFT UNDERLYING EVENT IF NEEDED
CALL HWMEVT
C--USER'S EVENT ANALYSIS
CALL HWANAL
100 CONTINUE
C--TERMINATE ELEMENTARY PROCESS
CALL HWEND
STOP
END
```

Various phases of the simulation can be suppressed by deleting the corresponding subroutine calls, or different subroutines may be substituted. For example, in studies at the parton level everything from CALL HWCFDR to CALL HMMEVT can be omitted.

The following is a full list of subroutines and functions, which are classified according to their initial letters. The starred sections may be omitted for certain event types provided the corresponding subroutine calls are suppressed in HWIGPR or HWBGEN. Brief subroutine descriptions may be found in Sect. 20.

Main program and initialisation:

HWIGPR **HWGIN**

User-provided routines to initialise, terminate and perform user's analysis of event data:

HWABEG **HWEND** **HWANAL**

Parton branching generation with interfering gluons:

HWB2ZF	HWBCON	HWBFIN	HWBGEN	HWBJCD	HWBNAS	HWBMAN	HWBSPA
HWDSPN	HWBSUD	HWBSUG	HWBSU1	HWBSU2	HWBTIM	HWBTMC	

Cluster hadronization model:

HACUT	HWDEC	HWCLIA	HWCFOR	HWCGSP	HWCHAD		
HWDSDS	HWDBZ	HWDDAD	HWDHIG	HWDFVY	HWDPID	HWDPMT	HWDTMR
HWDTWO	HWDMRT						

Elementary subprocess generation:

HWEFIN	HWE GAM	HWEINI	HWEONE	HWEPRO	HWE TWO		
HWHEFW	HWHBK1	HWIBRN	HWIBSG	HWIDTS	HWIDYP	HWIEPA	HWIEPG
HWIGCF	HWHEHO	HWHEW1	HWHEW2	HWHEW3	HWHEW4	HWHEWY	HWHTGC
HWHQCD	HWHGCM	HWHTGS	HWHTIG	HWHTGW	HWHTGY	HWHTGZ	HWHPHO
	HWHQCP	HWHWEK	HWHWP	HWHWH1			

*Soft minimum-bias hadron-hadron collision or underlying event:

HMMEVT	HWMLPS	HWNBBI	HWMULT				
---------------	---------------	---------------	---------------	--	--	--	--

Random number generators:

HWRAZM	HWREXP	HWREXT	HWREGAU	HWREGEN	HWREINT	HWRLOG	HWROPW
HWRDNG	HWRUNI						

*Space-like branching of incoming partons:

HWSBRN	HWSDGG	HWSDGQ	HWSFBR	HWSFUN	HWSGAM	HWSGEN	HWSGQQ
HWSSPC	HWSUD	HWSTAB	HWSVAL				

Miscellaneous utilities:

HWALF	HWBPR	HWDAT	HWDEC	HWDEEQ	HWEPR	HWGAU	HWIDT
HWINC	HWJINE	HWLDOD	HWLLOB	HWLROL	HWLUSM	HWPCM	HWUSQR
HWURAP	HWURES	HWUROB	HWUROF	HWUROT	HWUSQR	HWUSTA	
HWUTAB	HWUTIN						

Warning messages and error handling:

HWWARN

In addition there are dummy versions of the CERN PDFLIB structure function routines

PDFSET and **STRUCTF**, which should be deleted if the PDFLIB package is to be used.

5 Beams and Processes

As indicated above, a number of variables must be set in the main program to specify what is to be simulated:

Name	Description	Default
PART1	Type of particle in beam 1	'p'
PART2	Type of particle in beam 2	'p'
PBEAM1	Momentum of beam 1	20000.0
PBEAM2	Momentum of beam 2	20000.0
IPROC	Type of process to generate	1500
MAXEV	Number of events to generate	100

Note that in HERWIG the beam momenta **PBEAM1** and **PBEAM2** are both assumed to be large. Therefore fixed-target experiments should be simulated in a moving frame, such as the overall centre-of-mass frame, and then boosted back into the laboratory.

The beam particle types **PART1**, **PART2** can take any of the values **NAME** listed below.

Beam	NAME
e^+	e^+
e^-	e^-
μ^+	μ^+
μ^-	μ^-
γ	GAMA
p	p
\bar{p}	$p\bar{p}$
n	n
\bar{n}	$n\bar{n}$
π^+	π^+
π^-	π^-

The currently available processes IPRUC are as follows:

6 Input Parameters

The quantities that may be regarded as adjustable parameters are

IPROC	Process
100	$e^+e^- \rightarrow q\bar{q}$ (all flavours)
100+IQ	$e^+e^- \rightarrow q\bar{q}$ ($iQ = 1, 2, 3, 4, 5, 6$ for $q = d, u, s, c, b, t$)
110	$e^+e^- \rightarrow q\bar{q}g$ (all flavours)
110+IQ	$e^+e^- \rightarrow q\bar{q}g$ (iQ as above)
110+IL	$e^+e^- \rightarrow t\bar{t}$ ($iL = 2, 3$ for $\ell = \mu, \tau$)
200	$e^+e^- \rightarrow W^+W^-$ (see Sect. 15 on control of W/Z decays)
300	$e^+e^- \rightarrow Z^0H^0 \rightarrow Z^0q\bar{q}$ (all flavours)
300+IQ	$e^+e^- \rightarrow Z^0H^0 \rightarrow Z^0q\bar{q}$ (iQ as above)
306+IL	$e^+e^- \rightarrow Z^0H^0 \rightarrow Z^0t\bar{t}$ ($iL = 1, 2, 3$ for $\ell = e, \mu, \tau$)
310, 311	$e^+e^- \rightarrow Z^0H^0 \rightarrow Z^0W^+W^-$, $Z^0Z^0Z^0$
312	$e^+e^- \rightarrow Z^0H^0 \rightarrow Z^0\gamma\gamma$
399	$e^+e^- \rightarrow Z^0H^0 \rightarrow Z^0$ anything
400+ID	$e^+e^- \rightarrow \nu\bar{\nu}H^0 + e^+e^-H^0$ (ID as in IPROC = 300 + ID)
1350	$q\bar{q} \rightarrow Z^0/\gamma \rightarrow t\bar{t}$ (all lepton species)
1350+IL	$q\bar{q} \rightarrow Z^0/\gamma \rightarrow t\bar{t}$ ($iL = 1, 2, 3$ for $\ell = e, \mu, \tau$)
1400	$q\bar{q} \rightarrow W^\pm \rightarrow q'\bar{q}''$ (all flavours)
1400+IQ	$q\bar{q} \rightarrow W^\pm \rightarrow q'\bar{q}''$ (q' or q'' as above)
1450	$q\bar{q} \rightarrow W^\pm \rightarrow t\bar{\nu}_t$ (all lepton species)
1450+IL	$q\bar{q} \rightarrow W^\pm \rightarrow t\bar{\nu}_t$ (iL as above)
1499	$q\bar{q} \rightarrow W^\pm \rightarrow$ anything
1500	QCD 2 → 2 hard parton scattering
	After generation, IPROD is subprocess (see Sect. 13)
1600+ID	$gg \rightarrow H^0$ (ID as in IPROC = 300 + ID)
1700+IQ	QCD heavy quark production (iQ as above)
	After generation, IPROD is subprocess (see Sect. 13)
1800	QCD direct photon + jet production
	After generation, IPROD is subprocess (see Sect. 14)
1900+ID	$W^+W^- \rightarrow H^0$ (ID as in IPROC = 300 + ID)
2000	t production via W exchange (sum of 2001–2008)
2001–4	$i\bar{b} \rightarrow i\bar{t}$, $i\bar{b} \rightarrow i\bar{t}$, $i\bar{b} \rightarrow i\bar{t}$, $ub \rightarrow dt$
2005–8	$i\bar{b} \rightarrow i\bar{t}$, $ib \rightarrow ct$, $ib \rightarrow ct$, $db \rightarrow st$
2100	$W^\pm +$ jet production
2110	$W^\pm +$ jet production (Compton only: $q\bar{q} \rightarrow Wg$)
2120	$W^\pm +$ jet production (Annihilation only: $q\bar{q} \rightarrow Wg$)
8000	Minimum bias soft hadron-hadron event
9000	Deep inelastic lepton scattering (neutral current)
9000+IQ	Deep inelastic lepton scattering (NC on flavour iQ)
9010	Deep inelastic lepton scattering (charged current)
9010+IQ	Deep inelastic lepton scattering (CC on flavour iQ)
N.B.	Charged incoming leptons only
9100+IP	Heavy quark production by boson-gluon fusion in charged lepton-nucleon NC processes. IP: $1 = c\bar{c}$, $2 = b\bar{b}$, $3 = t\bar{t}$
9110+IP	CC processes. IP: $1 = s\bar{c}$, $2 = b\bar{c}$, $3 = s\bar{t}$, $4 = b\bar{t}$ (+ ch.conj.)
10000+IP	as IPROC = IP but with soft underlying event (soft remnant fragmentation in lepton-hadron) suppressed

Notes on parameters:

- QCDLAM can be identified at high momentum fractions (x or z) with the fundamental 5-flavour QCD scale $\Lambda_{MS}^{(5)}$. However, this relation does not necessarily hold in other regions of phase space, since higher order corrections are not treated precisely enough to remove renormalization scheme ambiguities [6].
- RMASS(1, 2, 3, 13) are effective light quark and gluon masses used in the hadronization phase of the program. They can be set to zero provided the parton shower cut-offs VQCUR and VGCUR are large enough to prevent divergences (see below).
- For cluster hadronization, it must be possible to split gluons into $q\bar{q}$, ie. RMASS(13) must be at least twice the lightest quark mass. Similarly it may be impossible for heavy-flavoured clusters to decay if RMASS(4, 5) are too low.
- VQCUR and VGCUR are needed if the quark and gluon effective masses become small. The condition to avoid divergences in parton showers is

$$1/Q_i + 1/Q_j < 1/\text{QCPL3} \quad (6.1)$$

for either i or j or both gluons, where $Q_i = \text{RMASS}(i) + \text{VQCUR}$ for quarks, $\text{RMASS}(13) + \text{VGCUR}$ for gluons, and QCPL3 is the equivalent $\Lambda^{(3)}$ computed from QCPLAM. In the notation of Ref. [6] and Sect. 2, QCPL3 is the 3-flavour equivalent of QCPL5 where

$$\text{QCPL5} = \text{QCPLAM} \times \exp\left(\frac{151 - 9\pi^2}{138}\right)/\sqrt{2} = 1.109 \times \text{QCPLAM}.$$

We have approximately

$$QCDL3 \simeq QCDL5 \times L_b^{107/1725} L_c^{107/2025} \exp\left(\frac{L_b + L_c}{27}\right) \quad (6.3)$$

where $L_{b,c} = 2 \ln(m_{b,c}/QCDL5)$, giving a default value of $QCDL3 \simeq 0.40$.

- $CLMAX$ determines the maximum allowed mass of a cluster made from quarks i and j as follows

$$M^2 < CLMAX^2 + (RMASS(i) + RMASS(j))^2. \quad (6.4)$$

Since the cluster mass spectrum falls rapidly at high mass, results become insensitive to $CLMAX$ at large values.

- $PSPLT$ determines the mass distribution in the cluster splitting $C\ell_1 \rightarrow C\ell_2 + C\ell_3$ when $C\ell_1$ is above the maximum allowed mass. The masses of $C\ell_2$ and $C\ell_3$ are generated uniformly in M_{PSPLT} . As long as the number of split clusters is small, dependence on $PSPLT$ is weak.

$QDIQK$ greater than twice the lightest diquark mass enables non-perturbative gluon splitting into diquarks as well as quarks. The probability of this is $PDIQK \times dQ/Q$ for scales Q below $QDIQK$. The diquark masses are taken to be the sum of constituent quark masses. Thus the default value $QDIQK = 0$ suppresses gluon \rightarrow diquark splitting.

- $QSPAC$ is the scale below which the structure functions of incoming hadrons are frozen and non-valence constituent partons are forced to evolve to valence partons.

$PTRMS$ is the width of the (Gaussian) intrinsic transverse momentum distribution of valence partons in incoming hadrons at scale $QSPAC$.

- $ENSOF$ is the enhancement factor used in choosing the multiplicity of the underlying soft event. The multiplicity distribution is taken to be that of a $p\bar{p}$ collision at c.m. energy $ENSOF \propto \sqrt{s}$.

In practice, the parameters that have been found most effective in fitting data are $QCDLAM$, the gluon effective mass $RMASS(13)$, and the cluster mass parameter $CLMAX$. Note that $QSPAC$, $PTRMS$ and $ENSOF$ do not affect lepton-lepton collisions.

The default parameter values have been found to give good agreement with event shape distributions at LEP [32,34].

A number of further parameters are needed to control the program and to turn various options on or off:

Name	Description	Default
IPRINT	Printout option	1
MAXPR	Number of events to print out	1
MAXER	Maximum number of errors	10
LNEWT	Unit for writing output events	0
LRSUD	Unit for reading Sudakov table	0
LNSUD	Unit for writing Sudakov table	7
NRN(1)	Random number seed 1	17673
NRN(2)	Random number seed 2	63565
WTMAX	Max weight (0 to search for it)	0.0
NWGT	Generate unweighted events	.TRUE.
AZSOFT	Soft gluon azimuthal correlations	.TRUE.
AZSPIN	Gluon spin azimuthal correlations	.TRUE.
NCOLJO	Number of colours	3
NFLAV	Number of (producible) flavours	6
PDFLIB	PDFLIB structure function set (if negative do not use PDFLIB)	-1
NSTRU	Input structure function set (1,2=Duke-Owens,1,2; 3,4=EHLQ1,2)	1
EPOLN	Electron beam polarization in DIS (± 1 for fully rh/lh polarized)	0.0
BGSHT	Scale= \hat{s} for boson-gluon fusion	.TRUE.
PTMIN	Min p_T in hadronic jet production	10.0
PTMAX	Max p_T in hadronic jet production	10^8
PTPW	$1/p_T^{PPW}$ for jet sampling	4.0
YJMIN	Min jet rapidity	-8.0
YJMAX	Max jet rapidity	8.0
EMMIN	Min dilepton mass in Drell-Yan	10.0
EMMAX	Max dilepton mass in Drell-Yan	10^8
EMPW	$1/m^{PPW}$ for Drell-Yan sampling	4.0
Q2MIN	Min Q^2 in deep inelastic	9999.5
Q2MAX	Max Q^2 in deep inelastic	10000.5
Q2PW	$1/Q^{PPW}$ for sampling	2.5
THMAX	Max thrust in 3 parton production	0.9

Printout options are:

- ```
IPRINT = 0 Print program title only
 1 Print selected input parameters
 2 1 + table of particle codes and properties
 3 2 + tables of Sudakov form factors
```

See Sect. 8 on form factors for details of  $LNSUD$  and  $LUSUD$ . If  $BGSHT$  is .FALSE., the scale used for heavy quark production via boson-gluon fusion in lepton-hadron collisions will be

$$\frac{2 \hat{s} \hat{t} \hat{u}}{\hat{s}^2 + \hat{t}^2 + \hat{u}^2}. \quad (6.5)$$



## 8 Form Factor File

HERWIG uses look-up tables of Sudakov form factors for the evolution of initial- and final-state parton showers. These can be read from an input file rather than being recomputed each time. The reading, writing and computing of form factor tables is controlled by integer parameters LRSUD and LWSUD:

```
LRSUD = N > 0 Read form factors for this run from unit N
LRSUD = 0 Compute new form factor tables for this run
LRSUD < 0 Form factor tables are already loaded
LWSUD = N > 0 Write form factors on unit N for future use
LWSUD = 0 Do not write new form factor tables
```

The option LRSUD < 0 allows the program to be initialized several times in the same run (eg. to generate various event types) without recomputing or rereading form factors.

Note that the Sudakov form factors depend on the parameters QCDLAM, VGCUT, NCFLD, NFLAV, RMASS(i3) and RMASS(i) for  $i = 1, \dots, NFLAV$ . Consequently form factor tables must be recomputed every time any of these parameters is changed. In version 5.1, these parameters are written/read with the form factor tables and checks are performed to ensure consistency.

## 9 Event Data

/HEPEVNT/ is the LEP standard common block containing current event data:

|              |                                                                                    |
|--------------|------------------------------------------------------------------------------------|
| NEVHEP       | event number                                                                       |
| NHEP         | number of entries for this event                                                   |
| ISTHEP(I)    | status of entry I (see below)                                                      |
| IDHEP(I)     | identity of entry I (Particle Data Group code)                                     |
| JMOHEP(1, I) | pointer to first mother of entry I (see below)                                     |
| JMOHEP(2, I) | pointer to second mother of entry I (see below)                                    |
| JDAHEP(1, I) | pointer to first daughter of entry I (see below)                                   |
| JDAHEP(2, I) | pointer to last daughter of entry I (see below)                                    |
| PHEP(*, I)   | $(p_x, p_y, p_z, E, M)$ of entry I: $M = \text{sign}(\sqrt{\text{abs}(m^2)}, m^2)$ |
| WHEP(*, I)   | $(x, y, z, t)$ of production vertex of entry I (not yet used)                      |

All momenta are given in GeV/c in the laboratory frame, in which the input beam momenta are PBEM1 and PBEM2 as specified by the user and point along the +z and -z directions respectively. Final state particles have ISTHEP(I) = 1. See Sect. 10 for a complete list of the special status codes used by HERWIG.

The identity codes IDHEP are as recommended by the LEP Working Group [31], ie. as defined by the Particle Data Group [39] plus IDHEP = 91 for clusters, 94 for jets, and 0 for others with no PDG code. HERWIG also has its own internal identity codes IDHW(I), stored in /HEBEVNT/. The utility subroutine HWUDT translates between HERWIG and PDG identity codes. See Sect. 16 for further details.

The mother/daughter pointers are standard, except that JMOHEP(2, I) and JDAHEP(2, I) for a parton are its colour mother and colour daughter, ie., the partons to which its colour and anticolour are connected, respectively. For this purpose the primary partons from a

hard subprocess are all regarded as outgoing (see examples in Sect. 13, 16 and 19). Since a quark has no anticolour, JDAHEP(2, I) is used to point to its flavour partner. Similarly for JMOHEP(2, I) in the case of an antiquark.

In addition to entries representing partons, particles, clusters etc., /HEPEVNT/ contains purely informational entries representing the total centre-of-mass momentum, hard and soft subprocess momenta, etc. See Sect. 10 for the corresponding status codes.

Information from all stages of event processing is retained in /HEPEVNT/ so the same particle may appear several times with different status codes. For example, an outgoing parton from a hard scattering (entered initially with status 113 or 114) will appear after processing as an on-mass-shell parton before QCD branching (status 123,124), an off-mass-shell entry representing the flavour and momentum of the outgoing jet (status 143,144), and a jet constituent (157). It might also appear again in other contexts, eg. as a spectator in a heavy flavour decay (status 154,160).

Incoming partons (entered with status 111,112, changed to 121,122 after branching) give rise to spacelike jets (status 141,142), with  $m^2 < 0$ , indicated by PHEP(5, IHEP) < 0, due to the loss of momentum via initial-state bremsstrahlung. The same applies in principle to incoming leptons, but QED radiative corrections are not yet included.

Each parton jet begins with a status 141–144 jet entry giving the total flavour and momentum of the jet. The first mother pointer of this entry gives the location of the parent hard parton, while the second gives that of the subprocess centre-of-mass momentum. If QCD branching has occurred, this is followed by a lightlike CDNE entry, which fixes the angular extent of the jet and its azimuthal orientation relative to the parton with which it interferes. The interfering parton is listed as the second mother of the cone. Next come the actual constituents of the jet. If no branching has occurred, there is no cone and the single jet constituent is the same as the jet. However, the jet and constituent momenta may not coincide if some transfer of momentum has occurred during cluster hadronization. See the sample output in Sect. 18 and 19 for illustration of these points.

## 10 Status Codes

A complete list of currently-used HERWIG status codes is given below. Many are used only in intermediate stages of event processing. The most important for users are probably 1 (final-state particle), 101–3 (initial state), 141–4 (jets), and 199 (decayed b- and t-flavoured hadrons).

The event status ISTAT in common /HEBEVNT/ is roughly ISTHEP – 100 where ISTHEP is the status of entries being processed. For completed events, ISTAT = 100.

| ISTHEP | Description                             |
|--------|-----------------------------------------|
| 1      | final state particle                    |
| 2      | gluon split to $q\bar{q}$               |
| 3      | documentation line                      |
| 100    | cone limiting jet evolution             |
| 101    | 'beam' (beam 1)                         |
| 102    | 'target' (beam 2)                       |
| 103    | overall centre of mass                  |
| 110    | unprocessed hard process c.m.           |
| 111    | unprocessed beam parton                 |
| 112    | unprocessed target parton               |
| 113    | unproc. first outgoing parton           |
| 114    | unproc. other outgoing parton           |
| 115    | unprocessed spectator parton            |
| 120-25 | as 110-15, after processing             |
| 130    | lepton in jet (unboosted)               |
| 131-34 | as 141-44, unboosted to c.m.            |
| 135    | space-like parton (beam, unboosted)     |
| 136    | space-like parton (target, unboosted)   |
| 137    | spectator (beam, unboosted)             |
| 138    | spectator (target, unboosted)           |
| 139    | parton from branching (unboosted)       |
| 140    | parton from gluon splitting (unboosted) |
| 141-44 | jet from parton type 111-14             |
| 145-50 | as 135-40 boosted, unclustered          |
| 151    | as 159, not yet clustered               |
| 152    | as 160, not yet clustered               |
| 153    | heavy quark before decay                |
| 154    | spectator before heavy decay            |
| 155    | spectator from beam                     |
| 156    | spectator from target                   |
| 157    | parton from QCD branching               |
| 158    | parton from gluon splitting             |
| 159    | parton from cluster splitting           |
| 160    | spectator after heavy decay             |
| 161    | beam cluster before soft process        |
| 162    | target cluster before soft process      |
| 163    | other cluster before soft process       |
| 167    | unhadronized beam cluster               |
| 168    | unhadronized target cluster             |
| 170    | soft process centre of mass             |
| 171    | soft cluster (beam, unhadronized)       |
| 172    | soft cluster (target, unhadronized)     |
| 173    | soft cluster (other, unhadronized)      |

## 11 Event Weights

The default is to generate unweighted events (EWWT = AWWT). Then event distributions are generated by computing a weight proportional to the cross section and comparing it with a random number times the maximum weight. Set WGTMAX to the maximum weight, or to zero for the program to compute it. If a weight greater than WGTMAX is generated during execution, a warning is printed and WGTMAX is reset. Similarly if the efficiency is too low (WGTMAX too large). If these errors occur too often, output event distributions could be distorted.

To generate weighted events, set NOWGT = .FALSE. in common /HERWIGNT/.

In QCD hard scattering and heavy flavour, direct photon, and W + jet production (IPROC = 1500, 1700, 1800, 2100 etc) the transverse energy distribution of weighted events (or the efficiency for unweighted events) can be varied using the parameters PTMIN, PTMAX and PPTOW.

Similarly in Drell-Yan processes (IPROC = 1350 etc) the lepton pair mass distribution is controlled by the parameters EMIN, EMAX and EMPOW, and in deep inelastic scattering (IPROC = 9000 etc) the  $Q^2$  distribution depends on Q2MIN, Q2MAX and Q2POW.

Data on weights generated are output at the end of the run. The mean weight is an estimate of the cross section (in nanobarns) integrated over the region used for event generation. Note that the mean weight is the sum of weights divided by the total number of *weights* generated, not the total number of *events*.

In version 5.1, the estimated cross section (in picobarns) and its error are also printed.

## 12 Heavy Flavour Decays

Heavy quark decays ( $b, t$  and higher generations) are treated as secondary hard subprocesses. Heavy flavoured hadrons are split into collinear heavy quark and spectator and the former decays independently. After decay, parton showers may be generated from coloured decay products, in the usual way. See Ref. [5] for details of the treatment of colour coherence in these showers.

As we discussed in Sect. 2, for high values of the top quark mass, the  $t$  quark is expected to decay before it has time to form a hadron. At present this possibility is *not* taken into account in HERWIG.

Users may wish to change the decay fractions for heavy quarks, for example to force a leptonic decay in every event. This can be done by modifying the contents of `COMMON/RWDPHV/`: `FBTM(1,1)`, `FBTM(2,1)`, ..., `FBTM(6,1)` are the  $b$ -quark decay fractions into the 6 doublets ( $d,u$ )  $(s,c)$   $(b,t)$   $(e,\nu_e)$   $(\mu,\nu_\mu)$  respectively. `FBTM(1,2)` etc are the corresponding  $\bar{b}$  fractions. Thus to get all  $b$ -hadrons to decay to muons, while keeping the default decay fractions for  $\bar{b}$ -hadrons, one would set `FBTM(J,1) = 0` for  $J \neq 5$  and `FBTM(5,1) = 1`, without changing `FBTM(J,2)`. `FRTP` is the corresponding array for the top quark, while `FRHY` is for quarks heavier than top. All these quantities can be changed from event to event.

### 13 QCD Hard Subprocesses

At present only  $2 \rightarrow 2$  subprocesses are implemented. They are classified as shown below.

| TINPRO | $1+2 \rightarrow 3+4$                           | c/f conn. |
|--------|-------------------------------------------------|-----------|
| 1      | $q+q \rightarrow q+q$                           | 3 4 2 1   |
| 2      | $q+q \rightarrow q+q$                           | 4 3 1 2   |
| 3      | $q+q' \rightarrow q+q'$                         | 3 4 2 1   |
| 4      | $q+\bar{q} \rightarrow q'+\bar{q}'$             | 2 4 1 3   |
| 5      | $q+\bar{q} \rightarrow q+\bar{q}$               | 3 1 4 2   |
| 6      | $q+\bar{q} \rightarrow q+\bar{q}$               | 2 4 1 3   |
| 7      | $q+\bar{q} \rightarrow g+g$                     | 2 4 1 3   |
| 8      | $q+\bar{q} \rightarrow g+g$                     | 2 3 4 1   |
| 9      | $g+\bar{q}' \rightarrow g+\bar{q}'$             | 3 1 4 2   |
| 10     | $g+g \rightarrow q+g$                           | 3 1 4 2   |
| 11     | $g+g \rightarrow q+g$                           | 3 4 2 1   |
| 12     | $\bar{q}+q \rightarrow \bar{q}'+q'$             | 3 1 4 2   |
| 13     | $\bar{q}+q \rightarrow \bar{q}+q$               | 2 4 1 3   |
| 14     | $\bar{q}+q \rightarrow \bar{q}+q$               | 3 1 4 2   |
| 15     | $\bar{q}+q \rightarrow g+g$                     | 3 1 4 2   |
| 16     | $\bar{q}+q \rightarrow g+g$                     | 4 1 2 3   |
| 17     | $\bar{q}+q' \rightarrow \bar{q}+q'$             | 2 4 1 3   |
| 18     | $\bar{q}+\bar{q} \rightarrow \bar{q}+\bar{q}$   | 4 3 1 2   |
| 19     | $\bar{q}+\bar{q} \rightarrow \bar{q}+\bar{q}$   | 3 4 2 1   |
| 20     | $\bar{q}+\bar{q}' \rightarrow \bar{q}+\bar{q}'$ | 4 3 1 2   |
| 21     | $\bar{q}+g \rightarrow \bar{q}+g$               | 2 4 1 3   |
| 22     | $\bar{q}+g \rightarrow \bar{q}+g$               | 4 3 1 2   |
| 23     | $g+q \rightarrow g+q$                           | 2 4 1 3   |
| 24     | $g+q \rightarrow g+q$                           | 3 4 2 1   |
| 25     | $g+\bar{q} \rightarrow g+\bar{q}$               | 3 1 4 2   |
| 26     | $g+\bar{q} \rightarrow g+\bar{q}$               | 4 3 1 2   |
| 27     | $g+g \rightarrow g+g$                           | 2 4 1 3   |
| 28     | $g+g \rightarrow g+g$                           | 4 1 2 3   |
| 29     | $g+g \rightarrow g+g$                           | 4 1 2 3   |
| 30     | $g+g \rightarrow g+g$                           | 4 3 1 2   |
| 31     | $g+g \rightarrow g+g$                           | 2 4 1 3   |

When different colour flows are possible, they are listed as separate subprocesses. This separation is not exact but is normally a good approximation [1,24]. The sum of the colour flows is the exact lowest-order cross section.

### 14 QCD Direct Photon Subprocesses

| TINPRO | $1+2 \rightarrow 3+4$                  | c/f conn. |
|--------|----------------------------------------|-----------|
| 41     | $q+\bar{q} \rightarrow g+\gamma$       | 2 3 1     |
| 42     | $q+g \rightarrow q+\gamma$             | 3 1 2     |
| 43     | $\bar{q}+q \rightarrow g+\gamma$       | 3 1 2     |
| 44     | $\bar{q}+g \rightarrow \bar{q}+\gamma$ | 2 3 1     |
| 45     | $g+q \rightarrow g+\gamma$             | 2 3 1     |
| 46     | $g+\bar{q} \rightarrow \bar{q}+\gamma$ | 3 1 2     |
| 47     | $g+g \rightarrow g+\gamma$             | 2 3 1     |

Note that at present the gluon fusion process 47, which proceeds via quark loops and is very small, is neglected.


### 15 Electroweak Subprocesses

`HERWIG` generates Higgs bosons through gluon fusion and  $W$  fusion in hadron-hadron collisions (IPROC=1600+ID and 1900+ID), and in lepton-lepton collisions through the Bjorken process (ie.  $Z^{(-)} \rightarrow Z^{(+)}H$  with one or both  $Z$ s off-shell) and  $W$  fusion (IPROC=300+ID and 400+ID). Each process is generated according to the exact leading order matrix element in the  $s$ -channel approximation. This results in unitarity violation for  $m_H \gg m_W$ ,  $s \gtrsim a$  few  $m_H^2$ , (where  $s = q_i^2$ ), so to regularize this, the  $m_H \Gamma_H$  in the propagator can be replaced by  $\sqrt{s} \Gamma_H(s)$ . The variable `IOPHIG` controls this procedure:

| IOPHIG | Choose $s$ according to            | Reweight? |
|--------|------------------------------------|-----------|
| 0      | $s^2/((s-m_H^2)^2 + m_H \Gamma_H)$ | Yes       |
| 1      | $1/((s-m_H^2)^2 + m_H \Gamma_H)$   | Yes       |
| 2      | $s^2/((s-m_H^2)^2 + m_H \Gamma_H)$ | No        |
| 3      | $1/((s-m_H^2)^2 + m_H \Gamma_H)$   | No        |

'c/f conn.' refers to the colour/flavour connections between the partons: ' $i j k l$ ' means that the colour of parton 1 comes from parton  $i$ , that of 2 from  $j$ , etc. For antiquarks, which have no colour (only anticolour), the label shows instead to which parton the flavour

is connected. For this colour/flavour labelling all partons are defined as outgoing. Thus, for example, process 10 has colour connections 3 1 4 2, corresponding to the colour flow diagram:



where reweighting means weighting the distribution back to

$$\frac{\sqrt{s} \Gamma_{\pi}(s)}{(s - m_H^2)^2 + \sqrt{s} \Gamma_{\pi}(s)} \quad (15.1)$$

The default is **IOPHIG** = 1. The difference between options 0 and 1 is purely in the weight distribution produced. Options 2 and 3 are intended primarily for users who wish to supply their own unitarity conserving reweighting function at the point indicated in routine **HWHIGM**. In all cases, the distribution is restricted to the range  $[m_H - \text{GAMMAX} \times \Gamma_H, m_H + \text{GAMMAX} \times \Gamma_H]$ . **GAMMAX** defaults to 10, but in the non-perturbative region  $m_H \gtrsim 1$  TeV should be reduced to protect against poor weight distributions. These considerations do not affect the distribution noticeably for  $m_H \lesssim 500$  GeV, and **GAMMAX** can safely be increased if necessary.

For each process, ID controls the Higgs decay: ID=1–6 for quarks, 7–9 for leptons, 10/11 for  $WW/ZZ$  pairs, and 12 for photons. In addition ID=0 gives quarks of all flavours, and ID=99 gives all decays. For each process, the average event weight is the cross section in nb times the branching fraction to the requested decay. The branching ratios to quarks use the next-to-leading logarithm corrections, those to  $WW/ZZ$  pairs allow for one or both bosons off-shell.

Gauge bosons formed by lepton-antilepton or quark-antiquark annihilation (**IPROC** = 100–199, 1300–1499) have their decay determined by the process code **IPROC**. Gauge bosons are also generated through the processes of  $W+1$  parton production in hadron-hadron collisions, and  $WW$  pair production in lepton-lepton collisions, as well as in the Higgs processes mentioned above. In these cases their decay is controlled by the variable **MODBOS(i)**. This controls the decay of the *i*th gauge boson per event:

| <b>MODBOS(i)</b> | <b>W Decay</b> | <b>Z Decay</b>       |
|------------------|----------------|----------------------|
| 0                | all            | all                  |
| 1                | $q\bar{q}$     | $q\bar{q}$           |
| 2                | $e\nu$         | $e^+e^-$             |
| 3                | $\mu\nu$       | $\mu^+\mu^-$         |
| 4                | $\tau\nu$      | $\tau^+\tau^-$       |
| 5                | $ev\&\mu\nu$   | $e^+e^-\&\mu^+\mu^-$ |
| 6                | all            | $\nu\nu$             |
| > 6              | all            | all                  |

All entries of **MODBOS** default to 0. Bosons which are produced in pairs (ie. from  $WW$  pair production, or Higgs decay) are symmetrized in **MODBOS(i)** and **MODBOS(i+1)**. For processes which directly produce gauge bosons, the event weight includes the branching fraction to the requested decay, but this is only true for Higgs production if decay to  $WW/ZZ$  is forced (ID=10/11) and not if ID=99. The spin correlations in the decays are handled in one of two ways:

- (a) the diagonal members of the spin density matrix are stored in **RHOHEP(i, IHEP)**, where  $i = 1, 2, 3$  for helicity =  $i-2$  in the centre-of-mass frame of their production, for processes where this matrix is diagonal (ie. there is no interference between spin states).
- (b) the correlations in the decay are handled directly by the production routine where (a) is not possible.

## 16 Including New Subprocesses

It should not be difficult for users to include further subprocesses in this version of the program if required. The parton and hard sub-process 4-momenta, masses and identity codes need to be entered in **COMMON/HEPEVT/** with the appropriate status codes **ISTHEP(I) = 110–114** to tell the program which is which (see the table in Sect. 10). The colour/flavour structure should be specified by the second mother and daughter pointers as explained in Sect. 13 (see also the sample output and guide, Sect. 18 and 19).

The **HERWIG** identity codes **IDHW(I)** in **COMMON/HEPEVT/** also need to be set correctly. The **IDHW** codes can be listed in a run with **IPRINT = 2**: the most important are the quarks 1–6 (as **IDHEP**), antiquarks 7–12, gluon 13, overall centre-of-mass 14, hard centre-of-mass 15, soft centre-of-mass 16, photon 59, leptons 121–126, antileptons 127–132.


The utility subroutine **BWULDT(IOPF, IPFG, IHWG, NAME)** is provided to translate between Particle Data Group code **IPFG**, **HERWIG** code **IHWG**, and **HERWIG CHARACTER+4 NAME**, with **IOPF = 1, 2, 3** depending on which of **IPFG**, **IHWG** and **NAME** is the input argument.

Consider for example the process of virtual photon-gluon fusion to make  $b + \bar{b}$  in proton-

electron collisions (in fact this process is now included as **IPROC = 9102**). We assume the user

provides a subroutine to generate the momenta **PHEP** for the hard subprocess  $e + g \rightarrow e\bar{b}\bar{b}$ .

The colour structure is



Thus the momenta generated, together with those of the initial beams and the overall centre of mass, could be entered in the following sequence:

| <b>IHEP</b> | <b>Entry</b> | <b>ISTHEP</b> | <b>IDHEP</b> | <b>JMOHEP</b> | <b>JDAHEP</b> | <b>IDHW</b> |
|-------------|--------------|---------------|--------------|---------------|---------------|-------------|
| 1           | e beam       | 101           | 11           | 0 0           | 0 0           | 121         |
| 2           | p beam       | 102           | 221 2        | 0 0           | 0 0           | 73          |
| 3           | ep c.m.      | 103           | 0            | 0 0           | 0 0           | 14          |
| 4           | e in         | 111           | 11           | 6 7           | 0 7           | 121         |
| 5           | gluon        | 112           | 21           | 6 9           | 0 8           | 13          |
| 6           | hard cm      | 110           | 0            | 4 5           | 7 9           | 15          |
| 7           | e out        | 113           | 11           | 6 4           | 0 4           | 121         |
| 8           | b            | 114           | 5            | 6 5           | 0 9           | 5           |
| 9           | b            | 114           | -5           | 6 8           | 0 5           | 11          |

Note that if there are more than two outgoing partons, the first has status 113 and all the others 114. Each parton has **JMOHEP(1, I) = 6** to indicate the location of the hard c.m. for this subprocess, while **JMOHEP(2, I)** gives the location of the colour mother (treating the incoming gluon as outgoing) or the connected electron. **JDAHEP(1, I)** will be set by the jet

generator HBBGEN, while JDABEP(2,I) points to the anticolour mother (or connected electron). Finally the HERWIG identifiers IDHW(I) could be set to the indicated values by means of the translation subroutine HWUDT as follows:

```
CHARACTER*4 NAME
.....
NHEP=9
IDBEP(1)=11
IDBEP(2)=2212
.....
IDBEP(9)=-5
DO 10 I=1,NHEP
10 CALL HWUDT(1,IDBEP(I),IDHW(I),NAME)
IDHW(6)=15
```

The last statement is needed because IDPDG(I) = 0 returns IDHW(I) = 14. If subroutine HBBGEN is now called, it will find the coloured partons and generate QCD jets from them. Subsequent calls to HWCRTR etc can then be used to form clusters and hadronize them. If the hard subprocess routine is called from HWEPRO, like those already provided, it should have two options controlled by the logical variable GENEV in COMMON/HWHARD/. For GENEV = .FALSE., an event weight (normally the cross section in nanobarns) is generated and stored as EWWT in COMMON/HWEVENT/. If this weight is accepted by HWEPRO, the subroutine is called a second time with GENEV = .TRUE. and the corresponding event data should then be generated and stored as explained above. On certain computers it will be necessary to SAVE those variables that determine event characteristics between the two subroutine calls.

## 17 Error Conditions

Certain combinations of input parameters may lead to problems in execution. HERWIG tries to detect these and print a warning. Errors during execution are dealt with by HWMARN which prints the calling subroutine and a code and takes appropriate action. In general, the larger the code the more serious the problem. Refer to the source program to find out why HWMARN was called. Events can be rerun by setting the random number seeds MRN(1) and MRN(2) to the values given in the error message or event dump. The contents of /HEPEVT/ can be printed by calling HWUPR, those of /HWPART/ (the last parton shower) by calling HWUPR.

Examples of error messages are:

```
HWMARN CALLED FROM SUBPROGRAM HWSBRN: CODE = 101
EVENT 31: SEEDS = 422399901 & 771980111 WEIGHT = 0.3893E-08
EVENT KILLED. EXECUTION CONTINUES
```

```
HWMARN CALLED FROM SUBPROGRAM HWCMD: CODE = 102
EVENT 51: SEEDS = 1033784787 & 1428957533 WEIGHT = 0.3893E-08
EVENT KILLED. EXECUTION CONTINUES
```

Spacelike (initial-state) parton branching had no phase space. This can happen due to cutoffs which are slightly different in the hard subprocess and the parton shower.  
Action taken: program throws away this event and starts a new one.

A cluster has been formed with too low a mass to represent any hadron of the correct flavour, and there is no colour-connected cluster from which the necessary additional mass could be transferred.  
Action taken: program throws away this event and starts a new one.

```
HWMARN CALLED FROM SUBPROGRAM HMUINE: CODE= 200
EVENT SURVIVES. RUN ENDS GRACEFULLY
```

```
HWMARN CALLED FROM SUBPROGRAM HMEV: CODE= 200
CPU TIME LIMIT LIABLE TO BE REACHED BEFORE GENERATING MAXEV EVENTS.
ACTION TAKEN: SKIPS TO TERMINAL CALCULATIONS USING EXISTING EVENTS.
```

```
HWMARN CALLED FROM SUBPROGRAM HBBSUD: CODE= 500
RUN CANNOT CONTINUE
```

The table of Sudakov form factors read on unit LRSUD does not extend to the maximum momentum scale QLIM specified for this run.  
Action taken: run aborted. The user must either reduce QLIM or set LRSUD=0 to make a bigger table (set LWSUD nonzero to write it).

```
HWMARN CALLED FROM SUBPROGRAM HMBSUD: CODE= 515
RUN CANNOT CONTINUE
```

The table of Sudakov form factors read on unit LRSUD is for a different value of a relevant parameter (in this case the b quark mass).  
Action taken: run aborted. The user must make a new table (set LWSUD nonzero to write it).

## 18 Sample Output

Below we give a complete listing of output from the program, set up for  $e^+ e^- \rightarrow b\bar{b}g$  at a c.m. energy of 91.2 GeV. The main features of the output are discussed in Sect. 19.

|                               |            |
|-------------------------------|------------|
| HERWIG 5.1                    | APRIL 1991 |
| INPUT CONDITIONS FOR THIS RUN |            |
| BEAM 1 (E+ ) MOMENTUM =       | 45.60      |
| BEAM 2 (E- ) MOMENTUM =       | 45.60      |
| PROCESS CODE (IPROC) =        | 115        |
| NUMBER OF FLAVOURS =          | 6          |
| STRUCTURE FUNCTION SET =      | 1          |
| AZIM SPIN CORRELATIONS =      | T          |
| AZIM SOFT CORRELATIONS =      | T          |
| QCD LAMBDA (GET) =            | 0.2000     |
| DOWN QUARK MASS =             | 0.3200     |
| UP QUARK MASS =               | 0.3200     |
| STRANGE QUARK MASS =          | 0.5000     |
| CHARMED QUARK MASS =          | 1.8000     |
| BOTTOM QUARK MASS =           | 5.2000     |
| TOP QUARK MASS =              | 100.0000   |
| GLUON EFFECTIVE MASS =        | 0.7500     |
| EXTRA SHOWER CUTOFF (Q) =     | 0.4800     |
| EXTRA SHOWER CUTOFF (G) =     | 0.0600     |

CLUSTER MASS PARAMETER = 3.5000  
 SPACELIKE EVOLN CUTOFF = 2.5000  
 INTRINSIC P-TRAN (RMS) = 0.0000  
 MAX THRUST FOR 2->3 = 0.9000

#### PDFLIB STRUCT FUNCTIONS NOT USED

NO EVENTS WILL BE WRITTEN TO DISK  
 WRITING SUDAKOV TABLE ON UNIT 7  
 PARTICLE TYPE 21=PIO SET STABLE  
 INITIAL SEARCH FOR MAX WEIGHT

|                      |            |
|----------------------|------------|
| PROCESS CODE IPROC = | 115        |
| RANDOM NO. SEED 1 =  | 1246579    |
| SEED 2 =             | 8147766    |
| NUMBER OF SHOTS =    | 2000       |
| NEW MAXIMUM WEIGHT = | 0.4702E+01 |
| NEW MAXIMUM WEIGHT = | 0.6034E+01 |
| NEW MAXIMUM WEIGHT = | 0.9074E+01 |

INITIAL SEARCH FINISHED

#### OUTPUT ON ELEMENTARY PROCESS

|                      |            |
|----------------------|------------|
| NUMBER OF EVENTS =   | 0          |
| NUMBER OF WEIGHTS =  | 2000       |
| MEAN VALUE OF WGT =  | 0.7198E+00 |
| RMS SPREAD IN WGT =  | 0.7733E+00 |
| ACTUAL MAX WEIGHT =  | 0.8239E+01 |
| ASSUMED MAX WEIGHT = | 0.9074E+01 |

|                      |          |
|----------------------|----------|
| PROCESS CODE IPROC = | 115      |
| CROSS SECTION (PB) = | 719.8381 |
| ERROR IN C-S (PB) =  | 17.2916  |
| EFFICIENCY PERCENT = | 7.9329   |

EVENT 1: 45.60 GEV/C E+ 0N 45.60 GEV/C E- PROCESS: 115  
 SEEDS: 17673 & 63565 STATUS: 100 ERROR: 0 WEIGHT: 0.7198E+00

#### --INITIAL STATE---

|         |                           |      |      |        |        |      |
|---------|---------------------------|------|------|--------|--------|------|
| IHEP ID | IDPDG IST M01 M02 DA1 DA2 | P-X  | P-Y  | P-Z    | ENERGY | MASS |
| 1 E+    | -11 101 0 0 0 0           | 0.00 | 0.00 | 45.60  | 45.60  | 0.00 |
| 2 E-    | 11 102 0 0 0 0            | 0.00 | 0.00 | -45.60 | 45.60  | 0.00 |
| 3 CMF   | 0 103 0 0 0 0             | 0.00 | 0.00 | 91.20  | 91.20  | 0.00 |

#### --HARD SUBPROCESS---

|                          |                           |        |        |       |        |       |
|--------------------------|---------------------------|--------|--------|-------|--------|-------|
| IHEP ID                  | IDPDG IST M01 M02 DA1 DA2 | P-X    | P-Y    | P-Z   | ENERGY | MASS  |
| 4 Z0/G                   | 23 120 1 2 5 7            | 0.00   | 0.00   | 0.00  | 0.00   | 91.20 |
| ---H/W/Z BOSON DECAYS--- |                           |        |        |       |        |       |
| IHEP ID                  | IDPDG IST M01 M02 DA1 DA2 | P-X    | P-Y    | P-Z   | ENERGY | MASS  |
| 5 BOTM                   | 5 123 4 6 8 7             | 35.40  | 12.90  | -8.32 | 38.04  | 5.20  |
| 6 GLUE                   | 21 124 4 7 12 5           | 3.67   | -11.51 | 2.89  | 12.45  | 0.76  |
| 7 BBAR                   | -5 124 4 5 18 6           | -39.07 | -1.39  | 5.43  | 39.81  | 5.20  |

#### --PARTON SHOWERS---

|         |                           |        |        |       |        |       |
|---------|---------------------------|--------|--------|-------|--------|-------|
| IHEP ID | IDPDG IST M01 M02 DA1 DA2 | P-X    | P-Y    | P-Z   | ENERGY | MASS  |
| 8 BOTM  | 94 143 5 4 10 11          | 33.41  | 12.18  | -7.85 | 37.11  | 7.17  |
| 9 CONE  | 0 100 5 6 0 0             | 0.85   | -0.73  | 0.10  | 1.13   | 0.00  |
| 10 BOTM | 5 157 8 22 34 21          | 31.72  | 12.16  | -6.97 | 35.07  | 5.20  |
| 11 GLUE | 21 2 8 14 22 23           | 1.69   | 0.02   | -0.88 | 2.05   | 0.75  |
| 12 GLUE | 94 144 6 4 14 17          | 3.46   | -10.87 | 2.73  | 16.86  | 10.67 |
| 13 CONE | 0 100 6 7 0 0             | -1.11  | 0.26   | 0.07  | 1.14   | 0.00  |
| 14 GLUE | 21 2 12 15 24 25          | 0.79   | 0.81   | 0.03  | 1.36   | 0.75  |
| 15 GLUE | 21 2 12 16 26 27          | -0.08  | -1.17  | 2.80  | 3.12   | 0.75  |
| 16 GLUE | 21 2 12 17 28 29          | 0.74   | -1.59  | 0.96  | 2.13   | 0.75  |
| 17 GLUE | 21 2 12 20 30 31          | 2.01   | -8.92  | -1.06 | 9.24   | 0.75  |
| 18 BBAR | 94 144 7 4 20 21          | -36.87 | -1.31  | 5.12  | 38.23  | 8.63  |
| 19 CONE | 0 100 7 6 0 0             | -0.40  | -0.98  | 0.34  | 1.11   | 0.00  |
| 20 GLUE | 21 2 18 21 32 33          | -1.42  | -0.26  | 1.49  | 2.21   | 0.75  |
| 21 BBAR | -5 157 18 10 40 33        | -35.45 | -1.05  | 3.63  | 36.02  | 5.20  |

#### --GLUON SPLITTING---

|         |                           |       |       |       |        |      |
|---------|---------------------------|-------|-------|-------|--------|------|
| IHEP ID | IDPDG IST M01 M02 DA1 DA2 | P-X   | P-Y   | P-Z   | ENERGY | MASS |
| 22 UBAR | -2 158 8 23 34 10         | 1.01  | -0.02 | -0.31 | 1.11   | 0.32 |
| 23 UP   | 2 158 8 24 35 22          | 0.67  | 0.04  | -0.57 | 0.94   | 0.32 |
| 24 UBAR | -2 158 12 25 35 23        | 0.20  | 0.45  | -0.07 | 0.59   | 0.32 |
| 25 UP   | 2 158 12 26 36 24         | 0.59  | 0.36  | 0.10  | 0.77   | 0.32 |
| 26 UBAR | -2 158 12 27 36 25        | -0.10 | -0.86 | 1.11  | 1.33   | 0.32 |
| 27 UP   | 2 158 12 28 37 26         | 0.02  | -0.51 | 1.69  | 1.79   | 0.32 |
| 28 UBAR | -2 158 12 29 37 27        | 0.44  | -1.02 | 0.76  | 1.39   | 0.32 |
| 29 UP   | 2 158 12 30 38 28         | 0.30  | -0.57 | 0.18  | 0.74   | 0.32 |
| 30 UBAR | -2 158 12 31 38 29        | 1.14  | -5.22 | -0.80 | 5.42   | 0.32 |
| 31 UP   | 2 158 12 32 39 30         | 0.87  | -3.70 | -0.26 | 3.82   | 0.32 |
| 32 DBAR | -1 158 16 33 39 31        | -0.59 | -0.16 | 0.89  | 1.13   | 0.32 |
| 33 DOWN | 1 158 16 21 40 32         | -0.84 | -0.10 | 0.60  | 1.08   | 0.32 |

#### --CLUSTER FORMATION---

|         |                           |       |       |       |        |      |
|---------|---------------------------|-------|-------|-------|--------|------|
| IHEP ID | IDPDG IST M01 M02 DA1 DA2 | P-X   | P-Y   | P-Z   | ENERGY | MASS |
| 34 CLUS | 91 183 10 22 41 42        | 32.73 | 12.14 | -7.28 | 36.17  | 6.06 |

| ---CLUSTER DECAYS---       |       |       |     |     |     |     |     |        |       |       |        |      |  |  |
|----------------------------|-------|-------|-----|-----|-----|-----|-----|--------|-------|-------|--------|------|--|--|
| IHEP                       | ID    | IDPDG | IST | M01 | M02 | DA1 | DA2 | P-X    | P-Y   | P-Z   | ENERGY | MASS |  |  |
| 41                         | BOBR  | -511  | 199 | 34  | 8   | 70  | 71  | 27.90  | 10.32 | -5.49 | 30.71  | 5.28 |  |  |
| 42                         | PI-   | -211  | 1   | 34  | 8   | 0   | 0   | 4.84   | 1.81  | -1.79 | 5.47   | 0.14 |  |  |
| 43                         | PI+   | 211   | 1   | 35  | 8   | 0   | 0   | 0.02   | -0.09 | -0.12 | 0.21   | 0.14 |  |  |
| 44                         | RHO-  | -213  | 197 | 35  | 8   | 55  | 56  | 0.86   | 0.40  | -0.51 | 1.32   | 0.77 |  |  |
| 45                         | K+    | 321   | 1   | 36  | 12  | 0   | 0   | 0.53   | 0.11  | 0.62  | 0.96   | 0.49 |  |  |
| 46                         | K*-   | -323  | 197 | 36  | 12  | 57  | 58  | -0.04  | -0.40 | 0.58  | 1.14   | 0.89 |  |  |
| 47                         | RHO+  | 213   | 197 | 37  | 12  | 59  | 60  | 0.50   | -1.39 | 2.47  | 2.98   | 0.77 |  |  |
| 48                         | PI-   | -211  | 1   | 37  | 12  | 0   | 0   | -0.04  | -0.15 | -0.01 | 0.21   | 0.14 |  |  |
| 49                         | RHO+  | 213   | 197 | 38  | 12  | 61  | 62  | 0.50   | -2.55 | 0.06  | 2.71   | 0.77 |  |  |
| 50                         | PI-   | -211  | 1   | 38  | 12  | 0   | 0   | 0.93   | -3.24 | -0.67 | 3.44   | 0.14 |  |  |
| 51                         | DL++  | 2224  | 197 | 39  | 12  | 63  | 64  | 0.23   | -1.25 | -0.52 | 1.84   | 1.23 |  |  |
| 52                         | DLB-  | -2224 | 197 | 39  | 12  | 65  | 66  | 0.05   | -2.61 | 1.15  | 3.11   | 1.23 |  |  |
| 53                         | RHO-  | -213  | 197 | 40  | 18  | 67  | 68  | -4.61  | -0.57 | -0.05 | 4.71   | 0.77 |  |  |
| 54                         | B+    | 521   | 199 | 40  | 18  | 72  | 73  | -31.67 | -0.59 | 4.27  | 32.40  | 5.27 |  |  |
| ---STRONG HADRON DECAYS--- |       |       |     |     |     |     |     |        |       |       |        |      |  |  |
| IHEP                       | ID    | IDPDG | IST | M01 | M02 | DA1 | DA2 | P-X    | P-Y   | P-Z   | ENERGY | MASS |  |  |
| 55                         | PI0   | 111   | 1   | 44  | 8   | 0   | 0   | 0.15   | 0.29  | -0.47 | 0.59   | 0.13 |  |  |
| 56                         | PI-   | -211  | 1   | 44  | 8   | 0   | 0   | 0.71   | 0.10  | -0.04 | 0.73   | 0.14 |  |  |
| 57                         | PI-   | -211  | 1   | 46  | 12  | 0   | 0   | -0.23  | -0.09 | 0.40  | 0.50   | 0.14 |  |  |
| 58                         | KBAR  | -311  | 198 | 46  | 12  | 69  | 69  | 0.20   | -0.31 | 0.18  | 0.64   | 0.50 |  |  |
| 59                         | PI0   | 111   | 1   | 47  | 12  | 0   | 0   | 0.48   | -1.32 | 1.80  | 2.29   | 0.13 |  |  |
| 60                         | PI+   | 211   | 1   | 47  | 12  | 0   | 0   | 0.02   | -0.06 | 0.67  | 0.69   | 0.14 |  |  |
| 61                         | PI0   | 111   | 1   | 49  | 12  | 0   | 0   | 0.08   | -1.80 | -0.16 | 1.81   | 0.13 |  |  |
| 62                         | PI+   | 211   | 1   | 49  | 12  | 0   | 0   | 0.42   | -0.75 | 0.21  | 0.90   | 0.14 |  |  |
| 63                         | PI+   | 211   | 1   | 51  | 12  | 0   | 0   | 0.22   | -0.40 | -0.29 | 0.56   | 0.14 |  |  |
| 64                         | P     | 2212  | 1   | 51  | 12  | 0   | 0   | 0.02   | -0.85 | -0.22 | 1.29   | 0.94 |  |  |
| 65                         | PI0   | 111   | 1   | 52  | 12  | 0   | 0   | 0.18   | -0.46 | 0.38  | 0.64   | 0.13 |  |  |
| 66                         | PBAR  | -2212 | 1   | 52  | 12  | 0   | 0   | -0.13  | -2.14 | 0.78  | 2.47   | 0.94 |  |  |
| 67                         | PI0   | 111   | 1   | 53  | 18  | 0   | 0   | -0.52  | -0.07 | 0.20  | 0.58   | 0.13 |  |  |
| 68                         | PI-   | -211  | 1   | 53  | 18  | 0   | 0   | -4.09  | -0.50 | -0.25 | 4.13   | 0.14 |  |  |
| 69                         | KOL   | 130   | 1   | 58  | 12  | 0   | 0   | -0.31  | 0.20  | -0.18 | 0.64   | 0.50 |  |  |
| ---HEAVY FLAVOUR DECAYS--- |       |       |     |     |     |     |     |        |       |       |        |      |  |  |
| IHEP                       | ID    | IDPDG | IST | M01 | M02 | DA1 | DA2 | P-X    | P-Y   | P-Z   | ENERGY | MASS |  |  |
| 70                         | BOTM  | 5     | 153 | 41  | 71  | 74  | 76  | 26.21  | 9.70  | -5.16 | 28.84  | 4.95 |  |  |
| 71                         | DBAR  | -1    | 125 | 41  | 76  | 80  | 76  | 1.69   | 0.63  | -0.33 | 1.86   | 0.32 |  |  |
| 72                         | UP    | 2     | 125 | 54  | 79  | 81  | 79  | -1.92  | -0.04 | 0.26  | 1.97   | 0.32 |  |  |
| ---PARTON SHOWERS---       |       |       |     |     |     |     |     |        |       |       |        |      |  |  |
| IHEP                       | ID    | IDPDG | IST | M01 | M02 | DA1 | DA2 | P-X    | P-Y   | P-Z   | ENERGY | MASS |  |  |
| 73                         | BBAR  | -6    | 163 | 54  | 72  | 77  | 79  | -29.75 | -0.55 | 4.01  | 30.43  | 4.95 |  |  |
| 74                         | NUEB  | -12   | 123 | 70  | 75  | 82  | 75  | 7.41   | 2.16  | -2.02 | 7.98   | 0.00 |  |  |
| 75                         | E-    | 11    | 124 | 70  | 74  | 83  | 74  | 2.86   | -0.01 | 0.04  | 2.86   | 0.00 |  |  |
| 76                         | CHARM | 4     | 124 | 70  | 74  | 80  | 70  | 15.94  | 7.54  | -3.18 | 18.01  | 1.80 |  |  |
| 77                         | UP    | 2     | 123 | 73  | 78  | 86  | 78  | -18.31 | -0.94 | 2.49  | 18.51  | 0.32 |  |  |
| 78                         | DBAR  | -1    | 124 | 73  | 77  | 88  | 77  | -5.82  | 0.17  | 0.02  | 5.83   | 0.32 |  |  |
| 79                         | CBAR  | -4    | 124 | 73  | 73  | 90  | 73  | -5.62  | 0.22  | 1.50  | 6.09   | 1.80 |  |  |
| 80                         | DBAR  | -1    | 160 | 71  | 85  | 93  | 85  | 1.69   | 0.63  | -0.33 | 1.86   | 0.32 |  |  |
| 81                         | UP    | 2     | 160 | 72  | 91  | 92  | 91  | -1.92  | -0.04 | 0.26  | 1.97   | 0.32 |  |  |
| 82                         | NUEB  | -12   | 1   | 74  | 70  | 0   | 0   | 7.41   | 2.16  | -2.02 | 7.98   | 0.00 |  |  |
| 83                         | E-    | 11    | 1   | 75  | 70  | 0   | 0   | 2.86   | -0.01 | 0.04  | 2.86   | 0.00 |  |  |
| ---CLUSTER FORMATION---    |       |       |     |     |     |     |     |        |       |       |        |      |  |  |
| IHEP                       | ID    | IDPDG | IST | M01 | M02 | DA1 | DA2 | P-X    | P-Y   | P-Z   | ENERGY | MASS |  |  |
| 92                         | CLUS  | 91    | 183 | 81  | 91  | 95  | 96  | -7.54  | 0.18  | 1.76  | 8.06   | 2.22 |  |  |
| 93                         | CLUS  | 91    | 183 | 85  | 90  | 97  | 98  | 17.63  | 8.17  | -3.51 | 19.87  | 2.22 |  |  |
| 94                         | CLUS  | 91    | 183 | 87  | 89  | 99  | 100 | -24.13 | -0.77 | 2.51  | 24.34  | 1.76 |  |  |
| ---CLUSTER DECAYS---       |       |       |     |     |     |     |     |        |       |       |        |      |  |  |
| IHEP                       | ID    | IDPDG | IST | M01 | M02 | DA1 | DA2 | P-X    | P-Y   | P-Z   | ENERGY | MASS |  |  |
| 92                         | CLUS  | 91    | 183 | 81  | 91  | 95  | 96  | -7.54  | 0.18  | 1.76  | 8.06   | 2.22 |  |  |
| 93                         | CLUS  | 91    | 183 | 85  | 90  | 97  | 98  | 17.63  | 8.17  | -3.51 | 19.87  | 2.22 |  |  |
| 94                         | CLUS  | 91    | 183 | 87  | 89  | 99  | 100 | -24.13 | -0.77 | 2.51  | 24.34  | 1.76 |  |  |
| ---CLUSTER DECAYS---       |       |       |     |     |     |     |     |        |       |       |        |      |  |  |
| IHEP                       | ID    | IDPDG | IST | M01 | M02 | DA1 | DA2 | P-X    | P-Y   | P-Z   | ENERGY | MASS |  |  |
| 95                         | PI+   | 211   | 1   | 92  | 72  | 0   | 0   | -0.62  | -0.11 | 0.22  | 0.68   | 0.14 |  |  |
| 96                         | DC+-  | -413  | 197 | 92  | 72  | 101 | 102 | -6.92  | 0.30  | 1.54  | 7.38   | 2.01 |  |  |
| 97                         | DC*+  | 413   | 197 | 93  | 84  | 103 | 104 | 15.59  | 7.08  | -3.15 | 17.53  | 2.01 |  |  |
| 98                         | PIO   | 111   | 1   | 93  | 84  | 0   | 0   | 2.04   | 1.09  | -0.36 | 2.34   | 0.13 |  |  |
| 99                         | PI+   | 211   | 1   | 94  | 86  | 0   | 0   | -10.09 | -0.29 | 0.92  | 10.14  | 0.14 |  |  |
| 100                        | A20   | 115   | 197 | 94  | 86  | 105 | 106 | -14.04 | -0.48 | 1.58  | 14.20  | 1.32 |  |  |
| ---STRONG HADRON DECAYS--- |       |       |     |     |     |     |     |        |       |       |        |      |  |  |
| IHEP                       | ID    | IDPDG | IST | M01 | M02 | DA1 | DA2 | P-X    | P-Y   | P-Z   | ENERGY | MASS |  |  |
| 101                        | DCBR  | -424  | 198 | 96  | 72  | 107 | 108 | -6.37  | 0.29  | 1.45  | 6.80   | 1.86 |  |  |
| 102                        | PI-   | -211  | 1   | 96  | 72  | 0   | 0   | -0.56  | 0.01  | 0.09  | 0.58   | 0.14 |  |  |
| 103                        | DCO   | 421   | 198 | 97  | 84  | 109 | 111 | 14.66  | 6.68  | -2.94 | 16.48  | 1.86 |  |  |
| 104                        | PI+   | 211   | 1   | 97  | 84  | 0   | 0   | 0.93   | 0.41  | -0.21 | 1.04   | 0.14 |  |  |
| 105                        | PI+   | 211   | 1   | 100 | 86  | 0   | 0   | -5.59  | -0.12 | 0.22  | 5.60   | 0.14 |  |  |
| 106                        | RHO-  | -213  | 198 | 100 | 86  | 112 | 113 | -8.45  | -0.36 | 1.36  | 8.60   | 0.77 |  |  |

```

PARTICLE TYPE 21=PIO SET STABLE
107 KO 311 198 101 72 114 114 -2.67 0.52 1.02 2.95 0.50
108 ETPR 331 198 101 72 115 116 -3.69 -0.23 0.43 3.85 0.96
109 KBAR -311 198 103 84 117 117 4.99 2.37 -0.95 5.62 0.50
110 RH00 113 198 103 84 118 119 5.11 2.33 -1.00 5.76 0.77
111 ETA 221 198 103 84 120 122 4.57 1.98 -0.99 5.11 0.55
112 PIO 111 1 106 86 0 0 -3.45 0.20 0.61 3.51 0.13
113 PI- -211 1 106 86 0 0 -5.00 -0.56 0.76 5.09 0.14
114 K0L 130 1 107 72 0 0 -2.67 0.52 1.02 2.95 0.50
115 GAMMA 22 1 108 72 0 0 -0.82 0.04 -0.05 0.82 0.00
116 RN00 113 198 108 72 123 124 -2.88 -0.27 0.48 3.03 0.77
117 K0L 130 1 109 84 0 0 4.99 2.37 -0.95 5.62 0.50
118 PI- -211 1 110 84 0 0 0.57 0.40 0.05 0.71 0.14
119 PI+ 211 1 110 84 0 0 4.54 1.94 -1.05 5.04 0.14
120 PI0 111 1 111 84 0 0 1.22 0.53 -0.25 1.36 0.13
121 PI- -211 1 111 84 0 0 1.57 0.53 -0.40 1.71 0.14
122 PI+ 211 1 111 84 0 0 1.79 0.92 -0.33 2.04 0.14
123 PI- -211 1 116 72 0 0 -2.39 -0.25 0.65 2.49 0.14
124 PI+ 211 1 116 72 0 0 -0.49 -0.02 -0.17 0.54 0.14

```

#### OUTPUT ON ELEMENTARY PROCESS

```

NUMBER OF EVENTS = 100
NUMBER OF WEIGHTS = 1339
MEAN VALUE OF WGT = 0.7323E+00
RMS SPREAD IN WGT = 0.8725E+00
ACTUAL MAX WEIGHT = 0.7355E+01
ASSUMED MAX WEIGHT = 0.9074E+01

PROCESS CODE IPROC = 115
CROSS SECTION (PB) = 732.2993
ERROR IN C-S (PB) = 23.8432
EFFICIENCY PERCENT = 8.0702

```

## 19 Guide to Sample Output

After listing the more important input parameter values, the program prints the message

PDFLIB STRUCT FUNCTIONS NOT USED

to tell us that the built-in structure functions have been selected, rather than the PDFLIB library (although of course they are not relevant for this process). The message

NO EVENTS WILL BE WRITTEN TO DISK

reminds the user that LNEVT = 0 for this run. Then a table of Sudakov form factors for parton branching is computed, since none has been provided on an input file (LRSUD = 0).

The table is written on an output file (LWSUD = 7) for use in future runs:

WRITING SUDAKOV TABLE ON UNIT 7

The default particle data table is modified by calling ENUSTA('PIO') to suppress  $\pi^0$  decays:

```

Next the program searches for the maximum weight, ie. the maximum cross section in the available phase space, as implied by the default value WGTMAX = 0. The parameter with means that $b\bar{b}$ events with a thrust of less than 0.9 will be generated. After this search, the result tells us that the initial estimate of the cross section for this sub-process in this region is 720 ± 17 picobarns, and the fraction of weights accepted for event generation is likely to be about 8%.
```

Since the default is MAXPR = 1, the first generated event is printed. The heading

```

EVENT 1: 45.60 GEV/c E+ ON 45.60 GEV/c E- PROCESS: 115
SEEDS: 17673 & 63565 STATUS: 100 ERROR: 0 WEIGHT: 0.7198E+00

```

tells us the beam and target, the random number seeds at the start of the event (so that it can be regenerated by setting NRN(1&2) to these values), and the process code IPROC. The status 100 means a complete event was generated and the zero error code means no problems were encountered. Since NOWGT = .TRUE. (unweighted event generation), each event has the mean weight computed earlier.

Next come the contents of COMMON/HEPEVT/ and related quantities. The various parts of this particular event are located as follows:

| Entry  | Description                                      |
|--------|--------------------------------------------------|
| 1-3    | Initial state                                    |
| 4      | Hard subprocess: $Z^0/\gamma$ production         |
| 5-7    | $Z^0/\gamma$ decay to $b\bar{b}g$                |
| 8-21   | Parton showers                                   |
| 22-33  | Gluon splitting                                  |
| 34-40  | Cluster formation                                |
| 41-69  | Cluster and hadron decays                        |
| 70-83  | Weak decays of $\overline{B^0}$ and $B^+$ mesons |
| 84-124 | Hadronization of $B^+/B^0$ decay products        |

We discuss each part in turn.

---INITIAL STATE---

| IRNP ID | IDPBG IST | M01 M02 DAI DA2 | P-X   | P-Y  | P-Z  | ENERGY | MASS   |
|---------|-----------|-----------------|-------|------|------|--------|--------|
| 1 E+    | -11       | 101 0 0 0       | 0 0 0 | 0.00 | 0.00 | 45.60  | 45.60  |
| 2 E-    | 11        | 102 0 0 0       | 0 0 0 | 0.00 | 0.00 | -45.60 | -45.60 |
| 3 CMF   | 0 103     | 0 0 0 0         | 0 0 0 | 0.00 | 0.00 | 91.20  | 91.20  |

**CMP** represents the overall centre-of-mass of the initial state. The ‘mother’ pointer  $M0i = JM0HEP(i, IHEP)$  and ‘daughter’ pointer  $DAi = JDHEP(i, IHEP)$  are set to zero for these entries.


---HARD SUBPROCESS---

| IHEP ID | IDPDG  | IST | M01 | M02 | DA1 | DA2  | P-X  | P-Y  | P-Z   | ENERGY | MASS |
|---------|--------|-----|-----|-----|-----|------|------|------|-------|--------|------|
| 4 ZO/G  | 23 120 | 1   | 2   | 5   | 7   | 0.00 | 0.00 | 0.00 | 91.20 | 91.20  |      |

---H/W/Z BOSON DECAYS---

| IHEP ID | IDPDG  | IST | M01 | M02 | DA1 | DA2    | P-X    | P-Y   | P-Z   | ENERGY | MASS |
|---------|--------|-----|-----|-----|-----|--------|--------|-------|-------|--------|------|
| 5 BOTM  | 6 123  | 4   | 6   | 8   | 7   | 35.40  | 12.90  | -8.32 | 38.94 | 5.20   |      |
| 6 GLUE  | 21 124 | 4   | 7   | 12  | 5   | 3.67   | -11.51 | 2.89  | 12.45 | 0.75   |      |
| 7 BBAR  | -5 124 | 4   | 5   | 18  | 6   | -39.07 | -1.39  | 5.43  | 39.81 | 5.20   |      |

Z0/G is the hard subprocess centre-of-mass (which is equal to the overall centre-of-mass because initial state QED radiation is not yet included). Its mother and daughter pointers give the locations of the incoming and outgoing partons. The status code 123 corresponds to give the locations of the incoming and outgoing partons. The status code 124 to all other outgoing partons in the hard subprocess. The first mother pointers show the location of the hard centre-of-mass, and the second mother of each parton is the ‘colour mother’, as explained above. Thus the colours of partons 5,6,7 are connected to 6,7,5 respectively. Since antiquarks carry no colour M02 for the  $b$  is used for the flavour connection to the  $b$ . Similarly, the first daughter points to the associated jet but the second daughter is the colour daughter, ie. the parton to which this one’s anticolour (or its flavour, in the case of the  $b$ ) is connected. Thus the anticolour connections of 5,6,7 in this case are to 7,5,6. The colour flow diagram is



---PARTON SHOWERS---

The QCD cascade from each hard parton is generated in sequence. First there is a jet entry ( $IHEP = 94$ ) giving the total jet momentum, flavour and mass. In the case of initial-state jets the mass would represent  $-|q^2|^{\frac{1}{2}}$ . M01 gives the parent hard parton and M02 the hard centre-of-mass. DA1 and DA2 point to the first and last parton in the jet after perturbative branching. If branching occurs, the next entry (CONE), is a lightlike 4-vector defining the radiation cone and the orientation of the radiation pattern.

The partons in the jet ( $ISTHEP = 157$ , plus 155,156 for spectators, with gluons reset to 2 by gluon splitting subroutine HWCGSP) have their colour and anticolour connections specified by M02 and DA2 respectively, as described above for the hard subprocess. The  $b$  jet is:

| IHEP ID | IDPDG  | IST | M01 | M02 | DA1 | DA2   | P-X   | P-Y   | P-Z   | ENERGY | MASS |
|---------|--------|-----|-----|-----|-----|-------|-------|-------|-------|--------|------|
| 8 BOTM  | 94 143 | 5   | 4   | 10  | 11  | 33.41 | 12.18 | -7.86 | 37.11 | 7.17   |      |
| 9 CONE  | 0 100  | 5   | 6   | 0   | 0   | 0.85  | -0.73 | 0.10  | 1.13  | 0.00   |      |
| 10 BOTM | 5 157  | 8   | 22  | 34  | 21  | 31.72 | 12.16 | -6.97 | 35.07 | 5.20   |      |
| 11 GLUE | 21 2   | 8   | 14  | 22  | 23  | 1.69  | 0.02  | -0.88 | 2.95  | 0.75   |      |

The gluon jet is:

| IHEP ID | IDPDG  | IST | M01 | M02 | DA1 | DA2   | P-X    | P-Y   | P-Z   | ENERGY | MASS |
|---------|--------|-----|-----|-----|-----|-------|--------|-------|-------|--------|------|
| 12 GLUE | 94 144 | 6   | 4   | 14  | 17  | 3.46  | -10.87 | 2.73  | 15.86 | 10.67  |      |
| 13 CONE | 0 100  | 6   | 7   | 0   | 0   | -1.11 | 0.26   | 0.07  | 1.14  | 0.00   |      |
| 14 GLUE | 21 2   | 12  | 15  | 24  | 25  | 0.79  | 0.81   | 0.03  | 1.36  | 0.75   |      |
| 15 GLUE | 21 2   | 12  | 16  | 26  | 27  | -0.08 | -1.17  | 2.80  | 3.12  | 0.75   |      |
| 16 GLUE | 21 2   | 12  | 17  | 28  | 29  | 0.74  | -1.59  | 0.96  | 2.13  | 0.75   |      |
| 17 GLUE | 21 2   | 12  | 20  | 30  | 31  | 2.01  | -8.92  | -1.06 | 9.24  | 0.75   |      |

The  $\bar{b}$  jet is:

| IHEP ID | IDPDG  | IST | M01 | M02 | DA1 | DA2    | P-X   | P-Y  | P-Z   | ENERGY | MASS |
|---------|--------|-----|-----|-----|-----|--------|-------|------|-------|--------|------|
| 18 BBAR | 94 144 | 7   | 4   | 20  | 21  | -36.87 | -1.31 | 5.12 | 38.23 | 8.63   |      |
| 19 CONE | 0 100  | 7   | 6   | 0   | 0   | -0.40  | -0.98 | 0.34 | 1.11  | 0.00   |      |
| 20 GLUE | 21 2   | 18  | 21  | 32  | 33  | -1.42  | -0.26 | 1.49 | 2.21  | 0.75   |      |
| 21 BBAR | -5 157 | 18  | 10  | 40  | 33  | -35.45 | -1.05 | 3.63 | 36.02 | 5.20   |      |

---GLUON SPLITTING---

As the first step in the cluster hadronization model, any gluons in the jets are split into light quark-antiquark pairs. The flavours of the pairs are chosen at random amongst those allowed by kinematics. The colour connections are remade accordingly.

| IHEP ID | IDPDG  | IST | M01 | M02 | DA1 | DA2   | P-X   | P-Y   | P-Z  | ENERGY | MASS |
|---------|--------|-----|-----|-----|-----|-------|-------|-------|------|--------|------|
| 22 UBAR | -2 158 | 8   | 23  | 34  | 10  | 1.01  | -0.02 | -0.31 | 1.11 | 0.32   |      |
| 23 UP   | 2 158  | 8   | 24  | 35  | 22  | 0.67  | 0.04  | -0.57 | 0.94 | 0.32   |      |
| 24 UBAR | -2 158 | 12  | 25  | 35  | 23  | 0.20  | 0.45  | -0.07 | 0.59 | 0.32   |      |
| 25 UP   | 2 158  | 12  | 26  | 36  | 24  | 0.59  | 0.36  | 0.10  | 0.77 | 0.32   |      |
| 32 DBAR | -1 158 | 18  | 33  | 39  | 31  | -0.59 | -0.16 | 0.89  | 1.13 | 0.32   |      |
| 33 DOWN | 1 158  | 18  | 21  | 40  | 32  | -0.84 | -0.10 | 0.60  | 1.08 | 0.32   |      |

Gluon radiation from the  $b$  and the  $\bar{b}$  will be limited by interference with the gluon, and that from the gluon by the  $b$  or  $\bar{b}$ . The momenta and masses of the partons are the raw on-shell values generated before QCD radiative corrections.

If the incoming partons were coloured, then they would have status 121 and 122, and in specifying their colour connections, they would be regarded as outgoing.

Each quark (or antiquark) is combined with its colour mother antiquark (or diquark) to make a cluster with the sum of their 4-momenta. Non-bean clusters with masses above the maximum specified by Eq. (2.2) would be split by creating new quark-antiquark pairs with ISHEP = 159 (there are no such pairs in this event). In a hadron collision, the beam remnant clusters, ie., those containing the spectator quarks from the incident hadrons (ISHEP = 167, 168) would be split into 2 ‘soft’ clusters each (see new features of version 4.3, above), making another two pairs. The clusters themselves are then listed.

#### ---CLUSTER FORMATION---

| IHEP ID | IDPDG  | IST   | M01   | M02    | DA1   | DA2   | P-X   | P-Y  | P-Z | ENERGY | MASS |
|---------|--------|-------|-------|--------|-------|-------|-------|------|-----|--------|------|
| 34 CLUS | 91 183 | 10 22 | 41 42 | 32 73  | 12 14 | -7.28 | 36.17 | 6.06 |     |        |      |
| 35 CLUS | 91 183 | 23 24 | 43 44 | 0.87   | 0.49  | -0.63 | 1.53  | 0.97 |     |        |      |
| 36 CLUS | 91 183 | 25 26 | 45 46 | 0.49   | -0.29 | 1.21  | 2.10  | 1.62 |     |        |      |
| 37 CLUS | 91 183 | 27 28 | 47 48 | 0.46   | -1.53 | 2.47  | 3.18  | 1.23 |     |        |      |
| 38 CLUS | 91 183 | 29 30 | 49 50 | 1.44   | -5.79 | -0.62 | 6.15  | 1.37 |     |        |      |
| 39 CLUS | 91 183 | 31 32 | 51 52 | 0.29   | -3.86 | 0.64  | 4.95  | 3.03 |     |        |      |
| 40 CLUS | 91 183 | 33 21 | 53 54 | -36.28 | -1.15 | 4.23  | 37.10 | 6.41 |     |        |      |

#### ---CLUSTER DECAYS---

The clusters, including the  $b$ -flavoured clusters 34 and 40, now decay isotropically into pairs of hadrons chosen according to density of states. Particles with ISHEP = 1 are stable. ISHEP = 199 indicates a heavy ( $b$  or  $t$ ) hadron which will later decay weakly via a secondary hard subprocess.

| IHEP ID  | IDPDG     | IST   | M01   | M02    | DA1   | DA2   | P-X   | P-Y  | P-Z | ENERGY | MASS |
|----------|-----------|-------|-------|--------|-------|-------|-------|------|-----|--------|------|
| 41 BOBR  | -511 199  | 34 8  | 70 71 | 27.90  | 10.32 | -5.49 | 30.71 | 5.28 |     |        |      |
| 42 PI-   | -211 1    | 34 8  | 0 0   | 4.84   | 1.81  | -1.79 | 5.47  | 0.14 |     |        |      |
| 43 PI+   | 211 1     | 35 8  | 0 0   | 0.02   | 0.09  | -0.12 | 0.21  | 0.14 |     |        |      |
| 44 RH0-  | -213 197  | 35 8  | 55 56 | 0.86   | 0.40  | -0.51 | 1.32  | 0.77 |     |        |      |
| 45 K+    | 321 1     | 36 12 | 0 0   | 0.53   | 0.11  | 0.62  | 0.96  | 0.49 |     |        |      |
| 46 K*-   | -323 197  | 36 12 | 57 58 | -0.04  | -0.40 | 0.58  | 1.14  | 0.89 |     |        |      |
| ... DL++ | 2224 197  | 39 12 | 63 64 | 0.23   | -1.25 | -0.52 | 1.84  | 1.23 |     |        |      |
| 52 DLB-  | -2214 197 | 39 12 | 65 66 | 0.05   | -2.61 | 1.15  | 3.11  | 1.23 |     |        |      |
| 53 RH0-  | -213 197  | 40 18 | 67 68 | -4.61  | -0.57 | -0.05 | 4.71  | 0.77 |     |        |      |
| 54 B+    | 521 199   | 40 18 | 72 73 | -31.67 | -0.59 | 4.27  | 32.40 | 5.27 |     |        |      |
| 55 PI0   | 111 1     | 44 8  | 0 0   | 0.15   | 0.29  | -0.47 | 0.59  | 0.13 |     |        |      |
| 56 PI-   | -211 1    | 44 8  | 0 0   | 0.71   | 0.10  | -0.04 | 0.73  | 0.14 |     |        |      |
| 57 PI-   | -211 1    | 46 12 | 0 0   | -0.23  | -0.09 | 0.40  | 0.50  | 0.14 |     |        |      |
| 58 KBAR  | -311 198  | 46 12 | 69 69 | 0.20   | -0.31 | 0.18  | 0.64  | 0.50 |     |        |      |
| 66 PBAR  | -2212 1   | 52 12 | 0 0   | -0.13  | -2.14 | 0.78  | 2.47  | 0.94 |     |        |      |
| 67 PI0   | 111 1     | 53 18 | 0 0   | -0.52  | -0.07 | 0.20  | 0.58  | 0.13 |     |        |      |
| 68 PI-   | -211 1    | 53 18 | 0 0   | -4.09  | -0.50 | -0.25 | 4.13  | 0.14 |     |        |      |

#### ---STRONG HADRON DECAYS---

Next come the decay products of unstable hadrons generated in the cluster decays. Remember that the  $\pi^0$  was set stable in the initialization phase.

| IHEP ID | IDPDG  | IST   | M01    | M02    | DA1   | DA2   | P-X   | P-Y  | P-Z | ENERGY | MASS |
|---------|--------|-------|--------|--------|-------|-------|-------|------|-----|--------|------|
| 92 CLUS | 91 183 | 81 91 | 95 96  | -7.54  | 0.18  | 1.76  | 8.06  | 2.22 |     |        |      |
| 93 CLUS | 91 183 | 85 89 | 97 98  | 17.63  | 8.17  | -3.51 | 19.87 | 2.22 |     |        |      |
| 94 CLUS | 91 183 | 87 89 | 99 100 | -24.13 | -0.77 | 2.51  | 24.34 | 1.76 |     |        |      |

#### ---CLUSTER FORMATION---

The weak decay products are thus the lepton pair already processed plus three clusters, which decay as in a primary hard process:

| IHEP ID | IDPDG  | IST   | M01    | M02    | DA1   | DA2   | P-X   | P-Y  | P-Z | ENERGY | MASS |
|---------|--------|-------|--------|--------|-------|-------|-------|------|-----|--------|------|
| 92 CLUS | 91 183 | 81 91 | 95 96  | -7.54  | 0.18  | 1.76  | 8.06  | 2.22 |     |        |      |
| 93 CLUS | 91 183 | 85 89 | 97 98  | 17.63  | 8.17  | -3.51 | 19.87 | 2.22 |     |        |      |
| 94 CLUS | 91 183 | 87 89 | 99 100 | -24.13 | -0.77 | 2.51  | 24.34 | 1.76 |     |        |      |

## 20 Subroutine Descriptions

-->CLUSTER DECAYS--

| HEP ID   | IDPDG IST | M01   | M02   | DA1   | DA2   | P-X    | P-Y    | P-Z   | ENERGY | MASS  |
|----------|-----------|-------|-------|-------|-------|--------|--------|-------|--------|-------|
| 95 PI+   | 211       | 1     | 92    | 72    | 0     | -0.62  | -0.11  | 0.22  | 0.68   | 0.14  |
| 96 DC*-  | -413      | 197   | 92    | 72    | 102   | -6.92  | 0.30   | 1.54  | 7.38   | 2.01  |
| 97 DC**+ | 413       | 197   | 93    | 84    | 103   | 104    | 15.59  | 7.08  | -3.15  | 17.53 |
| 98 PI0   | 111       | 1     | 93    | 84    | 0     | 2.04   | 1.09   | -0.36 | 2.34   | 0.13  |
| 99 PI+   | 211       | 1     | 94    | 86    | 0     | -10.09 | -0.29  | 0.92  | 10.14  | 0.14  |
| 100 A20  | 115       | 197   | 94    | 86    | 105   | 106    | -14.04 | -0.48 | 1.58   | 14.20 |
| .....    | .....     | ..... | ..... | ..... | ..... | .....  | .....  | ..... | .....  | ..... |
| 107 KO   | 311       | 198   | 101   | 72    | 114   | 114    | -2.67  | 0.52  | 1.02   | 2.95  |
| 108 ETPR | 331       | 198   | 101   | 72    | 115   | 116    | -3.69  | -0.23 | 0.43   | 3.85  |
| 109 KBAR | -311      | 198   | 103   | 84    | 117   | 117    | 4.99   | 2.37  | -0.95  | 5.62  |
| 110 RH00 | 113       | 198   | 103   | 84    | 118   | 119    | 5.11   | 2.33  | -1.00  | 5.76  |
| 111 ETA  | 221       | 198   | 103   | 84    | 120   | 122    | 4.57   | 1.98  | -0.99  | 5.11  |
| .....    | .....     | ..... | ..... | ..... | ..... | .....  | .....  | ..... | .....  | ..... |
| 123 PI-  | -211      | 1     | 116   | 72    | 0     | 0      | -2.39  | -0.25 | 0.65   | 2.49  |
| 124 PI+  | 211       | 1     | 116   | 72    | 0     | -0.49  | -0.02  | -0.17 | 0.54   | 0.14  |

The subsequent charmed particle decays are generated according to the simplified particle data tables. In  $t \rightarrow b \rightarrow c$  decays, the secondary  $b$ 's would be hadronized, undergo strong decays and then weak decay in the same way as above.

HEP ID IDPDG IST M01 M02 DA1 DA2 P-X P-Y P-Z ENERGY MASS

101 DCBR -421 198 96 72 107 108 -6.37 0.29 1.45 6.80 1.96

102 PI- -211 1 96 72 0 0 -0.56 0.01 0.09 0.58 0.14

103 DCO 421 198 97 84 109 111 14.66 6.68 -2.94 16.48 1.86

104 PI+ 211 1 97 84 0 0 0.93 0.41 -0.21 1.04 0.14

..... 311 198 101 72 114 114 -2.67 0.52 1.02 2.95 0.50

108 ETPR 331 198 101 72 115 116 -3.69 -0.23 0.43 3.85 0.96

109 KBAR -311 198 103 84 117 117 4.99 2.37 -0.95 5.62 0.50

110 RH00 113 198 103 84 118 119 5.11 2.33 -1.00 5.76 0.77

111 ETA 221 198 103 84 120 122 4.57 1.98 -0.99 5.11 0.56

..... -211 1 116 72 0 0 -2.39 -0.25 0.65 2.49 0.14

123 PI- 211 1 116 72 0 0 -0.49 -0.02 -0.17 0.54 0.14

Generation of this event is now complete. In the case of hadron collisions, this would be followed by the treatment of the soft underlying event. The undecayed beam clusters would undergo a soft collision with the remaining centre-of-mass energy in parallel with the hard subprocesses. This would produce soft clusters with a flat rapidity distribution and low transverse momenta, which would decay to provide the low  $p_T$  hadrons of the soft underlying event.

After the 100 events requested have been generated, an analysis of the associated weight distribution is printed. The information

$$\text{NUMBER OF WEIGHTS} = 1339$$

and

$$\begin{aligned} \text{CROSS SECTION (PB)} &= 732.2993 \\ \text{ERROR IN C-S (PB)} &= 23.8432 \end{aligned}$$

gives us a new, independent estimate of the cross section (which is less accurate in this case since the number of weights is less than the 2000 generated in the initial search). Finally we are told that the Monte Carlo efficiency of elementary subprocess generation during the run was 8%, as was predicted in the initial search:

$$\text{EFFICIENCY PERCENT} = 8.0702$$

| Name    | Description                                  |
|---------|----------------------------------------------|
| HWIGPR  | Main program                                 |
| HWIGIN  | Default initializations                      |
| HWABEG  | Initializes user's analysis                  |
| HWAEND  | Terminates user's analysis                   |
| HWANAL  | Performs user's analysis on event            |
| HWBZCF  | Computes azimuthal correlation functions     |
| HWBCON  | Makes colour connections between jets        |
| HWBFNM  | Transfers external lines of jet to /HSEPEVT/ |
| HWBGEM  | Finds unevolved partons and generates jets   |
| HWBJC0  | Combines jets with correct kinematics        |
| HWBMAS  | Computes masses and trans. momenta in jet    |
| HWBRAAN | Generates a timelike parton branching        |
| HWBSFA  | Computes momenta in spacelike jet            |
| HWBSFM  | Computes spin density/decay matrices         |
| HWBSUD  | Computes (or reads) Sudakov form factors     |
| HWBSUNG | Integrand in gluon Sudakov form factor       |
| HWBSU1  | First term in quark Sudakov form factor      |
| HWBSU2  | Second term in quark Sudakov form factor     |
| HWBTM   | Computes momenta in timelike jet             |
| HWBYMC  | Virtual mass cutoff for parton type ID       |
| HWCCUT  | Cuts a massive cluster in two                |
| HWCBEC  | Decays clusters into primary hadrons         |
| HWCFIA  | Sets up flavours for HNCIAHAD                |
| HWCFOR  | Forms clusters                               |
| HWCGSP  | Splits gluons                                |
| HWCHAD  | Decays a cluster into one or two hadrons     |
| HWDBOS  | Finds and decays W and Z bosons              |
| HWDBDZ  | Chooses decay mode of W and Z bosons         |
| HWDHAD  | Generates decays of unstable hadrons         |
| HWDTIG  | Finds and decays Higgs bosons                |
| HWDVVY  | Finds and decays heavy flavours              |
| HWDDPP  | Chooses a parton for HADHVV                  |
| HWDPWT  | Phase space decay weight                     |
| HWDTHR  | Generates a three-body decay                 |
| HWDTFO  | Generates a two-body decay                   |
| HWDTWT  | Weak ( $V - A$ ) decay weight                |
| HWETNM  | Final calculations on elementary subprocess  |
| HWEGAM  | Generates Weizsaecker-Williams photon        |
| HWEINI  | Initializes elementary subprocess            |
| HWEOME  | Sets up a $2 \rightarrow 1$ hard subprocess  |
| HWEPRO  | Generates elementary subprocess              |
| HWEITW0 | Sets up a $2 \rightarrow 2$ hard subprocess  |

| Name          | Description                                                         |
|---------------|---------------------------------------------------------------------|
| <b>HWHBGF</b> | Hard subprocess: boson-gluon fusion (BGF)                           |
| <b>HWHBK1</b> | Computes kinematics for BGF                                         |
| <b>HWHBRW</b> | Returns a phase-space point for BGF                                 |
| <b>HWHBSG</b> | Computes cross section for BGF                                      |
| <b>HWHDIS</b> | Hard subprocess: deep inelastic $e/\mu$ quark                       |
| <b>HWHDTP</b> | Hard subprocess: Drell-Yan $Z^0/\gamma$ prodn                       |
| <b>HWHFPA</b> | Hard subprocess: $e^+e^- \rightarrow q\bar{q}$                      |
| <b>HWHFPG</b> | Hard subprocess: $e^+e^- \rightarrow q\bar{q}g$                     |
| <b>HWHFW</b>  | Hard subprocess: $e^+e^- \rightarrow W^+W^-$                        |
| <b>HWHENO</b> | $e^+e^- \rightarrow W^+W^-$ subroutine                              |
| <b>HWHEM1</b> | $e^+e^- \rightarrow W^+W^-$ subroutine                              |
| <b>HWHEW2</b> | $e^+e^- \rightarrow W^+W^-$ subroutine                              |
| <b>HWHEN3</b> | $e^+e^- \rightarrow W^+W^-$ subroutine                              |
| <b>HWHEN4</b> | $e^+e^- \rightarrow W^+W^-$ subroutine                              |
| <b>HWHHVV</b> | Hard subprocess: heavy quark production                             |
| <b>HWHIGC</b> | Higgs $\rightarrow \gamma\gamma$ decay                              |
| <b>HWHIGF</b> | Higgs $\rightarrow W^+W^-$ decay                                    |
| <b>HWHIGH</b> | Choose Higgs mass for production routines                           |
| <b>HWHIGS</b> | Hard subprocess: $gg \rightarrow$ Higgs                             |
| <b>HWHIGT</b> | Computes $gg \rightarrow$ Higgs cross section                       |
| <b>HWHIGN</b> | $W+W^- \rightarrow$ Higgs cross section                             |
| <b>HWHIGY</b> | Computes $e^+e^- \rightarrow Z^0 \rightarrow Z^0 H^0$ cross section |
| <b>HWHIGZ</b> | Hard subprocess: $e^+e^- \rightarrow Z^0 \rightarrow Z^0 H^0$       |
| <b>HWHPHD</b> | Hard subprocess: direct photon production                           |
| <b>HWHQCD</b> | Hard subprocess: QCD $2 \rightarrow 2$                              |
| <b>HWHQCP</b> | Identifies QCD $2 \rightarrow 2$ hard subprocess                    |
| <b>HWHWEX</b> | Top production by W exchange                                        |
| <b>HWHWPR</b> | Hard subprocess: W production                                       |
| <b>HWHWJ</b>  | Hard subprocess: W + jet production                                 |
| <b>HWHWTT</b> | Generates min bias or soft underlying event                         |
| <b>HWHWPS</b> | Generates longitudinal phase space                                  |
| <b>HWHWBI</b> | Computes negative binomial probability                              |
| <b>HWHWLT</b> | Chooses min bias charged multiplicity                               |
| <b>HWHAZN</b> | Randomly rotated azimuth                                            |
| <b>HWREXP</b> | Random number: exponential distribution                             |
| <b>HWREXT</b> | Random number: exponential transverse mass                          |
| <b>HWRGAU</b> | Random number: Gaussian                                             |
| <b>HWRGEN</b> | Random number generator (1'Ecuyer's method)                         |
| <b>HWRINT</b> | Random integer                                                      |
| <b>HWRLOG</b> | Random logical                                                      |
| <b>HWRPOW</b> | Random number: power distribution                                   |
| <b>HWRUNG</b> | Random number: uniform + Gaussian tails                             |
| <b>HWRUND</b> | Random number: uniform                                              |

| Name            | Description                                                |
|-----------------|------------------------------------------------------------|
| <b>HWSRBN</b>   | Generates spacelike parton branching                       |
| <b>HWSRGG</b>   | Drees-Grassie gluon distribution in photon                 |
| <b>HWSRQ</b>    | Drees-Grassie quark distribution in photon                 |
| <b>HWSFR</b>    | Chooses a spacelike branching                              |
| <b>HWSFUN</b>   | Hadron structure functions                                 |
| <b>HWSAM</b>    | Gamma function (for structure functions)                   |
| <b>HWSSEN</b>   | Generates $\pi$ values for spacelike partons               |
| <b>HWSQQ</b>    | Inserts $g \rightarrow q\bar{q}$ part of gluon form factor |
| <b>HWSPC</b>    | Replaces spacelike partons by spectators                   |
| <b>HWSUD</b>    | Sudakov form factor/structure function                     |
| <b>HWSTAB</b>   | Interpolates in function table (for HWSUD)                 |
| <b>HWSVAL</b>   | Checks for valence parton                                  |
| <b>HWAULF</b>   | Two-loop QCD running coupling constant                     |
| <b>HWAUPR</b>   | Prints branching data for last parton shower               |
| <b>HWADAT</b>   | Particle properties (N.B. BLOCK DATA)                      |
| <b>HWAEC</b>    | Computes $e^+e^-$ cross section and asymmetry (given $q$ ) |
| <b>HWAEEQ</b>   | $e^+e^-$ cross section and asymmetry (given $q$ )          |
| <b>HWAEPR</b>   | Prints event data                                          |
| <b>HWAIAU</b>   | Adaptive Gaussian integration                              |
| <b>HWAIDT</b>   | Translates particle identity codes                         |
| <b>HWAUTNC</b>  | Initial parameter-dependent calculations                   |
| <b>HWAUNE</b>   | Initializes an event                                       |
| <b>HWAUDO</b>   | Lorentz 4-vector dot product                               |
| <b>HWAUDOB</b>  | Lorentz transformation: rest frame $\rightarrow$ lab       |
| <b>HWAUOF</b>   | Lorentz transformation: lab $\rightarrow$ rest frame       |
| <b>HWAUOR</b>   | Multiples by Lorentz matrix                                |
| <b>HWAUMAS</b>  | Puts mass in 5th component of vector                       |
| <b>HWAUCM</b>   | Centre-of-mass momentum                                    |
| <b>HWAUPAP</b>  | Rapidity                                                   |
| <b>HWAUES</b>   | Computes/print resonance data                              |
| <b>HWAUROB</b>  | Rotation by inverse of matrix $R$                          |
| <b>HWAUDDF</b>  | Rotation by matrix $R$                                     |
| <b>HWAUROT</b>  | Computes rotation $R$ from vector to z-axis                |
| <b>HWAUSOR</b>  | Sorts an array in ascending order                          |
| <b>HWAUSQR</b>  | Square root with sign retention                            |
| <b>HWAUSTA</b>  | Makes a particle type stable                               |
| <b>HWAUTAB</b>  | Interpolates in a table                                    |
| <b>HWAUTIM</b>  | Checks time remaining (N.B. VAX Fortran)                   |
| <b>HWAUDIF</b>  | Vector difference                                          |
| <b>HWAUDOT</b>  | Vector dot product                                         |
| <b>HWAUEQU</b>  | Vector equality                                            |
| <b>HWAUSCA</b>  | Vector times scalar                                        |
| <b>HWAUSUM</b>  | Vector sum                                                 |
| <b>HWAUZERO</b> | Vector zero                                                |
| <b>HWAURN</b>   | Issues warnings and deals with errors                      |

## Acknowledgements

Two of us (G.A. and I.S.) are grateful for the hospitality of DESY while a part of this work was performed.

## References

- [1] G. Marchesini and B.R. Webber, Nucl. Phys. B310 (1988) 461.
- [2] J.G. Knowles, Nucl. Phys. B304 (1988) 794.
- [3] J.G. Knowles, Nucl. Phys. B310 (1988) 571.
- [4] J.G. Knowles, Comp. Phys. Comm. 58 (1990) 271.
- [5] G. Marchesini, and B.R. Webber, Nucl. Phys. B330 (1990) 261.
- [6] S. Catani, G. Marchesini and B.R. Webber, Nucl. Phys. B349 (1991) 635.
- [7] G. Abbiendi and L. Stanco, DESY preprint 90-103, to be published in Comp. Phys. Comm.
- G. Abbiendi and L. Stanco, DESY preprint 90-112, to be published in Zeit. Phys. C.
- [8] A. Bassetto, M. Ciafaloni and G. Marchesini, Phys. Rep. 100 (1983) 201.
- [9] G. Marchesini and B.R. Webber, Nucl. Phys. B238 (1984) 1.
- [10] B.R. Webber, Nucl. Phys. B238 (1984) 492.
- [11] B.R. Webber, Ann. Rev. Nucl. Part. Sci. 36 (1986) 253.
- [12] G. Marchesini and B.R. Webber, Phys. Rev. D 38 (1988) 3419.
- [13] S. Catani and L. Trentadue, Phys. Lett. 217B (1989) 539; Nucl. Phys. B327 (1989) 353; Nucl. Phys. B353 (1991) 183.
- [14] Yu.I. Dokshitzer, V.A. Khoze and S.I. Troyan, in Proc. 6th Int. Conf. on Physics in Collision, ed. M. Derrick (World Scientific, Singapore, 1987).
- [15] Yu.L. Dokshitzer, V.A. Khoze, G. Marchesini and B.R. Webber, in Proc. XXV Rencontre de Moriond, March 1990, ed. J. Tran Thanh Van (Editions Frontieres, Gif sur Yvette).
- [16] JADE Collaboration, W. Bartel et al., Phys. Lett. 134B (1983) 275; Phys. Lett. 157B (1985) 340;
- TPC Collaboration, H. Alibra et al., Zeit. Phys. C28 (1985) 31;
- OPAL Collaboration, M.Z. Akrawy et al., CERN/PPE/91/31 (1991).
- [17] Yu.L. Dokshitzer, V.A. Khoze, S.I. Troyan and A.H. Mueller, Rev. Mod. Phys. 60 (1988) 373.
- [18] Yu.I. Dokshitzer, V.A. Khoze and S.I. Troyan, in Perturbative Quantum Chromodynamics, ed. A.H. Mueller (World Scientific, Singapore, 1989), p. 241.
- [19] C.P. Fong and B.R. Webber, Phys. Lett. 229B (1989) 289; Phys. Lett. 241B (1990) 255; Nucl. Phys. B355 (1991) 54.
- [20] OPAL Collaboration, M.Z. Akrawy et al., Phys. Lett. 247B (1990) 617.
- [21] S. Catani, F. Fiorani and G. Marchesini, Nucl. Phys. B336 (1990) 18.
- [22] L.N. Lipatov, Yad. Fiz. 23 (1976) 642 [Sov. J. Nucl. Phys. 23 (1976) 338]; E.A. Kuraev, I.N. Lipatov and V.S. Fadin, Zh. Eksp. Teor. Fiz. 72 (1977) 373 [Sov. Phys. JETP 45 (1977) 199]; Ya.Ya. Balitskii and L.N. Lipatov, Yad. Fiz. 28 (1978) 1597 [Sov. J. Nucl. Phys. 28 (1978) 822]; J. Bartels, Nucl. Phys. B151 (1979) 293; T. Jaroszewicz, Acta Phys. Pol. B11 (1980) 965; Phys. Lett. 116B (1982) 291.
- [23] G. Marchesini, and B.R. Webber, Nucl. Phys. B349 (1991) 617.
- [24] R.K. Ellis, G. Marchesini and B.R. Webber, Nucl. Phys. B286 (1987) 643.
- [25] D. Amati and G. Veneziano, Phys. Lett. 83B (1979) 87.
- [26] Yu.L. Dokshitzer and S.I. Troyan, Leningrad Nuclear Physics Institute preprint N922 (1984); Ya.I. Azimov, Yu.L. Dokshitzer, V.A. Khoze and S.I. Troyan, Phys. Lett. 165B (1985) 147; Zeit. Phys. C27 (1985) 65.
- [27] B. Andersson, G. Gustafsson, G. Ingelman and T. Sjöstrand, Phys. Rep. 97 (1983) 33.
- [28] UA1 Collaboration, C. Albajar et al., Nucl. Phys. B309 (1988) 405.
- [29] UA5 Collaboration, G.J. Alner et al., Nucl. Phys. B291 (1987) 445.
- [30] HERWIG version 3.4; CERN Program Library long write-up W5037.
- [31] T. Sjöstrand et al., in "Z Physics at LEP 1", ed. G. Altarelli, R. Kleiss and C. Verzegnassi, CERN Yellow Report 89-08, v.3, p.325.
- [32] OPAL Collaboration, M.Z. Akrawy et al., Zeit. Phys. C49 (1991) 375.
- [33] Z. Kunszt, P. Nason, G. Marchesini and B.R. Webber, in "Z Physics at LEP 1", ed. G. Altarelli, R. Kleiss and C. Verzegnassi, CERN Yellow Report 89-08, v.1, p.373.
- [34] J.W. Gary, private communication.
- [35] Z. Kunszt, private communication.
- [36] M. Drees and K. Grässle, Zeit. Phys. C28 (1985) 451.
- [37] PDFLIB: CERN Program Library long write-up W5051.
- [38] F. James, Comp. Phys. Comm. 60 (1990) 329.
- [39] Particle Data Group, J.J. Hernández et al., Review of Particle Properties, Phys. Lett. 239B (1990) 1.