

Cofinalities of Marczewski-like ideals

Jörg Brendle*
Graduate School of System Informatics
Kobe University
Rokko-dai 1-1, Nada-ku
Kobe 657-8501, Japan
email: brendle@kobe-u.ac.jp

and

Yurii Khomskii†
Hamburg University
Bundesstraße 55 (Geomatikum)
20146 Hamburg, Germany
email: yurii@deds.nl

and

Wolfgang Wohofsky†
Hamburg University
Bundesstraße 55 (Geomatikum)
20146 Hamburg, Germany
email: wolfgang.wohofsky@gmx.at

October 12, 2016

*Partially supported by Grants-in-Aid for Scientific Research (C) 24540126 and (C) 15K04977, Japan Society for the Promotion of Science, by JSPS and FWF under the Japan-Austria Research Cooperative Program *New developments regarding forcing in set theory*, and by the Isaac Newton Institute for Mathematical Sciences in the programme *Mathematical, Foundational and Computational Aspects of the Higher Infinite* (HIF) funded by EPSRC grant EP/K032208/1.

†Partially supported by the Isaac Newton Institute for Mathematical Sciences in the programme *Mathematical, Foundational and Computational Aspects of the Higher Infinite* (HIF) funded by EPSRC grant EP/K032208/1.

2010 Mathematics Subject Classification. Primary 03E17; Secondary 03E40, 03E50

Key Words. Laver forcing, Miller forcing, tree forcing, Marczewski ideal, nowhere Ramsey ideal, Laver ideal, Miller ideal, tree ideal, cofinality of an ideal

Abstract

We show that the cofinalities of both the Miller ideal m^0 (the σ -ideal naturally related to Miller forcing \mathbb{M}) and the Laver ideal ℓ^0 (related to Laver forcing \mathbb{L}) are larger than the size of the continuum \mathfrak{c} in ZFC.

1 Introduction

The purpose of this note is to prove (in ZFC) that the ideals naturally related to Laver forcing \mathbb{L} and to Miller forcing \mathbb{M} , the *Laver ideal* ℓ^0 and the *Miller ideal* m^0 , have cofinality strictly larger than \mathfrak{c} , the size of the continuum (Corollary 18 below). We will phrase our result in a more general framework and show that $\text{cof}(t^0) > \mathfrak{c}$ holds for all tree ideals t^0 derived from tree forcings \mathbb{T} satisfying a certain property (Theorem 13 in Section 3). This was known previously for the Marczewski ideal s^0 [JMS] and the nowhere Ramsey ideal r^0 [Ma], but it is unclear whether the method of proof for these two ideals works for ℓ^0 and m^0 (see the discussion in Section 2), and our approach is more general.

For a subtree $T \subseteq \omega^{<\omega}$ (or $T \subseteq 2^{<\omega}$), $[T] = \{x \in \omega^\omega : x \upharpoonright n \in T \text{ for all } n\}$ denotes the set of branches through T .

Definition 1 (Combinatorial tree forcing). A collection \mathbb{T} of subtrees of $\omega^{<\omega}$ is a *combinatorial tree forcing* if

1. $\omega^{<\omega} \in \mathbb{T}$,
2. (closure under subtrees) if $T \in \mathbb{T}$ and $s \in T$, then the tree $T_s = \{t \in T : s \subseteq t \text{ or } t \subseteq s\}$ also belongs to \mathbb{T} ,
3. (large disjoint antichains) there is a continuous function $f : \omega^\omega \rightarrow 2^\omega$ such that for all $x \in 2^\omega$, $f^{-1}(\{x\})$ is the set of branches of a tree in \mathbb{T} ,
4. (homogeneity) if $T \in \mathbb{T}$, then there is an order-preserving injection $i : \omega^{<\omega} \rightarrow T$ such that the map $g : \omega^\omega \rightarrow [T]$ given by $g(x) = \bigcup \{i(x \upharpoonright n) : n \in \omega\}$ is a homeomorphism and for any subtree $S \subseteq \omega^{<\omega}$, $S \in \mathbb{T}$ iff the downward closure of $i(S)$ belongs to \mathbb{T} .

\mathbb{T} is ordered by inclusion, that is, for $S, T \in \mathbb{T}$, $S \leq T$ if $S \subseteq T$.

Homogeneity says that the forcing looks the same below each condition. In view of homogeneity, “large disjoint antichains” implies that

5. each $T \in \mathbb{T}$ splits into continuum many trees with pairwise disjoint sets of branches, that is, there are $T_\alpha \in \mathbb{T}$, $\alpha < \mathfrak{c}$, with $T_\alpha \subseteq T$ such that $[T_\alpha] \cap [T_\beta] = \emptyset$ for $\alpha \neq \beta$.

In particular, there are \mathfrak{c} -sized antichains so that \mathbb{T} is not ccc and forcing notions like Cohen and random forcing do not fit into this framework.

For forcing notions whose conditions are subtrees of $2^{<\omega}$ like Sacks forcing \mathbb{S} , an analogous definition applies, with $\omega^{<\omega}$ and ω^ω replaced by $2^{<\omega}$ and 2^ω , respectively.

Definition 2 (Tree ideal). The *tree ideal* t^0 associated with the combinatorial tree forcing \mathbb{T} consists of all $X \subseteq \omega^\omega$ such that for all $T \in \mathbb{T}$ there is $S \leq T$ with $X \cap [S] = \emptyset$.

When $\mathbb{T} = \mathbb{S}$ is Sacks forcing, $t^0 = s^0$ is the well-known *Marczewski ideal*, and for Mathias forcing $\mathbb{T} = \mathbb{R}$, $t^0 = r^0$ is the ideal of *nowhere Ramsey* sets. *Laver forcing* \mathbb{L} [La] consists of trees $T \subseteq \omega^{<\omega}$ such that for all $t \in T$ containing the stem of T , $\text{stem}(T)$, the set of successors $\text{succ}_T(t) = \{n \in \omega : t \hat{\ } n \in T\}$ is infinite. *Miller forcing* \mathbb{M} [Mi] consists of trees $T \subseteq \omega^{<\omega}$ such that for all $t \in T$ there is $s \supseteq t$ in T such that $\text{succ}_T(s)$ is infinite. Note that \mathbb{L} and \mathbb{M} are combinatorial tree forcings in the above sense for $f : \omega^\omega \rightarrow 2^\omega$ given by $f(x)(n) = x(n) \bmod 2$ for all $x \in \omega^\omega$ and $n \in \omega$ witnesses “large disjoint antichains”. The Laver and Miller ideals ℓ^0 and m^0 are the corresponding tree ideals. For basic facts about such tree ideals, like non-inclusion between different ideals, see e.g. [Br].

Definition 3 (Cofinality of an ideal). Given an ideal \mathcal{I} , its *cofinality* $\text{cof}(\mathcal{I})$ is the smallest cardinality of a family $\mathcal{J} \subseteq \mathcal{I}$ such that every member of \mathcal{I} is contained in a member of \mathcal{J} .

A family like \mathcal{J} in this definition is said to be a *basis* of \mathcal{I} (or: *cofinal* in \mathcal{I}).

While the topic of our work are cofinalities of tree ideals, we note that other cardinal invariants of tree ideals t^0 , such as the *additivity* $\text{add}(t^0)$ (the least size of a subfamily $\mathcal{J} \subseteq t^0$ whose union is not in t^0) and the *covering number* $\text{cov}(t^0)$ (the least size of a subfamily $\mathcal{J} \subseteq t^0$ whose union is ω^ω) have been studied as well. If there is a fusion argument for \mathbb{T} , t^0 is a σ -ideal, and one has $\omega_1 \leq \text{add}(t^0) \leq \text{cov}(t^0) \leq \mathfrak{c}$, while the exact value of these two cardinals depends on the model of set theory. Furthermore, by “large disjoint antichains”, the *uniformity* $\text{non}(t^0)$ of a tree ideal t^0 (the smallest cardinality of a subset of ω^ω not belonging to t^0) is always equal to \mathfrak{c} . Since $\text{cof}(\mathcal{I}) \geq \text{non}(\mathcal{I})$ for any non-trivial ideal \mathcal{I} , $\text{cof}(t^0) \geq \mathfrak{c}$ follows, and the main problem about cofinalities of tree ideals is whether they can be equal to \mathfrak{c} or must be strictly above \mathfrak{c} .

The question whether $\text{cof}(\ell^0)$ and $\text{cof}(m^0)$ are larger than \mathfrak{c} was discussed in private communication with M. Dečo and M. Repický, and Repický [Re] in the meantime used our result to obtain a characterization of $\text{cof}(\ell^0)$ as $\mathfrak{d}((\ell^0)^\mathfrak{c})$.

Acknowledgments. This research was started in February 2015 while the first author was visiting the other two authors at the KGRC of the University of Vienna and at the Vienna University of Technology. He thanks the JSPS for its support and the universities in Vienna for their hospitality. Our work was continued in November 2015 at the University of East Anglia (first and second authors) and at the Isaac Newton Institute for Mathematical Sciences (INI) (first and third authors). We thank the INI for its support and both institutions for their hospitality.

2 The disjoint maximal antichain property

Definition 4. Let \mathbb{T} be a combinatorial tree forcing. \mathbb{T} has the *disjoint maximal antichain property* if there is a maximal antichain $(T_\alpha : \alpha < \mathfrak{c})$ in \mathbb{T} such that $[T_\alpha] \cap [T_\beta] = \emptyset$ for all $\alpha \neq \beta$.

The following has been known for some time (see also [Re, Theorem 1.2]).

Proposition 5. *Assume \mathbb{T} has the disjoint maximal antichain property. Then $\text{cf}(\text{cof}(t^0)) > \mathfrak{c}$.*

Proof. Let $(T_\alpha : \alpha < \mathfrak{c})$ be a disjoint maximal antichain in \mathbb{T} . Also let $\kappa = \text{cf}(\text{cof}(t^0))$ and assume $\kappa \leq \mathfrak{c}$. We shall derive a contradiction. Assume $\mathcal{X}_\alpha \subseteq t^0$, $\alpha < \kappa$, are of size $< \text{cof}(t^0)$. We shall show that $\mathcal{X} = \bigcup \{\mathcal{X}_\alpha : \alpha < \kappa\}$ is not cofinal in t^0 . By homogeneity of the tree forcing \mathbb{T} , we know that \mathcal{X}_α is not cofinal below T_α , that is, there is $X_\alpha \in t^0$, $X_\alpha \subseteq [T_\alpha]$, such that $X_\alpha \not\subseteq Y$ for all $Y \in \mathcal{X}_\alpha$. Let $X = \bigcup \{X_\alpha : \alpha < \kappa\}$. By disjointness of the maximal antichain, we see that $X \in t^0$. Obviously $X \not\subseteq Y$ for all $Y \in \mathcal{X}$, and we are done. \square

Note that for only showing $\text{cof}(t^0) > \mathfrak{c}$, the homogeneity of the forcing is not needed (that is, properties 1, 2, and 5 of Definition 1 are enough).

Definition 6. Let \mathbb{T} be a combinatorial tree forcing. \mathbb{T} has the *incompatibility shrinking property* if for any $T \in \mathbb{T}$ and any family $(S_\alpha : \alpha < \mu)$, $\mu < \mathfrak{c}$, in \mathbb{T} such that S_α is incompatible with T for all α , one can find $T' \leq T$ such that $[T']$ is disjoint from all the $[S_\alpha]$.

Proposition 7. *Let \mathbb{T} be a combinatorial tree forcing. The incompatibility shrinking property for \mathbb{T} implies the disjoint maximal antichain property for \mathbb{T} . In fact, it implies that any maximal antichain can be refined to a disjoint maximal antichain.*

Proof. Let $(T_\alpha : \alpha < \mathfrak{c})$ be a dense set of conditions in \mathbb{T} all of which lie below a given maximal antichain of size \mathfrak{c} . We construct $A \subseteq \mathfrak{c}$ of size \mathfrak{c} and $\{S_\alpha : \alpha \in A\} \subseteq \mathbb{T}$ such that

- $S_\alpha \leq T_\alpha$ for $\alpha \in A$,
- if $\alpha \notin A$, then T_α is compatible with some S_β for $\beta < \alpha$ with $\beta \in A$,
- $[S_\alpha] \cap [S_\beta] = \emptyset$ for $\alpha \neq \beta$ from A .

Clearly, these conditions imply that $(S_\alpha : \alpha \in A)$ is a disjoint maximal antichain. Also A must necessarily have size \mathfrak{c} .

Suppose we are at stage $\alpha < \mathfrak{c}$ of the construction. If T_α is compatible with some S_β where $\beta < \alpha$, $\beta \in A$, let $\alpha \notin A$, and we are done. If this is not the case, let $\alpha \in A$. By the incompatibility shrinking property we find $T' = S_\alpha$ as required. \square

Example 8. *Sacks forcing \mathbb{S} , Mathias forcing \mathbb{R} , and Silver forcing \mathbb{V} have the incompatibility shrinking property and thus also the disjoint maximal antichain property.*

To see this, simply use that for any two incompatible $S, T \in \mathbb{S}$, $[S] \cap [T]$ is at most countable, while in the case of \mathbb{V} , this intersection is finite and for \mathbb{R} , even empty.

From this we obtain that $cf(\text{cof}(s^0)) > \mathfrak{c}$ [JMS, Theorem 1.3] where s^0 is the Marczewski ideal, that $cf(\text{cof}(r^0)) > \mathfrak{c}$ [Ma] where r^0 is the ideal of nowhere Ramsey sets, and that $cf(\text{cof}(v^0)) > \mathfrak{c}$ where v^0 is the Silver ideal.

We also note that if there is a fusion argument for \mathbb{T} , then the continuum hypothesis CH implies the incompatibility shrinking property and thus also the disjoint maximal antichain property. For Laver and Miller forcings, a weaker hypothesis is sufficient.

Proposition 9. *Assume $\mathfrak{b} = \mathfrak{c}$. Then Laver forcing \mathbb{L} has the incompatibility shrinking property and thus also the disjoint maximal antichain property.*

Proof. Fix $T \in \mathbb{L}$, $\mu < \mathfrak{c}$, and any family $(S_\alpha : \alpha < \mu)$ in \mathbb{L} such that S_α is incompatible with T for all α . Since $T \cap S_\alpha$ does not contain a Laver tree, by [GRSS, Lemma 2.3], there is a function $g_\alpha : \omega^{<\omega} \rightarrow \omega$ such that if $x \in [T] \cap [S_\alpha]$, then there are infinitely many n with $x(n) < g_\alpha(x \upharpoonright n)$. By $\mathfrak{b} = \mathfrak{c}$, there is $f : \omega^{<\omega} \rightarrow \omega$ eventually dominating all g_α . Let $T' = \{s \in T : s(n) > f(s \upharpoonright n) \text{ for all } n \in \text{dom}(s) \text{ beyond the stem of } T\}$. Clearly T' is still a Laver tree with the same stem as T . Furthermore, $[T'] \cap [S_\alpha] = \emptyset$ for if x belonged to the intersection, we would have $x(n) < g_\alpha(x \upharpoonright n)$ for infinitely many n and $x(n) > f(x \upharpoonright n)$ for all n beyond the stem of T' , a contradiction. \square

A similar argument which we leave to the reader shows:

Proposition 10. *Assume $\mathfrak{d} = \mathfrak{c}$. Then Miller forcing \mathbb{M} has the incompatibility shrinking property and thus also the disjoint maximal antichain property.*

Question 11. *Do \mathbb{L} or \mathbb{M} have the disjoint maximal antichain property in ZFC?*

3 The selective disjoint antichain property

We now consider a property weaker than the disjoint maximal antichain property which is sufficient to show that the cofinalities of the Laver ideal ℓ^0 and the Miller ideal m^0 are larger than \mathfrak{c} in ZFC.

Definition 12. Let \mathbb{T} be a combinatorial tree forcing. \mathbb{T} has the *selective disjoint antichain property* if there is an antichain $(T_\alpha : \alpha < \mathfrak{c})$ in \mathbb{T} such that

- $[T_\alpha] \cap [T_\beta] = \emptyset$ for all $\alpha \neq \beta$,
- for all $T \in \mathbb{T}$ there is $S \leq T$ such that
 - either $S \leq T_\alpha$ for some $\alpha < \mathfrak{c}$,

– or $|[S] \cap [T_\alpha]| \leq 1$ for all $\alpha < \mathfrak{c}$.

We note that for our applications, it would be enough to have $|[S] \cap [T_\alpha]| \leq \aleph_0$ in the last clause.

Theorem 13. *Assume \mathbb{T} has the selective disjoint antichain property. Then $\text{cf}(\text{cof}(t^0)) > \mathfrak{c}$.*

Proof. Let $(T_\alpha : \alpha < \mathfrak{c})$ be a selective disjoint antichain in \mathbb{T} . Also assume that $(S_\beta : \beta < \mathfrak{c})$ is a list of all trees S in \mathbb{T} such that $|[S] \cap [T_\alpha]| \leq 1$ for all $\alpha < \mathfrak{c}$. Put $\kappa = \text{cf}(\text{cof}(t^0))$ and assume $\kappa \leq \mathfrak{c}$. Also assume $\mathcal{X}_\alpha \subseteq t^0$, $\alpha < \kappa$, are of size $< \text{cof}(t^0)$. As in the proof of Proposition 5, we shall show that $\mathcal{X} = \bigcup \{\mathcal{X}_\alpha : \alpha < \kappa\}$ is not cofinal in t^0 .

By “large disjoint antichains”, we find $T'_\alpha \leq T_\alpha$ such that $[T'_\alpha] \cap \bigcup_{\beta < \alpha} [S_\beta] = \emptyset$. By homogeneity, there is $X_\alpha \in t^0$ with $X_\alpha \subseteq [T'_\alpha]$ such that $X_\alpha \not\subseteq Y$ for all $Y \in \mathcal{X}_\alpha$. Let $X = \bigcup \{X_\alpha : \alpha < \kappa\}$. Obviously $X \not\subseteq Y$ for all $Y \in \mathcal{X}$. We need to show that X belongs to t^0 . To this end, let $T \in \mathbb{T}$.

First assume there is $S \leq T$ such that $S \leq T_\alpha$ for some $\alpha < \mathfrak{c}$. Then $[S] \cap X \subseteq X_\alpha$. Since $X_\alpha \in t^0$, there is $S' \leq S$ such that $[S'] \cap X_\alpha = \emptyset$, and $[S'] \cap X = \emptyset$ follows.

Next assume there is $S \leq T$ such that $|[S] \cap [T_\alpha]| \leq 1$ for all $\alpha < \mathfrak{c}$. Then $S = S_\beta$ for some $\beta < \mathfrak{c}$. By construction, we know that $X_\alpha \cap [S_\beta] = \emptyset$ for $\alpha > \beta$. Hence $[S_\beta] \cap X \subseteq \bigcup_{\alpha \leq \beta} [S_\beta] \cap [T_\alpha]$ and therefore $|[S_\beta] \cap X| < \mathfrak{c}$. Using again “large disjoint antichains”, we see that there is $S' \leq S_\beta$ such that $[S'] \cap X = \emptyset$, as required. This completes the proof of the theorem. \square

Again note that for only showing $\text{cof}(t^0) > \mathfrak{c}$, the homogeneity of the forcing is not needed (that is, properties 1, 2, and 5 of Definition 1 are enough).

The next property of a combinatorial tree forcing \mathbb{T} implies that \mathbb{T} adds a minimal real and, in fact, standard proofs of minimality go via this property.

Definition 14. Let \mathbb{T} be a combinatorial tree forcing. \mathbb{T} has the *constant or one-to-one property* if for all $T \in \mathbb{T}$ and all continuous $f : [T] \rightarrow 2^\omega$, there is $S \leq T$ such that $f \upharpoonright [S]$ is either constant or one-to-one.

It is known that both Miller forcing and Laver forcing have the constant or one-to-one property. For the former, this is implicit in work of Miller [Mi, Section 2], for the latter, in work of Gray [Gra] (see also [Gro, Theorems 2 and 7] for similar arguments). These results are formulated in terms of minimality. For completeness’ sake, we include a proof of the more difficult case of Laver forcing in our formulation. Note also that the result for \mathbb{M} is a trivial consequence of the result for \mathbb{L} .

Theorem 15 (Miller). *Miller forcing \mathbb{M} has the constant or one-to-one property.*

Theorem 16 (Gray). *Laver forcing \mathbb{L} has the constant or one-to-one property.*

Proof. Fix f and T . The pure decision property of Laver forcing implies:

Claim 16.1. Let $n \in \omega$ and $\tau \in T$ with $\text{stem}(T) \subseteq \tau$. There are $T' \leq_0 T_\tau$ and $s \in 2^n$ such that $[T'] \subseteq f^{-1}([s])$.

Claim 16.2. Let $\tau \in T$ with $\text{stem}(T) \subseteq \tau$. There are $T' \leq_0 T_\tau$ and $x = x_\tau \in 2^\omega$ such that if $(k_\tau^n : n \in \omega)$ is the increasing enumeration of $\text{succ}_{T'}(\tau)$ then $[T'_{\tau \hat{\ } k_\tau^n}] \subseteq f^{-1}([x \upharpoonright (|\tau| + n)])$.

Proof. Using Claim 16.1, construct a \leq_0 -decreasing sequence $(S^n : n \in \omega)$ with $S^0 \leq_0 T_\tau$ and a \subset -increasing sequence $(s^n \in 2^{n+|\tau|} : n \in \omega)$ such that $[S^n] \subseteq f^{-1}([s^n])$ for all n . Let $k_\tau^n = \min(\text{succ}_{S^n}(\tau) \setminus (k_\tau^{n-1} + 1))$ where we put $k_\tau^{-1} = -1$. Let T' be such that $\text{succ}_{T'}(\tau) = \{k_\tau^n : n \in \omega\}$ and $T'_{\tau \hat{\ } k_\tau^n} = S^n_{\tau \hat{\ } k_\tau^n}$. Also let $x = \bigcup_n s^n \in 2^\omega$. Then $T' \leq_0 T_\tau$ and $[T'_{\tau \hat{\ } k_\tau^n}] = [S^n_{\tau \hat{\ } k_\tau^n}] \subseteq [S^n] \subseteq f^{-1}([s^n]) = f^{-1}([x \upharpoonright (|\tau| + n)])$. \square

By Claim 16.2 and a fusion argument we see

Claim 16.3. There are $T' \leq_0 T$, $(x_\tau : \tau \in T', \text{stem}(T) \subseteq \tau)$, and $((k_\tau^n : n \in \omega) : \tau \in T', \text{stem}(T) \subseteq \tau)$ such that $(k_\tau^n : n \in \omega)$ is the increasing enumeration of $\text{succ}_{T'}(\tau)$ for all τ and $[T'_{\tau \hat{\ } k_\tau^n}] \subseteq f^{-1}([x_\tau \upharpoonright (|\tau| + n)])$ for all n and all τ . In particular $[T'_\tau] \subseteq f^{-1}([x_\tau \upharpoonright |\tau|])$ for all τ .

The properties of the x_τ imply in particular that $x_{\tau \hat{\ } k_\tau^n}$ converges to x_τ as n goes to infinity. Now define a rank function for $\tau \in T'$ as follows.

- $\rho(\tau) = 0 \iff \exists^\infty k \in \text{succ}_{T'}(\tau)$ such that $x_{\tau \hat{\ } k} \neq x_\tau$,
- for $\alpha > 0$, $\rho(\tau) = \alpha \iff \neg \rho(\tau) < \alpha \wedge \exists^\infty k \in \text{succ}_{T'}(\tau) (\rho(\tau \hat{\ } k) < \alpha)$.

By the convergence property of the x_τ , we see that $\rho(\tau) = 0$ implies in particular that the set $\{x_{\tau \hat{\ } k_\tau^n} : n \in \omega\}$ is infinite.

Case 1. $\rho(\tau) = \infty$ for some $\tau \in T'$ (i.e., the rank is undefined).

Then we can easily construct a Laver tree $S \leq T'$ such that $\text{stem}(S) = \tau$ and $x_\sigma = x_\tau$ for all $\sigma \in S$ with $\sigma \supseteq \tau$. We claim that $f \upharpoonright [S]$ is constant with value x_τ . Indeed let $y \in [S]$. Fix $k \geq |\tau|$. By construction $y \in [S_{y \upharpoonright k}] \subseteq f^{-1}([x_\tau \upharpoonright k])$. Since this holds for all k , $f(y) = x_\tau$, and we are done.

Case 2. $\rho(\tau)$ is defined for all $\tau \in T'$.

Recall that $F \subseteq T'$ is a *front* if for all $y \in [T']$ there is a unique n with $y \upharpoonright n \in F$. We build a subtree S of T' by specifying fronts F_n , $n \in \omega$, such that for every $\sigma \in F_{n+1}$ there is a (necessarily unique) $\tau \in F_n$ with $\tau \subset \sigma$. That is, S will be the tree generated by the fronts: $\sigma \in S$ iff there are $n \in \omega$ and $\tau \in F_n$ with $\sigma \subseteq \tau$. Additionally, we shall guarantee that there are $s_\tau \subseteq x_\tau$ for $\tau \in \bigcup_n F_n$ such that

- if $\sigma \neq \sigma'$ both are in F_n then $[s_\sigma] \cap [s_{\sigma'}] = \emptyset$,
- $[S_\tau] \subseteq f^{-1}([s_\tau])$ for $\tau \in \bigcup_n F_n$,
- if $\sigma \subset \tau$ with $\sigma \in F_n$ and $\tau \in F_{n+1}$ then $s_\sigma \subset s_\tau$,
- if $\sigma \subset \tau$ with $\sigma \in F_n$ and $\tau \in F_{n+1}$ then for every k with $|\sigma| \leq k < |\tau|$, $x_{\tau \upharpoonright k} = x_\sigma$ and $\rho(\sigma) > \rho(\tau \upharpoonright |\sigma| + 1) > \dots > \rho(\tau \upharpoonright |\tau| - 1) = 0$.

We first verify that this is enough to guarantee that $f \upharpoonright [S]$ is one-to-one. If $y, y' \in [S]$ are distinct, then there are $n, i, i' \in \omega$ such that $y \upharpoonright i$ and $y' \upharpoonright i'$ are distinct elements of F_n . Then $y \in f^{-1}([s_{y \upharpoonright i}])$, $y' \in f^{-1}([s_{y' \upharpoonright i'}])$ by the second clause, and $[s_{y \upharpoonright i}]$ and $[s_{y' \upharpoonright i'}]$ are disjoint by the first clause. Hence $f(y) \neq f(y')$ as required. Thus it suffices to construct the F_n and s_τ .

$n = 0$. We let $F_0 = \{\text{stem}(T')\} = \{\text{stem}(S)\}$. Also let $s_{\text{stem}(S)} = x_{\text{stem}(S)} \upharpoonright \text{stem}(S)$.

Suppose F_n and s_σ for $\sigma \in F_n$ have been constructed. We shall construct F_{n+1} , s_σ for $\sigma \in F_{n+1}$, as well as the part of the tree S in between F_n and F_{n+1} . Fix $\sigma \in F_n$. By A_σ^n we denote the part of S between σ and F_{n+1} , that is, $A_\sigma^n = \{\tau \in S : \sigma \subseteq \tau \text{ and } \tau \subset v \text{ for some } v \text{ in } F_{n+1}\}$. A_σ^n will be constructed recursively so as to satisfy the fourth clause above.

Put σ into A_σ^n . Suppose some $\tau \supseteq \sigma$ has been put into A_σ^n , $x_\tau = x_\sigma$ and, in case $\tau \supset \sigma$, $\rho(\sigma) > \rho(\tau)$. In case $\rho(\tau) = 0$, no successor of τ will be in A_σ^n and the successors of τ will belong to F_{n+1} , as explained below. If $\rho(\tau) > 0$, then $x_{\tau \hat{\ } k} = x_\tau$ for almost all $k \in \text{succ}_{T'}(\tau)$ and $\rho(\tau \hat{\ } k) < \rho(\tau)$ for infinitely many $k \in \text{succ}_{T'}(\tau)$. Hence we can prune the successor level of τ to $\text{succ}_S(\tau)$ such that $x_{\tau \hat{\ } k} = x_\tau$ and $\rho(\tau \hat{\ } k) < \rho(\tau)$ for all $k \in \text{succ}_S(\tau)$. The fourth clause is clearly satisfied. This completes the construction of A_σ^n .

Now fix $\tau \in A_\sigma^n$ with $\rho(\tau) = 0$. By pruning $\text{succ}_{T'}(\tau)$, if necessary, we may assume without loss of generality that the $x_{\tau \hat{\ } k^m}$, $m \in \omega$, are all pairwise distinct and converge to $x_\tau = x_\sigma$ and that, in fact, there is a strictly increasing sequence $(i_\tau^m : m \in \omega)$ such that $i_\tau^m = \min\{i : x_{\tau \hat{\ } k^m}(i) \neq x_\tau(i)\}$. Unfixing τ , we may additionally assume that if $\tau \neq \tau'$ are both in A_σ^n of rank 0 and $m, m' \in \omega$, then $i_\tau^m \neq i_{\tau'}^{m'}$. Finally we may assume that all such i_τ^m are larger than $|s_\sigma|$. This means in particular that $s_\sigma \subseteq x_{\tau \hat{\ } k^m}$ for all τ and m because $s_\sigma \subseteq x_\sigma = x_\tau$. Now choose $s_{\tau \hat{\ } k^m} \subseteq x_{\tau \hat{\ } k^m}$ such that $|s_{\tau \hat{\ } k^m}| > i_\tau^m$. Then $s_\sigma \subset s_{\tau \hat{\ } k^m}$ and the $s_{\tau \hat{\ } k^m}$ for distinct pairs (τ, m) with $\tau \in A_\sigma^n$ of rank 0 and $m \in \omega$ are pairwise incompatible.

Unfix $\sigma \in F_n$. Let $F_{n+1} = \{\tau \hat{\ } k^m : \tau \in A_\sigma^n \text{ for some } \sigma \in F_n, \rho(\tau) = 0, \text{ and } m \in \omega\}$. The third clause is immediate. To see the first clause, take distinct $\tau, \tau' \in F_{n+1}$. There are $\sigma, \sigma' \in F_n$ such that $\sigma \subset \tau$ and $\sigma' \subset \tau'$. If $\sigma \neq \sigma'$, then $[s_\tau] \cap [s_{\tau'}] = \emptyset$ because $[s_\sigma] \cap [s_{\sigma'}] = \emptyset$ and $s_\sigma \subset s_\tau$ and $s_{\sigma'} \subset s_{\tau'}$. If $\sigma = \sigma'$, then $[s_\tau] \cap [s_{\tau'}] = \emptyset$ by the construction in the previous paragraph. Finally, to see the second clause, by pruning T'_τ for $\tau \in F_{n+1}$ if necessary, we may assume $[T'_\tau] \subseteq f^{-1}([s_\tau])$ (see Claim 16.3). This completes the recursive construction and the proof of the theorem. \square

Proposition 17. *Assume \mathbb{T} is a combinatorial tree forcing with the constant or one-to-one property. Then \mathbb{T} has the selective disjoint antichain property.*

Proof. Let $f : \omega^\omega \rightarrow 2^\omega$ be a continuous function witnessing “large disjoint antichains” of \mathbb{T} . Let $\{x_\alpha : \alpha < \mathfrak{c}\}$ be an enumeration of 2^ω . Let $T_\alpha \in \mathbb{T}$ be such that $[T_\alpha] = f^{-1}(\{x_\alpha\})$. We check that $(T_\alpha : \alpha < \mathfrak{c})$ witnesses the selective disjoint antichain property. Clearly $[T_\alpha] \cap [T_\beta] = \emptyset$ for $\alpha \neq \beta$. Given $T \in \mathbb{T}$, find $S \leq T$ such that $f \upharpoonright [S]$ is constant or one-to-one. In the first case, $S \leq T_\alpha$ for some α , and in the second case, $|[S] \cap [T_\alpha]| \leq 1$ for all α , and we are done. \square

We are finally ready to complete the proof of the main result of this note.

Corollary 18. $cf(\text{cof}(\ell^0)) > \mathfrak{c}$ and $cf(\text{cof}(m^0)) > \mathfrak{c}$.

Proof. This follows from Theorem 16, Theorem 15, Proposition 17, and Theorem 13. \square

4 Problems

For some natural tree forcings, we still do not know whether the cofinality of the corresponding tree ideal is larger than \mathfrak{c} in ZFC. Let \mathbb{FM} denote *full splitting Miller forcing*, originally introduced by [NR] (see also [KL]), that is, conditions are Miller trees $T \subseteq \omega^{<\omega}$ such that whenever $s \in T$ is a splitting node, then $s \hat{\ } n \in T$ for all $n \in \omega$. fm^0 is the *full splitting Miller ideal*.

Question 19. *Is $\text{cof}(fm^0) > \mathfrak{c}$?*

By the discussion in Section 2 (before Proposition 9), we know this is true under CH.

More generally, one may ask:

Question 20. *Are there combinatorial tree forcings \mathbb{T} which consistently fail to have the disjoint maximal antichain property? Which consistently fail to satisfy $\text{cof}(t^0) > \mathfrak{c}$? For which t^0 consistently has a Borel basis?*

Note that the existence of a Borel basis implies $\text{cof}(t^0) = \mathfrak{c}$. By the above comment fm^0 has no Borel basis under CH, but this is open in ZFC. Question 20 is also of interest for tree forcings which do not necessarily satisfy all the clauses of Definition 1, e.g., for non-homogeneous forcing notions.

By [JMS, Theorems 1.4 and 1.5], we know that $\text{cof}(s^0)$ can consistently assume arbitrary values $\leq 2^{\mathfrak{c}}$ whose cofinality is larger than \mathfrak{c} and it is easy to see that the same arguments work for other tree ideals like m^0 and ℓ^0 . (In these models CH holds.)

Question 21. *Can we consistently separate the cofinalities of different tree ideals? E.g., are $\text{cof}(s^0) < \text{cof}(m^0)$ or $\text{cof}(m^0) < \text{cof}(s^0)$ consistent?*

References

- [Br] J. Brendle, *Strolling through paradise*, Fund. Math. **148** (1995), 1-25.
- [GRSS] M. Goldstern, M. Repický, S. Shelah, and O. Spinas, *On tree ideals*, Proc. Amer. Math. Soc. **123** (1995), 1573-1581.
- [Gra] C. Gray, *Iterated forcing from the strategic point of view*, Ph.D. thesis, University of California, Berkeley, California, 1980.
- [Gro] M. Groszek, *Combinatorics on ideals and forcing with trees*, J. Symbolic Logic **52** (1987), 582-593.

- [JMS] H. Judah, A. Miller, and S. Shelah, *Sacks forcing, Laver forcing, and Martin's axiom*, Arch. Math. Logic **31** (1992), 145-161.
- [KL] Y. Khomskii and G. Laguzzi, *Full-splitting Miller trees and infinitely often equal reals*, preprint.
- [La] R. Laver, *On the consistency of Borel's conjecture*, Acta Math. **137** (1976), 151-169.
- [Ma] P. Matet, unpublished.
- [Mi] A. Miller, *Rational perfect set forcing*, Contemporary Mathematics **31** (1984), 143-159.
- [NR] L. Newelski and A. Roslanowski, *The ideal determined by the unsymmetric game*, Proc. Amer. Math. Soc. **117** (1993), 823-831.
- [Re] M. Repický, *Cofinality of the Laver ideal*, preprint.