

HYPERGRAPHS WITH VANISHING TURÁN DENSITY IN UNIFORMLY DENSE HYPERGRAPHS

CHRISTIAN REIHER, VOJTĚCH RÖDL, AND MATHIAS SCHACHT

ABSTRACT. P. Erdős [*On extremal problems of graphs and generalized graphs*, Israel Journal of Mathematics **2** (1964), 183–190] characterised those hypergraphs F that have to appear in any sufficiently large hypergraph H of positive density. We study related questions for 3-uniform hypergraphs with the additional assumption that H has to be *uniformly dense* with respect to vertex sets. In particular, we characterise those hypergraphs F that are guaranteed to appear in large uniformly dense hypergraphs H of positive density. We also review the case when the density of the induced subhypergraphs of H may depend on the proportion of the considered vertex sets.

§1. INTRODUCTION

Unless said otherwise, all hypergraphs considered here are 3-uniform. For such a hypergraph $H = (V, E)$ the set of vertices is denoted by $V = V(H)$ and we refer to the set of hyperedges by $E = E(H)$. Moreover, we denote by $\partial H \subseteq V^{(2)}$ the subset of all two element subsets of V , that contains all pairs covered by some hyperedge $e \in E$. For a hyperedge $\{x, y, z\} \in E$ we sometimes simply write $xyz \in E$.

A classical extremal problem introduced by Turán [17] asks to study for a given hypergraph F its extremal function $\text{ex}(n, F)$ sending each positive integer to the maximum number of edges that a hypergraph of order n can have without containing F as a subhypergraph. In particular, one often focuses on the *Turán density* $\pi(F)$ of F defined by

$$\pi(F) = \lim_{n \rightarrow \infty} \frac{\text{ex}(n, F)}{\binom{n}{3}}.$$

The problem to determine the Turán densities of all hypergraphs is known to be very hard and so far it has been solved for a few hypergraphs only. A general result in this area due to Erdős [1] asserts that a hypergraph F satisfies $\pi(F) = 0$ if and only if it is tripartite in the sense that there is a partition $V(F) = X \cup Y \cup Z$ such that every edge of F contains precisely one vertex from each of X , Y , and Z .

2010 *Mathematics Subject Classification.* 05C35 (primary), 05C65, 05C80 (secondary).

Key words and phrases. quasirandom hypergraphs, extremal graph theory, Turán's problem.

The second author is supported by NSF grant DMS 1301698.

Following a suggestion by Erdős and Sós [3] we studied variants of Turán’s problem for *uniformly dense* hypergraphs [10–13]. Instead of finding the desired hypergraph F in an arbitrary “host” hypergraph H of sufficiently large density one assumes in these problems that there are no “sparse spots” in the edge distribution of H . There are various ways to make this precise and we refer to [11, Section 4] and [13, Section 2] for a more detailed discussion. Here we consider two closely related concepts, where the hereditary density condition pertains to large sets of vertices (see Sections 1.1 and 1.2 below).

1.1. Uniformly dense hypergraphs with positive density. The first concept we discuss here continues our work from [10–13]. Roughly speaking, this notion guarantees density d for all hypergraphs induced on sufficiently large vertex sets of linear size.

Definition 1.1. For real numbers $d \in [0, 1]$ and $\eta > 0$ we say that a hypergraph $H = (V, E)$ is $(d, \eta, 1)$ -dense if for all $U \subseteq V$ the estimate

$$|U^{(3)} \cap E| \geq d \binom{|U|}{3} - \eta |V|^3$$

holds, where $U^{(3)}$ denotes the set of all three element subsets of U .

The Turán densities associated with this concept are defined by

$$\pi_1(F) = \sup\{d \in [0, 1] : \text{for every } \eta > 0 \text{ and } n \in \mathbb{N} \text{ there exists} \\ \text{an } F\text{-free, } (d, \eta, 1)\text{-dense hypergraph } H \text{ with } |V(H)| \geq n\}.$$

Our main result characterises all hypergraphs F with $\pi_1(F) = 0$.

Theorem 1.2. *For a 3-uniform hypergraph F , the following are equivalent:*

- (a) $\pi_1(F) = 0$.
- (b) *There is an enumeration of the vertex set $V(F) = \{v_1, \dots, v_f\}$ and there is a three-colouring $\varphi: \partial F \rightarrow \{\text{red, blue, green}\}$ of the pairs of vertices covered by hyperedges of F such that every hyperedge $\{v_i, v_j, v_k\} \in E(F)$ with $i < j < k$ satisfies*

$$\varphi(v_i, v_j) = \text{red}, \quad \varphi(v_i, v_k) = \text{blue}, \quad \text{and} \quad \varphi(v_j, v_k) = \text{green}.$$

It is easy to see that tripartite hypergraphs F satisfy condition (b). Moreover, it follows from the work in [8] that every linear hypergraph F satisfies $\pi_1(F) = 0$. Linear hypergraphs have the property that every element of ∂F is contained in precisely one hyperedge of F . Consequently, we may consider an arbitrary vertex enumeration of F and then a colouring of ∂F satisfying condition (b) is forced. However, there are hypergraphs displaying condition (b), that are neither tripartite nor linear. For example, one can




FIGURE 1.1. Colouring of $\partial C_5^{(3)-}$ showing that $\pi_1(C_5^{(3)-}) = 0$. The ordering demanded by Theorem 1.2 (b) is from left to right, i.e., $x < w < v < z < y$, whereas on the cycle the vertices are ordered alphabetically with edges vw, wx, xy, yz, zv .

check that the hypergraph obtained from the tight cycle on five vertices by removing one hyperedge is such a hypergraph F (see Figure 1.1).

The easier implication of Theorem 1.2 is “(a) \implies (b).” For its proof we exhibit a “universal” hypergraph H all of whose subhypergraphs obey condition (b) and all of whose linear sized induced subhypergraphs have density $\frac{1}{27} - o(1)$. In other words, our argument establishing this implication does actually yield the following strengthening.

Fact 1.3. *If a hypergraph F does not have property (b) from Theorem 1.2, then $\pi_1(F) \geq \frac{1}{27}$.*

Proof. Given a positive integer n consider a three-colouring $\varphi: [n]^{(2)} \rightarrow \{\text{red, blue, green}\}$ of the pairs of the first n positive integers. We define a hypergraph H_φ with vertex set $[n]$ by regarding a triple $\{i, j, k\}$ with $1 \leq i < j < k \leq n$ as being a hyperedge if and only if $\varphi(i, j) = \text{red}$, $\varphi(i, k) = \text{blue}$, and $\varphi(j, k) = \text{green}$. Standard probabilistic arguments show that when φ is chosen uniformly at random, then for any fixed $\eta > 0$ the probability that H_φ is $(1/27, \eta, 1)$ -dense tends to 1 as n tends to infinity. On the other hand, as F does not satisfy condition (b) from Theorem 1.2, it is in a deterministic sense the case that F is never a subgraph of H_φ no matter how large n becomes. Thus we have indeed $\pi_1(F) \geq \frac{1}{27}$. \square

The combination of Theorem 1.2 and Fact 1.3 leads immediately to the following consequence, which shows that π_1 “jumps” from 0 to at least $\frac{1}{27}$.

Corollary 1.4. *If a hypergraph F satisfies $\pi_1(F) > 0$, then $\pi_1(F) \geq \frac{1}{27}$.*

1.2. Uniformly dense hypergraphs with vanishing density. The second concept we discuss here is closely related to the one from Definition 1.1. It was introduced by Erdős and Sós in [3] (see also [2, page 24]). To prepare its definition we need a concept of being

d -dense when d can be a function rather than just a single number and we shall consider *sequences* of hypergraphs instead of just one individual hypergraph.

Definition 1.5. (a) Let $\vec{H} = (H_n)_{n \in \mathbb{N}}$ be a sequence of hypergraphs with $|V(H_n)| \rightarrow \infty$ as $n \rightarrow \infty$ and let $d: (0, 1) \rightarrow (0, 1)$ be a function. We say that \vec{H} is d -dense provided that for every $\eta \in (0, 1)$ there is an $n_0 \in \mathbb{N}$ such that for $n \geq n_0$ every $U \subseteq V(H_n)$ with $|U| \geq \eta |V(H_n)|$ satisfies

$$|U^{(3)} \cap E(H_n)| \geq d(\eta) \binom{|U|}{3}.$$

(b) A hypergraph F is called *frequent* if for every function $d: (0, 1) \rightarrow (0, 1)$ and every d -dense sequence $\vec{H} = (H_n)_{n \in \mathbb{N}}$ of hypergraphs there is an integer n_0 such that F is a subhypergraph of every H_n with $n \geq n_0$.

Erdős and Sós [3, Proposition 3] described the following instructive example $(T_n)_{n \in \mathbb{N}}$ of a sequence of *ternary hypergraphs* that is d -dense for some function $d(\cdot)$, but not uniformly dense in the sense of Definition 1.1. Take the vertex set of T_n to be the set $\{0, 1, 2\}^n$ of all sequences with length n all of whose entries are 0, 1, or 2. Given three distinct vertices of T_n , say $\vec{x} = (x_1, \dots, x_n)$, $\vec{y} = (y_1, \dots, y_n)$, and $\vec{z} = (z_1, \dots, z_n)$ there is a least integer $i \in [n]$ for which $x_i = y_i = z_i$ is not the case and we put a hyperedge $\{\vec{x}, \vec{y}, \vec{z}\}$ into $E(T_n)$ if and only if this index i satisfies $\{x_i, y_i, z_i\} = \{0, 1, 2\}$. It was stated in [3] that the sequence of ternary hypergraphs is d -dense for some appropriate function $d(\cdot)$ and a short proof of this fact appeared in [4]. In Section 5 we obtain the following improvement.

Proposition 1.6. *The sequence of ternary hypergraphs $(T_n)_{n \in \mathbb{N}}$ is d -dense for any function $d: (0, 1] \rightarrow (0, 1]$ with $d(\eta) < \frac{1}{4} \eta^{\frac{2}{\log_2(3)-1}}$.*

Considering subsets $U \subseteq V(T_n)$ of the form $U = \{0, 1\}^r \times \{0, 1, 2\}^{n-r}$ shows that Proposition 1.6 is optimal whenever $\eta = (2/3)^r$ for some $r \in \mathbb{N}$. Since ternary hypergraphs are d -dense for some function $d(\cdot)$, it follows that every frequent hypergraph must be contained in some ternary hypergraph and Erdős wondered in [2] whether the converse of this holds as well. This was indeed verified by Frankl and Rödl in [4] and the following characterisation can be viewed as an analogue of Theorem 1.2 for d -dense hypergraphs.

Theorem 1.7. *A hypergraph F is frequent if, and only if it occurs as a subhypergraph of a ternary hypergraph.* □

It is not hard to show (see Lemma 5.3) that if F is a subhypergraph of some ternary hypergraph, then $F \subseteq T_{|V(F)|}$ and, consequently, Theorem 1.7 entails, that it is decidable whether a given hypergraph is frequent or not.

Organisation. The proof of the implication “(b) \implies (a)” of Theorem 1.2 utilises the hypergraph regularity method that is revisited in Section 2. This method allows us in Section 3 to reduce the problem of embedding hypergraphs satisfying the condition (b) in Theorem 1.2 into uniformly dense hypergraphs to a problem concerning so-called *reduced hypergraphs*. This reduction will be carried out in Section 3 and the main argument will then be given in Section 4. In Section 5 we prove Proposition 1.6, which implies the forward implication of Theorem 1.7.

For a more complete presentation we include a short proof of the backward implication of Theorem 1.7 as well, which follows the lines of the proof in [4]. In contrast to the proof of the implication “(b) \implies (a)” of Theorem 1.2 this proof is somewhat simpler and is based on a supersaturation argument. Extensions of our results to k -uniform hypergraphs with $k > 3$ will be discussed in the concluding remarks.

§2. HYPERGRAPH REGULARITY

A key tool in the proof of Theorem 1.2 is the regularity lemma for 3-uniform hypergraphs. We follow the approach from [15, 16] combined with the results from [7] and [9].

For two disjoint sets X and Y we denote by $K(X, Y)$ the complete bipartite graph with that vertex partition. We say that a bipartite graph $P = (X \cup Y, E)$ is (δ_2, d_2) -regular if for all subsets $X' \subseteq X$ and $Y' \subseteq Y$ we have

$$|e(X', Y') - d_2|X'||Y'|| \leq \delta_2|X||Y|,$$

where $e(X', Y')$ denotes the number of edges of P with one vertex in X' and one vertex in Y' . Moreover, for $k \geq 2$ we say a k -partite graph $P = (X_1 \cup \dots \cup X_k, E)$ is (δ_2, d_2) -regular, if all its $\binom{k}{2}$ naturally induced bipartite subgraphs $P[X_i, X_j]$ are (δ_2, d_2) -regular. For a tripartite graph $P = (X \cup Y \cup Z, E)$ we denote by $\mathcal{K}_3(P)$ the triples of vertices spanning a triangle in P , i.e.,

$$\mathcal{K}_3(P) = \{\{x, y, z\} \subseteq X \cup Y \cup Z : xy, xz, yz \in E\}.$$

If the tripartite graph P is (δ_2, d_2) -regular, then the *triangle counting lemma* implies

$$|\mathcal{K}_3(P)| \leq d_2^3|X||Y||Z| + 3\delta_2|X||Y||Z|. \tag{2.1}$$

We say a 3-uniform hypergraph $H = (V, E_H)$ is regular w.r.t. a tripartite graph P if it matches approximately the same proportion of triangles for every subgraph $Q \subseteq P$.

Definition 2.1. A 3-uniform hypergraph $H = (V, E_H)$ is (δ_3, d_3) -regular w.r.t. a tripartite graph $P = (X \cup Y \cup Z, E_P)$ with $V \supseteq X \cup Y \cup Z$ if for every tripartite subgraph $Q \subseteq P$ we have

$$||E_H \cap \mathcal{K}_3(Q)| - d_3|\mathcal{K}_3(Q)|| \leq \delta_3|\mathcal{K}_3(P)|.$$

Moreover, we simply say H is δ_3 -regular w.r.t. P , if it is (δ_3, d_3) -regular for some $d_3 \geq 0$. We also define the *relative density* of H w.r.t. P by

$$d(H|P) = \frac{|E_H \cap \mathcal{K}_3(P)|}{|\mathcal{K}_3(P)|},$$

where we use the convention $d(H|P) = 0$ if $\mathcal{K}_3(P) = \emptyset$. If H is not δ_3 -regular w.r.t. P , then we simply refer to it as δ_3 -irregular.

The regularity lemma for 3-uniform hypergraphs, introduced by Frankl and Rödl in [5], provides for a hypergraph H a partition of its vertex set and a partition of the edge sets of the complete bipartite graphs induced by the vertex partition such that for appropriate constants δ_3 , δ_2 , and d_2

- (1) the bipartite graphs given by the partitions are (δ_2, d_2) -regular and
- (2) H is δ_3 -regular for “most” tripartite graphs P given by the partition.

In many proofs based on the regularity method it is convenient to “clean” the regular partition provided by the lemma. In particular, we shall disregard hyperedges of H that belong to $\mathcal{K}_3(P)$ where H is not δ_3 -regular or where $d(H|P)$ is very small. These properties are rendered in the following somewhat standard corollary of the regularity lemma.

Theorem 2.2. *For every $d_3 > 0$, $\delta_3 > 0$ and $m \in \mathbb{N}$, and every function $\delta_2: \mathbb{N} \rightarrow (0, 1]$, there exist integers T_0 and n_0 such that for every $n \geq n_0$ and every n -vertex 3-uniform hypergraph $H = (V, E)$ the following holds.*

There exists a subhypergraph $\hat{H} = (\hat{V}, \hat{E}) \subseteq H$, an integer $\ell \leq T_0$, a vertex partition $V_1 \cup \dots \cup V_m = \hat{V}$, and for all integers i, j with $1 \leq i < j \leq m$ there exists a partition $\mathcal{P}^{ij} = \{P_\alpha^{ij} = (V_i \cup V_j, E_\alpha^{ij}): 1 \leq \alpha \leq \ell\}$ of $K(V_i, V_j)$ satisfying the following properties

- (i) $|V_1| = \dots = |V_m| \geq (1 - \delta_3)n/T_0$,
- (ii) for every $1 \leq i < j \leq m$ and $\alpha \in [\ell]$ the bipartite graph P_α^{ij} is $(\delta_2(\ell), 1/\ell)$ -regular,
- (iii) \hat{H} is δ_3 -regular w.r.t. all triads

$$P_{\alpha\beta\gamma}^{ijk} = P_\alpha^{ij} \cup P_\beta^{ik} \cup P_\gamma^{jk} = (V_i \cup V_j \cup V_k, E_\alpha^{ij} \cup E_\beta^{ik} \cup E_\gamma^{jk}), \quad (2.2)$$

- with $1 \leq i < j < k \leq m$ and $\alpha, \beta, \gamma \in [\ell]$, and $d(\hat{H}|P_{\alpha\beta\gamma}^{ijk})$ is either 0 or at least d_3 ,*
- (iv) and for every $1 \leq i < j < k \leq m$ we have

$$e_{\hat{H}}(V_i, V_j, V_k) \geq e_H(V_i, V_j, V_k) - (d_3 + \delta_3)|V_i||V_j||V_k|. \quad \square$$

The standard proof of Theorem 2.2 based on a refined version of the regularity lemma from [15, Theorem 2.3] can be found in [10, Corollary 3.3].

We shall use the *counting/embedding lemma*, which allows us to embed hypergraphs of fixed isomorphism type into appropriate and sufficiently regular and dense triads of the partition provided by Theorem 2.2. It is a direct consequence of [9, Corollary 2.3].

Theorem 2.3 (Embedding Lemma). *Let a hypergraph F with vertex set $[f]$ and $d_3 > 0$ be given. Then there exist $\delta_3 > 0$ and functions $\delta_2: \mathbb{N} \rightarrow (0, 1]$ and $N: \mathbb{N} \rightarrow \mathbb{N}$ such that the following holds for every $\ell \in \mathbb{N}$.*

Suppose $P = (V_1 \cup \dots \cup V_f, E_P)$ is a $(\delta_2(\ell), \frac{1}{\ell})$ -regular, f -partite graph whose vertex classes satisfy $|V_1| = \dots = |V_f| \geq N(\ell)$ and suppose H is an f -partite, 3-uniform hypergraph such that for all edges ijk of F we have

- (a) H is δ_3 -regular w.r.t. to the tripartite graph $P[V_i \cup V_j \cup V_k]$ and
- (b) $d(H|P[V_i \cup V_j \cup V_k]) \geq d_3$,

then H contains a copy of F . In fact, there is a monomorphism q from F to H with $q(i) \in V_i$ for all $i \in [f]$. \square

In an application of Theorem 2.3 the tripartite graphs $P[V_i \cup V_j \cup V_k]$ in (a) and (b) will be given by triads $P_{\alpha\beta\gamma}^{ijk}$ from the partition given by Theorem 2.2. For the proof of the direction “(b) \implies (a)” of Theorem 1.2 we consider for a fixed hypergraph F obeying condition (b) and fixed $\varepsilon > 0$ a sufficiently large uniformly dense hypergraph H of density ε . We will apply the regularity lemma in the form of Theorem 2.2 to H . The main part of the proof concerns the appropriate selection of dense and regular triads, that are ready for an application of the embedding lemma. In Section 3 we formulate a statement about reduced hypergraphs telling us that such a selection is indeed possible and in Section 4 we give its proof.

§3. MOVING TO REDUCED HYPERGRAPHS

In our intended application of the hypergraph regularity method we need to keep track which triads are dense and regular and natural structures for encoding such information are so-called reduced hypergraphs. We follow the terminology introduced in [12, Section 3].

Consider any finite set of indices I , suppose that associated with any two distinct indices $i, j \in I$ we have a finite nonempty set of vertices \mathcal{P}^{ij} , and that for distinct pairs of indices the corresponding vertex classes are disjoint. Assume further that for any three distinct indices $i, j, k \in I$ we are given a tripartite hypergraph \mathcal{A}^{ijk} with vertex classes \mathcal{P}^{ij} , \mathcal{P}^{ik} , and \mathcal{P}^{jk} . Under such circumstances we call the $\binom{|I|}{2}$ -partite hypergraph \mathcal{A} defined by

$$V(\mathcal{A}) = \bigcup_{\{i,j\} \in I^{(2)}} \mathcal{P}^{ij} \quad \text{and} \quad E(\mathcal{A}) = \bigcup_{\{i,j,k\} \in I^{(3)}} E(\mathcal{A}^{ijk})$$

a *reduced hypergraph*. We also refer to I as the *index set* of \mathcal{A} , to the sets \mathcal{P}^{ij} as the *vertex classes* of \mathcal{A} , and to the hypergraphs \mathcal{A}^{ijk} as the *constituents* of \mathcal{A} . The order of the indices appearing in the pairs and triples of the superscripts of the vertex classes and constituents

of \mathcal{A} plays no rôle here, i.e., $\mathcal{P}^{ij} = \mathcal{P}^{ji}$ and $\mathcal{A}^{ijk} = \mathcal{A}^{kij}$ etc. For $\mu > 0$ such a reduced hypergraph \mathcal{A} is said to be μ -dense if

$$|E(\mathcal{A}^{ijk})| \geq \mu |\mathcal{P}^{ij}| |\mathcal{P}^{ik}| |\mathcal{P}^{jk}|$$

holds for every triple $\{i, j, k\} \in I^{(3)}$.

In the light of the hypergraph regularity method, the proof of Theorem 1.2 reduces to the following statement whose proof will be given in the next section.

Lemma 3.1. *Given $\mu > 0$ and $f \in \mathbb{N}$ there exists an integers m such that the following holds. If \mathcal{A} is a μ -dense reduced hypergraph with index set $[m]$, vertex classes \mathcal{P}^{ij} , and constituents \mathcal{A}^{ijk} , then there are*

- (i) indices $\lambda(1) < \dots < \lambda(f)$ in $[m]$ and
- (ii) for each pair $1 \leq r < s \leq f$ there are three vertices $P_{\text{red}}^{\lambda(r)\lambda(s)}$, $P_{\text{blue}}^{\lambda(r)\lambda(s)}$, and $P_{\text{green}}^{\lambda(r)\lambda(s)}$ in $\mathcal{P}^{\lambda(r)\lambda(s)}$

such that for every triple of indices $1 \leq r < s < t \leq m$ the three vertices $P_{\text{red}}^{\lambda(r)\lambda(s)}$, $P_{\text{blue}}^{\lambda(r)\lambda(t)}$, and $P_{\text{green}}^{\lambda(s)\lambda(t)}$ form a hyperedge in $\mathcal{A}^{\lambda(r)\lambda(s)\lambda(t)}$.

At the end of this section we will prove that this lemma does indeed imply Theorem 1.2. For this purpose it will be more convenient to work with an alternative definition of π_1 that we denote by π_{\bullet} . In contrast to Definition 1.1 it speaks about being dense with respect to three subsets of vertices rather than just one.

Definition 3.2. A hypergraph $H = (V, E)$ of order $n = |V|$ is (d, η, \bullet) -dense if for every triple of subsets $X, Y, Z \subseteq V$ the number $e_{\bullet}(X, Y, Z)$ of triples $(x, y, z) \in X \times Y \times Z$ with $xyz \in E$ satisfies

$$e_{\bullet}(X, Y, Z) \geq d |X| |Y| |Z| - \eta n^3.$$

Accordingly, we set

$$\pi_{\bullet}(F) = \sup\{d \in [0, 1] : \text{for every } \eta > 0 \text{ and } n \in \mathbb{N} \text{ there exists} \\ \text{an } F\text{-free, } (d, \eta, \bullet)\text{-dense hypergraph } H \text{ with } |V(H)| \geq n\}. \quad (3.1)$$

Applying [13, Proposition 2.5] to $k = 3$ and $j = 1$ we deduce that every hypergraph F satisfies

$$\pi_{\bullet}(F) = \pi_1(F). \quad (3.2)$$

Consequently it is allowed to imagine that in clause (a) of Theorem 1.2 we would have written $\pi_{\bullet}(F) = 0$ instead of $\pi_1(F) = 0$.

Proof of Theorem 1.2 assuming Lemma 3.1. The implication “(a) \implies (b)” is implicit in Fact 1.3, meaning that we just need to consider the reverse direction. Suppose to this end that a hypergraph F satisfying condition (b) and some $\varepsilon > 0$ are given. We need to check that for $\varepsilon \gg \eta \gg n^{-1}$ every $(\varepsilon, \eta, \bullet)$ -dense hypergraph H of order n contains a copy of F .

Of course, we may assume that $V(F) = [f]$ holds for some $f \in \mathbb{N}$. Plugging F and $d_3 = \frac{\varepsilon}{4}$ into the embedding lemma we get a constant $\delta_3 > 0$, a function $\delta_2: \mathbb{N} \rightarrow (0, 1]$, and a function $N: \mathbb{N} \rightarrow \mathbb{N}$. Evidently we may assume that $\delta_3 \leq \frac{\varepsilon}{4}$, that $\delta_2(\ell) \ll \ell^{-1}$, and that N is increasing. Applying Lemma 3.1 with $\mu = \frac{\varepsilon}{8}$ and f we obtain an integer m . Given d_3 , δ_3 , m , and $\delta_2(\cdot)$ we get integers T_0 and n_0 from Theorem 2.2. Finally we choose

$$\eta = \frac{\varepsilon(1 - \delta_3)^3}{4T_0^3} \quad \text{and} \quad n_1 = 2T_0 \cdot N(T_0).$$

Now consider any $(\varepsilon, \eta, \bullet)$ -dense hypergraph H of order $n \geq n_1$. We contend that F appears as a subhypergraph of H . To see this we take

- a subhypergraph $\hat{H} = (\hat{V}, \hat{E}) \subseteq H$,
- a vertex partition $V_1 \cup \dots \cup V_m = \hat{V}$,
- an integer $\ell \leq T_0$,
- and pair partitions $\mathcal{P}^{ij} = \{P_\alpha^{ij} = (V_i \cup V_j, E_\alpha^{ij}): 1 \leq \alpha \leq \ell\}$ of $K(V_i, V_j)$ for all $1 \leq i < j \leq m$

satisfying the conditions (i)–(iv) from Theorem 2.2. The reduced hypergraph \mathcal{A} corresponding to this situation has index set $[m]$, vertex classes \mathcal{P}^{ij} and a triple $\{P_\alpha^{ij}, P_\beta^{ik}, P_\gamma^{jk}\}$ is defined to be an edge of the constituent \mathcal{A}^{ijk} if and only if $d(\hat{H}|P_{\alpha\beta\gamma}^{ijk}) \geq d_3$. As we shall verify below,

$$\mathcal{A} \text{ is } \mu\text{-dense.} \tag{3.3}$$

Due to Lemma 3.1 this means that there are

- indices $\lambda(1) < \dots < \lambda(f)$ in $[m]$ and
- for each pair $1 \leq r < s \leq f$ there are vertices $P_{\text{red}}^{\lambda(r)\lambda(s)}, P_{\text{blue}}^{\lambda(r)\lambda(s)}, P_{\text{green}}^{\lambda(r)\lambda(s)} \in \mathcal{P}^{\lambda(r)\lambda(s)}$

such that for every triple of indices $1 \leq r < s < t \leq m$ the three vertices $P_{\text{red}}^{\lambda(r)\lambda(s)}, P_{\text{blue}}^{\lambda(r)\lambda(t)}$, and $P_{\text{green}}^{\lambda(s)\lambda(t)}$ form a hyperedge in $\mathcal{A}^{\lambda(r)\lambda(s)\lambda(t)}$. These vertices are, of course, actually bipartite graphs forming dense regular triads. Since we have

$$|V_{\lambda(1)}| = \dots = |V_{\lambda(f)}| \geq \frac{(1 - \delta_3)n}{T_0} \geq \frac{n_1}{2T_0} = N(T_0) \geq N(\ell),$$

the embedding lemma is applicable to the hypergraph \hat{H} and to the f -partite graph with vertex partition $\bigcup_{r \in [f]} V_{\lambda(r)}$ and edge set $\bigcup_{rs \in \partial F} P_{\varphi(\lambda(r), \lambda(s))}^{\lambda(r)\lambda(s)}$, where $\varphi: \partial F \rightarrow \{\text{red, blue, green}\}$

denotes any colouring exemplifying that F does indeed possess property (b) from Theorem 1.2. Consequently, the monomorphism guaranteed by Theorem 2.3 yields a copy of F in $\hat{H} \subseteq H$.

So to conclude the proof it only remains to verify (3.3). Suppose to this end that some triple $\{i, j, k\} \in [m]^3$ is given. We have to verify that

$$|E(\mathcal{A}^{ijk})| \geq \mu |\mathcal{P}^{ij}| |\mathcal{P}^{ik}| |\mathcal{P}^{jk}| = \frac{\varepsilon}{8} \ell^3. \quad (3.4)$$

Using that H is $(\varepsilon, \eta, \bullet)$ -dense we infer

$$e_H(V_i, V_j, V_k) \geq \varepsilon |V_i| |V_j| |V_k| - \eta n^3$$

and by our choice of η it follows that

$$|V_i| |V_j| |V_k| \geq \left(\frac{(1 - \delta_3)}{T_0} \right)^3 n^3 = \frac{4\eta}{\varepsilon} n^3.$$

So altogether we have

$$e_H(V_i, V_j, V_k) \geq \frac{3}{4} \varepsilon |V_i| |V_j| |V_k|.$$

In combination with $\delta_3 \leq \frac{\varepsilon}{4} = d_3$ and condition (iv) from Theorem 2.2 this entails

$$e_{\hat{H}}(V_i, V_j, V_k) \geq \frac{1}{4} \varepsilon |V_i| |V_j| |V_k|. \quad (3.5)$$

On the other hand, by the triangle counting lemma (2.1) and $\delta_2 \ll \ell^{-1}$ each triad $P_{\alpha\beta\gamma}^{ijk}$ satisfies

$$\mathcal{K}_3(P_{\alpha\beta\gamma}^{ijk}) \leq (\ell^{-3} + 3\delta_2(\ell)) |V_i| |V_j| |V_k| \leq 2\ell^{-3} |V_i| |V_j| |V_k|,$$

for which reason

$$e_{\hat{H}}(V_i, V_j, V_k) \leq |E(\mathcal{A}^{ijk})| \cdot 2\ell^{-3} |V_i| |V_j| |V_k|.$$

Together with (3.5) this proves (3.4) and, hence, the implication from Lemma 3.1 to Theorem 1.2. \square

§4. PROOF OF THEOREM 1.2

This entire section is devoted to the proof of Lemma 3.1. We begin by outlining the main ideas of this proof. The argument proceeds in three stages. In the first of them we will choose a subset $X \subseteq [m]$ and for any two indices $r < s$ from X some vertex $P_{\text{red}}^{rs} \in \mathcal{P}^{rs}$ such that if $r < s < t$ are from X , then P_{red}^{rs} has large degree in \mathcal{A}^{rst} , where “large” means at least $\mu' |\mathcal{P}^{rt}| |\mathcal{P}^{st}|$ for some μ' depending only on μ . This argument will have the property that for fixed f and μ the size of X can be made as large as we wish by starting from a sufficiently large m . Then, in the next stage, we shrink the set X further to some $Y \subseteq X$ and select vertices $P_{\text{blue}}^{rt} \in \mathcal{P}^{rt}$ for all indices $r < t$ from Y such that if $r < s < t$ are from Y then the pair-degree of P_{red}^{rs} and P_{blue}^{rt} in \mathcal{A}^{rst} is still reasonably large, i.e., at least $\mu'' |\mathcal{P}^{st}|$

for some μ'' that depends again only on μ . Finally for some $Z \subseteq Y$ of size f we will manage to pick vertices P_{green}^{st} for $s < t$ from Z such that whenever $r < s < t$ are from Z the triple $P_{\text{red}}^{rs} P_{\text{blue}}^{rt} P_{\text{green}}^{st}$ appears in \mathcal{A}^{rst} . For this to succeed we just need $|Y|$ and hence also $|X|$ and m to be large enough depending on f and μ . We then enumerate $Z = \{\lambda(1), \dots, \lambda(f)\}$ in increasing order to conclude the argument.

The construction we use for the first stage proceeds in $m^* = |X|$ steps. In the first step we just select $1 \in X$. In the second step we put 2 into X and we will also make a decision concerning P_{red}^{12} . For that we ask every candidate $k \in [3, m]$ that might be put into X in the future to propose suitable choices for P_{red}^{12} . This leads us to consider for each such k the set $\mathcal{P}_{k,\text{red}}^{12} \subseteq \mathcal{P}^{12}$ of vertices with degree $\frac{\mu}{2} \cdot |\mathcal{P}^{1k}| |\mathcal{P}^{2k}|$ in \mathcal{A}^{12k} . Since \mathcal{A} is μ -dense we have $|\mathcal{P}_{k,\text{red}}^{12}| \geq \frac{\mu}{2} \cdot |\mathcal{P}^{12}|$ for each $k \geq 3$. Thus we can choose a vertex P_{red}^{12} in such a manner that it belongs to $\mathcal{P}_{k,\text{red}}^{12}$ for many k 's. From now on we restrict our attention to such k 's only. The third step begins by putting the smallest such k into X . If this happens to be, e.g., 7 then we ask each still relevant $k > 7$ for an opinion about the possible choices for the pair $(P_{\text{red}}^{17}, P_{\text{red}}^{27})$ and then we choose these two vertices in such a way that there are sufficiently many possibilities to continue. The general situation after h such steps is described in Lemma 4.1 below and the simpler Corollary 4.2 contains all that is needed for our intended application.

When reading the statement of the following lemma it might be helpful to think of M , m , and ε there as being m , m^* , and $\frac{\mu}{2}$ from the outline above. Also, n_1, \dots, n_h correspond to the indices which were already put into X whilst n_{h+1}, \dots, n_m are the indices that still have a chance of being put into X in the future.

Lemma 4.1. *Given $\varepsilon \in (0, 1)$ and positive integers $m \geq h$ there exists a positive integer $M = M(\varepsilon, m, h)$ for which the following is true. Suppose that we have*

- *nonempty sets \mathcal{P}^{rs} for $1 \leq r < s \leq M$ and*
- *further sets $\mathcal{P}_{t,\text{red}}^{rs} \subseteq \mathcal{P}^{rs}$ with $|\mathcal{P}_{t,\text{red}}^{rs}| \geq \varepsilon |\mathcal{P}^{rs}|$ for $1 \leq r < s < t \leq M$,*

then there are indices $n_1 < \dots < n_m$ in $[M]$ and there are elements $P_{\text{red}}^{n_r n_s} \in \mathcal{P}^{n_r n_s}$ for $1 \leq r < s \leq h$ such that

$$P_{\text{red}}^{n_r n_s} \in \bigcap_{t \in (s, m]} \mathcal{P}_{n_t, \text{red}}^{n_r n_s}.$$

Proof. We argue by induction on h . For the base case $h = 1$ we may take $M(\varepsilon, m, 1) = m$ and $n_r = r$ for all $r \in [m]$; because no vertices P_{red}^{rs} have to be chosen, the conclusion holds vacuously.

Now suppose that the result is already known for some integer h and all relevant pairs of ε and m , and that an integer $m \geq h + 1$ as well as a real number $\varepsilon \in (0, 1)$ are given. Set

$$m' = h + 1 + \left\lceil \frac{m - h - 1}{\varepsilon^h} \right\rceil \quad \text{and} \quad M = M(\varepsilon, m, h + 1) = M(\varepsilon, m', h).$$

To see that M is as desired, let sets \mathcal{P}^{rs} and $\mathcal{P}_{t,\text{red}}^{rs}$ as described above be given. Due to the definition of M , there are indices $n_1 < \dots < n_{m'}$ in $[M]$ and certain $P_{\text{red}}^{n_r n_s} \in \mathcal{P}^{n_r n_s}$ such that $P_{\text{red}}^{n_r n_s} \in \mathcal{P}_{n_t, \text{red}}^{n_r n_s}$ holds whenever $1 \leq r < s < t \leq m'$ and $s \leq h$. We set

$$\mathcal{P} = \mathcal{P}^{n_1 n_{h+1}} \times \dots \times \mathcal{P}^{n_h n_{h+1}}.$$

For each h -tuple $(P_1, \dots, P_h) \in \mathcal{P}$ we write

$$Q(P_1, \dots, P_h) = \{t \in [h + 2, m'] : P_r \in \mathcal{P}_{n_t, \text{red}}^{n_r n_{h+1}} \text{ for every } r \in [h]\}. \quad (4.1)$$

By counting the elements of

$$\{(t, P_1, \dots, P_h) : t \in Q(P_1, \dots, P_h)\}$$

in two different ways and using the lower bounds $|\mathcal{P}_{n_t, \text{red}}^{n_r n_{h+1}}| \geq \varepsilon |\mathcal{P}^{n_r n_{h+1}}|$ we get

$$\sum_{(P_1, \dots, P_h) \in \mathcal{P}} |Q(P_1, \dots, P_h)| = \sum_{t=h+2}^{m'} \prod_{r=1}^h |\mathcal{P}_{n_t, \text{red}}^{n_r n_{h+1}}| \geq (m' - h - 1) \varepsilon^h |\mathcal{P}|.$$

Hence, we may fix an h -tuple $(P_1, \dots, P_h) \in \mathcal{P}$ with

$$|Q(P_1, \dots, P_h)| \geq (m' - h - 1) \varepsilon^h \geq m - h - 1.$$

Now let $\ell_{h+2} < \dots < \ell_m$ be any elements from

$$Q = \{n_t : t \in Q(P_1, \dots, P_h)\}$$

in increasing order. Set

$$\ell_r = n_r \text{ for all } r \in [h + 1] \text{ as well as } P_{\text{red}}^{n_r, n_{h+1}} = P_r \text{ for all } r \in [h].$$

We claim that the indices $\ell_1 < \dots < \ell_m$ and the elements $P_{\text{red}}^{n_r n_s}$ with $1 \leq r < s \leq h + 1$ satisfy the conclusion. To see this let any $1 \leq r < s < t \leq m$ with $s \leq h + 1$ be given. We have to verify $P_{\text{red}}^{\ell_r \ell_s} \in \mathcal{P}_{\ell_t, \text{red}}^{\ell_r \ell_s}$. If $s \leq h$ this follows directly from $\ell_r = n_r$, $\ell_s = n_s$, $\ell_t \in \{n_{s+1}, \dots, n_{m'}\}$, and the inductive choice of the latter set. For the case $s = h + 1$ it follows from $t \geq h + 2$, that there is some $q \in Q(P_1, \dots, P_h)$ with $\ell_t = n_q$. The first property of q entails in view of (4.1) that $P_r \in \mathcal{P}_{n_q, \text{red}}^{n_r n_{h+1}}$ and, as $P_{\text{red}}^{n_r, n_{h+1}} = P_r$, this is exactly what we wanted. \square

The reason for having the two parameters m and h in this lemma is just that this facilitates the proof by induction on h . In applications one may always set $h = m$, since this gives the strongest possible conclusion for fixed m . Thus it might add to the clarity of exposition if we restate this case again, using the occasion to eliminate some double indices as well.

Corollary 4.2. *Suppose that for $M \gg \max(m, \varepsilon^{-1})$ we have*

- *nonempty sets \mathcal{P}^{rs} for $1 \leq r < s \leq M$ and*
- *further sets $\mathcal{P}_{t,\text{red}}^{rs} \subseteq \mathcal{P}^{rs}$ with $|\mathcal{P}_{t,\text{red}}^{rs}| \geq \varepsilon |\mathcal{P}^{rs}|$ for $1 \leq r < s < t \leq M$,*

then there is a subset $X \subseteq [M]$ of size m and there are elements $P_{\text{red}}^{rs} \in \mathcal{P}^{rs}$ for $r < s$ from X such that

$$P_{\text{red}}^{rs} \in \bigcap_t \{ \mathcal{P}_{t,\text{red}}^{rs} : t > s \text{ and } t \in X \}. \quad \square$$

As discussed above, this statement will be used below for choosing the vertices P_{red}^{rs} . The selection principle we use for choosing the P_{green}^{st} is essentially the same, but we have to apply the symmetry $r \mapsto M + 1 - r$ to the indices throughout. To prevent confusion when this happens within another argument, it might thus be helpful to restate the foregoing result as follows.

Corollary 4.3. *Suppose that for $M \gg \max(m, \varepsilon^{-1})$ we have*

- *nonempty sets \mathcal{P}^{st} for $1 \leq s < t \leq M$ and*
- *further sets $\mathcal{P}_{r,\text{green}}^{st} \subseteq \mathcal{P}^{st}$ with $|\mathcal{P}_{r,\text{green}}^{st}| \geq \varepsilon |\mathcal{P}^{st}|$ for $1 \leq r < s < t \leq M$,*

then there is a subset $Z \subseteq [M]$ of size m and there are elements $P_{\text{green}}^{st} \in \mathcal{P}^{st}$ for $s < t$ from Z such that

$$P_{\text{green}}^{st} \in \bigcap_r \{ \mathcal{P}_{r,\text{green}}^{st} : r < s \text{ and } r \in Z \}.$$

Proof. Set $\mathcal{P}_*^{rs} = \mathcal{P}^{M+1-s, M+1-r}$ for $1 \leq r < s \leq M$ and $\mathcal{P}_{t,\text{red}}^{rs} = \mathcal{P}_{M+1-t, \text{green}}^{M+1-s, M+1-r}$ for $1 \leq r < s < t \leq M$. Then apply Corollary 4.2, thus getting a certain set X and some elements P_{red}^{rs} . It is straightforward to check that

$$Z = \{M + 1 - x : x \in X\}$$

and $P_{\text{green}}^{st} = P_{\text{red}}^{M+1-t, M+1-s}$ are as desired. \square

The statement that follows coincides with [13, Lemma 7.1], where a short direct proof is given. For reasons of self-containment, however, we will show here that it follows easily from the above Corollary 4.3. Subsequently it will be used in the proof of a lemma playing a rôle similar to that of Lemma 4.1, but preparing the selection of the vertices P_{blue}^{rt} rather than P_{red}^{rs} . Specifically, the statement that follows will be used in that step of the proof of the next lemma that corresponds to choosing P_1, \dots, P_h in the proof of Lemma 4.1.

Corollary 4.4. *Suppose that for $M \gg \max(m, \varepsilon^{-1})$ we have*

- *nonempty sets W_1, \dots, W_M and*
- *further sets $D_{rs} \subseteq W_s$ with $|D_{rs}| \geq \varepsilon |W_s|$ for $1 \leq r < s \leq M$,*

then there is a subset $Z \subseteq [M]$ of size m and there are elements $d_s \in W_s$ for $s \in Z$ such that

$$d_s \in \bigcap_r \{D_{rs} : r < s \text{ and } r \in Z\}.$$

Proof. Let M be so large that the conclusion of Corollary 4.3 holds with $m + 1$ in place of m and with the same ε . Now let the sets W_s and D_{rs} as described above be given.

Set $\mathcal{P}^{st} = W_s$ for $1 \leq s < t \leq M$ and $\mathcal{P}_{r,\text{green}}^{st} = D_{rs}$ for $1 \leq r < s < t \leq M$. By hypothesis $\mathcal{P}_{r,\text{green}}^{st}$ is a sufficiently large subset of \mathcal{P}^{st} , so by our choice of M there is a set $Z^* \subseteq [M]$ of size $m + 1$ together with certain elements $P_{\text{green}}^{st} \in \mathcal{P}^{st}$ for $s < t$ from Z^* such that $P_{\text{green}}^{st} \in \mathcal{P}_{r,\text{green}}^{st}$ holds whenever $r < s < t$ are from Z^* . Set $z = \max(Z^*)$, $Z = Z^* \setminus \{z\}$, and $d_s = P_{\text{green}}^{sz}$ for all $s \in Z$. We claim that Z and the d_s are as demanded.

The condition $|Z| = m$ is clear, so now let any pair $r < s$ from Z be given. Then $r < s < z$ are from Z^* , whence $d_s = P_{\text{green}}^{sz} \in \mathcal{P}_{r,\text{green}}^{sz} = D_{rs}$. \square

The next lemma deals with the selection of “blue” vertices.

Lemma 4.5. *Given $\varepsilon \in (0, 1)$ and nonnegative integers $m \geq h$ there exists a positive integer $M = M(\varepsilon, m, h)$ for which the following is true. Suppose that we have*

- *nonempty sets \mathcal{P}^{rt} for $1 \leq r < t \leq M$ and*
- *further sets $\mathcal{P}_{s,\text{blue}}^{rt} \subseteq \mathcal{P}^{rt}$ with $|\mathcal{P}_{s,\text{blue}}^{rt}| \geq \varepsilon |\mathcal{P}^{rt}|$ for $1 \leq r < s < t \leq M$,*

then there are indices $n_1 < \dots < n_m$ in $[M]$ and there are elements $P_{\text{blue}}^{n_r n_t} \in \mathcal{P}^{n_r n_t}$ for all $1 \leq r < t \leq m$ with $r \leq h$ such that

$$P_{\text{blue}}^{n_r n_t} \in \bigcap_s \{\mathcal{P}_{n_s, \text{blue}}^{n_r n_t} : r < s < t\}.$$

Proof. Again we argue by induction on h with the base case $h = 0$ being trivial.

For the induction step we assume that the lemma is already known for some h and all possibilities for m and ε , and proceed to the case $m \geq h + 1$. We contend that $M = M(\varepsilon, m', h)$ is as desired when m' is chosen so large that the conclusion of Corollary 4.4 holds for $(m' - h - 1, m - h - 1)$ here in place of (M, m) there – with the same value of ε .

So let any sets \mathcal{P}^{rt} and $\mathcal{P}_{s,\text{blue}}^{rt}$ as described above be given. The choice of M guarantees the existence of some indices $n_1 < \dots < n_{m'}$ in $[M]$ together with certain elements $P_{\text{blue}}^{n_r n_t}$ satisfying the conclusion of Lemma 4.5 with m' in place of m . The m indices we are requested to find will be n_1, \dots, n_{h+1} and $(m - h - 1)$ members of the set $\{n_{h+2}, \dots, n_{m'}\}$, so in order to gain notational simplicity we may assume $n_r = r$ for all $r \in [m']$. Thus we have $P_{\text{blue}}^{rt} \in \mathcal{P}_{s,\text{blue}}^{rt}$ whenever $1 \leq r < s < t \leq m'$ and $r \leq h$.

Let us now define $W_j = \mathcal{P}^{h+1, h+j+1}$ for all $j \in [m' - h - 1]$ and $D_{ij} = \mathcal{P}_{h+i+1, \text{blue}}^{h+1, h+j+1}$ for all $i < j$ from $[m' - h - 1]$. Then the conditions of Corollary 4.4 are satisfied, meaning that there is a subset Z of $[m' - h - 1]$ of size $m - h - 1$ together with certain elements $d_j \in W_j$ for $j \in Z$ such that we have $d_j \in D_{ij}$ whenever $i < j$ are from Z .

We contend that the set of the m indices we are supposed to find can be taken to be

$$[h + 1] \cup ((h + 1) + Z).$$

To see this we may for simplicity assume $Z = [m - h - 1]$, so that the set of our m indices is simply $[m]$. Recall that we have already found above certain elements $P_{\text{blue}}^{rt} \in \mathcal{P}^{rt}$ for $1 \leq r < t \leq m$ with $r \leq h$ such that $P_{\text{blue}}^{rt} \in \mathcal{P}_{s, \text{blue}}^{rt}$ holds whenever $1 \leq r < s < t \leq m$ and $r \leq h$. So it remains to find further elements $P_{\text{blue}}^{h+1, t} \in \mathcal{P}^{h+1, t}$ for $t \in [h + 2, m]$ with $P_{\text{blue}}^{h+1, t} \in \mathcal{P}_{s, \text{blue}}^{h+1, t}$ whenever $h + 2 \leq s < t \leq m$. To this end, we use the vertices obtained by applying Corollary 4.4 and set $P_{\text{blue}}^{h+1, t} = d_{t-h-1}$ for all $t \in [m + 2, h]$. Observe that $P_{\text{blue}}^{h+1, t} \in W_{t-h-1} = \mathcal{P}^{h+1, t}$ holds for all relevant t . Moreover, if $h + 2 \leq s < t \leq m$, then we have indeed $P_{\text{blue}}^{h+1, t} = d_{t-h-1} \in D_{s-h-1, t-h-1} = \mathcal{P}_{s, \text{blue}}^{h+1, t}$. Thereby the proof by induction on h is complete. \square

For the same reasons as before we restate the case $h = m$ as follows.

Corollary 4.6. *Suppose that for $M \gg \max(m, \varepsilon^{-1})$ we have*

- *nonempty sets \mathcal{P}^{rt} for $1 \leq r < t \leq M$ and*
- *further sets $\mathcal{P}_{s, \text{blue}}^{rt} \subseteq \mathcal{P}^{rt}$ with $|\mathcal{P}_{s, \text{blue}}^{rt}| \geq \varepsilon |\mathcal{P}^{rt}|$ for $1 \leq r < s < t \leq M$,*

then there is a subset $Y \subseteq [M]$ of size m and there are elements $P_{\text{blue}}^{rt} \in \mathcal{P}^{rt}$ for $r < t$ from Y such that

$$P_{\text{blue}}^{rt} \in \bigcap_s \{ \mathcal{P}_{s, \text{blue}}^{rt} : r < s < t \text{ and } s \in Y \}. \quad \square$$

After these preparations we are ready to verify Lemma 3.1.

Proof of Lemma 3.1. Suppose

$$m \gg m_* \gg m_{**} \gg \max(f, \mu^{-1}).$$

Consider any three indices $1 \leq r < s < t \leq m$. For a vertex $P \in \mathcal{P}^{rs}$ we denote the degree of P in \mathcal{A}^{rst} by $d_t(P)$. In other words, this is the number of pairs $(Q, R) \in \mathcal{P}^{rt} \times \mathcal{P}^{st}$ with $\{P, Q, R\} \in E(\mathcal{A}^{rst})$. Further, we set

$$\mathcal{P}_{t, \text{red}}^{rs} = \{ P \in \mathcal{P}^{rs} : d_t(P) \geq \frac{\mu}{2} \cdot |\mathcal{P}^{rt}| |\mathcal{P}^{st}| \}.$$

Since

$$\begin{aligned} \mu |\mathcal{P}^{rs}| |\mathcal{P}^{rt}| |\mathcal{P}^{st}| &\leq |E(\mathcal{A}^{rst})| = \sum_{P \in \mathcal{P}^{rs}} d_t(P) = \sum_{P \in \mathcal{P}^{rs} \setminus \mathcal{P}_{t,\text{red}}^{rs}} d_t(P) + \sum_{P \in \mathcal{P}_{t,\text{red}}^{rs}} d_t(P) \\ &\leq \frac{\mu}{2} \cdot |\mathcal{P}^{rs}| |\mathcal{P}^{rt}| |\mathcal{P}^{st}| + |\mathcal{P}_{t,\text{red}}^{rs}| |\mathcal{P}^{rt}| |\mathcal{P}^{st}|, \end{aligned}$$

we have $|\mathcal{P}_{t,\text{red}}^{rs}| \geq \frac{\mu}{2} \cdot |\mathcal{P}^{rs}|$. So applying Corollary 4.2 with $(m, m^*, \frac{\mu}{2})$ here in place of (M, m, ε) there we get a set $X \subseteq [m]$ of size m_* together with some vertices P_{red}^{rs} satisfying the condition mentioned there. For simplicity we relabel our indices in such a way that $X = [m^*]$, intending to find the required indices $\lambda(1), \dots, \lambda(f)$ in $[m^*]$. This completes what has been called the first stage of the proof in the outline at the beginning of this section.

Next we look at any three indices $1 \leq r < s < t \leq m_*$. Recall that we just achieved $d_t(P_{\text{red}}^{rs}) \geq \frac{\mu}{2} \cdot |\mathcal{P}^{rt}| |\mathcal{P}^{st}|$. We write $p(P, Q)$ for the pair-degree of any two vertices $P \in \mathcal{P}^{rs}$ and $Q \in \mathcal{P}^{rt}$ in \mathcal{A}^{rst} , i.e., for the number of triples of this hypergraph containing both P and Q . Let us define

$$\mathcal{P}_{s,\text{blue}}^{rt} = \{Q \in \mathcal{P}^{rt} : p(P_{\text{red}}^{rs}, Q) \geq \frac{\mu}{4} \cdot |\mathcal{P}^{st}|\}.$$

Starting from the obvious formula

$$d(P_{\text{red}}^{rs}) = \sum_{Q \in \mathcal{P}^{rt}} p(P_{\text{red}}^{rs}, Q),$$

the same calculation as above discloses $|\mathcal{P}_{s,\text{blue}}^{rt}| \geq \frac{\mu}{4} \cdot |\mathcal{P}^{rt}|$. So we may apply Corollary 4.6 with $(m_*, m_{**}, \frac{\mu}{4})$ here instead of (M, m, ε) there in order to find a subset Y of $[m_*]$ of size m_{**} together with certain vertices P_{blue}^{rt} . As before it is allowed to suppose $Y = [m_{**}]$, in which case we have $p(P_{\text{red}}^{rs}, P_{\text{blue}}^{rt}) \geq \frac{\mu}{4} \cdot |\mathcal{P}^{st}|$ whenever $1 \leq r < s < t \leq m_{**}$.

Having thus completed the second stage we look at any three indices $1 \leq r < s < t \leq m_{**}$. Let $\mathcal{P}_{r,\text{green}}^{st}$ denote the set of all vertices R from \mathcal{P}^{st} for which the triple $\{P_{\text{red}}^{rs}, P_{\text{blue}}^{rt}, R\}$ belongs to \mathcal{A}^{rst} . Due to our previous choices we have $|\mathcal{P}_{r,\text{green}}^{st}| \geq \frac{\mu}{4} \cdot |\mathcal{P}^{st}|$. So we may apply Corollary 4.3 with $(m_{**}, f, \frac{\mu}{4})$ here rather than (M, m, ε) there, thus getting a certain set $Z \subseteq [m_{**}]$ and certain vertices $P_{\text{green}}^{st} \in \mathcal{P}^{st}$ for $s < t$ from Z . As always we may suppose that $Z = [f]$, so that $\{P_{\text{red}}^{rs}, P_{\text{blue}}^{rt}, P_{\text{green}}^{st}\}$ becomes a triple of \mathcal{A}^{rst} whenever $1 \leq r < s < t \leq f$. Now it is plain that the indices $\lambda(r) = r$ for $r \in [f]$ are as desired. \square

§5. UNIFORMLY DENSE WITH VANISHING DENSITY

We reprove Theorem 1.7 from [4] and we devote to each implication a separate section.

5.1. The forward implication. The statement that every frequent hypergraph is contained in one and, hence, eventually in all sufficiently large ternary hypergraphs, is a direct consequence of the fact that the sequence $(T_n)_{n \in \mathbb{N}}$ is itself d -dense for an appropriate function $d: (0, 1] \rightarrow (0, 1]$. This observation is due to Erdős and Sós [3] who left the verification to the reader. In [4, Proposition 3.1] it was shown that the sequence of ternary hypergraphs is d -dense for some function $d(\eta) = \eta^\varrho$ with $\varrho > 10$. Here we sharpen this estimate and establish Proposition 1.6, which gives the optimal exponent

$$\varrho = \frac{2}{\log_2(3) - 1} \approx 3.419\dots \quad (5.1)$$

More precisely, we prove the following lemma, which yields Proposition 1.6.

Lemma 5.1. *For ϱ given in (5.1), $\ell \geq 1$, $X \subseteq V(T_\ell)$, and $|X| = \eta \cdot 3^\ell$ we have*

$$e(X) \geq \frac{1}{4}\eta^\varrho \cdot \frac{|X|^3}{6} - \frac{3}{8} \cdot 3^\ell.$$

For the proof of this lemma we shall utilise the following inequality.

Fact 5.2. *If $x, y, z \in [0, 1]$ and $\tau = \varrho + 3$ for ϱ given in (5.1), then*

$$x^\tau + y^\tau + z^\tau + 24xyz \geq 3^{3-\tau}(x + y + z)^\tau.$$

Proof. In the proof the following identity will be handy to use

$$2^{\tau-1} = 3^{\tau-3}. \quad (5.2)$$

As the unit cube is compact, there is a point $(x_*, y_*, z_*) \in [0, 1]^3$ at which the continuous function $f: [0, 1]^3 \rightarrow \mathbb{R}$ given by

$$(x, y, z) \longmapsto x^\tau + y^\tau + z^\tau + 24xyz - 3^{3-\tau}(x + y + z)^\tau$$

attains its minimum value, say ξ . Due to symmetry we may suppose that $x_* \geq y_* \geq z_*$. Assume for the sake of contradiction that $\xi < 0$.

Since $\tau > 1$, convexity implies

$$x^\tau + y^\tau \geq 2\left(\frac{x+y}{2}\right)^\tau = 2^{1-\tau}(x+y)^\tau \stackrel{(5.2)}{=} 3^{3-\tau}(x+y)^\tau.$$

Consequently, $f(x, y, 0) \geq 0$ for all real $x, y \in [0, 1]$ and we have $x_*, y_*, z_* > 0$.

The minimality of ξ implies

$$\begin{aligned} x_*^\tau \xi &\leq x_*^\tau f\left(1, \frac{y_*}{x_*}, \frac{z_*}{x_*}\right) = x_*^\tau + y_*^\tau + z_*^\tau + 24x_*^{\tau-3} \cdot x_* y_* z_* - 3^{3-\tau}(x_* + y_* + z_*)^\tau \\ &= \xi + 24(x_*^{\tau-3} - 1)x_* y_* z_*, \end{aligned}$$

i.e., $24(1 - x_*^{\tau-3})x_*y_*z_* \leq \xi(1 - x_*^\tau)$, which due to the assumption $\xi < 0$ is only possible if $x_* = 1$. In other words, the function $x \mapsto f(x, y_*, z_*)$ from $[0, 1]$ to \mathbb{R} attains its minimum at the boundary point $x = 1$ and for this reason we have $\frac{df(x, y_*, z_*)}{dx}\big|_{x=1} \leq 0$, i.e.,

$$\tau + 24y_*z_* \leq \tau \cdot 3^{3-\tau}(1 + y_* + z_*)^{\tau-1}. \quad (5.3)$$

Next we observe that the function $z \mapsto f(1, 1, z)$ from $[0, 1]$ to \mathbb{R} is concave, because

$$\begin{aligned} \frac{d^2 f(1, 1, z)}{dz^2} &= (\tau - 1)\tau (z^{\tau-2} - 3^{3-\tau}(2 + z)^{\tau-2}) \\ &= (\tau - 1)\tau \left(\frac{(3z)^{\tau-2} - 3(2 + z)^{\tau-2}}{3^{\tau-2}} \right) < 0. \end{aligned}$$

Together with

$$f(1, 1, 0) = 2 - 3^{3-\tau} \cdot 2^\tau \stackrel{(5.2)}{=} 0 \quad \text{and} \quad f(1, 1, 1) = 27 - 3^{3-\tau} \cdot 3^\tau = 0$$

this proves that $f(1, 1, z) \geq 0$ holds for all $z \in [0, 1]$, which in view of $x_* = 1$ yields $y_* < 1$. Thus the function $y \mapsto f(1, y, z_*)$ from $[0, 1]$ to \mathbb{R} attains its minimum at the interior point $y = y_*$ and we infer $\frac{df(1, y, z_*)}{dy}\big|_{y=y_*} = 0$, i.e.,

$$\tau y_*^{\tau-1} + 24z_* = \tau \cdot 3^{3-\tau}(1 + y_* + z_*)^{\tau-1}.$$

In combination with (5.3) this proves $24(1 - y_*)z_* \geq \tau(1 - y_*^{\tau-1})$ and recalling $y_* \geq z_*$ we arrive at

$$24(1 - y_*)y_* \geq \tau(1 - y_*^{\tau-1}) > \frac{32}{5}(1 - y_*^5), \quad (5.4)$$

where we used $\tau = \varrho + 3 > 6.4$ for the last inequality (see (5.1)). Dividing by $(1 - y_*)y_*$ leads to

$$\frac{1 + y_* + y_*^2 + y_*^3 + y_*^4}{y_*} = \frac{1 - y_*^5}{(1 - y_*)y_*} \stackrel{(5.4)}{<} \frac{15}{4}. \quad (5.5)$$

Now for the function $h: (0, 1) \rightarrow \mathbb{R}$ given by $h(t) = \frac{1}{t} + 1 + t + t^2 + t^3$ we have

$$h'(t) < 0 \iff t^2(1 + 2t + 3t^2) < 1.$$

Consequently, there is a unique point $t_* \in (0, 1)$, at which h attains its global minimum and a short calculation reveals $t_* \in \left[\frac{5}{9}, \frac{4}{7}\right]$.

From (5.5) we may now deduce

$$\left(\frac{1}{t_*} + 1 + t_* \right) + t_*^2 + t_*^3 < \frac{15}{4}.$$

Since $t \mapsto \frac{1}{t} + 1 + t$ is decreasing on $(0, 1)$, this may be weakened to

$$\frac{7}{4} + 1 + \frac{4}{7} + \left(\frac{5}{9}\right)^2 + \left(\frac{5}{9}\right)^3 < \frac{15}{4},$$

which, however, is not the case. Thus $\xi \geq 0$ and Fact 5.2 is proved. \square

Lemma 5.1 follows by a simple inductive argument from the inequality from Fact 5.2.

Proof of Lemma 5.1. The case $\ell = 1$ is clear, since then the right-hand side cannot be positive. Proceeding inductively we assume from now on that the lemma holds for $\ell - 1$ in place of ℓ and look at an arbitrary set $X \subseteq V(T_\ell)$.

Let $V(T_\ell) = V_1 \cup V_2 \cup V_3$ be a partition of the vertex set of T_ℓ such that

- each of V_1 , V_2 , and V_3 induces a copy of $T_{\ell-1}$
- and all triples $v_1v_2v_3$ with $v_i \in V_i$ for $i = 1, 2, 3$ are edges of T_ℓ .

Setting $X_i = X \cap V_i$ and $\eta_i = |X_i|/3^{\ell-1}$ for $i = 1, 2, 3$ we get

$$\begin{aligned} e(X) &= e(X_1) + e(X_2) + e(X_3) + |X_1||X_2||X_3| \\ &\geq \left(\frac{\eta_1^{\ell+3} + \eta_2^{\ell+3} + \eta_3^{\ell+3} + 24\eta_1\eta_2\eta_3}{4} \right) \frac{(3^{\ell-1})^3}{6} - 3 \cdot \frac{3}{8} \cdot 3^{\ell-1} \end{aligned}$$

from the induction hypothesis. In view of Fact 5.2 it follows that

$$e(X) \geq \frac{27\eta^{\ell+3}}{4} \cdot \frac{(3^{\ell-1})^3}{6} - \frac{3}{8} \cdot 3^\ell, \tag{5.6}$$

where

$$\eta = \frac{\eta_1 + \eta_2 + \eta_3}{3} = \frac{|X_1| + |X_2| + |X_3|}{3^\ell} = \frac{|X|}{3^\ell},$$

meaning that (5.6) simplifies to the desired estimate

$$e(X) \geq \frac{\eta^\ell}{4} \cdot \frac{|X|^3}{6} - \frac{3}{8} \cdot 3^\ell. \quad \square$$

We conclude this subsection by observing that frequent hypergraphs on ℓ vertices must be contained in the ternary hypergraph on 3^ℓ vertices.

Lemma 5.3. *If a hypergraph F on ℓ vertices is frequent, then it is a subhypergraph of the ternary hypergraph T_ℓ .*

Proof. It follows from Lemma 5.1 that there is some $n \in \mathbb{N}$ with $F \subseteq T_n$. Thus it suffices to prove that if $F \subseteq T_n$ and $v(F) = \ell$, then $F \subseteq T_\ell$ holds as well. We do so by induction on ℓ , the base case $\ell \leq 3$ being clear.

Now let any hypergraph F appearing in some ternary hypergraph and with $\ell \geq 4$ vertices be given and choose $n \in \mathbb{N}$ minimal with $F \subseteq T_n$. Take a partition $V(T_n) = V_1 \cup V_2 \cup V_3$ such that each of V_1 , V_2 , and V_3 induces a copy of T_{n-1} and such that all further edges of T_n are of the form $v_1v_2v_3$ with $v_i \in V_i$ for $i = 1, 2, 3$. By the minimality of n each of the three sets $V_i \cap V(F)$ with $i = 1, 2, 3$ contains less than ℓ vertices, so by the induction hypothesis they induce subhypergraphs of T_n that appear already in $T_{\ell-1}$. Therefore we have indeed $F \subseteq T_\ell$. \square

5.2. The backward implication. For completeness we include a proof of the fact that subhypergraphs of ternary hypergraphs are indeed frequent. This proof follows the lines of the work in [4] and will be done by induction on the order of the hypergraph whose frequency we wish to establish. In order to carry the induction it will help us to address the corresponding supersaturation assertion directly. Let us recall to this end that a *homomorphism* from a hypergraph F to another hypergraph H is a map $\varphi: V(F) \rightarrow V(H)$ sending edges of F to edges of H ; explicitly, this means that $\{\varphi(x), \varphi(y), \varphi(z)\} \in E(H)$ is required to hold for every triple $xyz \in E(F)$. The set of these homomorphisms is denoted by $\text{Hom}(F, H)$ and $\text{hom}(F, H) = |\text{Hom}(F, H)|$ stands for the *number* of homomorphisms from F to H .

Proposition 5.4. *Given a hypergraph F which is a subhypergraph of some ternary hypergraph and a function $d: (0, 1) \rightarrow (0, 1)$, there are constants $\eta, \xi > 0$ such that*

$$\text{hom}(F, H) \geq \xi v(H)^{v(F)}$$

is satisfied by every hypergraph H with the property that $e(U) \geq d(\varepsilon)|U|^3/6$ holds whenever $U \subseteq V(H)$, $\varepsilon \in [\eta, 1]$, and $|U| \geq \varepsilon |V(H)|$.

Proof. We argue by induction on $v(F)$. The base case $v(F) \leq 2$ is clear, since then F cannot have any edge and $\eta = \xi = 1$ works. For $v(F) = 3$ we take $\eta = 1$ as well as $\xi = d(1)$. As every edge of H gives rise to six homomorphisms from F to H we get indeed $\text{hom}(F, H) \geq 6e(H) \geq d(1)v(H)^3$.

For the induction step let a hypergraph F with $v(F) \geq 4$ and a function $d: (0, 1) \rightarrow (0, 1)$ be given. Let $\ell \geq 2$ be minimal with $F \subseteq T_\ell$. For simplicity we will suppose that F is in fact an induced subhypergraph of T_ℓ .

Again we take a partition $V(T_\ell) = V_1 \cup V_2 \cup V_3$ such that V_i spans a copy of $T_{\ell-1}$ for $i = 1, 2, 3$ and all further edges of T_ℓ are of the form $v_1v_2v_3$ with $v_i \in V_i$ for $i = 1, 2, 3$. By symmetry we may suppose, after a possible renumbering of indices, that $|V(F) \cap V_3| \geq 2$ holds. Let F_{12} and F_3 be the restrictions of F to $V_1 \cup V_2$ and V_3 , respectively. Moreover, we will need the hypergraph F_* arising from F by deleting all but one vertex from $V(F) \cap V_3$. An alternative and perhaps helpful description of F_* is that can be obtained from F_{12} by adding a new vertex z and all triples v_1v_2z with $v_1 \in V(F) \cap V_1$ and $v_2 \in V(F) \cap V_2$.

Intuitively the reason why there should be many homomorphisms from F into an n -vertex hypergraph H satisfying some local density condition is the following. Due to $v(F_*) < v(F)$ we may assume by induction that $\text{hom}(F_*, H) = \Omega(n^{v(F_*)})$. This means that there is a collection of $\Omega(n^{v(F_{12})})$ homomorphisms φ from F_{12} to H that can be extended in $\Omega(n)$ many ways to a member of $\text{Hom}(F_*, H)$. For each such φ the set $A_\varphi \subseteq V(H)$ consisting of the possible images of the new vertex z in such an extension inherits a local density

condition, because its size is linear, and a further use of the induction hypothesis shows that there are $\Omega(n^{v(F_3)})$ homomorphisms from F_3 to A_φ . These homomorphisms can in turn be regarded as extensions of φ to members of $\text{Hom}(F, H)$. This argument can be performed for any φ and thus we get $\text{Hom}(F, H) \geq \Omega(n^{v(F_{12})}) \cdot \Omega(n^{v(F_3)}) = \Omega(n^{v(F)})$.

Proceeding now to the details of this derivation let η_* and ξ_* denote the constants obtained by applying the induction hypothesis to F_* and $d(\cdot)$. The minimality of ℓ implies $v(F_3) < v(F)$ and therefore we may apply the induction hypothesis to F_3 and the function $d': (0, 1) \rightarrow (0, 1)$ defined by $\varepsilon \mapsto d(\varepsilon \cdot \xi_*/2)$, thus obtaining two further constants η_3 and ξ_3 . We contend that

$$\eta = \min\left(\eta_*, \frac{1}{2}\xi_*\eta_3\right) \quad \text{and} \quad \xi = \frac{\xi_*^{v(F_3)+1}\xi_3}{2^{v(F_3)+1}}$$

have the requested properties.

Now let any hypergraph H with $e(U) \geq d(\varepsilon)|U|^3/6$ for all $\varepsilon \in [\eta, 1]$ $U \subseteq V(H)$ with $|U| \geq \varepsilon|V(H)|$ be given and put $n = v(H)$. Due to $\eta_* \geq \eta$ we have

$$\text{hom}(F_*, H) \geq \xi_* n^{v(F_*)}. \quad (5.7)$$

For every homomorphism $\varphi \in \text{Hom}(F_{12}, H)$ we consider the set

$$A_\varphi = \{v \in V(H) : \varphi \cup \{(z, v)\} \in \text{Hom}(F_*, H)\}$$

of vertices that can be used for extending φ to a homomorphism $\varphi \cup \{(z, v)\}$ from F_* to H . It will be convenient to identify these sets with the subhypergraphs of H they induce. Finally we define

$$\Phi = \{\varphi \in \text{Hom}(F_{12}, H) : |A_\varphi| \geq \frac{1}{2}\xi_* n\}$$

to be the set of those homomorphisms from F_{12} to H that admit a substantial number of such extensions.

Since $v(F_*) = v(F_{12}) + 1$ we obtain from (5.7)

$$\xi_* n^{v(F_{12})+1} \leq \sum_{\varphi \in \text{Hom}(F_{12}, H)} |A_\varphi| \leq |\Phi| \cdot n + n^{v(F_{12})} \cdot \frac{1}{2}\xi_* n,$$

whence

$$|\Phi| \geq \frac{1}{2}\xi_* n^{v(F_{12})}. \quad (5.8)$$

Moreover it is clear that

$$\text{hom}(F, H) = \sum_{\varphi \in \text{Hom}(F_{12}, H)} \text{hom}(F_3, A_\varphi) \quad (5.9)$$

and the next thing we show is that for every $\varphi \in \Phi$ we have

$$\text{hom}(F_3, A_\varphi) \geq \xi_3 \left(\frac{1}{2}\xi_* n\right)^{v(F_3)}. \quad (5.10)$$

Owing to our inductive choice of η_3 and ξ_3 it suffices for the verification of this estimate to show that if $\varepsilon \in [\eta_3, 1]$, $U \subseteq A_\varphi$, and $|U| \geq \varepsilon|A_\varphi|$, then $e(U) \geq d'(\varepsilon)|U|^3/6$. But since $\varphi \in \Phi$ leads to $|U| \geq \frac{1}{2}\varepsilon\xi_*n$, this follows immediately from $\frac{1}{2}\varepsilon\xi_* \geq \frac{1}{2}\xi_*\eta_3 \geq \eta$, the definition of d' , and from our choice of H .

Taken together (5.9), (5.10), and (5.8) yield

$$\text{hom}(F, H) \geq \sum_{\varphi \in \Phi} \text{hom}(F_3, A_\varphi) \geq |\Phi| \cdot \xi_3 \left(\frac{1}{2}\xi_*n\right)^{v(F_3)} \geq \frac{\xi_*^{v(F_3)+1} \xi_3}{2^{v(F_3)+1}} n^{v(F)},$$

as desired. \square

Proposition 5.4 implies that all subhypergraphs of ternary hypergraphs are frequent and combined with Lemma 5.3 this shows that being frequent is a decidable property.

§6. CONCLUDING REMARKS

6.1. Hypergraphs with uniformly positive density. In [13, Section 2] we defined for a given antichain $\mathcal{A} \subseteq \wp([k])$ and given real numbers $d \in [0, 1]$, $\eta > 0$ the concept of a k -uniform hypergraph being (d, η, \mathcal{A}) -dense. An obvious modification of (3.1) does then lead to corresponding *generalised Turán densities* $\pi_{\mathcal{A}}(F)$ of k -uniform hypergraphs F . Now the question presents itself to determine $\pi_{\mathcal{A}}(F)$ for all antichains \mathcal{A} and all hypergraphs F . At the moment this appears to be a hopelessly difficult task, as it includes, among many further variations, the original version of Turán's problem to determine the ordinary Turán density $\pi(F)$ of any hypergraph F .

For the time being it might be more reasonable to focus on the case $\mathcal{A} = [k]^{(k-2)}$ (or stronger density assumptions), as it might be that for this case one can establish a theory that resembles to some extent the classical theory for graphs initiated by Turán himself and developed further by Erdős, Stone, and Simonovits and many others.

Another possible direction is to characterise for given \mathcal{A} the hypergraphs F with $\pi_{\mathcal{A}}(F) = 0$ and here it seems natural to pay particular attention to the *symmetric case*, when $\mathcal{A} = [k]^{(j)}$ contains all j -element subsets of $[k]$. Let us now describe an extension of Theorem 1.2 to this setting. First of all, a k -uniform hypergraph $H = (V, E)$ is said to be (d, η, j) -dense, for real numbers $d \in [0, 1]$, $\eta > 0$, and $j \in [k-1]$, if for every j -uniform hypergraph G on V the collection $\mathcal{K}_k(G)$ of all k -subsets of V inducing a clique $K_k^{(j)}$ in G obeys the estimate

$$|E \cap \mathcal{K}_k(G)| \geq d|\mathcal{K}_k(G)| - \eta|V|^k.$$

One then defines for every k -uniform hypergraph F

$$\pi_j(F) = \sup\{d \in [0, 1]: \text{for every } \eta > 0 \text{ and } n \in \mathbb{N} \text{ there exists an } F\text{-free,} \\ (d, \eta, j)\text{-dense, } k\text{-uniform hypergraph } H \text{ with } |V(H)| \geq n\}$$

and [13, Proposition 2.5] shows that these densities $\pi_j(\cdot)$ agree with the densities $\pi_{[k]^{(j)}}(\cdot)$ alluded to in the first paragraph of this subsection.

For $j = k - 1$ it is known that every k -uniform hypergraph F satisfies $\pi_{k-1}(F) = 0$, which follows for example from the work in [6]. Theorem 1.2 address the case $j = k - 2$ for $k = 3$ and for general k we obtain the following characterisation.

Theorem 6.1. *For a k -uniform hypergraph F , the following are equivalent:*

- (a) $\pi_{k-2}(F) = 0$.
- (b) *There are an enumeration of the vertex set $V(F) = \{v_1, \dots, v_f\}$ and a k -colouring $\varphi: \partial F \rightarrow [k]$ of the $(k - 1)$ -sets of vertices covered by hyperedges of F such that every hyperedge $e = \{v_{i(1)}, \dots, v_{i(k)}\} \in E(F)$ with $i(1) < \dots < i(k)$ satisfies*

$$\varphi(e \setminus \{v_{i(\ell)}\}) = \ell \quad \text{for every } \ell \in [k]. \tag{6.1}$$

This can be established in the same way as Theorem 1.2, but using the hypergraph regularity lemma for k -uniform hypergraphs. For the corresponding notion of reduced hypergraphs we refer to [13, Definition 4.1] and for guidance on the reduction corresponding to Section 3 above we refer to the part of the proof of [13, Proposition 4.5] presented in Section 4 of that article.

For $j \in [k - 3]$ we believe Theorem 6.1 extends in the natural way, where the k -colouring φ in part (b) is replaced by a $\binom{k}{j+1}$ -colouring of the $(j + 1)$ -sets covered by an edge of F and condition (6.1) is replaced by a statement to the effect that the edges of F are rainbow and mutually order-isomorphic when one takes these colours into account.

For $j = 0$ such a characterisation leads to k -partite k -uniform hypergraphs F and, hence, such a result renders a common generalisation of Erdős' result from [1] and Theorem 6.1 and we shall return to this in the near future.

Despite this progress the problem to describe for an arbitrary (asymmetric) antichain $\mathcal{A} \subseteq \wp([k])$ the k -uniform class $\{F: \pi_{\mathcal{A}}(F) = 0\}$ remains challenging. In the 3-uniform case the investigation of $\{F: \pi_{\clubsuit}(F) = 0\}$ and $\{F: \pi_{\spadesuit}(F) = 0\}$, where $\clubsuit = \{1, 23\}$ and $\spadesuit = \{12, 13\}$, shows that algebraic structures enter the picture and this is currently work in progress of the authors.

We close this section with the following questions that compares $\pi_{\clubsuit}(F) = \pi_1(F)$ with $\pi(F)$ for 3-uniform hypergraphs.

Question 6.2. Is $\pi_1(F) < \pi(F)$ for every 3-uniform hypergraph F with $\pi(F) > 0$?

Roughly speaking, this question has an affirmative answer, if no 3-uniform hypergraph F with positive Turán density has an extremal hypergraph H that is uniformly dense with respect to large vertex sets $U \subseteq V(H)$ (see also [3, Problem 7] for a related assertion). In light of the fact, that all known extremal constructions for such 3-uniform hypergraphs F are obtained from blow-ups or iterated blow-ups of smaller hypergraphs, which fail to be $(d, \eta, 1)$ -dense for all $d > 0$ and sufficiently small $\eta > 0$, the answer to Question 6.2 might be affirmative. Recalling that $\pi(F) = \pi_0(F)$ may suggest many generalisations of Question 6.2 to k -uniform hypergraphs F of the form: *For which F do we have $\pi_j(F) < \pi_i(F)$ for $0 \leq i < j < k$?* At this point this is only known for $i = 0$ and $j = k - 1$ and Question 6.2 is the first interesting open case.

6.2. Hypergraphs with uniformly vanishing density. Definition 1.5 admits a straightforward generalisation to k -uniform hypergraphs: one just replaces all occurrences of the word “hypergraph” by “ k -uniform hypergraph” and all occurrences of the number 3 by k .

The sequence of ternary hypergraphs generalises to a sequence $(T_n^{(k)})_{n \in \mathbb{N}}$ of k -uniform hypergraphs that might be called *k -ary* and are defined as follows. The vertex set of $T_n^{(k)}$ is $[k]^n$ and given k vertices $\bar{x}_1, \dots, \bar{x}_k$, say with coordinates $\bar{x}_i = (x_{i1}, \dots, x_{in})$ for $i \in [k]$ one looks at the least number $m \in [n]$ for which $x_{1m} = \dots = x_{km}$ fails and declares $\{\bar{x}_1, \dots, \bar{x}_k\}$ to be an edge of $T_n^{(k)}$ if and only if $\{x_{1m}, \dots, x_{km}\} = [k]$ holds. The proof of Theorem 1.7 (and of Lemma 5.3) generalises in the following way (see [4]).

Theorem 6.3. *A k -uniform hypergraph F on ℓ vertices is frequent if, and only if it is a subhypergraph of the k -ary hypergraph $T_\ell^{(k)}$ on k^ℓ vertices. \square*

Some further questions concerning frequent hypergraphs arise naturally and below we discuss a few of them.

In the context of 3-uniform hypergraphs one may use three sets instead of one set in the definition of d -dense (see Definition 1.5 (a)) and this leads to a question that is somewhat different from the one answered by Theorem 1.7. This happens because the – perhaps on first sight expected – analogue of (3.2) does not hold. More explicitly, we say that a sequence $\vec{H} = (H_n)_{n \in \mathbb{N}}$ of 3-uniform hypergraphs with $v(H_n) \rightarrow \infty$ as $n \rightarrow \infty$ is $(d, \bullet\bullet)$ -dense for a function $d: (0, 1) \rightarrow (0, 1)$ provided that for every $\eta > 0$ there is some $n_0 \in \mathbb{N}$ such that for every $n \geq n_0$ and all choices of $X, Y, Z \subseteq V(H_n)$ with $|X||Y||Z| \geq \eta|V(H_n)|^3$ there are at least $d(\eta)|X||Y||Z|$ ordered triples $(x, y, z) \in X \times Y \times Z$ with $xyz \in E(H_n)$. Besides, a 3-uniform hypergraph F is called $\bullet\bullet$ -frequent if for every function $d: (0, 1) \rightarrow (0, 1)$ and every $(d, \bullet\bullet)$ -dense sequence $\vec{H} = (H_n)_{n \in \mathbb{N}}$ of 3-uniform hypergraphs there exists an $n_0 \in \mathbb{N}$ with $F \subseteq H_n$ for every $n \geq n_0$.

The relation of this concept to being d -dense is as follows: If a sequence \vec{H} of 3-uniform hypergraphs is $(d, \bullet\bullet)$ -dense, then, by looking only at the case $X = Y = Z$ in the definition above, one sees that \vec{H} is also d -dense. On the other hand, being d -dense does not even imply being $(d', \bullet\bullet)$ -dense for any function d' . As an example we mention that the sequence of ternary hypergraphs fails to be $(d, \bullet\bullet)$ -dense for every $d: (0, 1) \rightarrow (0, 1)$.

As a corollary of Theorem 1.7 subhypergraphs of ternary hypergraphs are $\bullet\bullet$ -frequent, but the converse implication may not hold. This leads to the following intriguing problem.

Problem 6.4. Characterise $\bullet\bullet$ -frequent 3-uniform hypergraphs.

Similar to studying $\pi_j(\cdot)$ for k -uniform hypergraphs for every $j < k$ one may study dense sequences with respect to different uniformities. More precisely, for a given integer $j \in [k - 1]$ and a function $d: (0, 1) \rightarrow (0, 1)$ we say that a sequence $\vec{H} = (H_n)_{n \in \mathbb{N}}$ of k -uniform hypergraphs with $v(H_n) \rightarrow \infty$ as $n \rightarrow \infty$ is (d, j) -dense if for every $\eta > 0$ there is an $n_0 \in \mathbb{N}$ such that for every $n \geq n_0$ and every j -uniform hypergraph G on $V(H_n)$ with $|\mathcal{K}_k(G)| \geq \eta |V(H_n)|^k$ the estimate

$$|E(H_n) \cap \mathcal{K}_k(G)| \geq d(\eta) |\mathcal{K}_k(G)|$$

holds. Moreover, a k -uniform hypergraph F is defined to be j -frequent if for every function $d: (0, 1) \rightarrow (0, 1)$ and every (d, j) -dense sequence $\vec{H} = (H_n)_{n \in \mathbb{N}}$ of k -uniform hypergraphs there exists an $n_0 \in \mathbb{N}$ with $F \subseteq H_n$ for every $n \geq n_0$. In particular, 1-frequent is the same as frequent in the sense of Theorem 6.3.

Similar as discussed above the k -ary hypergraphs show that there is a subtle difference between $(d, 1)$ -dense sequences and $(d, [k]^{(1)})$ -dense sequences (where we take k sets instead of one set). However, for $j \geq 2$ one can follow the argument presented in the proof of [13, Proposition 2.5] to show that a k -uniform hypergraph F is j -frequent if and only if it is $[k]^{(j)}$ -frequent (defined in the obvious way). As a result one can show that every k -uniform hypergraph F is $(k - 1)$ -frequent by following the inductive proof on the number of edges of the counting lemma for hypergraphs. This leaves open to characterise the j -frequent hypergraphs for $j \in [2, k - 2]$.

Finally, we mention that one may also consider (d, \mathcal{A}) -dense sequences of hypergraphs for asymmetric antichains \mathcal{A} and characterising \mathcal{A} -frequent hypergraphs is widely open.

REFERENCES

- [1] P. Erdős, *On extremal problems of graphs and generalized graphs*, Israel J. Math. **2** (1964), 183–190. MR0183654 (32 #1134) ↑1, 6.1
- [2] P. Erdős, *Problems and results on graphs and hypergraphs: similarities and differences*, Mathematics of Ramsey theory, Algorithms Combin., vol. 5, Springer, Berlin, 1990, pp. 12–28. MR1083590 ↑1.2, 1.2

- [3] P. Erdős and V. T. Sós, *On Ramsey-Turán type theorems for hypergraphs*, *Combinatorica* **2** (1982), no. 3, 289–295, DOI [10.1007/BF02579235](https://doi.org/10.1007/BF02579235). MR698654 (85d:05185) ↑1, 1.2, 1.2, 5.1, 6.1
- [4] P. Frankl and V. Rödl, *Some Ramsey-Turán type results for hypergraphs*, *Combinatorica* **8** (1988), no. 4, 323–332, DOI [10.1007/BF02189089](https://doi.org/10.1007/BF02189089). MR981890 ↑1.2, 1.2, 1.2, 5, 5.1, 5.2, 6.2
- [5] P. Frankl and V. Rödl, *Extremal problems on set systems*, *Random Structures Algorithms* **20** (2002), no. 2, 131–164, DOI [10.1002/rsa.10017.abs](https://doi.org/10.1002/rsa.10017.abs). MR1884430 (2002m:05192) ↑2
- [6] Y. Kohayakawa, V. Rödl, and J. Skokan, *Hypergraphs, quasi-randomness, and conditions for regularity*, *J. Combin. Theory Ser. A* **97** (2002), no. 2, 307–352, DOI [10.1006/jcta.2001.3217](https://doi.org/10.1006/jcta.2001.3217). MR1883869 (2003b:05112) ↑6.1
- [7] W. T. Gowers, *Quasirandomness, counting and regularity for 3-uniform hypergraphs*, *Combin. Probab. Comput.* **15** (2006), no. 1-2, 143–184, DOI [10.1017/S0963548305007236](https://doi.org/10.1017/S0963548305007236). MR2195580 (2008b:05175) ↑2
- [8] Y. Kohayakawa, B. Nagle, V. Rödl, and M. Schacht, *Weak hypergraph regularity and linear hypergraphs*, *J. Combin. Theory Ser. B* **100** (2010), no. 2, 151–160, DOI [10.1016/j.jctb.2009.05.005](https://doi.org/10.1016/j.jctb.2009.05.005). MR2595699 (2011g:05215) ↑1.1
- [9] B. Nagle, A. Poerschke, V. Rödl, and M. Schacht, *Hypergraph regularity and quasi-randomness*, *Proceedings of the Twentieth Annual ACM-SIAM Symposium on Discrete Algorithms*, SIAM, Philadelphia, PA, 2009, pp. 227–235. MR2809322 ↑2, 2
- [10] Chr. Reiher, V. Rödl, and M. Schacht, *On a Turán problem in weakly quasirandom 3-uniform hypergraphs*, available at [arXiv:1602.02290](https://arxiv.org/abs/1602.02290). Submitted. ↑1, 1.1, 2
- [11] ———, *Embedding tetrahedra into quasirandom hypergraphs*, *J. Combin. Theory Ser. B* **121** (2016), 229–247, DOI [10.1016/j.jctb.2016.06.008](https://doi.org/10.1016/j.jctb.2016.06.008). MR3548293 ↑1, 1.1
- [12] ———, *Some remarks on π_{\aleph}* , *Connections in Discrete Mathematics*, Cambridge University Press, Cambridge, available at [arXiv:1602.02299](https://arxiv.org/abs/1602.02299). To appear. ↑1, 1.1, 3
- [13] ———, *On a generalisation of Mantel’s theorem to uniformly dense hypergraphs*, *Int. Math. Res. Not. IMRN*, DOI [10.1093/imrn/rnx017](https://doi.org/10.1093/imrn/rnx017). To appear. ↑1, 1.1, 3, 4, 6.1, 6.1, 6.2
- [14] V. Rödl, *On universality of graphs with uniformly distributed edges*, *Discrete Math.* **59** (1986), no. 1-2, 125–134, DOI [10.1016/0012-365X\(86\)90076-2](https://doi.org/10.1016/0012-365X(86)90076-2). MR837962 (88b:05098) ↑
- [15] V. Rödl and M. Schacht, *Regular partitions of hypergraphs: regularity lemmas*, *Combin. Probab. Comput.* **16** (2007), no. 6, 833–885. MR2351688 (2008h:05083) ↑2, 2
- [16] ———, *Regular partitions of hypergraphs: counting lemmas*, *Combin. Probab. Comput.* **16** (2007), no. 6, 887–901. MR2351689 (2008j:05238) ↑2
- [17] P. Turán, *Eine Extremalaufgabe aus der Graphentheorie*, *Mat. Fiz. Lapok* **48** (1941), 436–452 (Hungarian, with German summary). MR0018405 (8,284j) ↑1

FACHBEREICH MATHEMATIK, UNIVERSITÄT HAMBURG, HAMBURG, GERMANY

E-mail address: Christian.Reiher@uni-hamburg.de

E-mail address: schacht@math.uni-hamburg.de

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, EMORY UNIVERSITY, ATLANTA, USA

E-mail address: rodl@mathcs.emory.edu