

Algebraic Flow Theory of Infinite Graphs ^{*†}

B. MIRAFTAB^a, M.J. MOGHADAMZADEH^b

^a*Fachbereich Mathematik, Universität Hamburg, Bundesstraße 55, 20146 Hamburg, Germany*

^b*Department of Mathematical Sciences, Sharif University of Technology, Tehran, Iran*

P.O. Box 64615-334,

babak.miraftab@uni-hamburg.de

javad_mz123@yahoo.com

Abstract

A problem by Diestel is to extend algebraic flow theory of finite graphs to infinite graphs with ends. In order to pursue this problem, we define an A -flow and non-elusive H -flow for arbitrary graphs and for abelian Hausdorff topological groups H and compact subsets $A \subseteq H$. We use these new definitions to extend several well-known theorems of flows in finite graphs to infinite graphs.

1 Introduction

The concept of flow is a main topic in graph theory and has various applications, as e.g. in electric networks. Algebraic flow theory for finite graphs is well studied, see [10, 11, 12, 14, 18]. But when it comes to infinite graphs, much less is known. There are some results for electrical networks, see [1, 7, 8, 9], but not for group-valued flows. In fact Diestel's problem [8, Problem 4.27] to extend flow theory to infinite graphs is still widely open. Here we are doing a first step towards its solution.

In Section 2, we give our main definition for flows in infinite graphs. Roughly speaking, a flow is a map from the edge set of a graph to an abelian Hausdorff topological group such that the sum over all edges in each finite cut is trivial. With this in mind, we shall extend the following theorems of finite graphs:

- A finite graph has a non-elusive \mathbb{Z}_2 -flow if and only if its degrees are even.
- A finite cubic graph has a non-elusive \mathbb{Z}_4 -flow if and only if it is 3-edge-colorable.

^{*}*Key Words:* Contraction, Flow, Infinite graph.

[†]2010 *Mathematics Subject Classification:* 05C21, 05C63, 22A05.

- Every finite graph containing a Hamilton cycle has a non-elusive \mathbb{Z}_4 -flow.

Our main tool to prove these results is Theorem 5, which offers some kind of compactness method to extend results for finite graphs to infinite graphs of arbitrary degree, i.e. that need not be locally finite. However it is worth remarking that not all theorems about flows in finite graphs have a straightforward analogue in the infinite case: a finite cubic graph G has a non-elusive \mathbb{Z}_3 -flow if and only if G is bipartite, see [7, Proposition 6.4.2]. This is wrong for infinite graphs. Figure 1.1 shows a cubic bipartite graph without any non-elusive \mathbb{Z}_3 -flow. Even further restrictions on the ends of that graph, e.g. requiring them to have edge- or vertex-degree 3, fails in our example. (For more about the ends of a graph and the topological setting, we refer readers to [8] and the references therein.)




Figure 1.1

In the Section 4, we define the concept of tension for infinite graphs. Heuristically, a tension is a map from the edge set of a graph to an abelian Hausdorff topological group such that the sum over all edges in each finite cycle is trivial.

2 Preliminaries

We refer readers to [7], for the standard terminology and notations in this paper. A 1-way infinite path is called a *ray*, a 2-way infinite path is a *double ray*, and the subrays of a ray or double ray are its tails. Two rays in a graph $G = (V, E)$ are equivalent if no finite set of vertices separates them. This is an equivalence relation whose classes are the *ends* of G . Now, consider a locally finite graph G as one-dimensional CW complex and compactify G by using the Freudenthal compactification method. We denote this new topological space by $|G|$, for more on $|G|$, see [6] and [8]. Let D be a subset of edges of G . We denote the closure of the point set $\cup_{d \in D} d$ in $|G|$ by \overline{D} . A *circle* in $|G|$ is a homeomorphic image of the unit circle S^1 . Analogously an *arc* in $|G|$ is a homeomorphic image of the closed interval $[0, 1]$. We denote the *cut space*, *finite cut space*, *topological cycle space* and *finite cycle space* of a graph G by $\mathcal{B}(G)$, $\mathcal{B}_{\text{fin}}(G)$, $\mathcal{C}(G)$ and $\mathcal{C}_{\text{fin}}(G)$, respectively. For more details about the equivalent definitions of topological cycle space and its properties, see [7, 8]. Note that $\mathcal{B}(G)$ is a vector space over \mathbb{Z}_2 . We now define the degree of an end of the graph G . The *edge-degree* of an end ω is the maximum number of edge-disjoint rays in ω . In addition, let D be a subset of the set of edges of G . Then we say that an end ω is *D-even* if there exists a finite vertex set S so that for all finite vertex sets $S' \supseteq S$ it holds that the maximal number of edge-disjoint arcs from S' to ω contained in \overline{D} is even. If D is all the edges of G , we

remove D from the notation and we only say that ω has an even edge-degree. For more about the degree of ends, see [3, 4]. The following theorem describes the elements of the cycle space for locally finite graphs. For the proof, see [7, Theorem 8.5.10] and [2, Theorem 5].

Theorem 1. *Let $G = (V, E)$ be a locally finite connected graph. Then an edge set $D \subseteq E$ lies in $\mathcal{C}(G)$ if and only if one of the following equivalent statements holds*

- (i) D meets every finite cut in an even number of edges.
- (ii) Every vertex and every end of G is D -even.

Let us review some notions of the compactness method for locally finite graphs. Suppose that v_0, v_1, \dots is an enumeration of V . We define $S_n = v_0, \dots, v_n$, for every $n \in \mathbb{N}$. Put G_n for the minor of G obtained by contracting each component of $G \setminus S_n$ to a vertex. Note that we delete any loop, but we keep multiple edges. The vertices of G_n outside S_n are called *dummy vertices* of G_n . Let $G = (V, E)$ be a graph. A *directed edge* is an ordered triple (e, x, y) , where $e = xy \in E$. So we can present each edge according to its direction by $\vec{e} = (e, x, y)$ or $\overleftarrow{e} = (e, y, x)$. We use \vec{E} for the set of all oriented edges of G . For two subsets X, Y (not necessarily disjoint) of V and a subset \vec{C} of \vec{E} , we define

$$\vec{C}(X, Y) := \{(e, x, y) \in \vec{C} \mid x \in X, y \in Y, x \neq y\}.$$

It is worth mentioning that we can express every finite cut of our graph by a pair (X, Y) , where X and $Y = V \setminus X$ are two subsets of the vertices. Thus for every finite cut (X, Y) , we have an oriented cut $\vec{E}(X, Y)$. The set $\vec{\mathcal{B}}_{\text{fin}}(G)$ denotes the set of all oriented finite cuts i.e. $\vec{\mathcal{B}}_{\text{fin}}(G) = \{\vec{E}(A, B) \mid (A, B) \in \mathcal{B}_{\text{fin}}(G)\}$. Let H be an abelian group(not necessarily finite). Then we denote all maps from \vec{E} to H such that $f(\vec{e}) = -f(\overleftarrow{e})$ for every non-loop $\vec{e} \in \vec{E}$ by $H^{\vec{E}}$ and we introduce the following notation only for $\vec{E}(A, B) \in \vec{\mathcal{B}}_{\text{fin}}(G)$

$$f(A, B) := \sum_{\vec{e} \in \vec{E}(A, B)} f(\vec{e}).$$

Also $H^{\vec{\mathcal{B}}_{\text{fin}}(G)}$ denotes all maps from $\mathcal{B}_{\text{fin}}(G)$ to H such that $f(A, B) = -f(B, A)$ for every $\vec{E}(A, B) \in \vec{\mathcal{B}}_{\text{fin}}(G)$. Let us review the definition of group-valued flows for finite graphs¹. A nowhere-zero H -flow of the graph G is a map $f \in H^{\vec{E}}$ with the following properties:

- C1: $f(\vec{e}) \neq 0$, for every $\vec{e} \in \vec{E}$.
- C2: $f(\{v\}, V) = 0$ for all vertices v of V .²

A drawback of the above definition is that it depends on degrees of vertices. So it is meaningless whenever our graph has a vertex with infinite degree. To concoct this definition, we switch every

¹Our approach is due to [7].

²This condition is known as the Kirchhoff's law.

vertex with every oriented cut of our graph in the condition C2 which means $f(A, B) = 0$ for all finite cuts (A, B) . More precisely we have the following definition:

Definition 1: Let H be an abelian Hausdorff topological group and let A be a compact subset of H . We define $\sigma: H^{\vec{E}} \rightarrow H^{\vec{\mathcal{B}}_{\text{fin}}(G)}$ such that

$$f(X, Y) = \sum_{\vec{e} \in \vec{E}(X, Y)} f(\vec{e})$$

for any finite oriented cut $\vec{E}(X, Y)$. Let M be a subset of $\vec{\mathcal{B}}_{\text{fin}}(G)$. Then we say that G has an A -flow with respect to M if $F_M = \{f \in A^{\vec{E}} \mid \sigma(f)(\vec{E}(X, Y)) = 0 \text{ for every } \vec{E}(X, Y) \in M\}$ is not empty and we say that G has an A -flow if G has an A -flow with respect to $\vec{\mathcal{B}}_{\text{fin}}(G)$. If f is an A -flow and $A \subseteq H \setminus \{0\}$, then we also call f a *non-elusive H -flow*.

Definition 2: With the above notation, suppose that G has an A -flow, where $H = \mathbb{Z}$ with the discrete topology and $A = \{-(k-1), \dots, k-1\} \setminus \{0\}$. Then we say that G has a k -flow.

If a graph G has more than one component, then G has an A -flow if and only if each of its components does. That is why we restrict ourselves to connected graphs from now on. So let G be a connected graph.

It is worth mentioning that if G is locally finite, then this definition coincides with the one in Section 4.3 of [8] for abelian groups. If the graph G is locally finite, then using the compactness method, we can generalize almost all theorems of finite flow theory to infinite.

Definition 3: Let $M = \{C_1, \dots, C_t\}$ be a finite subset of $\mathcal{B}_{\text{fin}}(G)$. Then we define a multigraph G_M according to M . Each cut $C_i \in M$ belongs to a bipartition (A_i, B_i) of V such that C_i are the $A_i - B_i$ edges. The vertices of G_M are the words $X_1 \cdots X_t$, where $X_i \in \{A_i, B_i\}$ for $i = 1, \dots, t$ in such a way that $\bigcap_{i=1}^t X_i \neq \emptyset$. Between two vertices $X_1 \cdots X_t$ and $X'_1 \cdots X'_t$ of G_M , there is an edge for each edge between $\bigcap_{i=1}^t X_i$ and $\bigcap_{i=1}^t X'_i$. We say that G_M is obtained from G by *contracting* with respect to M .

Remark 2. Let G be an infinite graph and let M be a finite subset of $\mathcal{B}_{\text{fin}}(G)$. Then throughout this paper we always first consider G_M without its loops and then we apply the corresponding result for finite multigraphs. Now we extend the flow in an arbitrary way to the loops. This is possible, as no cut of G contains a loop and so the assignments of loops do not influence whether our function is a flow or not.

Remark 3. The definition of G_M leads to a map $\phi: G \rightarrow G_M$, where every vertex u of G is mapped to a unique word $V_u \in V(G_M)$, it is contained in. Indeed, looking at each finite cut in M , we can construct the unique word $X_1 \cdots X_t$ in such a way that every X_i contains u , for each $i \in \{1, \dots, t\}$ and so $u \in \bigcap_{i=1}^t X_i$. We notice that each edge of G induces an edge of G_M . Indeed, it is not hard to see that ϕ defines a bijective map on the set of edges. Also, it is worth mentioning that $\phi^{-1}(U_1) \cap \phi^{-1}(U_2) = \emptyset$ for every two vertices U_1 and U_2 of $V(G_M)$. Thus the vertex set of G_M is a partition of V .

Our compactness method is more general than the ordinary compactness method for locally finite graphs as mentioned above. When the graph G is locally finite, for each G_n ³, we can choose a suitable subset M of the set of finite cuts such that G_M coincides with G_n .

3 Flows on Infinite Graphs

First, we start with the following lemma.

Lemma 4. *Let G be a graph and M be a finite subset of $\mathcal{B}_{\text{fin}}(G)$. Then we have $M \subseteq \mathcal{B}(G_M) \subseteq \mathcal{B}_{\text{fin}}(G)$.*

Proof. First, we show that $M \subseteq \mathcal{B}(G_M)$. Let $C = E(A, B) \in M$. Then consider the set of all words containing A and do the same for all words containing B , say \mathcal{A} and \mathcal{B} , respectively. The sets \mathcal{A} and \mathcal{B} form a partition of G_M and so we have C as a cut of G_M . Note that \mathcal{A} and \mathcal{B} are not empty, since every $uv \in C$ induces vertices $V_u \in \mathcal{A}$ and $V_v \in \mathcal{B}$. Now, assume that $C = E(A, B) \in \mathcal{B}(G_M)$. We deduce from Remark 3 that the edges between A and B in G_M are those between $\phi^{-1}(A)$ and $\phi^{-1}(B)$. Hence $(\phi^{-1}(A), \phi^{-1}(B))$ forms a partition of G and so C is a finite cut of G . \square

The following theorem plays a vital role in this paper and is a basic key to generalize flow theory of finite to infinite graphs.

Theorem 5. *Let G be a graph and H be an abelian Hausdorff topological group with compact subset A . Then G has an A -flow if and only if G_M has an A -flow for every finite subset M of $\mathcal{B}_{\text{fin}}(G)$.*

Proof. First, assume that G has an A -flow. By Lemma 4, every finite cut of G_M belongs to $\mathcal{B}_{\text{fin}}(G)$. So every A -flow of G is an A -flow of G_M . In particular, G_M has some A -flow. For the backward implication, since H is a topological group, the sets $H^{\vec{E}}$ and $H^{\vec{\mathcal{B}_{\text{fin}}}(G)}$ are endowed with the product topology. Let $M = \{C_1, \dots, C_t\}$ be a subset of $\vec{\mathcal{B}_{\text{fin}}}(G)$. We define $\sigma_i: H^{\vec{E}} \rightarrow H$ by $\sigma_i(f) = \sum_{e \in C_i} f(e)$. Since the sum operation is a continuous map, σ_i is continuous for each i . Therefore $\sigma_i^{-1}(0)$ is a closed subspace in $H^{\vec{E}}$, as H is Hausdorff. On the other hand, by Tychonoff's theorem (see [13, Theorem 37.3]), $A^{\vec{E}}$ is compact and so is $\sigma_i^{-1}(0) \cap A^{\vec{E}}$. It is clear that $F_M = \bigcap_{i=1}^t \sigma_i^{-1}(0) \cap A^{\vec{E}}$ and so F_M is compact. Since G_M has an A -flow, by definition, the set $F_{\mathcal{B}(G_M)}$ is not empty. Lemma 4 implies that F_M is not empty. Hence the intersection of every finite family of $F_{\{C_i\}}$ with $C_i \in \mathcal{B}_{\text{fin}}(G)$ is not empty. Since $A^{\vec{E}}$ is compact, we deduce that $F_{\mathcal{B}_{\text{fin}}(G)} = \bigcap_{C_i \in \mathcal{B}_{\text{fin}}(G)} F_{\{C_i\}}$ is not empty, see [13, Theorem 26.9]. Thus G has an A -flow. \square

³For definition of G_n , see Preliminaries.

For finite graphs, the existence of a nowhere-zero H -flow does not depend on the structure of H but only on its order, see [7, Corollary 6.3.2]. In the next corollary, we show that the same is true for infinite graphs.

Corollary 6. *Let H and H' be two finite abelian groups with equal order. Then G has a non-elusive H -flow if and only if G has a non-elusive H' -flow.*

Proof. We note that H and H' are endowed by the discrete topologies and so they are compact. Suppose G has a non-elusive H -flow. By Theorem 5, for every finite subset M of $\mathcal{B}_{\text{fin}}(G)$, the multigraph G_M has a non-elusive H -flow. We notice that G_M might have infinitely many loops. Since each loop appears twice, we can ignore them and so we only care the rest of edges which are finite. Thus we are able to apply [7, Corollary 6.3.2] and conclude that every G_M has a non-elusive H' -flow. Again, it follows from Theorem 5 that G has a non-elusive H -flow. The other direction follows from the symmetry of the statement. \square

There is a direct connection between k -flows and non-elusive \mathbb{Z}_k -flows in finite graphs which was discovered by Tutte, see [19]. In the next corollary, we use Theorem 5 and show that having a k -flow and a non-elusive \mathbb{Z}_k -flow are equivalent in infinite graphs.

Corollary 7. *A graph admits a k -flow if and only if it admits a non-elusive \mathbb{Z}_k -flow.*

Proof. The canonical homomorphism $\mathbb{Z} \rightarrow \mathbb{Z}_k$ implies the forward implication. For the converse, assume that G has a non-elusive \mathbb{Z}_k -flow. By Theorem 5, for every finite subset M of $\mathcal{B}_{\text{fin}}(G)$, the multigraph G_M has a non-elusive \mathbb{Z}_k -flow. We consider \mathbb{Z}_k with the discrete topology. It follows from Theorem 5 and [19] that every G_M has a k -flow. Again, we invoke Theorem 5 and we conclude that G has a k -flow. \square

Next up, we study non-elusive \mathbb{Z}_m -flows for some special values of m . First, we study non-elusive \mathbb{Z}_2 -flows for locally finite graphs. It is worth mentioning that if G is an arbitrary infinite graph and G has a non-elusive \mathbb{Z}_2 -flow, then one can see that all finite cuts of G are even and vice versa. First we need a notation. Suppose that $G = (V, E)$ is a graph and F is a subset of E . We define the indicate function $\delta_F: E \rightarrow \mathbb{Z}_2$ in the following way:

$$\delta_F(e) := \begin{cases} 1 & \text{for } e \in F \\ 0 & \text{for } e \notin F \end{cases}$$

Theorem 8. *Let $G = (V, E)$ be a locally finite graph and let F be a subset of E . Then δ_F is a \mathbb{Z}_2 -flow if and only if $F \in \mathcal{C}(G)$.*

Proof. First suppose that δ_F is a \mathbb{Z}_2 -flow. It is not hard to see that every vertex and every end of G is F -even. So it follows from Theorem 1 that F belongs to the cycle space of G . For the backward implication, since $F \in \mathcal{C}(G)$, we are able to invoke Theorem 1 and conclude that every vertex and every end of G is F -even. Thus it implies that δ_F is a \mathbb{Z}_2 -flow. \square

It is not hard to see that if a cubic graph G has a non-elusive \mathbb{Z}_3 -flow, then G is bipartite. For a cubic graph G , having a non-elusive \mathbb{Z}_3 -flow is equivalent to having an orientation of G in such a way that for every vertex v of G all incident edges of v are either directed outward or directed inward and moreover all assignments are one. Let G be a graph as depicted on Figure 1.1. Consider orientations with the above property. So we have two cases. In each case, we have a finite cut whose sum of assignments is not zero, see Figure 3.0.1.



Figure 3.0.1

Hence, we propose this question: When does a cubic graph has a non-elusive \mathbb{Z}_3 -flow?

Recently, Thomassen used S^1 and R_3 in flow theory of finite multigraphs and investigated the connection of such flows with \mathbb{Z}_3 -flows for finite multigraphs, see [16]. Now let us review these notations here. Let $G = (V, E)$ be a finite multigraph without loops. Then an S^1 -flow is the same as a flow whose flow values are complex numbers with absolute value 1. But we first choose an orientation for each $e \in E$ and then we assign elements of S^1 on the edges.⁴ Let R_k denote the set of k -th roots of unity, that is, the solutions to the equation $z^k = 1$.

Lemma 9. [16, Proposition 1] *Let G be a finite multigraph without loops. Then (i) and (ii) below are equivalent, and they imply the statement (iii)*

- (i) G has a non-elusive \mathbb{Z}_3 -flow.
- (ii) G has an R_3 -flow.
- (iii) G has an S^1 -flow.

If G is cubic, the three statements are equivalent, and G satisfies (i), (ii), (iii) if and only if G is bipartite.

We generalize Lemma 9. We replace the condition *cubic* with an edge dominating set H of vertices such that the degree of every vertex of H is 3. A subset H of vertices is an *edge dominating set* if every edge of the graph has an end vertex in H .

Lemma 10. *Let G be a finite multigraph without loops with a connected edge dominating set U of vertices such that every vertex of U has degree 3. Then the following three statements are equivalent.*

⁴We follow this approach only for the next three results.

- (i) G has a non-elusive \mathbb{Z}_3 -flow.
- (ii) G has an R_3 -flow.
- (iii) G has an S^1 -flow.

Proof. By Lemma 9, it is enough to show that (iii) \Rightarrow (ii). One may suppose that G has at least one edge. Assume that G has an S^1 -flow, say f . Choose an edge of G , say uv with $u \in U$. We notice that U contains at least two vertices. Because if U has only one vertex, then every vertex in $V \setminus U$ would have degree one and so we are not able to have an S^1 -flow. Let $f(uv) = z_1 \in S^1$. Since f is an S^1 -flow, there are $z_2, z_3 \in S^1$ such that $z_1 + z_2 + z_3 = 0$. Note that z_2 and z_3 are unique. Let w be a neighbour of u in U . Then degree of w is three and so the values of f on edges incident to w lie exactly in the set $\{z_1, z_2, z_3\}$. Since U is connected and meets every edge of G , we know that f assigns z_1, z_2 or z_3 to every edge of G . Thus f is a $\{z_1, z_2, z_3\}$ -flow on G . Since there is a bijection between $\{z_1, z_2, z_3\}$ and R_3 , we find an R_3 -flow for G . \square

Now, we are ready to answer this question: When does a cubic graph have a non-elusive \mathbb{Z}_3 -flow?

Theorem 11. *If G is a cubic graph, then the following statements are equivalent.*

- (i) G has a non-elusive \mathbb{Z}_3 -flow.
- (ii) G has an R_3 -flow.
- (iii) G has an S^1 -flow.

Proof. (i) \Rightarrow (ii) It follows from Theorems 5 and 9 that for every finite subset M of $\mathcal{B}_{\text{fin}}(G)$, the multigraph G_M has an R_3 -flow. So by Theorem 5, G has an R_3 -flow. (ii) \Rightarrow (iii) is trivial. (iii) \Rightarrow (i) By Theorem 5, the multigraph G_M has an S^1 -flow. We notice that as we mentioned in Remark 2, we ignore all loops of G_M . Let U be the set of all vertices that are incident with an edge from a cut of M . We note that U is finite. We add some paths of G to $G[U]$ until we get a connected graph N . Note that it suffices to take only finitely many paths, i.e. we may assume that N is finite. Let S_N be the set of vertices of N and assume that G_N is obtained by contracting the components of $G \setminus S_N$ to dummy vertices, similar to constructing of multigraph G_n for the compactness method. Obviously, S_N is an edge dominating set of vertices of G_N and moreover the degree of each vertex of N is 3. We notice that G_N has an S^1 -flow, as G has an S^1 -flow. By Lemma 10, the multigraph G_N has a non-elusive \mathbb{Z}_3 -flow. Since every element of M is a cut of G_N , the multigraph G_M has a non-elusive \mathbb{Z}_3 -flow. We invoke Theorem 5 and we conclude that G has a non-elusive \mathbb{Z}_3 -flow. \square

Next, we study non-elusive \mathbb{Z}_4 -flows.

Theorem 12. *Let $G = (V, E)$ be a locally finite graph. Then G has a non-elusive \mathbb{Z}_4 -flow if and only if E is the union of two elements of its topological cycle space.*

Proof. First, suppose that G has a non-elusive \mathbb{Z}_4 -flow. By Corollary 6, we can assume that G has a non-elusive $\mathbb{Z}_2 \oplus \mathbb{Z}_2$ -flow, say f . Set $E_i = \{e \in E(G) \mid \pi_i(f(e)) \neq 0\}$ for $i = 0, 1$, where π_1 and π_2 are the projection maps on the first and second coordinates, respectively. Since G has a $\mathbb{Z}_2 \oplus \mathbb{Z}_2$ -flow, each finite cut of G meets E_i evenly. We now invoke Theorem 1 and conclude that every E_i belongs to the topological cycle space of G , for $i = 0, 1$. For the backward implication, let $G = G_1 \cup G_2$ with $E(G_i) \in \mathcal{C}(G)$, for $i = 1, 2$. It follows from Theorem 1 and Theorem 8 that each G_i has a non-elusive \mathbb{Z}_2 -flow. Thus we can find a non-elusive $\mathbb{Z}_2 \oplus \mathbb{Z}_2$ -flow and by Theorem 6, we are done. \square

3.1 Edge-coloring for infinite graphs

If G is a cubic finite graph, then the conditions of having a non-elusive \mathbb{Z}_4 -flow and 3-edge-colorability of G are equivalent, but this is not true for infinite graphs. Let G be a graph as depicted in Figure 3.1.1. Suppose that G has a non-elusive \mathbb{Z}_4 -flow. On the other hand, we are able to contract the graph G to the Petersen graph. But it is known that the Petersen graph is not 3-edge-colorable. So we deduce that the Petersen graph does not admit a non-elusive \mathbb{Z}_4 -flow and it implies that G does not have a non-elusive \mathbb{Z}_4 -flow, either.

It seems that the notion of edge-coloring is not suitable for a characterization of when an infinite graph with ends admits a k -flow, but that a generalization of edge-colorability (“*semi- k -edge-colorability*”, to be defined below) is. We only need a definition of generalized edge-coloring for cubic graphs here which implies the existence of a non-elusive \mathbb{Z}_4 -flow. Hence we will define this concept under the name of semi-edge-coloring. Next, we define semi-edge-coloring for k -regular graphs where k is an odd number and we show that this definition for cubic graphs is equivalent to having a non-elusive \mathbb{Z}_4 -flow.




Figure 3.1.1

Before defining this new edge-colorability, note that we can define k -flow axiomatically for finite graphs. Our objective is to show that every graph which has a k -flow is a contraction of a cubic graph which has a k -flow. In order to show this, we need a definition. We call a map \mathcal{F} from the class of all finite graphs to \mathbb{Z}_2 a “*Boolean functor of having the property P* ” if G has the property P if and only if $\mathcal{F}(G) = 1$. For instance, having a k -flow is a Boolean functor. We denote it by \mathcal{F} . We notice that if $\mathcal{F}(G) = 1$ for a given graph G , then $\mathcal{F}(H) = 1$, where H is a contraction of G . The property of admitting a k -flow or equivalently a non-elusive \mathbb{Z}_k -flow can be characterized as follows:

Theorem 13. *Let $k > 2$ be an odd number and \mathcal{F} be the Boolean functor of having a non-elusive \mathbb{Z}_k -flow for every finite graph and \mathcal{F}' be another Boolean functor which satisfy the following three properties.*

- (i) \mathcal{F} and \mathcal{F}' are the same for cubic graphs.
- (ii) If $\mathcal{F}'(G) = 1$, then $\mathcal{F}'(H) = 1$ for every contraction⁵ H of G .
- (iii) If $\mathcal{F}'(G) = 1$, then there is a cubic graph H with $\mathcal{F}'(H) = 1$ such that G is a contraction of H .

Then \mathcal{F} and \mathcal{F}' are equal.

Proof. Assume that $\mathcal{F}(G) = 1$, for a given finite graph G and let f be a non-elusive \mathbb{Z}_k -flow of G . We now introduce a cubic graph H' such that G is a contraction of H' and $\mathcal{F}'(H') = 1$. Our strategy is to switch all vertices with degrees at least four with vertices with degrees at most three and then we eliminate all vertices with degrees two. Let $v \in V(G)$ with degree at least four. Suppose that the sum of values of two edges e_1 and e_2 that are incident with v is 0. First, we add a new vertex u . Then we separate these two edges from v and we join e_1 and e_2 to u . In other words, the degree of u is two and e_1 and e_2 are incident to u . So the degree of the vertex v reduces by 2 in the new graph. Now we assume that there are two edges which are incident to v and the sum of their flows is not 0, say e_1 and e_2 . We separate e_1 and e_2 from v with a new vertex u like in the previous case and join the new vertex u to v . In other words, we substitute these two edges with a claw i.e. $K_{1,3}$. We continue this process for all vertices of G until $\Delta(G) \leq 3$ is obtained. We call the new graph H . Next we are going to replace the vertices of degree two with $K_{3,3}$. Suppose that e_1 and e_2 are incident edges to the vertex v with $\deg(v) = 2$. Without loss of generality, we can assume that the orientation of e_1 is toward v . It is not hard to see that there are $a, b \in \mathbb{Z}_k \setminus \{0\}$ such that $f(e_1) + a + b = 0$. Consider the complete bipartite graph $K_{3,3}$. Since the degree of each vertex is 3, we can find a non-elusive \mathbb{Z}_k -flow on $K_{3,3}$ such that the value of all edges belong to the set $\{f(e_1), a, b\}$. Suppose that $e = v_1v_2$ of $K_{3,3}$ with the value $f(e_1)$ and the orientation from v_1 to v_2 . We remove the edge $e = v_1v_2$ from $K_{3,3}$ and the vertex v of G . Now, we join the edge e_1 to v_2 and e_2 to v_1 . We repeat this process for all vertices of degree 2. Hence we obtain a cubic graph H' with a non-elusive \mathbb{Z}_k -flow and so $\mathcal{F}(H') = \mathcal{F}'(H') = 1$. Therefore $\mathcal{F}'(G) = 1$, as G is contraction of H' . Hence we have shown that if $\mathcal{F}(G) = 1$, then $\mathcal{F}'(G) = 1$.

Now, if $\mathcal{F}'(G) = 1$, then the condition (iii) gives us an H with $\mathcal{F}'(H) = 1$, which G is a contraction of H and so $\mathcal{F}(H) = 1$. Thus we deduce that $\mathcal{F}(G) = 1$, as desired. \square

The proof of the preceding theorem implies the following corollary. We note that as we mentioned before “contraction” used in this paper is different from “minor”, see the footnote.

Corollary 14. *Every graph admitting a k -flow is a contraction of a cubic graph which has a k -flow.*

⁵The contracted vertex sets need not be connected.

We now are ready to state the definition of semi-edge-colorability which was mentioned above.

Definition 4: Let k be a positive integer. A *semi- k -edge-coloring* of a graph G is a map from $E(G)$ to $\{1, 2, \dots, k\}$, with the property that for every finite cut C of G , if the number of edges of C with the color i is c_i , then the all numbers c_1, \dots, c_k have the same parity. A graph G is *semi- k -edge-colorable* if G has a semi- k -edge-coloring.

We use flows to characterize semi-edge-colorings. First, let $V = \oplus_{i=1}^{k-1} \mathbb{Z}_2$ be the vector space over \mathbb{Z}_2 and e_i for $i = 1, \dots, k-1$ be the standard basis. Set $\mathcal{A} = \{e_1, \dots, e_{k-1}, \sum_{i=1}^{k-1} e_i\}$. Note that \mathcal{A} is compact with the discrete topology. We now use the notation of [16] and we state the following lemma.

Lemma 15. *Let G be a finite graph and k be a positive integer. Then with the above notation, the following statements are equivalent.*

- (i) G is semi- k -edge-colorable.
- (ii) G has an \mathcal{A} -flow.

Proof. The one to one correspondence between the color set $\{c_1, \dots, c_k\}$ and $\{e_1, \dots, e_{k-1}, \sum_{i=1}^{k-1} e_i\}$ induces a bijection between the set of semi- k -edge-colorings and the set of \mathcal{A} -flows. \square

Immediately, Theorem 8 implies the following remark:

Remark 16. Let G be a locally finite graph and k be a positive integer. Then the following statements are equivalent.

- (i) G is semi- $2k$ -edge-colorable.
- (ii) The degrees of all vertices and ends of G are even.
- (iii) G has a non-elusive \mathbb{Z}_2 -flow.

Our objective is to show that every 3-edge-colorable finite graph is a contraction of a cubic 3-edge-colorable graph. In order to show this, we show that the definition of semi-edge-coloring is the only definition which is compatible with the three properties of Theorem 13 for finite graphs, but instead of cubic graphs, we can have k -regular graphs. In other words, the Boolean functor having semi- k -edge-colorability is the unique Boolean functor which satisfies the conditions (i)-(iii) of Theorem 13.

Theorem 17. *Let k be an odd number, let \mathcal{F} be the Boolean functor of a finite graph being semi- k -edge-colorable and let \mathcal{F}' be another Boolean functor which satisfy the three following properties*

- (i) \mathcal{F} and \mathcal{F}' are the same for k -regular graphs.
- (ii) If $\mathcal{F}'(G) = 1$, then $\mathcal{F}'(H) = 1$ for every contraction H of G .

- (iii) If $\mathcal{F}'(G) = 1$, then there is a finite k -regular H such that G is a contraction of H with $\mathcal{F}'(H) = 1$.

Then \mathcal{F} and \mathcal{F}' are equal.

Proof. Assume that a graph G is semi- k -edge-colorable and so $\mathcal{F}(G) = 1$. We construct a k -regular graph H such that $\mathcal{F}'(H) = 1$ and moreover G is a contraction of H . We notice that as we mentioned before the contracted vertex sets need not be connected. Let v be an arbitrary vertex of G . If $\deg(v) = 2n$, then each color appears an even number of times, as the number of colors is odd and the degree is even. Thus we are able to form pairs $P_i = \{e_i^1, e_i^2\}$ of edges with the same color. Consider a k -edge-coloring of the complete graph K_{k+1} . We delete an edge e of the color of the edges of P_i , join the edges in P_i to the end vertices of e in K_{k+1} and we denote by L the union of $K_{k+1} \setminus \{e\}$ with edges e_i^1 and e_i^2 . We do this for every P_i for $i = 1, \dots, n$. If $\deg(v) = 2n + 1$, then each color appears an odd number of times. From each color, we choose an incident edge of v . We separate them and we attach them to a new vertex u . We notice that the degree of u is k . Thus the number of colors appears in the rest of incident edges of v is even. Again we are able to pair these edges. We do same for the paired edges as above. Hence the vertex v is replaced by the union of some copies of L and the vertex u . Now, we do the same for every vertex of G . Finally, we obtain a k -edge-colorable k -regular graph H which contains G as a contraction. Hence since $\mathcal{F}'(H) = \mathcal{F}(H) = \mathcal{F}(G) = 1$, we can conclude that $\mathcal{F}'(G) = 1$. If $\mathcal{F}'(G) = 1$, then we note that semi-edge-colorability is preserved by contraction. So the first and third conditions imply that $\mathcal{F}(G) = 1$. \square

The proof of the preceding theorem implies the following corollary.

Corollary 18. *Every 3-edge-colorable finite graph is a contraction of a k -regular 3-edge-colorable graph, where $k \geq 3$ is an odd number.*

In finite cubic graphs, the existence of non-elusive \mathbb{Z}_4 -flows and 3-edge-colorability are equivalent, see [7, Proposition 6.4.5]. Next, we generalize this fact to infinite graphs.

Theorem 19. *Let G be a graph. Then G has a non-elusive \mathbb{Z}_4 -flow if and only if G is semi-3-edge-colorable.*

Proof. First, assume that G is semi-3-edge-colorable. Since every contraction of G is semi-3-edge-colorable, we conclude that every G_M is semi-3-edge-colorable, for every finite subset M of $\mathcal{B}_{\text{fin}}(G)$. It follows from Corollary 18 that there is a cubic graph \widetilde{G}_M in such a way that \widetilde{G}_M is 3-edge-colorable and moreover G_M is a contraction of \widetilde{G}_M . We invoke Part (ii) of [7, Proposition 6.4.5] and we conclude that \widetilde{G}_M has a non-elusive \mathbb{Z}_4 -flow, as \widetilde{G}_M is a cubic graph and it is 3-edge-colorable. We notice that by the definition of \widetilde{G}_M , we deduce that G_M has a non-elusive \mathbb{Z}_4 -flow. Now, by Theorem 5, we deduce that G has a non-elusive \mathbb{Z}_4 -flow. For the forward implication, by Corollary 6, G has a non-elusive $\mathbb{Z}_2 \oplus \mathbb{Z}_2$ -flow, say f . We define a semi-3-edge-coloring $c: E(G) \rightarrow \mathbb{Z}_2 \oplus \mathbb{Z}_2 \setminus \{(0, 0)\}$ by $c(e) = f(e)$. Let F be a finite cut of G . Then since f is a non-elusive $\mathbb{Z}_2 \oplus \mathbb{Z}_2$,

the map f sums up to zero on the edges of F . In particular, the sum of all assignments of edges with the value $(1, 0)$ is zero. Thus we are able to deduce that the parity of every color of each edge of F is the same. Thus G is semi-3-edge-colorable, as desired. \square

3.2 Hamiltonicity

A graph is *Eulerian* if it is connected and all vertices have even degree. We call a finite graph *supereulerian* if it has a spanning Eulerian subgraph.

Lemma 20. *Every finite supereulerian graph has a non-elusive \mathbb{Z}_4 -flow.*

Proof. Let G be a supereulerian graph. Then by Corollary 6, it is enough to show that G has a non-elusive $\mathbb{Z}_2 \oplus \mathbb{Z}_2$ -flow. Let C be a spanning Eulerian subgraph of G . The degree of every vertex of G in C is even. Thus the constant function with the value $(0, 1)$ is a non-elusive \mathbb{Z}_2 -flow in C . We denote this $\mathbb{Z}_2 \oplus \mathbb{Z}_2$ -flow by F . Let e_1, \dots, e_k be an enumeration of the edges outside C . Suppose that u_i and v_i are the end vertices of e_i . Since C is a spanning Eulerian subgraph of G , we can find a walk P_i in C between u_i and v_i . We define a new flow F_i by assigning $(1, 0)$ to every edge of $P_i \cup \{e_i\}$. Note that $P_i \cup \{e_i\}$ is an Eulerian subgraph. So F_i is a $\mathbb{Z}_2 \oplus \mathbb{Z}_2$ -flow of G , for $i = \{1, \dots, k\}$. Then $\sum_{i=1}^k F_i + F$ is a $\mathbb{Z}_2 \oplus \mathbb{Z}_2$ -flow, too. Now, we claim that $\sum_{i=1}^k F_i + F$ is a non-elusive $\mathbb{Z}_2 \oplus \mathbb{Z}_2$ -flow. It is enough to show that $\sum_{i=1}^k F_i + F$ is non-zero for an arbitrary edge of C , as the value of e_i is $(1, 0)$, for $i = 1, \dots, k$. Since the second component of the map $\sum_{i=1}^k F_i + F$ is always 1 for every edge of C , the flow $\sum_{i=1}^k F_i + F$ is non-elusive and the claim is proved, as desired. \square

Remark 21. Catlin [5] showed that every finite 4-edge-connected graph is supereulerian. Thus it follows from Lemma 20 that every finite 4-edge-connected graph has a 4-flow. This result has been proved by Jaeger [10].

A *Hamiltonian circle* is a circle containing every vertex of an infinite graph. It is worth mentioning that every Hamiltonian circle contains all vertices and all ends precisely once.

Corollary 22. *Every graph containing a Hamiltonian circle has a non-elusive \mathbb{Z}_4 -flow.*

Proof. Let C be a Hamiltonian circle of $|G|$ and M be a finite subset of $\mathcal{B}_{\text{fin}}(G)$. Also, let $\phi : G \rightarrow G_M$ be the map which is defined in Remark 3. Then $\phi(C)$ is a spanning Eulerian subgraph of G_M and so G_M is supereulerian. It follows from Lemma 20 that G_M has a non-elusive \mathbb{Z}_4 -flow for every finite subset M of $\mathcal{B}_{\text{fin}}(G)$. Now, we invoke Theorem 5 and we conclude that G has a non-elusive \mathbb{Z}_4 -flow. \square

3.3 Conjectures

In the study of flow theory one main point of interest is the connection to the edge-connectivity. For example, if a finite graph is 2-edge-connected, then it has a non-elusive \mathbb{Z}_6 -flow, see [14].

Next up, we show that the connection between edge-connectivity and the existence of a non-elusive flow for infinite graphs admits exactly the same connection as for finite graphs.

Corollary 23. *If n -edge-connectivity implies the existence of an m -flow for finite graphs, then this implication holds for infinite graphs as well.*

Proof. Let M be a finite subset of $\mathcal{B}_{\text{fin}}(G)$. Note that since G is n -edge-connected, the multi-graph G_M is n -edge-connected. By assumption, the graph G_M has an m -flow. Now, we invoke Theorem 5 and conclude that G has an m -flow. \square

As a corollary of Remark 21 and Corollary 23, we obtain the following.

Corollary 24. *Every 4-edge-connected graph has a 4-flow.* \square

There are some famous conjectures in finite flow theory such as the four-flow conjecture and the three-flow conjecture. If these conjectures hold true for finite graphs, then they are true for infinite graphs and vice versa.

Five-flow conjecture: Every 2-edge-connected graph has a 5-flow.

Four-flow conjecture: Let G be a bridgeless graph. If for every finite subset M of $\mathcal{B}_{\text{fin}}(G)$, G_M does not contain the Petersen graph as a topological minor, then G has a non-elusive 4-flow.

Three-flow conjecture: Every 4-edge-connected graph has a 3-flow.

In 1961, Seymour [14] has shown that every finite bridgeless graph has a 6-flow. Immediately, Theorem 5 implies the following theorem.

Theorem 25. *Every bridgeless graph G has a 6-flow.* \square

4 Tension of Infinite Graphs

Another concept related to flows is tension. Let $G = (V, E)$ be a finite graph and K be a group that is not necessarily abelian. We call a map $f : \vec{E} \rightarrow K$ a K -tension if f satisfies $\sum_{e \in \vec{C}} f(e) = 0$ for every directed cycle \vec{C} of G . We note that we sum up the assignments of edges with respect to a cyclic order. If $f(\vec{e}) \neq 0$ for every $\vec{e} \in \vec{E}$ then G has a *nowhere-zero K -tension*. Since we are studying cycles, it does not matter where we start, and moreover, if G has a K -tension, the choice of our edge orientation is irrelevant, as every element of K has its inverse. So we can define our K -tension for infinite graphs G in an analogous manner with superseding finite cuts with finite cycles in the definition of a flow. Suppose that K is a Hausdorff topological group with a compact subset A of K . We define $\sigma : K^{\vec{E}} \rightarrow K^{\mathcal{C}_{\text{fin}}(G)}$ such that $\sigma(f)(\vec{C}) = \sum_{\vec{e} \in \vec{C}} f(\vec{e})$ for any finite oriented cycle C . Let M be a subset of $\mathcal{C}_{\text{fin}}(G)$. Then we say that G has an A -tension with respect to M if $F_M = \{f \in A^{\vec{E}} \mid \sigma(f)(C) = 0 \text{ for every } C \in M\}$ is not empty and we say that G has an

A-tension if G has an A -tension with respect to $\mathcal{C}_{\text{fin}}(G)$. If f is an A -tension and $K \setminus \{0\}$ then we say that f is a *non-elusive K -tension*.

If f is an A -flow and $A \subseteq K \setminus \{0\}$, then we also call f a *non-elusive H -flow*. Now, a natural question arises: When does an infinite graph G have a non-elusive K -tension? At first glance, it seems that we can use the concept of dual graphs. A pair of *dual graphs* is a pair of graphs (G, G^*) such that there is a bijection $\phi : E(G) \rightarrow E(G^*)$ with the property that a finite set $A \subseteq E(G)$ is the edge set of a cycle if and only if $\phi(A)$ is a bond (minimal edge cut) in G^* . Thomassen [15, Theorem 3.2] showed that a 2-connected graph G has a dual graph if and only if G is planar and any two vertices of G are separated by a finite edge cut. Moreover if G^* is a dual graph of G and $A \subseteq E(G)$, then G^*/A^* is a dual graph of $G - A$, see [17, Lemma 9.11]. For more details regarding the concept of duality with the topological approach, see [3]. We denoted by G^*/A^* the graph obtained from G^* by contracting all edges of A^* . Hence, for defining the similar graph like G_M in Definition 3, we have to delete some edges from G and it holds true only for planar graphs where every two of its vertices are separated by a finite edge cut. In the next theorem, we delete edges for an arbitrary graph and show that the above argument is still true.

Theorem 26. *Let G be a graph and \mathcal{C} be a finite subset of $\vec{\mathcal{C}}_{\text{fin}}(G)$. Then G has a non-elusive K -tension if and only if every finite subset \mathcal{C} of $\vec{\mathcal{C}}_{\text{fin}}(G)$ has a non-elusive K -tension.*

Proof. Set

$$F_{\mathcal{C}} = \{f \text{ is a } K\text{-tension of } G \mid f \text{ is a nowhere-zero } K\text{-tension with respect to } \mathcal{C}\}.$$

Then $F_{\mathcal{C}}$ is not empty for any finite subset \mathcal{C} of $\vec{\mathcal{C}}_{\text{fin}}(G)$. So using an analogous method as in the proof of Theorem 5, we conclude that G has a non-elusive K -tension. \square

Acknowledgements. The authors are deeply grateful to the referees for careful reading. Also the authors are grateful to Pascal Gollin, Matthias Hamann and Peter Christian Heinig for their comments.

References

- [1] R. Aharoni, E. Berger, A. Georgakopoulos, A. Perlstein, P. Sprüssel, The max-flow min-cut theorem for countable networks, J.Comb. Theory B 101 (2011), 1–17.
- [2] E. Berger, H. Bruhn, Eulerian edge sets in locally finite graphs, Combinatorica 31 (2011), 21–38.
- [3] H. Bruhn, R. Diestel, Duality in infinite graphs, Combin. Probab. Comput. 15 (2006), no. 1-2, 75–90.
- [4] H. Bruhn, M. Stein, On end degrees and infinite cycles in locally finite graphs, Combinatorica 27 (2007), 269–291 .
- [5] P.A. Catlin, A reduction method to find spanning Eulerian subgraphs, J. Graph Theory 12 (1988), 29–45.

- [6] R. Diestel, End spaces and spanning trees, *J. Comb. Theory, Ser. B*, 96 (2006), 846–854 .
- [7] R. Diestel, *Graph theory*, 4th edition, Springer-Verlag, (2010).
- [8] R. Diestel, Locally finite graphs with ends: a topological approach, *Discrete Math.* 311 (2011), no. 15, 1423–1447.
- [9] A. Georgakopoulos, Uniqueness of electrical currents in a network of finite total resistance, *J. Lond. Math. Soc. (2)* 82 (2010), no. 1, 256–272.
- [10] F. Jaeger, Flows and generalized coloring theorems in graphs, *J. Combin. Theory Ser. B* 26 (1979), no. 2, 205–216.
- [11] F. Jaeger, On circular flows in graphs, in *finite and infinite sets*, *Colloq. Math. Soc. János Bolyai* 37, North-Holland, Amsterdam, 1984, 391–402.
- [12] L.M. Lovasz, C. Thomassen, Y. Wu, C.Q. Zhang, *J. Combin. Theory Ser. B* 103 (2013), no. 5, 587–598.
- [13] J. Munkres, *Topology: a first course*, Prentice-Hall, Inc., Englewood Cliffs, N.J., (2000).
- [14] P.D. Seymour, Nowhere-zero 6-flows, *J. Combin. Theory Ser. B* 30 (1981), no. 2, 130–135.
- [15] C. Thomassen, Duality of infinite graphs, *J. Combin. Theory Ser. B* 33 (1982), no. 2, 137–160.
- [16] C. Thomassen, Group flow, complex flow, unit vector flow, and the $(2 + \epsilon)$ -flow conjecture, *J. Combin. Theory Ser. B* 108 (2014), 81–91.
- [17] C. Thomassen, Planarity and duality of finite and infinite graphs, *J. Combin. Theory Ser. B* 29 (1980), no. 2, 244–271.
- [18] C. Thomassen, The weak 3-flow conjecture and the weak circular flow conjecture, *J. Combin. Theory Ser. B* 102 (2012), no. 2, 521–529.
- [19] W.T. Tutte, A contribution to the theory of chromatic polynomial, *Canad. J. Math.*, 6 (1954), pp. 80–91.