

# FORCING QUASIRANDOMNESS WITH TRIANGLES

CHRISTIAN REIHER AND MATHIAS SCHACHT

**ABSTRACT.** We study *forcing pairs* for *quasirandom graphs*. Chung, Graham, and Wilson initiated the study of families  $\mathcal{F}$  of graphs with the property that if a large graph  $G$  has approximately homomorphism density  $p^{e(F)}$  for some fixed  $p \in (0, 1]$  for every  $F \in \mathcal{F}$ , then  $G$  is quasirandom with density  $p$ . Such families  $\mathcal{F}$  are said to be *forcing*. Several forcing families were found over the last three decades and characterising all bipartite graphs  $F$  such that  $(K_2, F)$  is a forcing pair is a well-known open problem in the area of quasirandom graphs, which is closely related to Sidorenko's conjecture. In fact, most of the known forcing families involve bipartite graphs only.

We consider forcing pairs containing the triangle  $K_3$ . In particular, we show that if  $(K_2, F)$  is a forcing pair, then so is  $(K_3, F^\triangleright)$ , where  $F^\triangleright$  is obtained from  $F$  by replacing every edge of  $F$  by a triangle (each of which introduces a new vertex). For the proof we first show that  $(K_3, C_4^\triangleright)$  is a forcing pair, which strengthens related results of Simonovits and Sós and of Conlon et al.

## §1. INTRODUCTION

The systematic study of quasirandom graphs was initiated by Thomason [52, 53] and Chung, Graham, and Wilson [12] and over the last 30 years many generalisations and extensions to directed graphs [23], tournaments [7], hypergraphs [1, 4–6, 16, 26, 33, 35, 54], set systems [8], permutations [19], groups [22], subsets of cyclic groups and finite fields [10, 55], and sparse graphs [11, 15, 31] were established by several researchers (see, e.g., the surveys [32, 34] for a more detailed discussion). Roughly speaking, a given discrete structure is quasirandom if it shares important properties with a “truly random” structure of the same size. In the context of graphs this is made precise by mimicking the uniform edge distribution of the random graph  $G(n, p)$ .

**Definition 1.1.** For  $\varepsilon > 0$  and  $p \in (0, 1]$  we say a graph  $G = (V, E)$  is  $(\varepsilon, p)$ -*quasirandom*, if for all subsets  $X, Y \subseteq V$  we have

$$|e_G(X, Y) - p|X||Y|| \leq \varepsilon|V|^2,$$

where edges contained in the intersection  $X \cap Y$  are counted twice in  $e_G(X, Y)$ .

In the context of quasirandom graphs we often consider sequences of graphs  $\vec{G} = (G_n)_{n \in \mathbb{N}}$  with  $|V(G_n)| \rightarrow \infty$ . Then we may say that the sequence  $\vec{G}$  is  $p$ -*quasirandom* if for every  $\varepsilon > 0$

---

2010 *Mathematics Subject Classification.* 05C80.

*Key words and phrases.* quasirandom graphs, forcing pairs.

Research of both author was supported by ERC Consolidator Grant 724903.

all but finitely many members of the sequence are  $(\varepsilon, p)$ -quasirandom. For a simpler discussion, we sometimes say that a graph  $G$  is  $p$ -quasirandom without any reference to the sequence, by which we mean that some sufficiently large graph  $G$  is  $(\varepsilon, p)$ -quasirandom for some small unspecified  $\varepsilon > 0$ . If the density  $p$  is of no particular importance, then we may just say  $G$  is quasirandom.

A large part of the theory on quasirandom graphs concerns equivalent characterisations of  $p$ -quasirandom graph sequences. Early results in that direction implicitly appeared in [2, 3, 20, 40] and Chung, Graham, and Wilson [12] gave six alternative characterisation. Since then many more such characterisations were found (see, e.g., [9, 16, 28, 29, 41–43, 46–49, 56]).

Here we focus on characterisations that rely on the densities of graph homomorphisms of given graphs  $F$  into large graphs  $G$ , which we denote by  $t(F, G)$ . Let us recall that a pair of graphs  $(F_1, F_2)$  is said to be *forcing* if for every  $p \in (0, 1]$  and  $\varepsilon > 0$  there is some  $\delta > 0$  such that the following holds: if a graph  $G$  satisfies

$$t(F_1, G) \geq (1 - \delta)p^{e(F_1)} \quad \text{and} \quad t(F_2, G) \leq (1 + \delta)p^{e(F_2)} \quad (1.1)$$

then it is  $(\varepsilon, p)$ -quasirandom. This notion goes back to [12] and has frequently been discussed in the literature. The probably most classical example of such a family is the pair  $(K_2, C_4)$ . The statement that the pair  $(K_2, F)$  is forcing for every bipartite graph  $F$  that does not happen to be a forest, called the *forcing conjecture*, can be traced back to Skokan and Thoma [49] (see also [14]). It has been the subject of intensive study that led to its verification in various cases. For the most recent contributions to the forcing conjecture and the closely related conjectures of Erdős and Simonovits [45] and of Sidorenko [44] we refer to [14, 17, 18, 25, 30, 36, 50].

Until recently all known forcing pairs contained bipartite graphs only. In fact, already Chung, Graham, and Wilson [12] noted that  $(K_2, K_3)$  (and also  $(K_{1,2}, K_3)$ ) is not a forcing pair, by giving an example of  $n$ -vertex graphs  $G$  with all vertices having degree close to  $n/2$  and with  $t(K_3, G) \approx 1/8$ , but containing independent sets and cliques of size  $\lfloor n/4 \rfloor$ . However, it was shown by Simonovits and Sós [47] that such a situation can be avoided by appealing to the *hereditary* nature of quasirandom graphs, i.e., if  $G = (V, E)$  is  $p$ -quasirandom, then the induced subgraphs  $G[U]$  are  $p$ -quasirandom for linear sized subsets  $U \subseteq V$ . Simonovits and Sós then showed that requiring



$$t(F, G[U]) = \left(1 \pm \delta \frac{|V|}{|U|}\right) p^{e(F)} \quad (1.2)$$

for a given graph  $F$  with at least one edge and for all  $U \subseteq V$  forces  $G$  to be  $p$ -quasirandom. Recently, Conlon, Hàn, Person, and Schacht [16] (see also [24]) observed that condition (1.2) gives rise to a forcing pair  $(F, M_F)$  for an appropriate graph  $M_F$  depending on  $F$ .

We study forcing pairs involving triangles. For that case it was shown in [16] that the pair  $(K_3, M)$  is forcing, where  $M$  denotes the line graph of the 3-dimensional Boolean cube, depicted on the right. The idea behind the proof is roughly as follows: Three successive applications of the Cauchy–Schwarz inequality yield  $t(M, G) \geq t(K_3, G)^8$  for any graph  $G = (V, E)$ . On the other hand, the assumption (1.1) for  $(K_3, M)$  tells us  $t(K_3, G) \gtrsim p^3$  and  $t(M, G) \lesssim p^{24}$  for some real  $p \in (0, 1]$ . Consequently, an approximate equality must hold in each of these three steps, and it may be argued that this is in turn only possible if for all subsets  $A, B, C \subseteq V$  we have

$$\Delta(A, B, C) \approx p^3 |A| |B| |C|,$$

where  $\Delta(A, B, C)$  denotes the number of triangles with a vertex in  $A$ , a vertex in  $B$ , and a vertex in  $C$ . This yields (1.2) for  $F = K_3$  and the Simonovits–Sós theorem implies that  $G$  is  $p$ -quasirandom.



Our main result shows that the same effect as above can be achieved with two applications of the Cauchy–Schwarz inequality only. This implies that the pair  $(K_3, C_4^\triangleright)$  is forcing, where  $C_4^\triangleright$  is obtained from the 4-cycle  $C_4$  where every edge is replaced by a triangle (each of which introduces a new vertex), i.e., the graph shown on the left. As we shall explain in more detail below, we have  $t(C_4^\triangleright, G) \geq t(K_3, G)^4$  for all graphs  $G$ , and if approximate equality holds for some graph  $G$ , then it satisfies the assumption of the following theorem, which weakens the assumption of the Simonovits–Sós theorem in the triangle case.

**Theorem 1.2.** *For every  $p \in (0, 1]$  and  $\varepsilon > 0$  there is an  $\eta > 0$  such that any graph  $G = (V, E)$  satisfying*

$$\Delta(A, B, V) = p^3 |A| |B| |V| \pm \eta |V|^3 \quad (1.3)$$

*for all  $A, B \subseteq V$  is  $(\varepsilon, p)$ -quasirandom.*

The following corollary renders the aforementioned connection between condition (1.3) and  $t(C_4^\triangleright, G) \approx t(K_3, G)^4$  and strengthens the result of Conlon et al. that  $(K_3, M)$  is a forcing pair.

**Corollary 1.3.** *The pair  $(K_3, C_4^\triangleright)$  is forcing.*

As it turns out Corollary 1.3 applied in the context of weighted graphs allows the following general result, which is our main result on forcing pairs involving triangles.

**Corollary 1.4.** *If  $(K_2, F)$  is a forcing pair, then so is  $(K_3, F^\triangleright)$ .*

**Organisation.** We prove Theorem 1.2 in Section 2 and Corollary 1.3 in Section 3. In Section 4 we switch to the analytical language of graphons and prove Corollary 1.4. We conclude by recording some further observations and problems for future research in Section 5.

## §2. THE TWO SETS CONDITION


The proof of Theorem 1.2 is based on the regularity method for graphs. This means that we use a regularity lemma and a counting lemma in order to reduce the problem at hand to a somewhat different one that speaks about a certain “reduced graph.” In the present situation we need to conceive this reduced graph as a *weighted graph*. Such objects may also be regarded as symmetric matrices with entries from the unit interval. The precise statement we shall require is stated as Lemma 2.1 below. The interested reader may check that this lemma could conversely also be deduced from Theorem 1.2.

**Lemma 2.1.** *Given any real numbers  $p \in (0, 1]$  and  $\varepsilon > 0$  there is a real  $\delta > 0$  such that the following holds: Let  $(d_{ij})_{i,j \in [t]} \in [0, 1]^{t \times t}$  be a symmetric matrix such that for all distinct indices  $i, j \in [t]$  we have*

$$d_{ij} \sum_{k \in [t]} d_{ik} d_{jk} = (p^3 \pm \delta)t. \quad (2.1)$$

Then  $d_{ij} = p \pm \varepsilon$  holds for all  $i, j \in [t]$ .

*Proof.* Throughout the proof we work with the hierarchy  $\delta \ll \varrho \ll p, \varepsilon$  for some auxiliary chosen constant  $\varrho$ .



Since the sum on the left-hand side of (2.1) is at most  $t$ , we have  $d_{ij} \geq p^3 - \delta \geq p^3/2$  for all  $i, j \in [t]$ . The main idea is to choose three indices  $x, y, z \in [t]$  such that the difference  $d_{xy} - d_{xz}$  is as large as possible. Set  $a = d_{xy}$  and  $b = d_{xz}$ . The maximality of  $a - b$  entails

$$d_{yi} \geq b \text{ and } d_{zi} \leq a \text{ for all } i \in [t]. \quad (2.2)$$

By (2.1) we have

$$a \cdot \sum_{i \in [t]} d_{xi} d_{yi} \leq (p^3 + \delta)t \quad \text{and} \quad b \cdot \sum_{i \in [t]} d_{xi} d_{zi} \geq (p^3 - \delta)t,$$

whence

$$\sum_{i \in [t]} d_{xi} (a d_{yi} - b d_{zi}) \leq 2\delta t.$$

Now (2.2) yields

$$a d_{yi} - b d_{zi} = a(d_{yi} - b) + b(a - d_{zi}) \geq ((d_{yi} - b) + (a - d_{zi}))p^3/2 \geq 0$$

for all  $i \in [t]$ , so it follows that

$$\sum_{i \in [t]} (d_{yi} - b) + \sum_{i \in [t]} (a - d_{zi}) \leq 8p^{-6}\delta t, \quad (2.3)$$

where neither sum on the left-hand side is negative. Thus there is an index  $w \in [t]$  with  $(d_{wy} - b) + (a - d_{wz}) \leq 8p^{-6}\delta$ . In particular, the numbers  $\bar{a} = d_{wz} \leq a$  and  $\bar{b} = d_{wy} \geq b$  (see (2.2)) satisfy

$$|a - \bar{a}| + |\bar{b} - b| \leq 8p^{-6}\delta. \quad (2.4)$$

Applying (2.1) to the pairs  $(w, y)$  and  $(w, z)$  in place of  $(i, j)$  and subtracting the resulting estimates we obtain

$$\left| \sum_{i \in [t]} d_{wi}(\bar{a}d_{zi} - \bar{b}d_{yi}) \right| \leq 2\delta t.$$

Thus the triangle inequality and (2.3) lead to

$$\begin{aligned} |a\bar{a} - b\bar{b}| \cdot p^3 t / 2 &\leq \left| \sum_{i \in [t]} d_{wi}(a\bar{a} - b\bar{b}) \right| \\ &\leq \left| \sum_{i \in [t]} d_{wi}\bar{a}(a - d_{zi}) \right| + \left| \sum_{i \in [t]} d_{wi}\bar{b}(d_{yi} - b) \right| + \left| \sum_{i \in [t]} d_{wi}(\bar{a}d_{zi} - \bar{b}d_{yi}) \right| \\ &\leq 2(1 + 4p^{-6})\delta t, \end{aligned}$$

i.e.,  $|a\bar{a} - b\bar{b}| \leq 4p^{-3}(1 + 4p^{-6})\delta$ . Consequently (2.4) yields

$$p^3(a - b) \leq a^2 - b^2 \leq |a\bar{a} - b\bar{b}| + a|a - \bar{a}| + b|\bar{b} - b| \leq 4p^{-3}(1 + 4p^{-6})\delta + 8p^{-6}\delta$$

and thus  $a - b \leq 4p^{-6}(1 + 2p^{-3} + 4p^{-6})\delta \leq \varrho$ .

Now for any four indices  $i, j, k, \ell \in [t]$  the extremal choice of  $a - b$  gives

$$|d_{ij} - d_{k\ell}| \leq |d_{ij} - d_{i\ell}| + |d_{i\ell} - d_{k\ell}| \leq 2(a - b) \leq 2\varrho.$$

In other words, there is an interval of length  $2\varrho$  containing all the  $d_{ij}$ . In the light of (2.1) and the smallness of  $\varrho$  this interval needs to be contained in  $(p - \varepsilon, p + \varepsilon)$ . Thereby Lemma 2.1 is proved.  $\square$

As we have already said, our proof of Theorem 1.2 depends on Szemerédi's regularity lemma [51], a version of which we would like to state next:

**Lemma 2.2.** *For every positive real number  $\delta$  there is a positive integer  $T$  such that every graph  $G = (V, E)$  admits a partition  $V = V_0 \cup V_1 \cup \dots \cup V_t$  of its vertex set obeying the following conditions:*

- (a)  $t \leq T$ ,  $|V_0| \leq \delta |V|$ , and  $|V_1| = \dots = |V_t| > 0$ .
- (b) For each  $i \in [t]$  there are at most  $\delta t$  many indices  $j \in [t]$  such that the pair  $(V_i, V_j)$  is not  $\delta$ -quasirandom.

Here a pair  $(A, B)$  of nonempty subsets of  $V$ , say with density  $d = \frac{e(A, B)}{|A||B|}$ , is said to be  $\delta$ -quasirandom if we have  $e(X, Y) = d|X||Y| \pm \delta|A||B|$  for all  $X \subseteq A$  and  $Y \subseteq B$ .

The above statement differs in several aspects from the “standard” regularity lemma and we briefly discuss those differences:

- The crucial property obtained for most pairs  $(V_i, V_j)$  is often taken to be something called  $\delta$ -regularity\* rather than  $\delta$ -quasirandomness. These two concepts are known to be equivalent up to polynomial losses in the involved constants, and in fact  $\delta$ -regularity implies  $\delta$ -quasirandomness. Our reason for working with this notion here is that allows a slightly cleaner presentation of the proof.

---

\*A pair  $(A, B)$  of subsets of  $G$  is said to be  $\delta$ -regular if  $|d(X, Y) - d(A, B)| \leq \delta$  holds for all  $X \subseteq A$  and  $Y \subseteq B$  with  $|X| \geq \delta|A|$  and  $|Y| \geq \delta|B|$ .

- Instead of the second condition one usually finds a weaker clause just stating that at most  $\delta t^2$  pairs  $(V_i, V_j)$  fail to be quasirandom in the literature. The above version has also been used and to obtain it, one may apply the standard version of the regularity lemma with some appropriate  $\delta' \ll \delta$  in place of  $\delta$  and then relocate all classes  $V_i$  with  $i > 0$  that violate (b) to  $V_0$ .
- Usually one requires also a lower bound  $t_0$  on the number of vertex classes  $t$  in advance and then one obtains  $T \geq t \geq \min(t_0, |V|)$  rather than just  $T \geq t$  in the first part of (a). The rationale behind this is that in many applications one has no intentions of “looking inside the individual  $V_i$ ,” wherefore it brings certain advantages to have these sets reasonably small. In our current situation, however, even the extreme outcome  $t = 1$  would be useful. In view of (b) it would mean that the pair  $(V_1, V_1)$  is  $\delta$ -quasirandom, and since, provided that  $\delta$  is small,  $V_1$  would be almost all of  $V(G)$ , this is essentially all we need to infer for the proof of Theorem 1.2.

In the course of proving Theorem 1.2 we will also need to be able to count triangles after regularising  $G$ . This will be rendered by the following strong, but well-known, form of the triangle counting lemma.

**Lemma 2.3.** *Let  $A$ ,  $B$ , and  $C$  denote three nonempty subsets of  $V(G)$  for some graph  $G$ . Suppose that the pairs  $(B, C)$ ,  $(C, A)$ , and  $(A, B)$  have densities  $\alpha$ ,  $\beta$ , and  $\gamma$  respectively, and that at least two of these three pairs are  $\delta$ -quasirandom. Then*

$$\Delta(A, B, C) = (\alpha\beta\gamma \pm 2\delta)|A||B||C|.$$

*Proof of Theorem 1.2.* We begin by choosing certain auxiliary constants obeying the hierarchy

$$\eta \ll T^{-1} \ll \delta \ll \varepsilon, p,$$

where  $T$  is the integer obtained by applying the regularity Lemma 2.2 with  $\delta$ . Now let any graph  $G = (V, E)$  satisfying

$$\Delta(A, B, V) = p^3 |A| |B| |V| \pm \eta |V|^3 \tag{2.5}$$

for all  $A, B \subseteq V$  be given. The regularity lemma yields a partition

$$V(G) = V_0 \cup V_1 \cup \dots \cup V_t$$

satisfying the above clauses (a) and (b). For  $i, j \in [t]$  we denote the density of the pair  $(V_i, V_j)$  by  $d_{ij}$ . The assumption (2.5) is only going to be used in the special case  $A, B \in \{V_1, \dots, V_t\}$ . It then discloses the following useful property of the numbers  $d_{ij}$ :

$$d_{ij} \sum_{k \in [t]} d_{ik} d_{jk} = (p^3 \pm 9\delta)t, \tag{2.6}$$

for all  $i, j \in [t]$ . To see this, we consider any two indices  $i, j \in [t]$ . Let  $R_i$  denote the set of all  $k \in [t]$  for which the pair  $(V_i, V_k)$  is not  $\delta$ -quasirandom, let  $R_j$  be defined similarly with respect to  $j$ , and set  $R = R_i \cup R_j$ . Owing to condition (b) from Lemma 2.2 we have  $|R_i| \leq \delta t$  and  $|R_j| \leq \delta t$ , whence  $|R| \leq 2\delta t$ . Let us write  $M = |V_1| = \dots = |V_t|$ .

Then  $Mt = |V| - |V_0| \geq (1 - \delta)|V|$ . As we may assume  $\delta \leq \frac{1}{2}$ , it follows that  $|V| \leq 2Mt$ , whence  $|V_0| \leq 2\delta Mt$ . Now we have

$$\left| \Delta(V_i, V_j, V) - M^3 d_{ij} \sum_{k \in [t]} d_{ik} d_{jk} \right| \leq \Delta(V_i, V_j, V_0) + \sum_{k \in [t]} \left| \Delta(V_i, V_j, V_k) - M^3 d_{ij} d_{ik} d_{jk} \right|.$$

Here the first term may be estimated trivially by

$$\Delta(V_i, V_j, V_0) \leq |V_0| |V_i| |V_j| \leq 2\delta M^3 t.$$

Moreover, for  $k \in [t] \setminus R$  the triangle counting lemma (Lemma 2.3) tells us that

$$\left| \Delta(V_i, V_j, V_k) - M^3 d_{ij} d_{ik} d_{jk} \right| \leq 2\delta M^3,$$

while for  $k \in R$  we still have the obvious bound

$$\left| \Delta(V_i, V_j, V_k) - M^3 d_{ij} d_{ik} d_{jk} \right| \leq M^3.$$

Due to  $|R| \leq 2\delta t$  all this combines to

$$\left| \Delta(V_i, V_j, V) - M^3 d_{ij} \sum_{k \in [t]} d_{ik} d_{jk} \right| \leq 6\delta M^3 t. \quad (2.7)$$

On the other hand, plugging  $A = V_i$  and  $B = V_j$  into (2.5) we learn

$$\left| \Delta(V_i, V_j, V) - p^3 M^2 |V| \right| \leq \eta |V|^3,$$

which in turn yields

$$\left| \Delta(V_i, V_j, V) - p^3 M^3 t \right| \leq p^3 M^2 (|V| - Mt) + \eta |V|^3.$$

In view of

$$M^2 (|V| - Mt) + \eta |V|^3 = M^2 |V_0| + \eta |V|^3 \leq M^3 (2\delta t + 8\eta t^3) \leq M^3 t (2\delta + 8\eta t^2)$$

a suitable choice of  $\eta$  leads to

$$\left| \Delta(V_i, V_j, V) - p^3 M^3 t \right| \leq 3\delta M^3 t.$$

Together with (2.7) this concludes the proof of (2.6).

We may assume that depending on  $\varepsilon$  and  $p$  the constant  $\delta$  has been chosen so small that Lemma 2.1 guarantees  $d_{ij} = p \pm \frac{\varepsilon}{2}$  for all  $i, j \in [t]$ . Let us write  $S$  for the set of all pairs  $(i, j) \in [t]^2$  such that the pair  $(V_i, V_j)$  is not  $\delta$ -quasirandom. Notice that condition (b) from Lemma 2.2 implies  $|S| \leq \delta t^2$ .

Now for any  $A, B \subseteq V$  we have

$$\left| e(A, B) - p |A| |B| \right| \leq \sum_{i=0}^t \sum_{j=0}^t \left| e(A \cap V_i, B \cap V_j) - p |A \cap V_i| |B \cap V_j| \right|. \quad (2.8)$$

Each term on the left-hand side having  $i = 0$ ,  $j = 0$ , or  $(i, j) \in S$  may be bounded from above by  $|V_i| |V_j|$ , so altogether these terms contribute at most

$$|V|^2 - (|V| - |V_0|)^2 + |S| M^2 \leq 2\delta |V|^2 + \delta M^2 t^2 \leq 3\delta |V|^2.$$

Owing to the quasirandomness, each of the remaining terms on the right hand side of (2.8) may be estimated as follows:

$$\begin{aligned} & \left| e(A \cap V_i, B \cap V_j) - p |A \cap V_i| |B \cap V_j| \right| \\ & \leq \left| e(A \cap V_i, B \cap V_j) - d_{ij} |A \cap V_i| |B \cap V_j| \right| + |d_{ij} - p| |V_i| |V_j| \\ & \leq \left( \delta + \frac{\varepsilon}{2} \right) |V_i| |V_j| \end{aligned}$$

So taken together these terms amount to at most  $(\delta + \frac{\varepsilon}{2}) |V|^2$ , and in view of  $\delta \ll \varepsilon$  we finally we arrive at

$$\left| e(A, B) - p |A| |B| \right| \leq (4\delta + \frac{\varepsilon}{2}) |V|^2 \leq \varepsilon |V|^2$$

for arbitrary  $A, B \subseteq V$ . This proves that  $G$  is indeed  $(\varepsilon, p)$ -quasirandom.  $\square$

### §3. PROOF OF COROLLARY 1.3

In this section we deduce Corollary 1.3. The only thing we need to check is the following proposition, which combined with Theorem 1.2 yields the corollary.

**Proposition 3.1.** *Suppose that a graph  $G = (V, E)$  satisfies*

$$t(K_3, G) \geq (1 - \delta)p^3 \quad \text{and} \quad t(C_4^\triangleright, G) \leq (1 + \delta)p^{12}$$

*for some  $p, \delta \in (0, 1]$ . Then*

$$\Delta(A, B, V) = p^3 |A| |B| |V| \pm 8\delta^{1/4} p^3 |V|^3$$

*holds for all  $A, B \subseteq V$ .*


Besides the Cauchy–Schwarz inequality itself the proof will also use the following known and easy to confirm result on situations where equality holds almost.

**Fact 3.2.** *Let  $x_1, \dots, x_n$ ,  $\alpha$ , and  $\nu$  denote any real numbers satisfying*

$$\sum_{i=1}^n x_i = \alpha n \quad \text{and} \quad \sum_{i=1}^n x_i^2 = (\alpha^2 + \nu^2)n.$$

*Then we have  $\sum_{i=1}^m x_i = \alpha m \pm \nu n$  for any  $m = 0, 1, \dots, n$ .*

*Proof of Proposition 3.1.* Let  $G = (V, E)$ ,  $p$ , and  $\delta$  be as in the hypothesis and let  $A, B \subseteq V$  be arbitrary.



We begin the proof by setting up some notation: Let  $S$  denote the graph obtained by gluing two triangles together at a vertex (see the picture on the left). We consider the homomorphism densities of  $K_3$ ,  $S$ , and  $C_4^\triangleright$  and define the real numbers  $a$ ,  $b$ , and  $c$  so as to obey

$$\text{hom}(K_3, G) = ap^3 n^3, \tag{3.1}$$

$$\text{hom}(S, G) = bp^6 n^5, \tag{3.2}$$



and

$$\text{hom}(C_4^{\triangleright}, G) = cp^{12}n^8, \quad (3.3)$$

where  $n = |V|$ . Notice that the assumption translates to  $a \geq 1 - \delta$  and  $c \leq 1 + \delta$ . Given a vertex  $x \in V$  we write  $T_x$  for the number of pairs  $(y, z) \in V^2$  such that  $xyz$  is a triangle in  $G$ . Moreover, for any two vertices  $u, v \in V$  we denote the number of triples  $(x, y, z) \in V^3$  with  $ux, uy, vx, vz, xy, xz \in E$  by  $S_{uv}$ . In terms of these numbers the equations (3.1), (3.2), and (3.3) rewrite as

$$\sum_{x \in V} T_x = ap^3n^3, \quad (3.4)$$

$$\sum_{x \in V} T_x^2 = \sum_{(u,v) \in V^2} S_{u,v} = bp^6n^5, \quad (3.5)$$

and

$$\sum_{(u,v) \in V^2} S_{u,v}^2 = cp^{12}n^8. \quad (3.6)$$

Thus the Cauchy–Schwarz inequality implies

$$(1 - \delta)^4 \leq a^4 \leq b^2 \leq c \leq 1 + \delta. \quad (3.7)$$

Due to Fact 3.2 (applied with  $\alpha = ap^3n^2$  and  $\nu = p^3n^2\sqrt{b - a^2}$ ), (3.4), and (3.5) we have

$$\Delta(A, V, V) = \sum_{x \in A} T_x = ap^3n^2 |A| \pm \sqrt{b - a^2} \cdot p^3n^3.$$

Owing to  $1 - \delta \leq a \leq 1 + \delta$  and  $\sqrt{b - a^2} \leq 3\sqrt{\delta}$  (see (3.7)), this leads to

$$\Delta(A, V, V) = p^3n^2 |A| \pm 4\delta^{1/2}p^3n^3. \quad (3.8)$$

Similarly Fact 3.2 (applied with  $\alpha = bp^6n^3$  and  $\nu = p^6n^3\sqrt{c - b^2}$ ), (3.5), and (3.6) give

$$\sum_{(u,v) \in A^2} S_{u,v}^2 \leq bp^6n^3 |A|^2 + \sqrt{c - b^2} \cdot p^6n^5 \leq ((1 + \delta) |A|^2 + 3\delta^{1/2}n^2)p^6n^3,$$

whence

$$\sum_{y \in V} \Delta(A, y, V)^2 = \sum_{(u,v) \in A^2} S_{u,v}^2 \leq (|A|^2 + 4\delta^{1/2}n^2)p^6n^3. \quad (3.9)$$

Now if  $Q$  and  $R$  denote the real numbers satisfying

$$\sum_{y \in V} \Delta(A, y, V) = p^3n^2Q \quad \text{and} \quad \sum_{y \in V} \Delta(A, y, V)^2 = p^6n^3R,$$

then (3.8) and (3.9) entail

$$Q = |A| \pm 4\delta^{1/2}n \quad \text{and} \quad R \leq |A|^2 + 4\delta^{1/2}n^2, \quad (3.10)$$

whilst a final application of Fact 3.2 (applied with  $\alpha = p^3nQ$  and  $\nu = p^3n\sqrt{R - Q^2}$ ) reveals

$$\Delta(A, B, V) = p^3nQ |B| \pm p^3n^2\sqrt{R - Q^2}. \quad (3.11)$$

It follows from (3.10) that  $R - Q^2 \leq 12\delta^{1/2}n^2$  and owing to (3.11) we obtain

$$\begin{aligned} \left| \Delta(A, B, V) - p^3 |A| |B| n \right| &\leq p^3 n |B| |Q - |A|| + 4\delta^{1/4} p^3 n^3 \\ &\leq p^3 n^3 (4\delta^{1/2} + 4\delta^{1/4}) \leq 8\delta^{1/4} p^3 n^3, \end{aligned}$$

as desired.  $\square$

#### §4. PROOF OF COROLLARY 1.4

**4.1. Notation.** We mostly follow the notation from Lovász's research monograph [37] and in this subsection we remind the reader of what we actually need. By  $\mathcal{W}$  we mean the space of all bounded symmetric measurable functions from the unit square  $[0, 1]^2$  to the set of reals. So  $\mathcal{W}$  is a linear space whose members are sometimes referred to as *kernels*. It is known that for each kernel  $W$  the maximum

$$\|W\|_{\square} = \max \left\{ \left| \int_{A \times B} W(x, y) \, dx \, dy \right| : A, B \subseteq [0, 1] \text{ measurable} \right\}$$

exists and that  $W \mapsto \|W\|_{\square}$  is a norm on  $\mathcal{W}$ , the so-called *cutnorm*. If  $\|W\|_{\square} = 0$  holds for  $W \in \mathcal{W}$ , then this kernel vanishes almost everywhere, i.e., the set  $\{(x, y) \in [0, 1]^2 : W(x, y) \neq 0\}$  has measure zero (see [37, Section 8.2.3]).

The group of measure preserving bijections from the unit interval onto itself is denoted by  $\mathcal{S}_{[0,1]}$ . This group acts in an obvious way on the space of kernels by

$$W^{\varphi}(x, y) = W(\varphi(x), \varphi(y))$$

for all  $W \in \mathcal{W}$ ,  $\varphi \in \mathcal{S}_{[0,1]}$ , and  $x, y \in [0, 1]$ . The *cut distance*  $\delta_{\square}(W_1, W_2)$  between two kernels  $W_1, W_2 \in \mathcal{W}$  is defined by

$$\delta_{\square}(W_1, W_2) = \inf \{ \|W_1 - W_2^{\varphi}\|_{\square} : \varphi \in \mathcal{S}_{[0,1]} \}.$$

Actually, this infimum is known to be a minimum, but this fact is rarely needed and we shall make no use of it.

Those  $W \in \mathcal{W}$  that satisfy  $W(x, y) \in [0, 1]$  for all  $x, y \in [0, 1]$  are called *graphons* and the set of all graphons is denoted by  $\mathcal{W}_0$ . With each graph  $G = (V, E)$  we can associate a graphon  $W_G$  by taking an arbitrary partition  $[0, 1] = \bigcup_{v \in V} P_v$  of the unit interval into measurable pieces of measure  $|V|^{-1}$  and defining for all  $x, y \in [0, 1]$

$$W_G(x, y) = \begin{cases} 1, & \text{if } uv \in E, \text{ where } x \in P_u, y \in P_v \\ 0, & \text{else.} \end{cases}$$

This graphon depends, of course, not only on  $G$  but also on the underlying partition, but modulo the action of  $\mathcal{S}_{[0,1]}$  mentioned above it is uniquely determined by  $G$ .

An important insight due to Lovász and Szegedy [38] is the compactness of the pseudometric space  $(\mathcal{W}_0, \delta_{\square})$ . In fact, the compactness easily implies the regularity lemma for graphs (see also [37, Theorem 9.23]). This result does actually occupy a central place in the limit

theory of dense graphs. Besides, it is beautifully complemented by the fact that the set  $\{W_G : G \text{ is a graph}\}$  is dense in  $(\mathcal{W}_0, \delta_\square)$ .

Given a graph  $F$  and a kernel  $W$  the *homomorphism density*  $t(F, W)$  is defined to be the multidimensional integral

$$t(F, W) = \int_{[0,1]^{V(F)}} \prod_{uv \in E(F)} W(x_u, x_v) \prod_{u \in V(F)} dx_u.$$

This stipulation extends the usual definition of homomorphism densities for graphs in the sense that  $t(F, W_G) = t(F, G)$  holds for all graphs  $F$  and  $G$ .

Analytically speaking, the *global counting lemma* asserts that for every graph  $F$  the map  $W \mapsto t(F, W)$  from  $(\mathcal{W}_0, \delta_\square)$  to  $[0, 1]$  is Lipschitz continuous with Lipschitz constant  $e(F)$  (see [37, Theorem 10.23]).

**4.2. Forcing families.** Let us write  $W \equiv p$  for a kernel  $W$  and a real number  $p$  if  $W$  agrees almost everywhere with the constant function whose value is always  $p$ .

**Lemma 4.1.** *A pair of graphs  $(F_1, F_2)$  is forcing if and only if we have  $W \equiv p$  for every real  $p \in (0, 1]$  and every graphon  $W$  with  $t(F_i, W) = p^{e(F_i)}$  for  $i = 1, 2$ .*

*Proof.* This is implicit in the discussion from [37, Section 16.7.1]. □

Theorem 1.2 is the discrete analogue of the following statement.

**Theorem 4.2.** *If a graphon  $W$  and a real  $p \in (0, 1]$  are such that*

$$W(x, y) \int_0^1 W(x, z)W(y, z) dz = p^3$$

*holds for almost all  $(x, y) \in [0, 1]^2$ , then  $W \equiv p$ .*

One way to show this proceeds by carefully repeating the proof of Lemma 2.1 in this analytical setting. This is not hard, but somewhat technical, and hence we would like to present an alternative argument here.

*Proof of Theorem 4.2.* Define a graphon  $U$  by

$$U(x, y) = W(x, y) \int_0^1 W(x, z)W(y, z) dz \tag{4.1}$$

for all  $x, y \in [0, 1]$ . Now the assumption  $U \equiv p^3$  leads to  $t(K_3, W) = t(K_2, U) = p^3$  and  $t(C_4^\triangleright, W) = t(C_4, U) = p^{12}$ . Since the pair  $(K_3, C_4^\triangleright)$  is forcing, it follows by Lemma 4.1 that we have indeed  $W \equiv p$ . □

We conclude this section with the proof of our main result on forcing pairs involving triangles.

*Proof of Corollary 1.4.* Suppose that  $W$  is a graphon and  $p \in (0, 1]$  is a real number such that

$$t(K_3, W) = p^3 \quad \text{and} \quad t(F^\triangleright, W) = p^{3e(F)}. \tag{4.2}$$

In view of Lemma 4.1 we have to prove that  $W \equiv p$ . To this end, we look again at the graphon  $U$  defined by (4.1). The hypothesis (4.2) rewrites in terms of  $U$  as

$$t(K_2, U) = p^3 \quad \text{and} \quad t(F, U) = p^{3e(F)}.$$

Since the pair  $(K_2, F)$  is forcing, it follows that  $U \equiv p^3$ , and in the light of Theorem 4.2 we get indeed  $W \equiv p$ .  $\square$

## §5. CONCLUDING REMARKS

We close with a few remarks and open problems for future research.

- Corollary 1.4 raises the general problem to characterise all graphs  $F$  with the property that the pair  $(K_3, F)$  is forcing. However, given our current state of knowledge and the fact that this is still open for  $(K_2, F)$  it appears even unclear to formulate a plausible conjecture in this regard.
- The proof of Theorem 1.2 presented in Section 2 is based on Szemerédi's regularity lemma and as a consequence this proof requires that  $\eta^{-1}$  behaves like an exponential tower of height  $\text{poly}(\varepsilon^{-1}, p^{-1})$ . We would like to thank L. M. Lovász for pointing out to us that a different argument utilising the Frieze–Kannan regularity lemma [21] shows that  $\eta^{-1} = 2^{(\varepsilon p)^{-\Theta(1)}}$  would suffice as well. To see this one exploits that the assumption of Theorem 1.2 implies  $t(K_3, G) \approx p^3$  and  $t(C_4^\triangleright, G) \approx p^{12}$ . Due to the global counting lemma this gives us two approximate equalities for the densities  $d_{ij}$  arising in a Frieze–Kannan regular partition of  $G$ . Similar as in the proof of Proposition 3.1 two reverse applications of the Cauchy–Schwarz inequality then entail that the assumption (2.1) of Lemma 2.1 holds with at most  $o(t^2)$  exceptions. Working a little bit harder in the proof of this lemma it can then be shown that this is enough to imply that there are at most  $o(t^2)$  pairs  $(i, j) \in [t]^2$  for which  $d_{ij} \approx p$  fails. This, however, is in turn equivalent to  $G$  being  $p$ -quasirandom.

Recently, it was shown by Conlon, Fox, and Sudakov [13] that the corresponding dependency of the parameters in the Simonovits–Sós theorem for the triangle is in fact linear (see also [27, 39] for further results). In view of these results, it seems an interesting open question, whether Theorem 1.2 holds also for  $\eta = \text{poly}(\varepsilon, p)$ .

- It appears to be an intriguing open problem to find the appropriate generalisation of Theorem 1.2 (and Corollary 1.3) for other graphs than the triangle. At this point even for cliques  $K_k$  with  $k \geq 4$  this is an open problem. For integers  $1 \leq \ell \leq k$  we say  $K_k$  is  $\ell$ -forcing, if every graph  $G = (V, E)$  satisfying for all subsets  $X_1, \dots, X_\ell \subseteq V$

$$\mathcal{K}_k(X_1, \dots, X_\ell) = p^{\binom{k}{2}} |V|^{k-\ell} \prod_{i=1}^{\ell} |X_i| + o(|V|^k)$$

for some  $p \in (0, 1]$  is  $p$ -quasirandom, where  $\mathcal{K}_k(X_1, \dots, X_\ell)$  denotes the number of  $k$ -tuples

$$(v_1, \dots, v_k) \in X_1 \times \dots \times X_\ell \times V^{k-\ell}$$

that span a  $K_k$  in  $G$ . The Simonovits–Sós theorem implies for every  $k \geq 2$  that  $K_k$  is  $k$ -forcing and it is not hard to show that no clique is 1-forcing. Theorem 1.2 tells us that  $K_3$  is 2-forcing and it would be interesting to determine for every  $k \geq 4$  the smallest  $\ell$  such that  $K_k$  is  $\ell$ -forcing. Currently, we are not aware of any reason that rules out the possibility that every clique  $K_k$  is 2-forcing or that there is a universal bound independent of  $k$ .

- One may also consider hypergraphs extensions of those results. For example, one may investigate, whether the tetrahedron  $K_4^{(3)}$  is 3-forcing for the notion of quasirandomness investigated in [6, 33].

## REFERENCES

- [1] E. Aigner-Horev, D. Conlon, H. Hàn, Y. Person, and M. Schacht, *Quasirandomness in hypergraphs*. Submitted. ↑1
- [2] N. Alon, *Eigenvalues and expanders*, Combinatorica **6** (1986), no. 2, 83–96, DOI [10.1007/BF02579166](#). Theory of computing (Singer Island, Fla., 1984). MR875835 ↑1
- [3] N. Alon and F. R. K. Chung, *Explicit construction of linear sized tolerant networks*, Proceedings of the First Japan Conference on Graph Theory and Applications (Hakone, 1986), 1988, pp. 15–19, DOI [10.1016/0012-365X\(88\)90189-6](#). MR975519 ↑1
- [4] F. R. K. Chung, *Quasi-random classes of hypergraphs*, Random Structures Algorithms **1** (1990), no. 4, 363–382, DOI [10.1002/rsa.3240010401](#). MR1138430 ↑1
- [5] ———, *Quasi-random hypergraphs revisited*, Random Structures Algorithms **40** (2012), no. 1, 39–48, DOI [10.1002/rsa.20388](#). MR2864651 ↑1
- [6] F. R. K. Chung and R. L. Graham, *Quasi-random hypergraphs*, Random Structures Algorithms **1** (1990), no. 1, 105–124, DOI [10.1002/rsa.3240010108](#). MR1068494 ↑1, 5
- [7] ———, *Quasi-random tournaments*, J. Graph Theory **15** (1991), no. 2, 173–198, DOI [10.1002/jgt.3190150206](#). MR1106530 ↑1
- [8] ———, *Quasi-random set systems*, J. Amer. Math. Soc. **4** (1991), no. 1, 151–196, DOI [10.2307/2939258](#). MR1077279 ↑1
- [9] ———, *Maximum cuts and quasirandom graphs*, Random graphs, Vol. 2 (Poznań, 1989), Wiley-Intersci. Publ., Wiley, New York, 1992, pp. 23–33. MR1166604 ↑1
- [10] ———, *Quasi-random subsets of  $Z_n$* , J. Combin. Theory Ser. A **61** (1992), no. 1, 64–86, DOI [10.1016/0097-3165\(92\)90053-W](#). MR1178385 ↑1
- [11] ———, *Sparse quasi-random graphs*, Combinatorica **22** (2002), no. 2, 217–244, DOI [10.1007/s004930200010](#). Special issue: Paul Erdős and his mathematics. MR1909084 ↑1
- [12] F. R. K. Chung, R. L. Graham, and R. M. Wilson, *Quasi-random graphs*, Combinatorica **9** (1989), no. 4, 345–362, DOI [10.1007/BF02125347](#). MR1054011 (91e:05074) ↑1, 1, 1
- [13] D. Conlon, J. Fox, and B. Sudakov, *Hereditary quasirandomness without regularity*, Math. Proc. Cambridge Philos. Soc., available at [arXiv:1611.02099](#). To appear. ↑5
- [14] ———, *An approximate version of Sidorenko’s conjecture*, Geom. Funct. Anal. **20** (2010), no. 6, 1354–1366. MR2738996 ↑1
- [15] D. Conlon, J. Fox, and Y. Zhao, *Extremal results in sparse pseudorandom graphs*, Adv. Math. **256** (2014), 206–290, DOI [10.1016/j.aim.2013.12.004](#). MR3177293 ↑1
- [16] D. Conlon, H. Hàn, Y. Person, and M. Schacht, *Weak quasi-randomness for uniform hypergraphs*, Random Structures Algorithms **40** (2012), no. 1, 1–38, DOI [10.1002/rsa.20389](#). MR2864650 ↑1, 1, 1

- [17] D. Conlon, J. H. Kim, C. Lee, and J. Lee, *Some advances on Sidorenko's conjecture*, available at [arXiv:1510.06533](#). Submitted. [↑1](#)
- [18] D. Conlon and J. Lee, *Finite reflection groups and graph norms*, Adv. Math. **315** (2017), 130–165, DOI [10.1016/j.aim.2017.05.009](#). MR3667583 [↑1](#)
- [19] J. N. Cooper, *Quasirandom permutations*, J. Combin. Theory Ser. A **106** (2004), no. 1, 123–143, DOI [10.1016/j.jcta.2004.01.006](#). MR2050120 [↑1](#)
- [20] P. Frankl, V. Rödl, and R. M. Wilson, *The number of submatrices of a given type in a Hadamard matrix and related results*, J. Combin. Theory Ser. B **44** (1988), no. 3, 317–328, DOI [10.1016/0095-8956\(88\)90040-8](#). MR941440 [↑1](#)
- [21] A. Frieze and R. Kannan, *Quick approximation to matrices and applications*, Combinatorica **19** (1999), no. 2, 175–220, DOI [10.1007/s004930050052](#). MR1723039 [↑5](#)
- [22] W. T. Gowers, *Quasirandom groups*, Combin. Probab. Comput. **17** (2008), no. 3, 363–387, DOI [10.1017/S0963548307008826](#). MR2410393 [↑1](#)
- [23] S. Griffiths, *Quasi-random oriented graphs*, J. Graph Theory **74** (2013), no. 2, 198–209, DOI [10.1002/jgt.21701](#). MR3090716 [↑1](#)
- [24] H. Hàn, Y. Person, and M. Schacht, *Note on forcing pairs*, The Sixth European Conference on Combinatorics, Graph Theory and Applications (EuroComb 2011), Electron. Notes Discrete Math., vol. 38, Elsevier Sci. B. V., Amsterdam, 2011, pp. 437–442, DOI [10.1016/j.endm.2011.09.071](#). [↑1](#)
- [25] H. Hatami, *Graph norms and Sidorenko's conjecture*, Israel J. Math. **175** (2010), 125–150, DOI [10.1007/s11856-010-0005-1](#). MR2607540 [↑1](#)
- [26] J. Haviland and A. Thomason, *Pseudo-random hypergraphs*, Discrete Math. **75** (1989), no. 1-3, 255–278, DOI [10.1016/0012-365X\(89\)90093-9](#). Graph theory and combinatorics (Cambridge, 1988). MR1001401 [↑1](#)
- [27] X. He, *Linear dependence between hereditary quasirandomness conditions*, available at [arXiv:1707.05396](#). Submitted. [↑5](#)
- [28] H. Huang and C. Lee, *Quasi-randomness of graph balanced cut properties*, Random Structures Algorithms **41** (2012), no. 1, 124–145, DOI [10.1002/rsa.20384](#). MR2943429 [↑1](#)
- [29] S. Janson and V. T. Sós, *More on quasi-random graphs, subgraph counts and graph limits*, European J. Combin. **46** (2015), 134–160, DOI [10.1016/j.ejc.2015.01.001](#). MR3305351 [↑1](#)
- [30] J. H. Kim, C. Lee, and J. Lee, *Two approaches to Sidorenko's conjecture*, Trans. Amer. Math. Soc. **368** (2016), no. 7, 5057–5074, DOI [10.1090/tran/6487](#). MR3456171 [↑1](#)
- [31] Y. Kohayakawa and V. Rödl, *Regular pairs in sparse random graphs. I*, Random Structures Algorithms **22** (2003), no. 4, 359–434, DOI [10.1002/rsa.10081](#). MR1980964 [↑1](#)
- [32] ———, *Szemerédi's regularity lemma and quasi-randomness*, Recent advances in algorithms and combinatorics, CMS Books Math./Ouvrages Math. SMC, vol. 11, Springer, New York, 2003, pp. 289–351, DOI [10.1007/0-387-22444-0\\_9](#). MR1952989 [↑1](#)
- [33] Y. Kohayakawa, V. Rödl, and J. Skokan, *Hypergraphs, quasi-randomness, and conditions for regularity*, J. Combin. Theory Ser. A **97** (2002), no. 2, 307–352, DOI [10.1006/jcta.2001.3217](#). MR1883869 [↑1](#), [5](#)
- [34] M. Krivelevich and B. Sudakov, *Pseudo-random graphs*, More sets, graphs and numbers, Bolyai Soc. Math. Stud., vol. 15, Springer, Berlin, 2006, pp. 199–262, DOI [10.1007/978-3-540-32439-3\\_10](#). MR2223394 [↑1](#)
- [35] J. Lenz and D. Mubayi, *Eigenvalues and linear quasirandom hypergraphs*, Forum Math. Sigma **3** (2015), e2, 26, DOI [10.1017/fms.2014.22](#). MR3324939 [↑1](#)
- [36] J. X. Li and B. Szegedy, *On the logarithmic calculus and Sidorenko's conjecture*, Combinatorica, available at [arXiv:1107.1153](#). To appear. [↑1](#)
- [37] L. Lovász, *Large networks and graph limits*, American Mathematical Society Colloquium Publications, vol. 60, American Mathematical Society, Providence, RI, 2012. MR3012035 [↑4.1](#), [4.2](#)
- [38] L. Lovász and B. Szegedy, *Szemerédi's lemma for the analyst*, Geom. Funct. Anal. **17** (2007), no. 1, 252–270, DOI [10.1007/s00039-007-0599-6](#). MR2306658 [↑4.1](#)



- [39] Chr. Reiher and M. Schacht, *Quasirandomness from hereditary subgraph densities*. Manuscript. [↑5](#)
- [40] V. Rödl, *On universality of graphs with uniformly distributed edges*, Discrete Math. **59** (1986), no. 1-2, 125–134, DOI [10.1016/0012-365X\(86\)90076-2](#). MR837962 [↑1](#)
- [41] A. Shapira, *Quasi-randomness and the distribution of copies of a fixed graph*, Combinatorica **28** (2008), no. 6, 735–745, DOI [10.1007/s00493-008-2375-0](#). MR2488748 [↑1](#)
- [42] A. Shapira and R. Yuster, *The effect of induced subgraphs on quasi-randomness*, Random Structures Algorithms **36** (2010), no. 1, 90–109, DOI [10.1002/rsa.20289](#). MR2591049 [↑1](#)
- [43] ———, *The quasi-randomness of hypergraph cut properties*, Random Structures Algorithms **40** (2012), no. 1, 105–131, DOI [10.1002/rsa.20364](#). MR2864654 [↑1](#)
- [44] A. F. Sidorenko, *Inequalities for functionals generated by bipartite graphs*, Diskret. Mat. **3** (1991), no. 3, 50–65 (Russian); English transl., Discrete Math. Appl. **2** (1992), no. 5, 489–504. MR1138091 [↑1](#)
- [45] M. Simonovits, *Extremal graph problems, degenerate extremal problems, and supersaturated graphs*, Progress in graph theory (Waterloo, Ont., 1982), Academic Press, Toronto, ON, 1984, pp. 419–437. MR776819 [↑1](#)
- [46] M. Simonovits and V. T. Sós, *Szemerédi’s partition and quasirandomness*, Random Structures Algorithms **2** (1991), no. 1, 1–10, DOI [10.1002/rsa.3240020102](#). MR1099576 [↑1](#)
- [47] ———, *Hereditarily extended properties, quasi-random graphs and not necessarily induced subgraphs*, Combinatorica **17** (1997), no. 4, 577–596, DOI [10.1007/BF01195005](#). MR1645698 [↑1](#), [1](#)
- [48] ———, *Hereditary extended properties, quasi-random graphs and induced subgraphs*, Combin. Probab. Comput. **12** (2003), no. 3, 319–344, DOI [10.1017/S0963548303005613](#). Combinatorics, probability and computing (Oberwolfach, 2001). MR1988980 [↑1](#)
- [49] J. Skokan and L. Thoma, *Bipartite subgraphs and quasi-randomness*, Graphs Combin. **20** (2004), no. 2, 255–262, DOI [10.1007/s00373-004-0556-1](#). MR2080111 (2005e:05130) [↑1](#), [1](#)
- [50] B. Szegedy, *An information theoretic approach to Sidorenko’s conjecture*, available at [arXiv:1406.6738](#). Preprint. [↑1](#)
- [51] E. Szemerédi, *Regular partitions of graphs*, Problèmes combinatoires et théorie des graphes (Colloq. Internat. CNRS, Univ. Orsay, Orsay, 1976), Colloq. Internat. CNRS, vol. 260, CNRS, Paris, 1978, pp. 399–401 (English, with French summary). MR540024 (81i:05095) [↑2](#)
- [52] A. Thomason, *Pseudorandom graphs*, Random graphs ’85 (Poznań, 1985), North-Holland Math. Stud., vol. 144, North-Holland, Amsterdam, 1987, pp. 307–331. MR930498 [↑1](#)
- [53] ———, *Random graphs, strongly regular graphs and pseudorandom graphs*, Surveys in combinatorics 1987 (New Cross, 1987), London Math. Soc. Lecture Note Ser., vol. 123, Cambridge Univ. Press, Cambridge, 1987, pp. 173–195. MR905280 [↑1](#)
- [54] H. Towsner,  *$\sigma$ -algebras for quasirandom hypergraphs*, Random Structures Algorithms **50** (2017), no. 1, 114–139, DOI [10.1002/rsa.20641](#). MR3583029 [↑1](#)
- [55] R. M. Wilson, *Cyclotomy and difference families in elementary abelian groups*, J. Number Theory **4** (1972), 17–47, DOI [10.1016/0022-314X\(72\)90009-1](#). MR0309755 [↑1](#)
- [56] R. Yuster, *Quasi-randomness is determined by the distribution of copies of a fixed graph in equicardinal large sets*, Combinatorica **30** (2010), no. 2, 239–246, DOI [10.1007/s00493-010-2496-0](#). MR2676838 [↑1](#)

FACHBEREICH MATHEMATIK, UNIVERSITÄT HAMBURG, HAMBURG, GERMANY

*E-mail address:* Christian.Reiher@uni-hamburg.de

*E-mail address:* schacht@math.uni-hamburg.de