

## Five-Particle Phase-Space Integrals in QCD\*

---

**Oleksandr Gituliar**<sup>†</sup>

*II. Institut für Theoretische Physik, Universität Hamburg*

*E-mail: [oleksandr@gituliar.net](mailto:oleksandr@gituliar.net)*

**Vitaly Magerya**

*II. Institut für Theoretische Physik, Universität Hamburg*

*E-mail: [vitaly.magerya@tx97.net](mailto:vitaly.magerya@tx97.net)*

**Andrey Pikelner**

*II. Institut für Theoretische Physik, Universität Hamburg*<sup>‡</sup>

*E-mail: [andrey.pikelner@desy.de](mailto:andrey.pikelner@desy.de)*

We present analytical expressions for the 31 five-particle phase-space master integrals in massless QCD as an  $\epsilon$ -series with coefficients being multiple zeta values of weight up to 12. In addition, we provide a computer code for the Monte-Carlo integration in higher dimensions, based on the RAMBO algorithm, that has been used to numerically cross-check the obtained results in 4, 6, and 8 dimensions.

*Loops and Legs in Quantum Field Theory (LL2018)*

*29 April 2018 - 04 May 2018*

*St. Goar, Germany*

---

\*Based on [1].

<sup>†</sup>Speaker.

<sup>‡</sup>On leave of absence from Joint Institute for Nuclear Research, Dubna, Russia.

## 1. Introduction

Nowadays, perturbative calculations play the key role in describing data from high-energy particle colliders, such as the LHC, as well as in improving the precision of numerical parameters in the Standard Model and other models. It is clear now that higher-order calculations will play an even more crucial role in processing data from future high-luminosity colliders, like the FCC or the ILC, where theoretical errors will dominate over experimental statistical errors. These arguments motivate us to make one step forward beyond available fully-inclusive phase-space integrals for a four-particle decay [2] and calculate a set of yet unknown integrals that corresponds to a five-particle decay of a color-neutral off-shell particle in Quantum Chromodynamics.

A particular application of these integrals we have in mind is the extraction of NNLO time-like splitting functions [3] from a semi-inclusive one-particle decay process, as for example discussed in [4, 5]: the integrals we will be looking at correspond to a fully inclusive cross section, and can be used to determine integration constants when calculating exclusive quantities using the method of differential equations.

In this article, we focus on the calculation of master integrals that can be used to express any other integral of the corresponding topology provided a set of integration-by-parts rules (IBP) [6] is known. Our approach is based on techniques for solving dimensional recurrence relations (DRR) [7] described in [8, 9]. In particular, we use the DREAM package [8] to obtain numerical results for the desired integrals with 2000-digit precision, and restore their analytical form in terms of multiple zeta values (MZV) [10, 11, 12] up to weight 12 using the PSLQ method [13] as implemented in Mathematica. We also present a Monte-Carlo code, based on the RAMBO algorithm [14], for numerical integration of the phase-space integrals in arbitrary (integer) number of dimensions that has been used to check consistency of the obtained results.

This article is organized as following. In Section 2 we introduce our notation and describe our calculational method in more detail. In Section 3 we provide complete results for four-particle integrals and discuss numerical cross-checks using Monte-Carlo integration. In Section 4 we make our final remarks.


Accompanying to this paper are the auxiliary files on the arXiv<sup>1</sup> containing the complete master integrals with MZV weight up to 12, as well as the Monte-Carlo integration routines with the corresponding results.

## 2. The Method

We start by identifying a set of five-particle phase-space master integrals using two different approaches for consistency. As the first approach we exploit the equivalence of IBP rules for cut and ordinary propagators, and obtain the complete basis of phase-space master integrals by taking all five-particle cuts of the 28 master integrals for four-loop massless propagators found in [15], discarding those that do not correspond to the squared matrix elements of the  $1 \rightarrow 5$  process (i.e. only leaving graphs which are bipartite), and reducing the remaining integrals with Laporta-style IBP reduction [16, 17] as implemented in FIRE5 [18]. As an alternative approach, we construct the complete expression for the total cross section of the  $1 \rightarrow 5$  process in QCD using QGRAF [19]

---

<sup>1</sup><https://arXiv.org/abs/1803.09084>



**Table 1:** Cut diagrams for five-particle phase-space master integrals in QCD. Dashed lines represent cut propagators and carry final-state momenta  $p_1, \dots, p_5$ . Labels represent propagators, so that "123" corresponds to  $p_1 + p_2 + p_3$  and "012" to  $q - p_1 - p_2$  (where  $q$  is the initial-state momentum, i.e.,  $q = p_1 + \dots + p_5$ ).

and FORM [20], and then reduce it with the help of FIRE5. Both methods give 31 master integrals listed in Table 1, with each having up to 6 unique propagators. Our notation for these integrals is

$$F_i = S_\Gamma \int \text{dPS}_5 \frac{1}{D_1^{(i)} \dots D_n^{(i)}}, \quad (2.1)$$

where  $D_j^{(i)}$  are propagators that take the form of invariant scalar products

$$s_{kl\dots q} = (p_k + p_l + \dots + p_q)^2, \quad (2.2)$$

$\text{dPS}_5$  is a five-particle phase-space element in  $D$  dimensions

$$\text{dPS}_N = \left( \prod_{i=1}^N \text{d}^D p_i \delta^+(p_i^2) \right) \delta^{(D)}(q - p_1 - \dots - p_N), \quad (2.3)$$

and  $S_\Gamma$  is a common normalization factor chosen for convenience<sup>2</sup> to be

$$S_\Gamma = (q^2)^{5-2D} \frac{(2\pi)^4}{\pi^{2D}} \Gamma\left(\frac{D}{2} - 1\right) \Gamma\left(3\frac{D}{2} - 3\right). \quad (2.4)$$

With this normalization and knowing the volume of the complete  $N$ -particle phase space<sup>3</sup>

$$\int \text{dPS}_N = (q^2)^{\frac{D}{2}(N-1)-N} \frac{\pi^{\frac{D}{2}(N-1)}}{(2\pi)^{N-1}} \frac{\Gamma\left(\frac{D}{2} - 1\right)^N}{\Gamma\left(\left(\frac{D}{2} - 1\right)(N-1)\right) \Gamma\left(\left(\frac{D}{2} - 1\right)N\right)}, \quad (2.5)$$

we can already fix the value of  $F_1$  as:

$$F_1 = S_\Gamma \int \text{dPS}_5 = \frac{\Gamma\left(\frac{D}{2} - 1\right)^6 \Gamma\left(3\frac{D}{2} - 3\right)}{\Gamma\left(4\frac{D}{2} - 4\right) \Gamma\left(5\frac{D}{2} - 5\right)} \quad (2.6)$$

Next, with the help of LiteRed [21] and FIRE5 [18] we derive a set of lowering dimensional recurrence relations which express master integrals in  $D+2$  dimensions in terms of master integrals in  $D$  dimensions:

$$F_i(D+2) = M_{ij}(D) F_j(D). \quad (2.7)$$

In the general case  $M(D)$  is expected to have a block-triangular structure, but in our case it can be shuffled into triangular form. The general structure of our  $M(D)$  can be visualized as:

<sup>2</sup>This way we prevent additional constants (e.g.  $\gamma_E$  or  $\ln \pi$ ) to appear in the final results, hence reducing its size as well as a size of the basis for PSLQ algorithm.

<sup>3</sup>The dependence on  $q^2$  is trivial here, and can be restored by power counting. We will omit it from now on, setting  $q^2$  to 1.



$F_i(D)$  in the limit of  $D$  going to imaginary infinity. Our case contains two important simplifications, and will not require such analysis.

The first simplification is that since  $M(D)$  is triangular, the homogeneous part of eq. (2.9) decouples into a set of first order difference equations:

$$H_i(D+2) = M_{ii}(D)H_i(D) \quad (2.12)$$

For rational  $M_{ii}(D)$  written in the following form (compare to eq. (2.10)):

$$M_{ii}(D) = C \frac{(\frac{D}{2} - a_1)(\frac{D}{2} - a_2) \dots (\frac{D}{2} - a_A)}{(\frac{D}{2} - b_1)(\frac{D}{2} - b_2) \dots (\frac{D}{2} - b_B)}, \quad (2.13)$$

the homogeneous solution can immediately be found as:

$$H_i(D) = C^{\frac{D}{2}} \frac{\Gamma(\frac{D}{2} - a_1) \Gamma(\frac{D}{2} - a_2) \dots \Gamma(\frac{D}{2} - a_A)}{\Gamma(\frac{D}{2} - b_1) \Gamma(\frac{D}{2} - b_2) \dots \Gamma(\frac{D}{2} - b_B)} \quad (2.14)$$

Explicitly, for  $H_{28}(D)$  we have:

$$H_{28}(D) = \left(-\frac{3^3}{4^3}\right)^{\frac{D}{2}} \frac{(\frac{D}{2} - 2)^2 (\frac{D}{2} - \frac{2}{3}) (\frac{D}{2} - \frac{1}{3})}{(\frac{D}{2} - \frac{3}{2})^2 (\frac{D}{2} - 1) (\frac{D}{2} - \frac{1}{2})} \quad (2.15)$$

The second simplification, is that as we will argue further, all  $\omega_i(D)$  for  $i > 1$  are zero. To see this, first let us look at the asymptotic behavior of  $F_i(D)$  at large  $D$ . Rewriting eq. (2.1) as an integral over invariants  $s_{ij}$  gives

$$F_i = S_\Gamma \left( \prod_{k=1}^{N-1} \frac{\Omega_{D-k}}{2} \right) \int \left( \prod_{l < m} \frac{ds_{lm}}{2} \right) (\Delta_N)^{\frac{D-N-1}{2}} \Theta(\Delta_N) \delta(1 - s_{1\dots N}) \frac{1}{D_1^{(i)} \dots D_n^{(i)}}, \quad (2.16)$$

where  $\Delta_N$  is the Gram determinant defined as

$$\Delta_N = \frac{(-1)^{N+1}}{2^N} \begin{vmatrix} s_{11} & s_{12} & \dots & s_{1N} \\ s_{12} & s_{22} & \dots & s_{2N} \\ \vdots & \vdots & \ddots & \vdots \\ s_{1N} & s_{2N} & \dots & s_{NN} \end{vmatrix}, \quad (2.17)$$

and  $\Omega_k$  is the surface area of a unit hypersphere in  $k$ -dimensional space

$$\Omega_k = 2\pi^{\frac{k}{2}} \Gamma\left(\frac{k}{2}\right)^{-1}. \quad (2.18)$$

If  $\Delta_N(s_{ij})$  has a unique global maximum inside the integration region, we can apply Laplace's method to eq. (2.16) and find its asymptotic as

$$F_i(D \rightarrow \infty) = S_\Gamma \left( \prod_{k=1}^{N-1} \Omega_{D-k} \right) (\Delta_N^{\max})^{\frac{D}{2}} \left( \frac{2\pi}{D} \right)^{\frac{1}{2} \left( \frac{N(N-1)}{2} - 1 \right)} (\mathcal{C}_i + \mathcal{O}(D^{-1})), \quad (2.19)$$

where  $\mathcal{C}_i$  is a constant that depends on the location of the maximum and the denominators  $D_j^{(i)}$ , but not on  $D$ .

The global maximum of  $\Delta_N$  is reached when all  $s_{ij}$  ( $i \neq j$ ) are identical and equal to  $\frac{2}{N(N-1)}$ . Geometrically this configuration corresponds to the vectors  $\vec{p}_i$  pointing to the vertices of a regular  $N$ -hedron embedded into Euclidean space of  $(N-1)$  dimensions. The maximum value is then

$$\Delta_N^{max} = \frac{1}{N^N(N-1)^{N-1}}, \quad (2.20)$$

and explicitly we get

$$F_i(D \rightarrow \infty) = \pi^{\frac{7}{2}} 2^{\frac{25}{2}} \frac{(4^4 5^5)^{-\frac{D}{2}} \Gamma(\frac{3D}{2} - 3)}{D^{\frac{9}{2}} \Gamma(\frac{D-4}{2}) \Gamma(\frac{D-3}{2}) \Gamma(\frac{D-1}{2})} (\mathcal{C}_i + \mathcal{O}(D^{-1})) \sim \frac{1}{D^{\frac{5}{2}}} \left( \frac{3^3}{4^4 5^5} \right)^{\frac{D}{2}}. \quad (2.21)$$

It follows that all  $F_i(D)$  have identical asymptotic behavior up to a constant  $\mathcal{C}_i$ . As a confirmation, it can be shown that eq. (2.21) is asymptotically the same expression as we had for  $F_1$  in eq. (2.6).

Now we can compare this asymptotic for  $F_i(D)$  to the asymptotic for  $H_i(D)$  from eq. (2.14). Indeed, for  $H_{28}(D)$  from eq. (2.15) we have:

$$H_{28}(D) \sim \frac{1}{D^{\frac{1}{2}}} \left( -\frac{3^3}{4^3} \right)^{\frac{D}{2}}. \quad (2.22)$$

This grows asymptotically exponentially faster than eq. (2.21). In fact, all  $H_i(D)$  for  $i > 1$  grow exponentially faster than  $F_i(D)$ , and since  $F$  and  $H$  are connected via eq. (2.11), this can only happen if the corresponding periodic functions  $\omega_i(D)$  are zero.

Thus, to find  $F_i(D)$  we only need to find  $R_i(D)$ , the inhomogeneous solutions to eq. (2.9). We compute them as a series in  $\varepsilon = (4-D)/2$  using DREAM with 2000-digit accuracy, and then restore the analytical form of the series coefficients in terms of MZVs using the PSLQ method [13]. This way we obtain the analytical result for all master integrals up to MZVs of weight 12 using the corresponding bases from [10] and the SummerTime package [12] for their numerical evaluation. Corresponding expressions are presented in the auxiliary files on the arXiv.

### 3. Crosschecks

#### 3.1 Four-Particle Integrals

As the first consistency check of our method we reproduce results for four-particle phase-space integrals reported in [2]. We perform all the steps described in Section 2. Generating the IBP rules with the help of LiteRed and then proceeding with DREAM we obtain the final result with 2000-digit accuracy. The series reconstructed with PSLQ, and truncated to MZV weight 6 are (using the original notation, and omitting  $S_\Gamma$  and  $q^2$  factors):

$$R_6 \equiv \text{---} \begin{array}{c} \bullet \\ \diagup \quad \diagdown \\ \bullet \quad \bullet \\ \diagdown \quad \diagup \\ \bullet \end{array} \text{---} = -1 + \zeta_2 + \varepsilon \left( -12 + 5\zeta_2 + 9\zeta_3 \right) + \varepsilon^2 \left( -91 + 27\zeta_2 \right. \quad (3.1) \\ \left. + 45\zeta_3 + \frac{61}{5}\zeta_2^2 \right) + \varepsilon^3 \left( -558 + 161\zeta_2 + 197\zeta_3 + 61\zeta_2^2 \right)$$





$i$	Numerical results			Analytic results		
	$D = 4$	$D = 6$	$D = 8$	$D = 4$	$D = 6$	$D = 8$
2	–	1708(2)00	4699(1)0	–	171085.62	47000.531
3	3.7823(4)	3.1704(2)	3.0221(1)	3.7823736	3.1704486	3.0221118
4	–	1504.7(8)	725.3(1)	–	1504.4507	725.26806
5	–	1007.4(5)	580.80(9)	–	1007.5235	580.76347
6	–	6191(5)00	14496(6)0	–	619633.25	144975.32
7	46.46(4)	18.533(2)	15.205(1)	46.435253	18.532303	15.205538
8	–	2031(2)0	5357(2)	–	20297.189	5355.3611
9	–	4313(3)	2406.7(4)	–	4312.8823	2406.7943
10	10.436(2)	7.1508(5)	6.5093(3)	10.435253	7.1507477	6.5092878
11	228.8(1)	62.67(1)	47.663(4)	229.11836	62.667046	47.663194
12	–	157.34(4)	102.26(1)	–	157.33521	102.26408
13	–	13729(8)	4000(1)	–	13732.166	4000.2779
14	–	268.46(8)	172.80(2)	–	268.45969	172.79805
15	–	6322(6)0	16048(5)	–	63316.356	16049.857
16	–	4414(3)0	12952(4)	–	44117.898	12951.443
17	–	1243.4(7)	709.6(1)	–	1243.1369	709.52840
18	–	3002(2)00	5899(3)0	–	300402.99	58965.517
19	–	4982(4)00	10637(4)0	–	498329.79	106357.81
20	–	2360(2)000	5777(2)00	–	2362594.9	577686.64
21	–	6312(5)0	20642(5)	–	63147.876	20642.071
22	–	8402(7)00	24407(8)0	–	840453.94	244075.75
23	–	1443(1)000	4556(1)00	–	1443198.3	455543.43
24	–	1391(1)00	3997(1)0	–	139263.92	39966.878
25	–	3347(3)00	8526(3)0	–	335128.10	85254.217
26	25.563(6)	15.376(1)	13.6042(7)	25.564747	15.376404	13.604247
27	–	697.3(3)	397.84(6)	–	697.18948	397.83514
28	143.9(1)	52.855(7)	42.917(3)	143.97886	52.853837	42.917424
29	–	4409(3)0	13702(3)	–	44117.898	13700.597
30	–	6327(6)0	16181(5)	–	63316.356	16178.566
31	–	8955(8)0	19055(8)	–	89611.062	19051.115

**Table 2:** Numerical results for the ratio  $F_i/F_1$  with the corresponding uncertainties (standard deviations) indicated in the parenthesis. Missing entries correspond to divergent integrals.

but does not give a flat distribution of points over the phase space. Still, the convergence of Vegas integration with this method appeared to be as good as with the RAMBO version. We do not supply a corresponding table of results, because it is very similar to Table 2.

We would like to note that such numerical integration is commonly done differently, e.g. via the method sector decomposition [26]. To apply that method to phase space integrals one needs to parameterize the phase space by a suitable rational mapping from a hypercube, and for five particles we found this to be challenging, whereas a four-particle phase-space mapping is known from [2].

## 4. Conclusions and outlook

In this article we present analytical expressions for five-particle phase-space integrals expressed in terms of multiple zeta values up to weight 12. The results are calculated using dimensional recurrence relation method with a 2000-digit accuracy using the DREAM package, and analytically restored via the PSLQ algorithm. We also present a computer code for the numerical integration of phase-space integrals in a higher-number of dimensions that has been used to cross-check the obtained results. The approach presented here shows excellent performance for calculating single-scale integrals without ultra-violet divergences and can be easily applied to other problems of this kind.

Further work in this direction includes the calculation of multi-scale phase-space integrals needed for the extraction of time-like splitting functions, with the presented integrals used as boundary conditions of differential equations. Also of interest is the calculation of the remaining unknown integrals with 2-, 3- and 4-particle cuts appearing in the optical theorem for the four-loop propagator.

## Acknowledgments

We are thankful to Sven Moch for numerous discussions and helpful suggestions concerning this work, and for proofreading this article. We were pleased to use Axodraw2 [27] to draw diagrams for this article.

This work was supported in part by the German Research Foundation DFG through the Collaborative Research Centre No. SFB 676 *Particles, Strings and the Early Universe: the Structure of Matter and Space-Time*.

## References

- [1] O. Gituliar, V. Magerya and A. Pikelner, *Five-Particle Phase-Space Integrals in QCD*, *JHEP* **06** (2018) 099[1803.09084].
- [2] A. Gehrmann-De Ridder, T. Gehrmann and G. Heinrich, *Four particle phase space integrals in massless QCD*, *Nucl.Phys.* **B682** (2004) 265[hep-ph/0311276].
- [3] A. A. Almasy, S. Moch and A. Vogt, *On the Next-to-Next-to-Leading Order Evolution of Flavour-Singlet Fragmentation Functions*, *Nucl. Phys.* **B854** (2012) 133[1107.2263].
- [4] O. Gituliar, *Master integrals for splitting functions from differential equations in QCD*, *JHEP* **02** (2016) 017[1512.02045].
- [5] O. Gituliar and S. Moch, *Towards three-loop QCD corrections to the time-like splitting functions*, *Acta Phys. Polon.* **B46** (2015) 1279[1505.02901].
- [6] K. G. Chetyrkin and F. V. Tkachov, *Integration by Parts: The Algorithm to Calculate beta Functions in 4 Loops*, *Nucl. Phys.* **B192** (1981) 159.
- [7] O. V. Tarasov, *Connection between Feynman integrals having different values of the space-time dimension*, *Phys. Rev.* **D54** (1996) 6479[hep-th/9606018].
- [8] R. N. Lee and K. T. Mingulov, *DREAM, a program for arbitrary-precision computation of dimensional recurrence relations solutions, and its applications*, 1712.05173.

- [9] R. N. Lee and K. T. Mingulov, *Meromorphic solutions of recurrence relations and DRA method for multicomponent master integrals*, 1712.05166.
- [10] H. Furusho, *The multiple zeta value algebra and the stable derivation algebra*, *Publications of the Research Institute for Mathematical Sciences* **39** (2003) 695[math/0011261].
- [11] J. Blumlein, D. J. Broadhurst and J. A. M. Vermaseren, *The Multiple Zeta Value Data Mine*, *Comput. Phys. Commun.* **181** (2010) 582[0907.2557].
- [12] R. N. Lee and K. T. Mingulov, *Introducing SummerTime: a package for high-precision computation of sums appearing in DRA method*, *Comput. Phys. Commun.* **203** (2016) 255[1507.04256].
- [13] H. Ferguson, D. Bailey and S. Arno, *Analysis of PSLQ, an integer relation finding algorithm*, *Mathematics of Computation of the American Mathematical Society* **68** (1999) 351.
- [14] R. Kleiss, W. J. Stirling and S. D. Ellis, *A New Monte Carlo Treatment of Multiparticle Phase Space at High-energies*, *Comput. Phys. Commun.* **40** (1986) 359.
- [15] P. A. Baikov and K. G. Chetyrkin, *Four Loop Massless Propagators: An Algebraic Evaluation of All Master Integrals*, *Nucl. Phys.* **B837** (2010) 186[1004.1153].
- [16] S. Laporta, *High precision calculation of multiloop Feynman integrals by difference equations*, *Int. J. Mod. Phys.* **A15** (2000) 5087[hep-ph/0102033].
- [17] S. Laporta, *High-precision calculation of the 4-loop contribution to the electron g-2 in QED*, *Phys. Lett.* **B772** (2017) 232[1704.06996].
- [18] A. V. Smirnov, *FIRE5: a C++ implementation of Feynman Integral REDuction*, *Comput. Phys. Commun.* **189** (2015) 182[1408.2372].
- [19] P. Nogueira, *Automatic Feynman graph generation*, *J. Comput. Phys.* **105** (1993) 279.
- [20] B. Ruijl, T. Ueda and J. Vermaseren, *FORM version 4.2*, 1707.06453.
- [21] R. N. Lee, *LiteRed 1.4: a powerful tool for reduction of multiloop integrals*, *J. Phys. Conf. Ser.* **523** (2014) 012059[1310.1145].
- [22] R. N. Lee, *Space-time dimensionality  $D$  as complex variable: Calculating loop integrals using dimensional recurrence relation and analytical properties with respect to  $D$* , *Nucl. Phys.* **B830** (2010) 474[0911.0252].
- [23] G. P. Lepage, *A New Algorithm for Adaptive Multidimensional Integration*, *J. Comput. Phys.* **27** (1978) 192.
- [24] T. Hahn, *CUBA: A Library for multidimensional numerical integration*, *Comput. Phys. Commun.* **168** (2005) 78[hep-ph/0404043].
- [25] B. Gough, *GNU Scientific Library Reference Manual*. Network Theory Ltd., 3rd ed., 2009.
- [26] T. Binoth and G. Heinrich, *An automatized algorithm to compute infrared divergent multiloop integrals*, *Nucl. Phys.* **B585** (2000) 741[hep-ph/0004013].
- [27] J. C. Collins and J. A. M. Vermaseren, *Axodraw Version 2*, 1606.01177.