

STABILITY OF LODAY CONSTRUCTIONS

AYELET LINDENSTRAUSS AND BIRGIT RICHTER

ABSTRACT. We study the question for which commutative ring spectra A the tensor of a simplicial set X with A , $X \otimes A$, is a stable invariant in the sense that it depends only on the homotopy type of ΣX . We prove several structural properties about different notions of stability, corresponding to different levels of invariance required of $X \otimes A$, and establish stability in important cases, such as complex and real periodic topological K-theory, KU and KO .

1. INTRODUCTION

For any simplicial set X and any commutative ring spectrum A one can form the tensor of X with A , $X \otimes A$. An important special case of this construction is topological Hochschild homology of A , $\mathrm{THH}(A)$, which is $S^1 \otimes A$. In the following we will often work with commutative R -algebras for some commutative ring spectrum R and we will take coefficients in a commutative A -algebra C . This requires working with *pointed* simplicial sets X and we denote the corresponding object by $\mathcal{L}_X^R(A; C)$ with $\mathcal{L}_X^S(A; A)$ just being $X \otimes A$ and $\mathcal{L}_{S^1}^R(A; C)$ being $\mathrm{THH}^R(A; C)$. We recall the definition of $\mathcal{L}_X^R(A; C)$ below in 1.1. As topological Hochschild homology is the target of a trace map from algebraic K-theory

$$(1.1) \quad K(A) \rightarrow \mathrm{THH}(A)$$

it has been calculated in many cases. Also, higher order topological Hochschild homology, which is $\mathcal{L}_{S^n}^R(A; C)$, has been determined in many important classes of examples, see for instance [3, 7, 11, 17, 19]. In [3] we develop several tools for calculating $\mathcal{L}_{\Sigma X}^R(A; C)$. However, if we want to determine the homotopy type of $\mathcal{L}_X^R(A; C)$ and X doesn't happen to be a suspension, then the range of methods is much sparser.

Rognes' redshift conjecture [1] predicts that applying algebraic K-theory raises chromatic level by one in good cases. In particular, higher chromatic phenomena could be detected by iterated algebraic K-theory of rings. If A is a commutative ring spectrum, then so are $K(A)$ and $\mathrm{THH}(A)$, and as the trace map is a map of commutative ring spectra, one can iterate the trace map from (1.1) to obtain

$$K(K(A)) \rightarrow \mathrm{THH}(\mathrm{THH}(A))$$

and one doesn't have to stop at two-fold iterations. As $X \otimes A$ is the tensor of A with X in the category of commutative ring spectra [9, chapter VII, §2, §3], one can identify

$$\mathrm{THH}(\mathrm{THH}(A)) = S^1 \otimes (S^1 \otimes A)$$

with $(S^1 \times S^1) \otimes A$ and this is *torus homology* of A . Similarly, any n -fold iteration of algebraic K-theory of A has an iterated trace map to $(S^1)^n \otimes A$. There are calculations of torus homology of $H\mathbb{F}_p$ for small n by Rognes, Veen [20] and Ausoni-Dundas, but a general result is missing. However, the homotopy type of $S^n \otimes H\mathbb{F}_p$ is known for every n and for small n $(S^1)^n \otimes A$ splits as follows: We have that $\Sigma(S^1)^n \simeq \Sigma(\bigvee_{i=1}^n \binom{n}{i} S^1)$ and one obtains for small n

$$(S^1)^n \otimes H\mathbb{F}_p \simeq \left(\bigvee_{i=1}^n \binom{n}{i} S^1 \right) \otimes H\mathbb{F}_p.$$

This gave rise to the question whether $\mathcal{L}_X^R(A; C)$ is a *stable invariant*, i.e., whether the homotopy type of $\mathcal{L}_X^R(A; C)$ only depends on the homotopy type of ΣX . There are positive results: $\mathcal{L}_X^{Hk}(HA)$ is a stable invariant if k is a field and A is a commutative Hopf algebra over k [2, Theorem 5.2] or if k is an arbitrary commutative ring and A is a smooth k -algebra [8, Example 2.6]. But Dundas and Tenti also show [8, §3.8] that $\mathcal{L}_X^{H\mathbb{Q}}(H\mathbb{Q}[t]/t^2)$ is *not* a stable invariant. They show that $\mathcal{L}_{S^1 \vee S^1 \vee S^2}^{H\mathbb{Q}}(H\mathbb{Q}[t]/t^2)$ and

Date: May 15, 2019.

We thank Bjørn Dundas for many helpful discussion and for spotting several dumb mistakes in earlier versions of this paper.

$\mathcal{L}_{S^1 \times S^1}^{H\mathbb{Q}}(H\mathbb{Q}[t]/t^2)$ differ and that reducing the coefficients from $H\mathbb{Q}[t]/t^2$ to $H\mathbb{Q}$ doesn't eliminate this discrepancy. This also implies that $\mathcal{L}_X^S(H\mathbb{Q}[t]/t^2)$ and $\mathcal{L}_X^S(H\mathbb{Q}[t]/t^2; H\mathbb{Q})$ are not stable invariants.

Our aim is to investigate the question of stability in a systematic manner. We start by giving several different notions of stability. Instead of asking for equivalent homotopy types of $\mathcal{L}_X^R(A; C)$ and $\mathcal{L}_Y^R(A; C)$ if $\Sigma X \simeq \Sigma Y$ we are asking when we actually get an equivalence $\mathcal{L}_X^R(A; C) \simeq \mathcal{L}_Y^R(A; C)$ of augmented commutative C -algebras. There are intermediate notions that ask for less structure to be preserved, for instance, that the equivalence $\mathcal{L}_X^R(A; C) \simeq \mathcal{L}_Y^R(A; C)$ is one of commutative R -algebras or of C - or R -modules.

We establish that stability is preserved by several constructions such as base-change and products but we also show which procedures do not preserve stability. For instance stability is not a transitive property: if $R \rightarrow A$ and $A \rightarrow B$ satisfy stability then this does not imply that $R \rightarrow B$ has this property.

The main purpose of this paper is to establish new cases where stability holds. For instance for any regular quotient $R \rightarrow R/(a_1, \dots, a_n)$ of a commutative ring R we obtain stability for the induced map of commutative ring spectra $HR \rightarrow HR/(a_1, \dots, a_n)$. Free commutative ring spectra generated by a module spectrum satisfy stability and we suggest a notion of *really smooth* maps of commutative ring spectra. These are maps $R \rightarrow A$ that can be factored as the canonical inclusion of R into a free commutative R -algebra spectrum followed by a map that satisfies étale descent. We show that really smooth maps satisfy stability. Other examples where stability holds are Thom spectra as well as $S \rightarrow KU$ and $S \rightarrow KO$.

For calculation one often doesn't really need stability, but the property of the suspension to decompose products is the crucial feature that one wants to have on the level of $\mathcal{L}_{(-)}^R(A; C)$. Therefore we say that $R \rightarrow A \rightarrow C$ decomposes products if

$$\mathcal{L}_{X \times Y}^R(A; C) \simeq \mathcal{L}_{X \vee Y \vee X \wedge Y}^R(A; C)$$

for all pointed simplicial sets X and Y . We use Greenlees' spectral sequence [10, Lemma 3.1] in the case $C = Hk$ for k a field to show that this decomposition property is preserved under forming suitable retracts.

1.1. Definition of $\mathcal{L}_X^R(A; C)$. Let X be a finite pointed simplicial set and let $R \rightarrow A \rightarrow C$ be a sequence of maps of commutative ring spectra. We assume that R is a cofibrant commutative S -algebra and that A and C are cofibrant commutative R -algebras. The cofibrancy assumptions on R , A and C will ensure that the homotopy type of $\mathcal{L}_X^R(A; C)$ is well-defined:

The *Loday construction with respect to X of A over R with coefficients in C* is the simplicial commutative augmented C -algebra spectrum $\mathcal{L}_X^R(A; C)$ whose p -simplices are

$$C \wedge \bigwedge_{x \in X_p \setminus *} A$$

where the smash products are taken over R . Here, $*$ denotes the basepoint of X and we place a copy of C at the basepoint. As the smash product over R is the coproduct in the category of commutative R -algebra spectra, the simplicial structure is straightforward: Face maps d_i on X induce multiplication in A or the A -action on C if the basepoint is involved. The degeneracy maps s_i on X cause the insertion of the unit map $\eta_A: R \rightarrow A$ over all n -simplices which are not hit by $s_i: X_{n-1} \rightarrow X_n$. As defined, $\mathcal{L}_X^R(A; C)$ is a simplicial commutative augmented C -algebra spectrum. We use the same symbol $\mathcal{L}_X^R(A; C)$ for its geometric realization. For $C = A$ we abbreviate $\mathcal{L}_X^R(A; A)$ by $\mathcal{L}_X^R(A)$. If $R = S$, then we omit it from the notation. Note that $\mathcal{L}_X^R(A)$ is by definition [9, VII, §2, §3] equal to $X \otimes A$ where $X \otimes A$ is formed in the category of commutative R -algebras.

If X is an arbitrary pointed simplicial set, then we can write it as the colimit of its finite pointed subcomplexes and the Loday construction with respect to X can then also be expressed as the colimit of the Loday construction for the finite subcomplexes.

2. NOTIONS OF STABILITY

The weakest notion of stability just asks for an abstract equivalence in the stable homotopy category:

Definition 2.1.

- (1) Let $R \rightarrow A$ be a cofibration of commutative S -algebras with R cofibrant. We call $R \rightarrow A$ *stable* if for every pair of pointed simplicial sets X and Y an equivalence $\Sigma X \simeq \Sigma Y$ implies that $\mathcal{L}_X^R(A) \simeq \mathcal{L}_Y^R(A)$.

- (2) Let $S \rightarrow R \rightarrow A \rightarrow B$ be a sequence of cofibrations of commutative S -algebras. Then we call (R, A, B) *stable*, if for every pair of pointed simplicial sets X and Y an equivalence $\Sigma X \simeq \Sigma Y$ implies that $\mathcal{L}_X^R(A; B) \simeq \mathcal{L}_Y^R(A; B)$.

Examples 2.2.

- Dundas and Tenti show that for any discrete smooth k -algebra A we have that $Hk \rightarrow HA$ is stable [8, Example 2.6].
- They show, however, that $H\mathbb{Q} \rightarrow H\mathbb{Q}[t]/t^2$ and $(H\mathbb{Q}, H\mathbb{Q}[t]/t^2, H\mathbb{Q})$ are not stable.
- If A is a commutative Hopf algebra over a field k , then Berest, Ramadoss and Yeung prove [2, §5] that $Hk \rightarrow HA$ and (Hk, HA, Hk) are stable.
- In [3] we show that for any sequence of cofibrations of commutative S -algebras $S \rightarrow A \rightarrow B \rightarrow A$ we get that

$$\mathcal{L}_X^B(A) \simeq \mathcal{L}_{\Sigma X}^A(B; A)$$

as augmented commutative A -algebras and hence $B \rightarrow A$ is stable if B is a cofibrant commutative augmented A -algebra.

In the above definition we just require an abstract weak equivalence, but one can also pose additional conditions on the equivalence $\mathcal{L}_X^R(A; B) \simeq \mathcal{L}_Y^R(A; B)$. A strong version of stability is the following:

Definition 2.3.

- (1) Let $R \rightarrow A$ be a cofibration of commutative S -algebras with R cofibrant. We call $R \rightarrow A$ *multiplicatively stable* if for every pair of pointed simplicial sets X and Y an equivalence $\Sigma X \simeq \Sigma Y$ implies that $\mathcal{L}_X^R(A) \simeq \mathcal{L}_Y^R(A)$ as commutative augmented A -algebra spectra.
- (2) Let $S \xrightarrow{\quad} R \xrightarrow{\alpha} A \xrightarrow{\beta} B$ be a sequence of cofibrations of commutative S -algebras. Then we call $R \rightarrow A \rightarrow B$ *multiplicatively stable* if for every pair of pointed simplicial sets X and Y an equivalence $\Sigma X \simeq \Sigma Y$ implies that $\mathcal{L}_X^R(A; B) \simeq \mathcal{L}_Y^R(A; B)$ and $\mathcal{L}_X^R(B) \simeq \mathcal{L}_Y^R(B)$ as commutative augmented B -algebras such that the diagram

$$\begin{array}{ccc} \mathcal{L}_X^R(A; B) & \xrightarrow{\simeq} & \mathcal{L}_Y^R(A; B) \\ \mathcal{L}_X^R(\beta) \downarrow & & \downarrow \mathcal{L}_X^R(\beta) \\ \mathcal{L}_X^R(B) & \xrightarrow{\simeq} & \mathcal{L}_Y^R(B) \end{array}$$

commutes.

Of course, there is a whole hierarchy of notions of stability. Instead of asking that the equivalence $\mathcal{L}_X^R(A) \simeq \mathcal{L}_Y^R(A)$ is one of augmented commutative A -algebras, we could ask for one of augmented commutative R -algebras or A - or just R -modules.

Definition 2.4. Let $R \rightarrow A$ be a cofibration of commutative S -algebras with R cofibrant. We call $R \rightarrow A$ *A -linearly stable* if for every pair of pointed simplicial sets X and Y an equivalence $\Sigma X \simeq \Sigma Y$ implies that $\mathcal{L}_X^R(A) \simeq \mathcal{L}_Y^R(A)$ as A -modules. Similarly, we call $R \rightarrow A$ *R -linearly stable* if for every pair of pointed simplicial sets X and Y an equivalence $\Sigma X \simeq \Sigma Y$ gives rise to an equivalence of R -modules $\mathcal{L}_X^R(A) \simeq \mathcal{L}_Y^R(A)$.

Remark 2.5. If $R \rightarrow A$ is A -linearly stable, then (R, A, B) is stable because

$$\mathcal{L}_X^R(A; B) \simeq \mathcal{L}_X^R(A) \wedge_A B.$$

If $R \rightarrow A$ is multiplicatively stable, then so is $R \rightarrow A \rightarrow B$.

A converse might not be true: Even if B is faithful as an A -module, we might not know that the equivalence $\mathcal{L}_X^R(A) \wedge_A B \simeq \mathcal{L}_Y^R(A) \wedge_A B$ is of the form $f \wedge_A B$, so we cannot deduce that $\mathcal{L}_X^R(A) \simeq \mathcal{L}_Y^R(A)$.

Let us start with several examples of multiplicative stability.

Proposition 2.6. *If B is an augmented commutative A -algebra, then $B \rightarrow A$ and $A \rightarrow \mathcal{L}_{\Sigma X}^A(B; A) \rightarrow A$ are multiplicatively stable.*

Proof. In the augmented case $A \rightarrow B \rightarrow A$, as an equivalence $\Sigma X \simeq \Sigma Y$ implies that $\mathcal{L}_{\Sigma X}^A(B; A) \simeq \mathcal{L}_{\Sigma Y}^A(B; A)$ as augmented commutative A -algebras, we also get that $\mathcal{L}_X^B(A) \simeq \mathcal{L}_Y^B(A)$ as augmented commutative A -algebras by applying [3, Theorem 3.3] to the sequence of maps $A = A \rightarrow B \rightarrow A$, so $B \rightarrow A$ is multiplicatively stable.

For the second claim observe that

$$\mathcal{L}_Y^A(\mathcal{L}_{\Sigma X}^A(B; A); A) \simeq \mathcal{L}_{Y \wedge \Sigma X}^A(B; A) = \mathcal{L}_{\Sigma Y \wedge X}^A(B; A).$$

Observe that for all X we have that $\mathcal{L}_X^A(A) \simeq A$ so $A \rightarrow \mathcal{L}_{\Sigma X}^A(B; A) \rightarrow A$ is multiplicatively stable. \square

Loday constructions for suspensions are stable:

Theorem 2.7. *Let $R \rightarrow A$ be a cofibration of commutative S -algebras with R cofibrant. Then $A \rightarrow \mathcal{L}_{\Sigma X}^R(A)$ is multiplicatively stable for all X .*

Proof. We have to show that $\mathcal{L}_Y^A(\mathcal{L}_{\Sigma X}^R(A))$ only depends on the homotopy type of ΣY . We first identify $\mathcal{L}_Y^A(\mathcal{L}_{\Sigma X}^R(A))$ with the help of [11, Remark 3.3] as

$$\begin{aligned} \mathcal{L}_Y^A(\mathcal{L}_{\Sigma X}^R(A)) &\simeq \mathcal{L}_Y^R(\mathcal{L}_{\Sigma X}^R(A)) \wedge_{\mathcal{L}_Y^R(A)} A \\ &\simeq \mathcal{L}_{Y \times \Sigma X}^R(A) \wedge_{\mathcal{L}_Y^R(A)} A \\ &\simeq \mathcal{L}_{(Y \times \Sigma X) \cup Y^*}^R(A) \\ &\simeq \mathcal{L}_{Y_+ \wedge \Sigma X}^R(A) \cong \mathcal{L}_{\Sigma(Y_+) \wedge X}^R(A). \end{aligned}$$

As $\Sigma(Y_+) \simeq \Sigma Y \vee S^1$, this depends only on ΣY . \square

Example 2.8. Applying Theorem 2.7 to $H\mathbb{F}_p$ and $\Sigma X = S^2$ gives that the map

$$H\mathbb{F}_p \rightarrow \mathrm{THH}^{[2]}(H\mathbb{F}_p) \simeq H\mathbb{F}_p \vee \Sigma^3 H\mathbb{F}_p$$

is multiplicatively stable for all primes p .

As we know from the algebraic setting that smooth algebras are stable, it is natural to consider free commutative A -algebra spectra. Let M be an A -module spectrum for some commutative S -algebra A . We consider the free commutative A -algebra on M ,

$$\mathbb{P}_A(M) = \bigvee_{n \geq 0} M^{\wedge n} / \Sigma_n$$

with the usual convention that $M^{\wedge 0} / \Sigma_0 = A$.

In the following we use several categories, so let's fix some notation. Let \mathcal{U} denote the category of unbased (compactly generated weak Hausdorff) spaces. For a commutative ring spectrum R , \mathcal{M}_R denotes the category of R -module spectra and \mathcal{C}_R denotes the category of commutative R -algebras.

Lemma 2.9. *For every simplicial set X there is a weak equivalence of commutative A -algebras*

$$\mathcal{L}_X^A(\mathbb{P}_A(M)) \simeq \mathbb{P}_A(X_+ \wedge M).$$

Proof. For the proof we use the fact that the category of commutative A -algebras is tensored over unpointed topological spaces and simplicial sets in a compatible way [9, VII §2, §3]. Note that $\mathcal{L}_X^A(\mathbb{P}_A(M)) = X \otimes_A \mathbb{P}_A(M)$ in the notation of [9].

We have the following chain of bijections for an arbitrary commutative A -algebra B :

$$\begin{aligned} \mathcal{C}_A(X \otimes_A \mathbb{P}_A(M), B) &\cong \mathcal{U}(X, \mathcal{C}_A(\mathbb{P}_A(M), B)) \\ &\cong \mathcal{U}(X, \mathcal{M}_A(M, B)) \\ &\cong \mathcal{M}_A(X_+ \wedge M, B) \\ &\cong \mathcal{C}_A(\mathbb{P}_A(X_+ \wedge M), B) \end{aligned}$$

where $X_+ \wedge M$ is the tensor of X with M in the category of A -modules. Hence the Yoneda lemma implies the claim. \square

Corollary 2.10. *In the setting above, if $\Sigma X \simeq \Sigma Y$, then $\mathcal{L}_X^A(\mathbb{P}_A(M)) \simeq \mathcal{L}_Y^A(\mathbb{P}_A(M))$ as commutative A -algebras.*

Proof. If $\Sigma X \simeq \Sigma Y$, then $\Sigma_+^\infty X \simeq \Sigma_+^\infty Y$ and as $X_+ \wedge M = \Sigma_+^\infty X \wedge M$ this implies that $\mathbb{P}_A(X_+ \wedge M) \simeq \mathbb{P}_A(Y_+ \wedge M)$ as commutative A -algebras. \square

The following example was also considered in [15, Lemma 5.5]. A cofibration $A \rightarrow B$ of commutative S -algebras with A cofibrant is called THH -étale if the canonical map $B \rightarrow \mathrm{THH}^A(B)$ is a weak equivalence.

Proposition 2.11. *If $A \rightarrow B$ is THH-étale, then for all connected pointed X the canonical map $B \rightarrow \mathcal{L}_X^A(B)$ is an equivalence. Hence, as this map is a map of augmented commutative B -algebras, $\mathcal{L}_X^A(B) \simeq \mathcal{L}_Y^A(B)$ for any pair of connected simplicial sets X and Y .*

Proof. The proof is by induction on the top dimension of a non-degenerate simplex in a finite connected simplicial set, and then by taking colimits in the infinite case. A connected 0-dimensional simplicial set consists of a point, where there is nothing to prove. Any 1-dimensional connected finite simplicial set is homotopy equivalent to a wedge of circles, so if $X \simeq S^1 \vee S^1 \vee \dots \vee S^1$ and $B \simeq \mathcal{L}_{S^1}^A B$,

$$\mathcal{L}_X^A B \simeq B \wedge_B B \wedge_B \dots \wedge_B B \simeq B.$$

Once we know the result for simplicial sets of dimension $\leq n - 1$, if we get a simplicial set X with a finite number of non-degenerate n -cells we proceed by induction on the number of those. As in the proof of Proposition 8-4 in [3], using the homotopy invariance of the construction and subdivision, if needed, we can assume that X can be constructed by adding a new non-degenerate simplex with an embedded boundary to a simplicial set homotopy equivalent to X with one non-degenerate n -cell deleted, for which the proposition holds by the induction on the number of non-degenerate n -cells. By the inductive hypothesis it also holds for the embedded boundary $\partial\Delta^n$, and since the new simplex being added is homotopy equivalent to a point, the proposition holds for it. By the connectivity and by homotopy invariance we can also assume that the basepoint of X is contained in the boundary of the new simplex being attached, so the identifications of all three Loday constructions with B are compatible. Then $\mathcal{L}_X^A(B) \simeq B \wedge_B B \simeq B$. \square

Remark 2.12. Examples of THH-étale maps $A \rightarrow B$ are Galois extensions in the sense of [16] but also étale maps in the sense of Lurie [13, Definition 7.5.1.4]. For a careful discussion of these notions and for comparison results see [14].

3. INHERITANCE PROPERTIES AND DESCENT

With the assumption of multiplicative stability we get a descent result:

Theorem 3.1. *If $R \rightarrow A \rightarrow B$ is multiplicatively stable, then $A \rightarrow B$ is multiplicatively stable.*

Proof. Let's assume that $\Sigma X \simeq \Sigma Y$. Then by assumption we get that $\mathcal{L}_X^R(B) \simeq \mathcal{L}_Y^R(B)$ and $\mathcal{L}_X^R(A; B) \simeq \mathcal{L}_Y^R(A; B)$ as commutative augmented B -algebras, compatibly with the module structure of the former over the latter. The Juggling Lemma [3, Lemma 3.1] yields an equivalence of augmented commutative B -algebras

$$\mathcal{L}_X^A(B) \simeq B \wedge_{\mathcal{L}_X^R(A; B)} \mathcal{L}_X^R(B) \text{ and } \mathcal{L}_Y^A(B) \simeq B \wedge_{\mathcal{L}_Y^R(A; B)} \mathcal{L}_Y^R(B).$$

Our assumptions guarantee that therefore $\mathcal{L}_X^A(B) \simeq \mathcal{L}_Y^A(B)$ as commutative augmented B -algebras. \square

One can upgrade this slightly and introduce coefficients:

Corollary 3.2. *If $S \rightarrow R \rightarrow A \rightarrow B \rightarrow C$ is a sequence of cofibrations of commutative A -algebras and both $R \rightarrow A \rightarrow C$ and $R \rightarrow B \rightarrow C$ are multiplicatively stable, then $A \rightarrow B \rightarrow C$ is multiplicatively stable as well.*

Lemma 3.3. *Let $A \leftarrow R \rightarrow B$ be cofibrations of commutative S -algebras with R cofibrant, then $\mathcal{L}_X^A(A \wedge_R B) \cong A \wedge_R \mathcal{L}_X^R(B)$ as simplicial commutative augmented $A \wedge_R B$ -algebras and hence on realizations as commutative augmented $A \wedge_R B$ -algebras.*

Proof. There is a direct isomorphism sending $A \wedge_R (B \wedge_R \dots \wedge_R B)$ to $(A \wedge_R B) \wedge_A \dots \wedge_A (A \wedge_R B)$ and this isomorphism is compatible with the multiplication. \square

This implies that stability is closed under base-change:

Proposition 3.4. *Let A and E be cofibrant commutative S -algebra spectra. If $R \rightarrow A$ is R -linearly stable, then so is $E \rightarrow E \wedge_R A$. If $R \rightarrow A$ is multiplicatively stable, then so is $E \rightarrow E \wedge_R A$.*

Proof. Assume that $\Sigma X \simeq \Sigma Y$. Then by assumption $\mathcal{L}_X^R(A) \simeq \mathcal{L}_Y^R(A)$ as R -modules or as augmented commutative A -algebras. But then also $E \wedge_R \mathcal{L}_X^R(A) \simeq E \wedge_R \mathcal{L}_Y^R(A)$ and by Lemma 3.3 this implies

$$\mathcal{L}_X^E(E \wedge_R A) \simeq \mathcal{L}_Y^E(E \wedge_R A).$$

\square

Remark 3.5. Note that the above implication cannot be upgraded to an equivalence: starting with the assumption that $\mathcal{L}_X^E(E \wedge_R A) \simeq \mathcal{L}_Y^E(E \wedge_R A)$, we get $E \wedge_R \mathcal{L}_X^R(A) \simeq E \wedge_R \mathcal{L}_Y^R(A)$. Even if $\mathcal{L}_X^R(A)$ and $\mathcal{L}_Y^R(A)$ are E -local in the category of R -modules, however, we don't know that the weak equivalence $E \wedge_R \mathcal{L}_X^R(A) \simeq E \wedge_R \mathcal{L}_Y^R(A)$ is of the form $E \wedge_R f$ (or a zigzag of such maps), but for the E -local Whitehead Theorem [4, Lemma 1.2] we have to have a map and not just an abstract isomorphism of E_* -homology groups.

Smashing with a fixed commutative R -algebra preserves stability:

Lemma 3.6. *Let A, B and C be cofibrant commutative R -algebras. Then there is an equivalence of commutative augmented $C \wedge_R B$ -algebras*

$$\mathcal{L}_X^{C \wedge_R A}(C \wedge_R B) \simeq C \wedge_R \mathcal{L}_X^A(B).$$

Hence if $f: A \rightarrow B$ is multiplicatively stable, then so is $C \wedge_R f: C \wedge_R A \rightarrow C \wedge_R B$.

Proof. The equivalence

$$\mathcal{L}_X^{C \wedge_R A}(C \wedge_R B) \simeq C \wedge_R \mathcal{L}_X^A(B)$$

is based on the equivalence

$$(C \wedge_R B) \wedge_{C \wedge_R A} (C \wedge_R B) \simeq C \wedge_R (B \wedge_A B).$$

□

Proposition 3.7. *Let R be a commutative ring and let $a \in R$ be a regular element. Then $HR \rightarrow HR/a$ is multiplicatively stable.*

Proof. We consider the pushout $HR \wedge_{HR[t]}^L HR$ where the right algebra map $R[t] \rightarrow R$ sends t to zero and the left algebra map sends t to a . Note that with respect to both of these maps $HR[t]$ is an augmented commutative HR -algebra spectrum. The Künneth spectral sequence for $\pi_*(HR \wedge_{HR[t]}^L HR)$ has as its E^2 -term $\mathrm{Tor}_{*,*}^{R[t]}(R, R)$ and we take the standard free $R[t]$ resolution

$$0 \longrightarrow R[t] \xrightarrow{t} R[t]$$

of R . Applying $(-)\otimes_{R[t]} R$ yields

$$\begin{array}{ccc} 0 & \longrightarrow & R[t] \otimes_{R[t]} R & \xrightarrow[\mathrm{id} \otimes a]{t \otimes \mathrm{id}} & R[t] \otimes_{R[t]} R \\ & & \cong \downarrow & & \downarrow \cong \\ 0 & \longrightarrow & R & \xrightarrow{a} & R. \end{array}$$

Note, that the regularity of a is needed to ensure injectivity on the left hand side.

We apply Lemma 3.3 and choose a cofibrant model of HR as a commutative $HR[t]$ -algebra and obtain

$$\mathcal{L}_X^{HR}(HR/a) \simeq \mathcal{L}_X^{HR}(HR \wedge_{HR[t]} HR) \simeq HR \wedge_{HR[t]} \mathcal{L}_X^{HR[t]}(HR)$$

where the right $HR[t]$ -module structure of $\mathcal{L}_X^{HR[t]}(HR)$ factors through the augmentation map sending t to 0. Assume that $\Sigma X \simeq \Sigma Y$. By Proposition 2.6 we have that $HR[t] \rightarrow HR$ is multiplicatively stable, so $\mathcal{L}_X^{HR[t]}(HR) \simeq \mathcal{L}_Y^{HR[t]}(HR)$ as commutative augmented HR -algebras. This yields an equivalence of commutative augmented $HR \wedge_{HR[t]} HR \simeq HR/a$ -algebras between $\mathcal{L}_X^{HR}(HR/a)$ and $\mathcal{L}_Y^{HR}(HR/a)$. □

Remark 3.8. The above result can be used for calculating torus homology for instance for $H\mathbb{Z} \rightarrow H\mathbb{Z}/p\mathbb{Z}$ for every prime p : We know the homotopy type of $\mathcal{L}_{S^k}^{H\mathbb{Z}}(H\mathbb{Z}/p\mathbb{Z})$ by [3, Proposition 5.3] for all k and therefore we get the homotopy type of $\mathcal{L}_{(S^1)^n}^{H\mathbb{Z}}(H\mathbb{Z}/p\mathbb{Z})$ as smash products over $H\mathbb{Z}/p\mathbb{Z}$ of $\binom{n}{k}$ copies of $\mathcal{L}_{S^k}^{H\mathbb{Z}}(H\mathbb{Z}/p\mathbb{Z})$ for $1 \leq k \leq n$.

Corollary 3.9. *For every commutative ring R and every regular element $a \in R$ the square-zero extension*

$$HR/a \rightarrow HR/a \vee \Sigma HR/a$$

is multiplicatively stable. In particular, for every commutative ring R , $HR \rightarrow HR \vee \Sigma HR$ is multiplicatively stable.

Proof. As $HR \rightarrow HR/a$ is multiplicatively stable we get by Lemma 3.3 that

$$\mathcal{L}_X^{HR/a}(HR/a \wedge_{HR} HR/a) \simeq HR/a \wedge_{HR} \mathcal{L}_X^{HR}(HR/a)$$

as augmented commutative $HR/a \wedge_{HR} HR/a$ -algebras and hence $HR/a \rightarrow HR/a \wedge_{HR} HR/a$ is multiplicatively stable. The Künneth spectral sequence yields that $\pi_*(HR/a \wedge_{HR} HR/a) \cong \Lambda_{R/a}(x)$ with $|x| = 1$. By [7, Proposition 2.1] this implies that

$$HR/a \wedge_{HR} HR/a \simeq HR/a \vee \Sigma HR/a$$

as a commutative augmented HR/a -algebra.

Considering the regular element $t \in R[t]$ gives that $HR \rightarrow HR \vee \Sigma HR$ is multiplicatively stable. \square

Stability is inherited by Loday constructions.

Proposition 3.10. *If $R \rightarrow A$ is multiplicatively stable, then so is $R \rightarrow \mathcal{L}_Z^R(A)$ for any Z .*

Proof. Assume that $\Sigma X \simeq \Sigma Y$. As $\Sigma(X \times Z) \simeq \Sigma X \vee \Sigma Z \vee \Sigma X \wedge Z$ we get that

$$\Sigma(X \times Z) \simeq \Sigma X \vee \Sigma Z \vee \Sigma X \wedge Z \simeq \Sigma Y \vee \Sigma Z \vee \Sigma Y \wedge Z \simeq \Sigma(Y \times Z)$$

and thus, as $R \rightarrow A$ is multiplicatively stable

$$\mathcal{L}_X^R(\mathcal{L}_Z^R(A)) \simeq \mathcal{L}_{X \times Z}^R(A) \simeq \mathcal{L}_{Y \times Z}^R(A) \simeq \mathcal{L}_Y^R(\mathcal{L}_Z^R(A)).$$

\square

Remark 3.11. One can interpret Proposition 3.10 as the statement that Loday constructions preserve stability because for all Z there is an equivalence of augmented commutative R -algebras $R \simeq \mathcal{L}_Z^R(R)$.

Note that we know that the square zero extensions $H\mathbb{F}_p \rightarrow H\mathbb{F}_p \vee \Sigma^3 H\mathbb{F}_p$ (Example 2.8) and $H\mathbb{F}_p \rightarrow H\mathbb{F}_p \vee \Sigma H\mathbb{F}_p$ (Corollary 3.9) are multiplicatively stable. However, if place the module $H\mathbb{F}_p$ in degree zero, then the following result shows that the square zero extension $H\mathbb{F}_p \rightarrow H\mathbb{F}_p \vee H\mathbb{F}_p$ is *not* multiplicatively stable for odd primes p . The proof is a direct adaptation of [8, §3.8].

Theorem 3.12. *Let p be an odd prime. Then $(H\mathbb{F}_p, H\mathbb{F}_p[t]/t^2, H\mathbb{F}_p)$ is not stable.*

Corollary 3.13. *The commutative $H\mathbb{F}_p$ -algebra $H\mathbb{F}_p[t]/t^2$ is neither multiplicatively stable nor $H\mathbb{F}_p[t]/t^2$ -linearly stable over $H\mathbb{F}_p$.*

Proof. If it were, then this would imply that $(H\mathbb{F}_p, H\mathbb{F}_p[t]/t^2, H\mathbb{F}_p)$ is stable. \square

Proof of Theorem 3.12. We know that

$$\pi_* \mathcal{L}_{S^1 \vee S^1 \vee S^2}^{\mathbb{F}_p}(\mathbb{F}_p[t]/t^2; \mathbb{F}_p) \cong \pi_* \mathcal{L}_{S^1}^{\mathbb{F}_p}(\mathbb{F}_p[t]/t^2; \mathbb{F}_p) \otimes_{\mathbb{F}_p} \pi_* \mathcal{L}_{S^1}^{\mathbb{F}_p}(\mathbb{F}_p[t]/t^2; \mathbb{F}_p) \otimes_{\mathbb{F}_p} \pi_* \mathcal{L}_{S^2}^{\mathbb{F}_p}(\mathbb{F}_p[t]/t^2; \mathbb{F}_p)$$

and by [3] we know what the tensor factors are:

$$\pi_* \mathcal{L}_{S^1}^{\mathbb{F}_p}(\mathbb{F}_p[t]/t^2; \mathbb{F}_p) \cong \mathbf{HH}_*^{\mathbb{F}_p}(\mathbb{F}_p[t]/t^2; \mathbb{F}_p) \cong \Lambda_{\mathbb{F}_p}(\varepsilon t) \otimes_{\mathbb{F}_p} \Gamma_{\mathbb{F}_p}(\varphi^0 t)$$

and

$$\pi_* \mathcal{L}_{S^2}^{\mathbb{F}_p}(\mathbb{F}_p[t]/t^2; \mathbb{F}_p) \cong \mathbf{HH}_*^{[2], \mathbb{F}_p}(\mathbb{F}_p[t]/t^2; \mathbb{F}_p) \cong \Gamma_{\mathbb{F}_p}(\varrho^0 \varepsilon t) \otimes \bigotimes_k (\Lambda_{\mathbb{F}_p}(\varepsilon \varphi^k t) \otimes \Gamma_{\mathbb{F}_p}(\varphi^0 \varphi^k t)).$$

Torus homology is the total complex of the bicomplex for $\mathcal{L}_{S^1 \times S^1}^{\mathbb{F}_p}(\mathbb{F}_p[t]/t^2; \mathbb{F}_p)$ as in [8]. In the bicomplex in bidegree (n, m) we have the term

$$\mathcal{L}_{[n] \times [m]}^{\mathbb{F}_p}(\mathbb{F}_p[t]/t^2; \mathbb{F}_p) \cong \mathbb{F}_p \otimes_{\mathbb{F}_p} (\mathbb{F}_p[t]/t^2)^{(n+1)(m+1)-1} \cong (\mathbb{F}_p[t]/t^2)^{(n+1)(m+1)-1}.$$

In total degree one we have contributions from $(0, 1)$ and $(1, 0)$ that we call y_1^v and y_1^h as in [8, §3.8].

Everything is a cycle here and these elements correspond to \otimes and $1 \otimes t$.

From now on we suppress the tensor signs from the notation and we denote the generators by matrices. In total degree two there are three possibilities $(0, 2)$, $(1, 1)$ and $(2, 0)$. There are the classes y_2^v in bidegree

$(0, 2)$, and y_2^h in bidegree $(2, 0)$ corresponding to the standard Hochschild generators $\begin{pmatrix} 1 \\ t \\ t \end{pmatrix}$ and $\begin{pmatrix} 1 & t \\ & t \end{pmatrix}$.

In bidegree (1, 1) there are the following possibilities for non-degenerate cycles:

$$\begin{pmatrix} 1 & t \\ t & t \end{pmatrix}, \begin{pmatrix} 1 & t \\ t & 1 \end{pmatrix}, \begin{pmatrix} 1 & t \\ 1 & t \end{pmatrix}, \begin{pmatrix} 1 & 1 \\ t & t \end{pmatrix}, \text{ and } \begin{pmatrix} 1 & 1 \\ 1 & t \end{pmatrix}.$$

As we are working over \mathbb{F}_p for an odd prime p , 2 is invertible. The boundary of $\frac{1}{2} \begin{pmatrix} 1 & 1 \\ 1 & t \end{pmatrix}$ is $\begin{pmatrix} 1 & t \\ 1 & t \end{pmatrix}$,

the boundary of $\frac{1}{2} \begin{pmatrix} 1 & 1 & 1 \\ 1 & t & t \end{pmatrix}$ is $\begin{pmatrix} 1 & 1 \\ t & t \end{pmatrix}$. Finally, we identify $\begin{pmatrix} 1 & t \\ t & t \end{pmatrix}$ as the boundary of $\begin{pmatrix} 1 & 1 \\ 1 & t \end{pmatrix}$.

The element $\begin{pmatrix} 1 & 1 \\ t & 1 \\ 1 & t \end{pmatrix}$ ensures that $\begin{pmatrix} 1 & t \\ t & 1 \end{pmatrix}$ is homologous to $\begin{pmatrix} 1 & 1 \\ t & t \end{pmatrix}$, so we are left with the generator

in bidegree (1, 1) given by $\begin{pmatrix} 1 & 1 \\ 1 & t \end{pmatrix}$.

So we get (at most) a 3-dimensional vector space in total degree 2.

In $\pi_2 \mathcal{L}_{S^1 \vee S^1 \vee S^2}^{\mathbb{F}_p}(\mathbb{F}_p[t]/t^2; \mathbb{F}_p)$ however, we get the generators $\varphi^0 t \otimes 1 \otimes 1$, $1 \otimes \varphi^0 t \otimes 1$, $\varepsilon t \otimes \varepsilon t \otimes 1$ and $1 \otimes 1 \otimes \rho^0 \varepsilon t$, so we have a 4-dimensional vector space. \square

Remark 3.14. For odd primes 2 is invertible and this reduces the number of generators in total degree 2 to 3 in the torus homology of $\mathbb{F}_p[t]/t^2$ over \mathbb{F}_p with \mathbb{F}_p -coefficients. For $p = 2$ one can check that there is an extra class coming from $\begin{pmatrix} 1 & t \\ t & 1 \end{pmatrix}$ which is homologous to $\begin{pmatrix} 1 & 1 \\ t & t \end{pmatrix}$ and to $\begin{pmatrix} 1 & t \\ 1 & t \end{pmatrix}$ so together with the class $\begin{pmatrix} 1 & 1 \\ 1 & t \end{pmatrix}$ this gives two generators in bidegree (1, 1) and the ones in (2, 0) and (0, 2) giving a total of dimension 4. As $\mathbb{F}_2[t]/t^2$ is a commutative Hopf algebra over \mathbb{F}_2 , we know that $\mathbb{F}_2 \rightarrow \mathbb{F}_2[t]/t^2$ and $\mathbb{F}_2 \rightarrow \mathbb{F}_2[t]/t^2 \rightarrow \mathbb{F}_2$ are stable.

We can model $S[t]/t^2$ as $S \wedge \Pi_+$ for the commutative pointed monoid $\Pi_+ = \{+, 1, t\}$.

Corollary 3.15. *The map $S \rightarrow S[t]/t^2$ is not multiplicatively stable.*

Proof. If it were multiplicatively stable, then by Lemma 3.3 $H\mathbb{F}_p \rightarrow H\mathbb{F}_p[t]/t^2$ would be stable, as well. \square

Remark 3.16. Neither stability nor multiplicative stability are transitive: for every commutative ring k the map $k \rightarrow k[t]$ is smooth, hence (multiplicatively) stable and $k[t] \rightarrow k[t]/t^2$ is stable by Proposition 3.7, but for $k = \mathbb{Q}$ Dundas and Tenti show [8] that $\mathbb{Q} \rightarrow \mathbb{Q}[t]/t^2$ is not stable and for $k = \mathbb{F}_p$ the know that $\mathbb{F}_p \rightarrow \mathbb{F}_p[t]/t^2$ is not multiplicatively stable.

The Loday construction behaves nicely with respect to pushouts:

Lemma 3.17. *If $C \leftarrow A \rightarrow B$ is a diagram of cofibrations of commutative R algebras and if A is cofibrant as a commutative R -algebra, then*

$$\mathcal{L}_X^R(C \wedge_A B) \simeq \mathcal{L}_X^R(C) \wedge_{\mathcal{L}_X^R(A)} \mathcal{L}_X^R(B).$$

Proof. This equivalence is proven using an exchange of priorities in a colimit diagram based on the equivalence

$$(C \wedge_A B) \wedge_R (C \wedge_A B) \simeq (C \wedge_R C) \wedge_{A \wedge_R A} (B \wedge_R B).$$

\square

Remark 3.18. Beware that the above identification does *not* imply that multiplicative stability is closed under pushouts in the category of commutative R -algebras, because if we know that $\mathcal{L}_X^R(D) \simeq \mathcal{L}_Y^R(D)$ as commutative augmented D -algebras (for $D = A, B$ and C), then this does not imply that $\mathcal{L}_X^R(C \wedge_A B)$ is equivalent to $\mathcal{L}_Y^R(C \wedge_A B)$ because we cannot guarantee that the equivalences $\mathcal{L}_X^R(D) \simeq \mathcal{L}_Y^R(D)$ commute with the structure maps in the pushout diagram.

For example we know that $H\mathbb{Q} \rightarrow H\mathbb{Q}[t]$ and $H\mathbb{Q} \rightarrow H\mathbb{Q}[t, x]$ are multiplicatively stable, but $H\mathbb{Q} \rightarrow H\mathbb{Q}[t]/t^2$ is not stable by [8], despite the fact that we can express the latter as a pushout $H\mathbb{Q}[t] \wedge_{H\mathbb{Q}[t, x]} H\mathbb{Q}[t]$ where x maps to t^2 on the left hand side and to 0 on the right hand side.

In the case of $A = R$ we *do* get a stability result:

Corollary 3.19. *Assume that $R \rightarrow B$ and $R \rightarrow C$ are multiplicatively stable. Then so is $R \rightarrow B \wedge_R C$.*

Proof. If $\Sigma X \simeq \Sigma Y$, then $\mathcal{L}_X^R(B) \simeq \mathcal{L}_Y^R(B)$ and $\mathcal{L}_X^R(C) \simeq \mathcal{L}_Y^R(C)$ by assumption and these equivalence are of commutative augmented B - and C -algebras, so in particular of commutative augmented R -algebras. Note that $\mathcal{L}_X^R(R) \simeq R$ for all pointed X . Hence by Lemma 3.17 we obtain

$$\mathcal{L}_X^R(B \wedge_R C) \simeq \mathcal{L}_X^R(B) \wedge_R \mathcal{L}_X^R(C) \simeq \mathcal{L}_Y^R(B) \wedge_R \mathcal{L}_Y^R(C) \simeq \mathcal{L}_Y^R(B \wedge_R C)$$

and this is an equivalence of commutative augmented $B \wedge_R C$ -algebras. \square

Example 3.20. We know from Proposition 3.7 that $HR \rightarrow HR/a$ is multiplicatively stable for every commutative ring R and every regular element $a \in R$. Corollary 3.19 implies that $HR \rightarrow HR/a \wedge_{HR} HR/a$ is multiplicatively stable and as before we know that $HR/a \wedge_{HR} HR/a \simeq HR/a \vee \Sigma HR/a$, so $HR \rightarrow HR/a \vee \Sigma HR/a$ is multiplicatively stable. For instance $H\mathbb{Z} \rightarrow H\mathbb{Z}/p \vee \Sigma H\mathbb{Z}/p$ is multiplicatively stable for all primes p .

Example 3.21. Taking the coproduct (with a cofibrant model of $H\mathbb{Z}[t]$ as a commutative $H\mathbb{Z}$ -algebra)

$$\begin{array}{ccc} H\mathbb{Z} & \longrightarrow & H\mathbb{Z}/p \\ \downarrow & & \downarrow \\ H\mathbb{Z}[t] & \longrightarrow & H\mathbb{Z}[t] \wedge_{H\mathbb{Z}}^L H\mathbb{Z}/p \simeq H\mathbb{Z}/p[t] \end{array}$$

shows that $H\mathbb{Z} \rightarrow H\mathbb{Z}/p[t]$ is multiplicatively stable.

Corollary 3.22. *Let R be a commutative ring and let (a_1, \dots, a_n) be a regular sequence in R , then $HR \rightarrow HR/(a_1, \dots, a_n)$ is multiplicatively stable.*

Proof. We prove the claim for two generators; the general case follows by induction. We take a cofibrant model of HR/a_1 as a commutative HR -algebra and we consider the standard free resolution $0 \longrightarrow R \xrightarrow{a_2} R$ of R/a_2 and obtain

$$0 \longrightarrow R/a_1 \otimes_R R \xrightarrow{\text{id} \otimes a_2} R/a_1 \otimes_R R$$

because multiplication by a_2 is injective on R/a_1 . Hence the coproduct $HR/a_1 \wedge_{HR}^L HR/a_2$ of

$$\begin{array}{ccc} HR & \longrightarrow & HR/a_1 \\ \downarrow & & \\ HR/a_2 & & \end{array}$$

is $HR/(a_1, a_2)$ and by Corollary 3.19 the claim follows. \square

Proposition 3.23. *Let k be a field and let Π_+ be a pointed commutative monoid. If $S \rightarrow Hk$ and $Hk \rightarrow Hk[\Pi_+]$ are multiplicatively stable, then $\Sigma X \simeq \Sigma Y$ implies that $\mathcal{L}_X(Hk[\Pi_+]) \simeq \mathcal{L}_Y(Hk[\Pi_+])$ as augmented commutative Hk -algebras.*

Proof. This follows from the splitting of $\mathcal{L}_X^S(Hk[\Pi_+])$ as a commutative augmented Hk -algebra [12, Theorem 7.1] as

$$\mathcal{L}_X^S(Hk[\Pi_+]) \simeq \mathcal{L}_X^S(Hk) \wedge_{Hk} \mathcal{L}_X^{Hk}(Hk[\Pi_+]).$$

\square

Proposition 3.24. *If $S \rightarrow A$ and $S \rightarrow B$ are cofibrations of commutative S -algebras and if A and B are (multiplicatively) stable, then if X and Y are connected and $\Sigma X \simeq \Sigma Y$, then*

$$\mathcal{L}_X(A \times B) \simeq \mathcal{L}_Y(A \times B)$$

as commutative S -algebras.

Proof. This follows from [3, Proposition 8.4] because $\mathcal{L}_X(A \times B) \simeq \mathcal{L}_X(A) \times \mathcal{L}_X(B)$ as commutative S -algebras. \square

The following notion is investigated in [15, 14].

Definition 3.25. Let $R \rightarrow A \rightarrow B$ be a sequence of cofibrations of commutative S -algebras with R cofibrant. Then this sequence *satisfies étale descent* if for all connected X the canonical map

$$\mathcal{L}_X^R(A) \wedge_A B \rightarrow \mathcal{L}_X^R(B)$$

is an equivalence.

If $R \rightarrow A \rightarrow B$ satisfies étale descent and if X is not connected, so for example $X = X_1 \sqcup X_2$ with X_i connected for $i = 1, 2$, then the formula becomes

$$\mathcal{L}_X^R(B) = \mathcal{L}_{X_1 \sqcup X_2}^R(B) \simeq \mathcal{L}_{X_1}^R(B) \wedge_R \mathcal{L}_{X_2}^R(B) \simeq \mathcal{L}_{X_1}^R(A) \wedge_A B \wedge_R \mathcal{L}_{X_2}^R(A) \wedge_A B.$$

The property of satisfying étale descent is closed under smashing with a fixed commutative S -algebra:

Lemma 3.26. *If $R \rightarrow A \rightarrow B$ satisfies étale descent and if C is a cofibrant commutative R -algebra, then $C \rightarrow C \wedge_R A \rightarrow C \wedge_R B$ satisfies étale descent.*

Proof. We know from Lemma 3.3 that $\mathcal{L}_X^C(C \wedge_R A) \simeq C \wedge_R \mathcal{L}_X^R(A)$. Therefore an exchange of pushouts yields

$$\begin{aligned} \mathcal{L}_X^C(C \wedge_R A) \wedge_{(C \wedge_R A)} (C \wedge_R B) &\simeq (C \wedge_R \mathcal{L}_X^R(A)) \wedge_{(C \wedge_R A)} (C \wedge_R B) \\ &\simeq (C \wedge_C C) \wedge_{R \wedge_R R} (\mathcal{L}_X^R(A) \wedge_A B) \\ &\simeq C \wedge_R \mathcal{L}_X^R(B) \simeq \mathcal{L}_X^C(C \wedge_R B). \end{aligned}$$

□

In the case of étale descent we can extend stable maps and get maps that are stable for connected X :

Proposition 3.27. *Let $R \rightarrow A \rightarrow B$ be a sequence of cofibrations of commutative S -algebras with R cofibrant. If $R \rightarrow A$ is multiplicatively stable and if $R \rightarrow A \rightarrow B$ satisfies étale descent, then if $\Sigma X \simeq \Sigma Y$ for connected X and Y we can conclude that there is a weak equivalence of augmented commutative B -algebras*

$$\mathcal{L}_X^R(B) \simeq \mathcal{L}_Y^R(B).$$

Proof. As X and Y are connected and as $R \rightarrow A$ is stable, the equivalence $\Sigma X \simeq \Sigma Y$ implies that $\mathcal{L}_X^R(A) \simeq \mathcal{L}_Y^R(A)$ and with étale descent we can upgrade this to

$$\mathcal{L}_X^R(B) \simeq \mathcal{L}_X^R(A) \wedge_A B \simeq \mathcal{L}_Y^R(A) \wedge_A B \simeq \mathcal{L}_Y^R(B).$$

□

Remark 3.28. We know that $H\mathbb{Q} \rightarrow H\mathbb{Q}[t]$ is stable and as $\mathbb{Q}[t]/t^2$ and $\mathbb{Q}[t]$ are commutative augmented \mathbb{Q} -algebras, we also know that $H\mathbb{Q}[t]/t^2 \rightarrow H\mathbb{Q}$ and $H\mathbb{Q}[t] \rightarrow H\mathbb{Q}$ are stable, but since $H\mathbb{Q} \rightarrow H\mathbb{Q}[t]/t^2$ and $H\mathbb{Q} \rightarrow H\mathbb{Q}[t]/t^2 \rightarrow H\mathbb{Q}$ are not stable, we won't have general descent results. For instance in the diagram

$$\begin{array}{ccc} & & \mathbb{Q}[t]/t^2 \\ & \nearrow \eta & \downarrow \varepsilon \\ \mathbb{Q} & \xlongequal{\quad} & \mathbb{Q} \end{array}$$

the maps $H(\varepsilon)$ and the identity on $H\mathbb{Q}$ are (even multiplicatively) stable, but $H\eta$ isn't.

For morphisms that are faithful Galois extensions and satisfy étale descent, we obtain a descent result for multiplicative stability:

Theorem 3.29. *Let $A \rightarrow B$ be a faithful Galois extension with finite Galois group G and assume that $A \rightarrow B$ satisfies étale descent. Assume that $\Sigma X \simeq \Sigma Y$ for connected X and Y implies that there is a G -equivariant equivalence $\mathcal{L}_X(B) \simeq \mathcal{L}_Y(B)$ as commutative B -algebras. Then also $\mathcal{L}_X(A) \simeq \mathcal{L}_Y(A)$ as commutative A -algebras.*

Proof. The base-change result for Galois extensions [16, Lemma 7.1.1] applied to the diagram

$$\begin{array}{ccc} A & \longrightarrow & B \\ \downarrow & & \downarrow \\ \mathcal{L}_X(A) & \longrightarrow & B \wedge_A \mathcal{L}_X(A) \end{array}$$

yields that $\mathcal{L}_X(A) \rightarrow B \wedge_A \mathcal{L}_X(A)$ is a G -Galois extension and by étale descent there is an equivalence of augmented commutative B -algebras $B \wedge_A \mathcal{L}_X(A) \simeq \mathcal{L}_X(B)$ which is G -equivariant where on the left hand side the only non-trivial G -action is on the B -factor and on the right hand side G -acts on $\mathcal{L}_X(B)$ by naturality in B . Hence we get a chain of G -equivariant equivalences of commutative B -algebras

$$B \wedge_A \mathcal{L}_X(A) \simeq \mathcal{L}_X(B) \simeq \mathcal{L}_Y(B) \simeq B \wedge_A \mathcal{L}_Y(A).$$

Taking G -homotopy fixed points then gives an equivalence of augmented commutative A -algebras

$$\mathcal{L}_X(A) \simeq \mathcal{L}_X(B)^{hG} \simeq \mathcal{L}_Y(B)^{hG} \simeq \mathcal{L}_Y(A).$$

□

There exist several definitions of smoothness in the literature (see for instance [16, 15]) using THH-étaleness and TAQ-étaleness. We suggest the following variant.

Definition 3.30. We call a map of cofibrant S -algebras $\varphi: R \rightarrow A$ *really smooth*, if it can be factored as $R \xrightarrow{i_R} \mathbb{P}_R(X) \xrightarrow{f} A$ where i_R is the canonical inclusion, X is an R -module and if $R \xrightarrow{i_R} \mathbb{P}_R(X) \xrightarrow{f} A$ satisfies étale descent.

Combining Proposition 3.27 and Corollary 2.10 we get:

Proposition 3.31. *If $R \rightarrow A$ is really smooth then $\Sigma X \simeq \Sigma Y$ for connected X and Y implies*

$$\mathcal{L}_X^R(A) \simeq \mathcal{L}_Y^R(A)$$

as commutative R -algebras.

The notion of being really smooth is transitive and closed under base change.

Lemma 3.32.

- If $\varphi: R \rightarrow A$ and $\psi: A \rightarrow B$ are really smooth, then so is $\psi \circ \varphi: R \rightarrow B$.
- If $\varphi: R \rightarrow A$ is really smooth and if C is a cofibrant commutative R -algebra, then $C \rightarrow C \wedge_A B$ is really smooth.

Proof. For transitivity we take the factorizations $\varphi = R \xrightarrow{i_R} \mathbb{P}_R(X) \xrightarrow{f} A$ and $\psi = A \xrightarrow{i_A} \mathbb{P}_A(Y) \xrightarrow{g} B$ and combine them to

$$R \xrightarrow{i_R} \mathbb{P}_R(X \vee Y) \simeq \mathbb{P}_R(X) \wedge_R \mathbb{P}_R(Y) \xrightarrow{f \wedge_R \text{id}} A \wedge_R \mathbb{P}_R(Y) \simeq \mathbb{P}_A(Y) \xrightarrow{g} B$$

So we have to show that for general maps $f: D \rightarrow A, g: A \rightarrow B, h: B \rightarrow C$ of commutative R -algebras:

- (1) If f satisfies étale descent, then so does $f \wedge_R \text{id}_C$ for every commutative R -algebra C .
- (2) If g and h satisfy étale descent, then so does $h \circ g$.

For (1) let X be connected. As $\mathcal{L}_X^R(-)$ commutes with pushouts (see Lemma 3.17), we get that $\mathcal{L}_X^R(A \wedge_R C) \simeq \mathcal{L}_X^R(A) \wedge_R \mathcal{L}_X^R(C)$. As f satisfies étale descent,

$$\mathcal{L}_X^R(A) \wedge_R \mathcal{L}_X^R(C) \simeq A \wedge_D \mathcal{L}_X^R(D) \wedge_R \mathcal{L}_X^R(C) \simeq A \wedge_D \mathcal{L}_X^R(D \wedge_R C)$$

and this in turn is equivalent to $(A \wedge_R C) \wedge_{(D \wedge_R C)} \mathcal{L}_X^R(D \wedge_R C)$.

The proof of (2) is straightforward because

$$\begin{aligned} C \wedge_A \mathcal{L}_X^R(A) &\simeq C \wedge_B (B \wedge_A \mathcal{L}_X^R(A)) \\ &\simeq C \wedge_B \mathcal{L}_X^R(B) \\ &\simeq \mathcal{L}_X^R(C). \end{aligned}$$

For the claim about base change consider the diagram

$$\begin{array}{ccc} R & \xrightarrow{i_R} & \mathbb{P}_R(X) \\ \eta \downarrow & & \downarrow f \\ C & \longrightarrow & C \wedge_R \mathbb{P}_R(X). \end{array}$$

Adjunction gives that $C \wedge_R \mathbb{P}_R(X) \simeq \mathbb{P}_C(C \wedge_R X)$. As $R \rightarrow \mathbb{P}_R(X) \rightarrow C$ satisfies étale descent we obtain with Lemma 3.26 that $C \rightarrow C \wedge_R \mathbb{P}_R(X) \rightarrow C \wedge_R A$ satisfies étale descent. □

4. THOM SPECTRA AND TOPOLOGICAL K-THEORY

Christian Schlichtkrull gives a closed formula for the Loday construction on Thom spectra [17, Theorem 1.1]: Let $f: W \rightarrow BF_{hI}$ be an E_∞ -map with W grouplike and let $T(f)$ denote the corresponding Thom spectrum. Then for any $T(f)$ -module spectrum M one has

$$(4.1) \quad \mathcal{L}_X(T(f); M) \simeq M \wedge \Omega^\infty(E_W \wedge X)_+$$

where E_W is the Omega spectrum associated to W (i.e., $W \simeq \Omega^\infty E_W$). If M is a commutative $T(f)$ -algebra spectrum, then the above equivalence is one of commutative $T(f)$ -algebras. For $M = T(f)$ the equivalence also respects the augmentation.

An immediate consequence of Schlichtkrull's result is the following:

Corollary 4.1. *If $T(f)$ is a Thom spectrum as above, then $S \rightarrow T(f)$ is multiplicatively stable.*

Proof. If $\Sigma X \simeq \Sigma Y$, then on the level of spectra we obtain

$$\Sigma(E_W \wedge X) \simeq E_W \wedge \Sigma X \simeq E_W \wedge \Sigma Y \simeq \Sigma(E_W \wedge Y),$$

but here suspension is invertible, hence $E_W \wedge X \simeq E_W \wedge Y$ and therefore

$$\mathcal{L}_X(T(f)) \simeq \mathcal{L}_Y(T(f)).$$

An equivalence of spectra induces an equivalence of infinite loop spaces and the $T(f)$ -algebra structure on $T(f) \wedge \Omega^\infty(E_W \wedge X)_+$ just comes from the one on $T(f)$ and the infinite loop structure on $\Omega^\infty(E_W \wedge X)$. This gives the multiplicativity of the stability. \square

The case of the suspension spectrum of an abelian topological group is a special case where we take $f: G \rightarrow BF_{hI}$ to be the trivial map. Then $T(f) \simeq \Sigma_+^\infty(G)$. Other examples are MU , MO , MSO , MSp or $MSpin$.

Remark 4.2. Note that by Corollary 3.15 spherical abelian monoid rings are *not* stable in general, whereas spherical abelian group rings are.

Bruno Stonek calculates higher THH of periodic complex topological K-theory, KU , and he determines topological André-Quillen homology of KU [19]. He uses Snaith's description of KU as the Bott localization of $\Sigma_+^\infty \mathbb{C}P^\infty$. The latter is a Thom spectrum because $\mathbb{C}P^\infty = BU(1)$ can be realized as a topological abelian group.

Theorem 4.3. *If X and Y are connected and $\Sigma X \simeq \Sigma Y$, then*

$$\mathcal{L}_X(KU) \simeq \mathcal{L}_Y(KU)$$

as commutative augmented KU -algebra spectra.

Proof. Let β denote the Bott element. Stonek uses Snaith's identification of KU as $\Sigma_+^\infty \mathbb{C}P^\infty[\beta^{-1}]$ to prove [19, Corollary 4.12] that there is a zigzag of equivalences

$$\begin{array}{ccc} \mathrm{THH}(KU) \simeq \mathrm{THH}(\Sigma_+^\infty \mathbb{C}P^\infty[\beta^{-1}]) & \xleftarrow{\simeq} & \mathrm{THH}(\Sigma_+^\infty \mathbb{C}P^\infty) \wedge_{\Sigma_+^\infty \mathbb{C}P^\infty} \Sigma_+^\infty \mathbb{C}P^\infty[\beta^{-1}] \\ & & \downarrow \simeq \\ & & (\mathrm{THH}(\Sigma_+^\infty \mathbb{C}P^\infty))[\beta^{-1}]. \end{array}$$

The same argument yields that for any connected X the localization of $\mathcal{L}_X(\Sigma_+^\infty \mathbb{C}P^\infty)$ at β is equivalent to $\mathcal{L}_X(\Sigma_+^\infty \mathbb{C}P^\infty[\beta^{-1}]) = \mathcal{L}_X(KU)$.

The localization map $\Sigma_+^\infty \mathbb{C}P^\infty \rightarrow \Sigma_+^\infty \mathbb{C}P^\infty[\beta^{-1}]$ satisfies étale descent, and therefore the composite $S \rightarrow \Sigma_+^\infty \mathbb{C}P^\infty \rightarrow \Sigma_+^\infty \mathbb{C}P^\infty[\beta^{-1}]$ identifies KU as an étale extension of a Thom spectrum. By Proposition 3.27 we obtain multiplicative stability for connected simplicial sets. \square

Corollary 4.4. *If X and Y are connected simplicial sets with $\Sigma X \simeq \Sigma Y$ then $\mathcal{L}_X(KO) \simeq \mathcal{L}_Y(KO)$ as commutative KO -algebras.*

Proof. Rognes shows [16, §5.3] that the complexification map $KO \rightarrow KU$ is a faithful C_2 -Galois extension of commutative ring spectra and Mathew [14, Example 4.6] deduces from [5, Example 5.9] that it satisfies étale descent. Schlichtkrull's equivalence from (4.1) is natural hence it preserves the C_2 -action. Therefore the result follows from Theorem 3.29. \square

5. THE GREENLEES SPECTRAL SEQUENCE

Let k be a field and let $A \rightarrow B$ be a morphism of connective commutative S -algebras with an augmentation to Hk satisfying some mild finiteness assumption. Then by [10, Lemma 3.1] there is a spectral sequence

$$E_{s,t}^2 = \pi_s(B \wedge_A Hk) \otimes_k \pi_t(A) \Rightarrow \pi_{s+t}(B).$$

Let p be an odd prime. We consider the cofibration $S^1 \vee S^1 \hookrightarrow S^1 \times S^1 \rightarrow S^2$ and the associated pushout diagram

$$\begin{array}{ccc} \mathcal{L}_{S^1 \vee S^1}^R(\mathbb{F}_p[t]/t^2; \mathbb{F}_p) & \longrightarrow & \mathcal{L}_{S^1 \times S^1}^R(\mathbb{F}_p[t]/t^2; \mathbb{F}_p) \\ \downarrow & & \downarrow \\ H\mathbb{F}_p & \longrightarrow & \mathcal{L}_{S^2}^R(\mathbb{F}_p[t]/t^2; \mathbb{F}_p). \end{array}$$

Here, R can be S or $H\mathbb{F}_p$. For $R = S$ we obtain a Greenlees spectral sequence

$$(5.1) \quad \pi_s(\mathcal{L}_{S^2}(\mathbb{F}_p[t]/t^2; \mathbb{F}_p)) \otimes_{\mathbb{F}_p} \pi_t(\mathcal{L}_{S^1 \vee S^1}(\mathbb{F}_p[t]/t^2; \mathbb{F}_p)) \Rightarrow \pi_{s+t}(\mathcal{L}_{S^1 \times S^1}(\mathbb{F}_p[t]/t^2; \mathbb{F}_p))$$

whereas for $R = H\mathbb{F}_p$ the spectral sequence is

$$(5.2) \quad \pi_s(\mathcal{L}_{S^2}^{\mathbb{F}_p}(\mathbb{F}_p[t]/t^2; \mathbb{F}_p)) \otimes_{\mathbb{F}_p} \pi_t(\mathcal{L}_{S^1 \vee S^1}^{\mathbb{F}_p}(\mathbb{F}_p[t]/t^2; \mathbb{F}_p)) \Rightarrow \pi_{s+t}(\mathcal{L}_{S^1 \times S^1}^{\mathbb{F}_p}(\mathbb{F}_p[t]/t^2; \mathbb{F}_p)).$$

In (5.1) every term $\mathcal{L}_X(\mathbb{F}_p[t]/t^2; \mathbb{F}_p)$ splits as

$$\mathcal{L}_X(H\mathbb{F}_p) \wedge_{H\mathbb{F}_p} \mathcal{L}_X^{\mathbb{F}_p}(\mathbb{F}_p[t]/t^2; \mathbb{F}_p)$$

naturally in X , and going from working over S to working over $H\mathbb{F}_p$ simply collapses the $\mathcal{L}_X(H\mathbb{F}_p)$ to $\mathcal{L}_X^{H\mathbb{F}_p}(H\mathbb{F}_p) \simeq H\mathbb{F}_p$. Therefore we get a surjection of the spectral sequence of (5.1) onto the one of (5.2), and if all the spectral sequence differentials vanish on the former, they have to vanish on the latter too. But we know that the rank of $\pi_2(\mathcal{L}_{S^1 \times S^1}^{\mathbb{F}_p}(\mathbb{F}_p[t]/t^2; \mathbb{F}_p))$ is less than the rank of the E^2 -term in total degree 2, hence there has to be a non-trivial differential in (5.2) and hence also in (5.1). This implies the following result.

Theorem 5.1. *For every odd prime p , $(S, H\mathbb{F}_p[t]/t^2, H\mathbb{F}_p)$ is not stable.*

Instead of stability we can consider the following property of Loday constructions.

Definition 5.2. Let R be a cofibrant commutative ring spectrum and let $R \rightarrow A \rightarrow C$ be a sequence of cofibrations of commutative R -algebras. We say that $R \rightarrow A \rightarrow C$ *decomposes products* if for all pointed simplicial sets X and Y

$$\mathcal{L}_{X \times Y}^R(A; C) \simeq \mathcal{L}_{X \vee Y \vee X \wedge Y}^R(A; C)$$

Note that the right hand side is equivalent to $\mathcal{L}_X^R(A; C) \wedge_C \mathcal{L}_Y^R(A; C) \wedge_C \mathcal{L}_{X \wedge Y}^R(A; C)$.

Proposition 5.3. *Let $R \rightarrow A \rightarrow B \rightarrow A \rightarrow Hk$ be a sequence of commutative S -algebras that turns B into an augmented commutative A -algebra. Assume that k is a field.*

If $R \rightarrow B \rightarrow Hk$ decomposes products then so does $R \rightarrow A \rightarrow Hk$.

Proof. The naturality of the Loday construction ensures that the vertical compositions in the diagram

$$\begin{array}{ccccc} \mathcal{L}_{X \vee Y}^R(A; Hk) & \longrightarrow & \mathcal{L}_{X \times Y}^R(A; Hk) & \longrightarrow & \mathcal{L}_{X \wedge Y}^R(A; Hk) \\ \downarrow & & \downarrow & & \downarrow \\ \mathcal{L}_{X \vee Y}^R(B; Hk) & \longrightarrow & \mathcal{L}_{X \times Y}^R(B; Hk) & \longrightarrow & \mathcal{L}_{X \wedge Y}^R(B; Hk) \\ \downarrow & & \downarrow & & \downarrow \\ \mathcal{L}_{X \vee Y}^R(A; Hk) & \longrightarrow & \mathcal{L}_{X \times Y}^R(A; Hk) & \longrightarrow & \mathcal{L}_{X \wedge Y}^R(A; Hk) \end{array}$$

are the identity. Therefore the spectral sequence

$$\pi_s(\mathcal{L}_{X \wedge Y}^R(A; Hk)) \otimes_k \pi_t(\mathcal{L}_{X \vee Y}^R(A; Hk)) \Rightarrow \pi_{s+t}(\mathcal{L}_{X \times Y}^R(A; Hk))$$

is a direct summand of the one for B . So if the spectral sequence for A had a non-trivial differential, then the one for B also had to have one, but as B decomposes products, this cannot happen. \square

Note that this gives an additive splitting, but we can't rule out multiplicative extensions.

If B does not decompose products, then this does *not* imply that A doesn't either. A concrete counterexample is $S \rightarrow H\mathbb{Q} \rightarrow H\mathbb{Q}[t]/t^2 \rightarrow H\mathbb{Q}$. Here, $S \rightarrow H\mathbb{Q}[t]/t^2$ does not decompose products, but $S \rightarrow H\mathbb{Q}$ is even multiplicatively stable.

6. TRANSITIVITY AND SPLITTINGS

Let k be a field. The case $k \rightarrow k[t] = R \rightarrow k[t]/t^m = R/t^m \rightarrow k = R/t$ for $m \geq 2$ is special in the sense that the quotient $k[t]/t^m$ is itself a commutative augmented k -algebra, so we can combine our splitting result for higher order Shukla homology [3, Proposition 7.5] with the juggling formula [3, Theorem 3.3]. We have [3, Theorem 7.6]:

$$(6.1) \quad \mathrm{THH}^{[n]}(k[t]/t^m; k) \simeq \mathrm{THH}^{[n]}(k[t]; k) \wedge_{Hk} \mathrm{THH}^{[n], k[t]}(k[t]/t^m; k)$$

for all $n \geq 1$ and for all $m \geq 2$ and in this special case we can get a Hk -version of this result.

Theorem 6.1. *Let k be a field and let m be greater or equal to 2. Then for all $n \geq 1$*

$$(6.2) \quad \mathrm{THH}^{[n], k}(k[t]/t^m; k) \simeq \mathrm{THH}^{[n], k}(k[t]; k) \wedge_{\mathrm{THH}^{[n-1], k[t]}(k)}^L \mathrm{THH}^{[n], k[t]}(k[t]/t^m; k).$$

This is a transitivity result for $k \rightarrow k[t] \rightarrow k[t]/t^m \rightarrow k$.

Proof. Consider the diagram

$$\begin{array}{ccccc} Hk & \longrightarrow & \mathrm{THH}^{[n-1], k[t]}(k) & \longrightarrow & \mathrm{THH}^{[n], k}(k[t]; k) \\ \downarrow & & \downarrow & & \downarrow \\ \mathrm{THH}^{[n], k[t]}(k[t]/t^m; k) & \longrightarrow & \mathrm{THH}^{[n-1], k[t]/t^m}(k) & \longrightarrow & \mathrm{THH}^{[n], k}(k[t]/t^m; k). \end{array}$$

□

The left-hand square is a homotopy pushout square by [3, Proposition 7.5] and the juggling formula [3, Theorem 3.3] applied to $Hk \rightarrow Hk[t] \rightarrow Hk[t]/t^m \rightarrow Hk$ ensures that the right-hand square is also a homotopy pushout square because for all $n \geq 1$

$$\mathrm{THH}^{[n], k}(k[t]/t^m; k) \simeq \mathrm{THH}^{[n], k}(k[t]; k) \wedge_{\mathrm{THH}^{[n-1], k[t]}(k)}^L \mathrm{THH}^{[n-1], k[t]/t^m}(k).$$

This yields that the outer rectangle is also a homotopy pushout square and this was the claim.

Remark 6.2. Note that there cannot be a version of (6.1) and (6.2) for arbitrary connected X : We know that $Hk[t] \rightarrow Hk[t]/t^2 \rightarrow Hk$ is multiplicatively stable for all fields k and we know that $Hk \rightarrow Hk[t] \rightarrow Hk$ is stable. But for any odd prime p we know that $H\mathbb{F}_p \rightarrow \mathbb{F}_p[t]/t^2 \rightarrow H\mathbb{F}_p$ is not stable and that there is an actual discrepancy between $\pi_2(\mathcal{L}_{S^1 \times S^1}^{H\mathbb{F}_p}(H\mathbb{F}_p[t]/t^2; H\mathbb{F}_p))$ and $\pi_2(\mathcal{L}_{S^1 \vee S^1 \vee S^2}^{H\mathbb{F}_p}(H\mathbb{F}_p[t]/t^2; H\mathbb{F}_p))$, so there cannot be an equivalence between $\mathcal{L}_{S^1 \times S^1}^{H\mathbb{F}_p}(H\mathbb{F}_p[t]/t^2; H\mathbb{F}_p)$ and $\mathcal{L}_{S^1 \vee S^1 \vee S^2}^{H\mathbb{F}_p}(H\mathbb{F}_p[t]/t^2; H\mathbb{F}_p)$.

REFERENCES

- [1] Christian Ausoni, John Rognes, *The chromatic red-shift in algebraic K-theory*, Enseign. Math. 54 (2), (2008), 9–11.
- [2] Yuri Berest, Ajay C. Ramadoss, Wai-kit Yeung, *Representation homology of spaces and higher Hochschild homology*, arXiv:1703.03505
- [3] Irina Bobkova, Eva Höning, Ayelet Lindenstrauss, Kate Poirier, Birgit Richter and Inna Zakharevich, *Splittings and calculational techniques for higher THH*, arXiv:1808.05440, to appear in *Algebr. Geom. Topol.*
- [4] Aldridge K. Bousfield, *The localization of spectra with respect to homology*, Topology 18 (1979), no. 4, 257–281.
- [5] Dustin Clausen, Akhil Mathew, Niko Naumann and Justin Noel, *Descent in algebraic K-theory and a conjecture of Ausoni-Rognes*, to appear in the Journal of the European Mathematical Society, arXiv:1606.03328.
- [6] Bjørn Ian Dundas, Thomas Goodwillie, Randy McCarthy, *The local structure of algebraic K-theory*, Algebra and Applications, 18. Springer-Verlag London, Ltd., London, 2013. xvi+435 pp.
- [7] Bjørn Ian Dundas, Ayelet Lindenstrauss, Birgit Richter, *On higher topological Hochschild homology of rings of integers*, Mathematical Research Letters 25, 2 (2018), 489–507.
- [8] Bjørn Ian Dundas, Andrea Tenti, *Higher Hochschild homology is not a stable invariant*, Mathematische Zeitschrift, Volume 290, Issue 1-2, 145–154.
- [9] Anthony D. Elmendorf, Igor Kriz, Michael A. Mandell, J. Peter May, *Rings, modules, and algebras in stable homotopy theory*. With an appendix by M. Cole. Mathematical Surveys and Monographs, 47. American Mathematical Society, Providence, RI, (1997), xii+249.
- [10] John P. C. Greenlees, *Ausoni-Bökstedt duality for topological Hochschild homology*, J. Pure Appl. Algebra 220 (2016), no. 4, 1382–1402.

- [11] Gemma Halliwell, Eva Höning, Ayelet Lindenstrauss, Birgit Richter, Inna Zakharevich, *Relative Loday constructions and applications to higher THH-calculations*, *Topology Appl.* 235 (2018), 523–545.
- [12] Lars Hesselholt, Ib Madsen, *On the K-theory of finite algebras over Witt vectors of perfect fields*, *Topology* 36 (1997), no. 1, 29–101.
- [13] Jacob Lurie, *Higher Algebra*, book draft available at <http://www.math.harvard.edu/~lurie/>
- [14] Akhil Mathew, *THH and base-change for Galois extensions of ring spectra*, *Algebr. Geom. Topol.* 17 (2017), no. 2, 693–704.
- [15] Randy McCarthy, Vahagn Minasian, *HKR theorem for smooth S-algebras*, *J. Pure Appl. Algebra* 185 (2003), no. 1-3, 239–258.
- [16] John Rognes, *Galois extensions of structured ring spectra. Stably dualizable groups*, *Mem. Amer. Math. Soc.* 192 (2008), no. 898, viii+137 pp.
- [17] Christian Schlichtkrull, *Higher topological Hochschild homology of Thom spectra*, *J. Topol.* 4 (2011), no. 1, 161–189.
- [18] Victor P. Snaith, *Algebraic cobordism and K-theory*, *Mem. Amer. Math. Soc.* 21 (1979), no. 221, vii+152 pp.
- [19] Bruno Stonek, *Higher topological Hochschild homology of periodic complex K-theory*, arXiv:1801.00156
- [20] Torleif Veen, *Detecting periodic elements in higher topological Hochschild homology*, *Geom. Topol.* 22 (2018), no. 2, 693–756.

MATHEMATICS DEPARTMENT, INDIANA UNIVERSITY, 831 EAST THIRD STREET, BLOOMINGTON, IN 47405, USA
E-mail address: `alindens@indiana.edu`

FACHBEREICH MATHEMATIK DER UNIVERSITÄT HAMBURG, BUNDESSTRASSE 55, 20146 HAMBURG, GERMANY
E-mail address: `birgit.richter@uni-hamburg.de`