

General Perturbations for Braneworld Compactifications and the Six Dimensional Case

S. L. Parameswaran^{a1}, S. Randjbar-Daemi^{b2} and A. Salvio^{c3}

^a *II Institute for Theoretical Physics,
University of Hamburg, DESY Theory Group, Notkestrasse 85, Bldg. 2a, D-22603
Hamburg, Germany*

^b *International Center for Theoretical Physics,
Strada Costiera 11, 34014 Trieste, Italy*

^c *Institut de Théorie des Phénomènes Physiques,
EPFL, CH-1015 Lausanne, Switzerland
and
IFAE, Universitat Autònoma de Barcelona,
08193 Bellaterra, Barcelona, Spain*


Abstract

Our main objective is to study how braneworld models of higher codimension differ from the 5D case and traditional Kaluza-Klein compactifications. We first derive the classical dynamics describing the physical fluctuations in a wide class of models incorporating gravity, non-Abelian gauge fields, the dilaton and two-form potential, as well as 3-brane sources. Next, we use these results to study braneworld compactifications in 6D supergravity, focusing on the bosonic fields in the minimal model; composed of the supergravity-tensor multiplet and the $U(1)$ gauge multiplet whose flux supports the compactification. For unwarped models sourced by positive tension branes, a harmonic analysis allows us to solve the large, coupled, differential system completely and obtain the full 4D spin-2,1 and 0 particle spectra, establishing (marginal) stability and a qualitative behaviour similar to the smooth sphere compactification. We also find interesting results for models with negative tension branes; extra massless Kaluza-Klein vector fields can appear in the spectra, beyond those expected from the isometries in the internal space. These fields imply an enhanced gauge symmetry in the low energy 4D effective theory obtained by truncating to the massless sector, which is explicitly broken as higher modes are excited, until the full 6D symmetries are restored far above the Kaluza-Klein scale. Remarkably, the low energy effective theory does not seem to distinguish between a compactification on a smooth sphere and these singular, deformed spheres.

¹Email: susha.louise.parameswaran@desy.de

²Email: seif@ict.p.trieste.it

³Email: salvio@ifae.es



Contents

1	Introduction	3
2	The Model	5
2.1	Field content	6
2.2	The action	6
2.3	The equations of motion (EOMs) and solutions	8
3	General Perturbations	11
3.1	Bilinear action	11
3.2	Local symmetries	13
4	Perturbations in the Light Cone Static Gauge	14
4.1	Gauge fixing	14
4.2	Bilinear action in the light cone static gauge	15
4.2.1	Spin-2 action	15
4.2.2	Spin-1 action	16
4.2.3	Spin-0 action and singularities due to backreacting, fluctuating branes	16
5	6D (and 5D) Braneworlds	18
5.1	Gravitational fluctuations	18
5.1.1	Randall-Sundrum	19
5.1.2	6D Brane Worlds	19
5.2	Vector fluctuations	21
5.2.1	Randall-Sundrum	21
5.2.2	6D Brane Worlds	21
5.3	Massless vectors and 4D gauge symmetries	28
5.3.1	Why there are three massless vector modes	28
5.3.2	The absence of enhanced gauge symmetries in the full 4D theory . .	29
5.3.3	The emergence of enhanced gauge symmetries at low energies	31
5.4	Massless scalars	32
5.4.1	Randall-Sundrum	32
5.4.2	6D Brane Worlds	32
6	Summary of Results	33
7	Conclusions	35
A	General ξ-Dependent Bilinear Action	37
B	Spin-0 Bilinear Action in the Light Cone Static Gauge	37
C	Spin-0 Spectrum for 6D Supergravity Compactification	39

1 Introduction

Almost two decades on, branes are evermore ubiquitous in the models constructed to understand particle physics and cosmology, with all their How?'s and Why?'s. As fundamental objects, they are the D-branes and NS-branes (or M-branes) of string (or M) theory, but within a low-energy effective field theory description, they are introduced as braneworlds. Often these braneworlds are considered as infinitely thin but finite tension objects, like for their more fundamental cousins, although sometimes it proves necessary to resolve their structure by adding some thickness.

A codimension one brane necessarily forms a boundary in the bulk space, since there is no path which can lead from one side to the other without traversing the brane. The gravitational backreaction of these objects is well understood; whilst the metric is continuous across the brane, its first derivative can have finite discontinuities. Branes with more than one transverse dimension are qualitatively different, and much harder, due to their sourcing of singularities in the transverse space. Still, codimension two branes can also be handled with some control; they backreact on the geometry in such a way as to produce relatively mild conical singularities.

The construction of solutions sourced by branes, with up to two codimensions, in various field theory models is by now a well-developed art. In 5D, the archetype is of course the construction of Randall and Sundrum [1, 2]. In 6D, we take the general warped braneworld compactifications (“conical-GGP solutions”) of 6D N=1 gauged supergravity [3] found in [4, 5] as representative. These solutions additionally invoke fluxes, which are also playing a dominant role in string compactifications today, and indeed models with two extra dimensions are the simplest in which flux compactifications can be studied. Having established the solutions, we can begin to ask about their physics: Are they stable to small perturbations? What are the symmetries and particle content of the low energy effective field theory? Is it chiral? What are the modifications to 4D Einsteinian gravity? What would be the effective vacuum energy measured by a 4D observer? What role do the branes play in these and other phenomena? And so on.

The first step towards answering these questions is to analyze the classical spectra of small fluctuations around the solution. A number of such studies have been made recently for the conical-GGP solutions. In [6] we worked out the spectra for certain 4D gauge fields and fermions present in the model and no tachyons or ghosts were found amongst them. A similar (marginal) stability was found in [7], where the axially symmetric modes for some of the scalar perturbations were calculated. The spectrum for the gravitino has also been analyzed in [8]. In [9], meanwhile, we studied the tachyonic instabilities that can arise from the non-axially symmetric, 4D scalar fluctuations descending from 6D gauge fields, and charged under the background fluxes⁴. Whether a given model with a given flux suffers from this instability turns out to depend on the tensions of the branes present.

We now intend to complete the spectral analysis for the bosonic fluctuations about the braneworld solutions of 6D supergravity. Our particular focus in this paper is on the so-called Salam-Sezgin sector – that arising from the supergravity-tensor multiplet and the

⁴The end point of this instability is studied in [10].

$U(1)$ gauge multiplet in which the background monopole lies – which was partially treated in [11, 7]. The remaining sectors have been completed elsewhere [6, 9]. We will calculate the corresponding spectra for the 4D spin-2 and – for unwarped backgrounds – spin-1 and spin-0 fields. The model that we are studying is complicated, and technically difficult. However, this goes hand in hand with its advantage of generality, and indeed the results for several simpler scenarios can be extracted from our work at its various stages.

Our approach will be that established in [12], where a formalism was developed to analyze the spectra of small perturbations about arbitrary solutions of Einstein, Yang-Mills and scalar systems. The first part of this paper can be considered as a generalization of that work, where we now include the presence of thin source 3-branes and extra bulk fields that are generically present in supergravity theories; the dilaton and anti-symmetric two-form potential. With little extra cost, we actually keep the number of dimensions transverse to the brane general.

We first derive the general form of the bilinear action that describes the behaviour of small fluctuations. For codimension-two or higher, we include fluctuations of the brane positions in the transverse directions, the so-called “branons”. We then apply the light-cone gauge (for bulk fields) and static gauge (for branons) to restrict to physical degrees of freedom, and decouple the dynamics for the spin-2, -1 and -0 fluctuations. The gauge-fixed bilinear action thus obtained provides the starting point to calculate the Kaluza-Klein (KK) spectra for the conical-GGP solutions, as well as, for example, the 5D Randall-Sundrum models and the non-supersymmetric Einstein-Yang Mills(-dilaton) model in any dimension.

In the second part of this paper, we use these general equations to study the behaviour of braneworld models in 6D (along the way also recover some of well-known aspects of the 5D scenarios). Here, since we include the backreaction of the branes, the dynamics of the branons are not well-defined⁵. Therefore, to study the spin-0 sector, we choose to truncate the branons by *e.g.* placing the branes at orbifold fixed points, or taking the brane tensions to be very large making the branes rigid within our range of validity. Meanwhile, the conical singularities in the curvature that are induced by the codimension two branes do not prevent us from understanding the behaviour of the bulk fluctuations.

We are able to derive the spectrum for the 4D spin-2 fields in the model’s full warped generality. The spin-1 and spin-0 sectors present large coupled differential systems, and by finding a set of harmonics on the 2D internal space (the “rugbyball”), we are also able to solve these systems analytically for the unwarped case. In this way, we obtain *all* the 4D modes for unwarped compactifications with positive tension brane sources, and qualitatively, we observe the same behaviour as in the smooth sphere compactification without branes – including marginal stability.

In the presence of negative tension codimension-two branes, meanwhile, the physics can surprise. Here, despite the fact that brane sources clearly break the $SU(2)$ isometries of the sphere to $U(1)$, three massless spin-1 fields⁶ can be found amongst the KK spectra for special values of the conical deficit angle. These special deficit angles, $\delta = -2\pi, -4\pi, \dots$,

⁵Indeed, the behaviour of the branons is usually considered under the probe brane approximation, in which the brane tension is much smaller than the bulk gravitational scale, so that the backreaction can be neglected [13, 14].

⁶In addition to any massless gauge fields arising from unbroken higher dimensional gauge symmetries.

allow three Killing vectors to be well-defined everywhere outside the branes, although only one of them can be globally integrated to an isometry.

Whether or not the massless vectors are gauge bosons of an enhanced gauge symmetry in the 4D theory can be understood by going beyond bilinear order and considering the interaction terms. We find the presence of KK modes that are not in well-defined representations of the $SU(2)$ generated by the Killing vectors, and therefore the full 4D theory does not enjoy an $SU(2)$ gauge symmetry. For this reason, we do not expect the classical masslessness of the vector fields to survive quantum corrections. Meanwhile, all our bosonic massless modes do fall into well-defined $SU(2)$ representations, and therefore we argue that the classical low energy 4D effective field theory – obtained by truncating to the massless sector – does enjoy an enhanced KK gauge symmetry beyond the isometries! Moreover, it appears that the low energy theory does not distinguish between compactifications on the smooth sphere and these singular, deformed spheres.

Let us now give an outline for the remainder of the paper. The first part presents a rather general analysis that determines the dynamics of perturbations in braneworld compactifications. In the next section, we introduce the model (both theories and background solutions) and discuss the scenarios to which our analysis can be applied. In Section 3, we introduce the perturbations about the background, obtain the bilinear action that describes their dynamics, and discuss the local symmetries of this action. In Section 4, we use these symmetries to fix to the “light cone static gauge”, and give the bilinear action in this gauge, in which the different spin sectors decouple.

Then begins the second part, which uses the previous results to study the 4D fields that emerge in various scenarios. In Section 5, our main interest is in the braneworld solutions of 6D supergravity, but we also discuss a non-supersymmetric 6D model and the 5D Randall-Sundrum models. In the main text we present the KK spectra for spin-2 and spin-1 fields and identify the massless spin-0 fields; the complete spin-0 sector can be found in the appendices. Finally, we understand in detail the physical significance of the extra massless 4D vector modes that can appear in the spectra, and the gauge invariance that emerges in the 4D theory.

We summarise our results in Section 6, before concluding in Section 7.

2 The Model

We begin with the definition of our model. The main focus of the present paper will be a class of bosonic 6D field theories with thin codimension-two branes. In particular we are interested in the bosonic part of 6D $N=1$ gauged supergravity [3]. However, throughout the article we shall keep a general space-time dimension D as far as possible, and certain truncations of the field content allow our analysis to be applied to several different scenarios, including the non-supersymmetric Einstein-Yang-Mills theory or the Randall-Sundrum Model.

2.1 Field content

The basic ingredients of our model are the higher dimensional metric G_{MN} , where the space-time indices run over $M, N, \dots = 0, \dots, D - 1$, and the gauge field \mathcal{A}_M of a compact Lie group \mathcal{G} . These are bulk fields in the sense that they depend on all the space-time coordinates X^M .

We also want to consider a certain number \mathcal{N} of 3-branes embedded in the D -dimensional space time. To do so we introduce, following Ref. [15], \mathcal{N} functions $Y_k^M(x_k)$, $k = 1, \dots, \mathcal{N}$, which represent the positions of the branes in the D -dimensional space time. The x_k represent the 4D coordinates on the brane, $x_k = \{x_k^\alpha\}$, where α, β, \dots are the 4D indices. Not all the space-time components of $Y_k^M(x_k)$ are physical degrees of freedom: 4 space-time components for each k can be gauged away by using the 4D (general) coordinate transformation invariance acting on x_k [15], as we will explicitly do in Subsection 4.1. We consider $Y_k^M(x_k)$ to be a brane field because it depends only on a 4D world-volume coordinate. These fields are important to introduce the branes in a covariant way, and indeed we can construct the induced metrics on the branes by means of

$$g_{\alpha\beta}^k = G_{MN}(Y_k(x_k))\partial_\alpha Y_k^M(x_k)\partial_\beta Y_k^M(x_k). \quad (2.1)$$

In order to complete the bosonic part of the 6D supergravity, one should add other bulk fields in addition to G_{MN} and \mathcal{A}_M , that is a dilaton ϕ and a 2-form field B_{MN} , which emerge from the graviton multiplet and an antisymmetric tensor multiplet [3]. We will refer to B_{MN} as the Kalb-Ramond field. Moreover, concerning the 6D supergravity, we shall assume that \mathcal{G} is a product of simple groups that include a $U(1)_R$ gauged R-symmetry. In general one can also add some hypermultiplets [3], which turn out to be important to cancel gauge and gravitational anomalies [16, 17]. In the bosonic sector this leads to additional scalar fields Φ^α (hyperscalars) in some representation of \mathcal{G} ; however, from now on we set $\Phi^\alpha = 0$. We do so because we are interested in the linear perturbations which mix with the D -dimensional gravitational fluctuations h_{MN} : indeed, for the class of backgrounds we are interested in (see Subsection 2.3), the Φ^α decouple from h_{MN} . Their inclusion should be straightforward.

Therefore the bulk and the brane field contents that we consider are respectively:

$$\{G_{MN}, \mathcal{A}_M, \phi, B_{MN}\} \quad \text{and} \quad \{Y_k^M(x_k), \dots\}. \quad (2.2)$$

The dots in the second set of (2.2) represent additional brane fields that we can always introduce, but which are not required by general covariance; for example they can be the fields of the Standard Model (SM).

2.2 The action

We split the action functional S into the bulk action S_B , which depends only on the bulk fields, and the brane action S_b that is a functional of the brane fields as well.

The bulk action is⁷[3]

$$S_B = \int d^D X \sqrt{-G} \left\{ \frac{1}{\kappa^2} \left[R - \frac{1}{4} (\partial\phi)^2 \right] - \frac{1}{4} e^{\phi/2} F^2 - \frac{\kappa^2}{48} e^\phi H_{MNP} H^{MNP} - \mathcal{V}(\phi) \right\}, \quad (2.3)$$

where G is the determinant of G_{MN} and κ is the D -dimensional Planck scale; also⁸ $F^2 \equiv F_{MN} F^{MN}$ and $(\partial\phi)^2 \equiv \partial_M \phi \partial^M \phi$. The explicit expression for the gauge field strength F_{MN} is⁹

$$F_{MN} = \partial_M \mathcal{A}_N - \partial_N \mathcal{A}_M + g \mathcal{A}_M \times \mathcal{A}_N, \quad (2.4)$$

where g is the gauge coupling, which in fact represents a collection of independent gauge couplings including that of the $U(1)_R$ subgroup, g_1 . H_{MNP} is the Kalb-Ramond field strength, which contains a Chern-Simons coupling as follows [18]:

$$H_{MNP} = \partial_M B_{NP} + F_{MN} \mathcal{A}_P - \frac{g}{3} \mathcal{A}_M (\mathcal{A}_N \times \mathcal{A}_P) + 2 \text{ cyclic perms.} \quad (2.5)$$

The function $\mathcal{V}(\phi)$ is the dilaton potential. In the supersymmetric model this is fixed to be $\mathcal{V}(\phi) = 8g_1^2 e^{-\phi/2} / \kappa^4$.

Meanwhile, we consider the following 3-brane action

$$S_b = \sum_k \left(-T_k \int d^4 x_k \sqrt{-g^k} \right) \equiv -T \int d^4 x \sqrt{-g}, \quad (2.6)$$

where g^k is the determinant of (2.1) and T_k are the tensions of the branes. From now on (unless otherwise stated) we suppress the index k , as we have done on the right hand side of (2.6). The reader may have noticed that we have not introduced the Gibbons-Hawking boundary term, which is generically necessary to treat codimension one branes [19]. Indeed, we shall apply our analysis only to those codimension one models whose branes are placed on orbifold fixed points, in which case the Gibbons-Hawking boundary term is not present [20].

We can summarise by saying that our analysis will apply to the following two types of models:

1. 6D N=1 gauged supergravity.
2. Einstein-Yang-Mills theories, with a dilaton or cosmological constant Λ , for a general space-time dimension.

The second case includes, for example, the RS models [1, 2] or the non-supersymmetric 6D Einstein-Yang-Mills- Λ (EYMA) model [13, 21]. They can be obtained by simply fixing the appropriate dimension and setting $H_{MNP} = 0$, $\phi = 0$ and $\mathcal{V}(0) = \Lambda$. Even if our main

⁷We choose signature $(-, +, \dots, +)$, and define $R_{MN}{}^R{}_S = \partial_M \Gamma_{NS}^R - \partial_N \Gamma_{MS}^R + \Gamma_{MP}^R \Gamma_{NS}^P - \Gamma_{NP}^R \Gamma_{MS}^P$ and $R_{MN} = R_{PM}{}^P{}_N$.

⁸A trace overall is understood when we write a product of Lie algebra valued objects: e.g. in Eq. (2.3) $F^2 \equiv \text{Tr}(F^2)$.

⁹We define the cross-product as $(\mathcal{A}_M \times \mathcal{A}_N)^I = f^{IJK} \mathcal{A}_M^J \mathcal{A}_N^K$, with f^{IJK} the structure constants of \mathcal{G} : $[T^I, T^J] = if^{IJK} T^K$, where T^I are the generators of \mathcal{G} .

interest is in models of Type 1 we will also consider the second class for several reasons. In this way, we will see that our results can be applied in quite general contexts, and it will also provide interesting additional ways to check our formulae. Moreover, in the future it should help us to figure out the role of supersymmetry in the linear perturbations.

Finally, it is important to note that the actions S_B and S_b are invariant with respect to both the D -dimensional and the 4D coordinate transformations (acting respectively on X^M and x^α). We will discuss the local symmetries of the present model and an explicit gauge fixing for the linear perturbations in Subsections 3.2 and 4.1.

2.3 The equations of motion (EOMs) and solutions

The EOMs that follow from the variation of the action $S_B + S_b$ are:

$$R^{MN} - \frac{1}{2}G^{MN}R = \frac{\kappa^2}{2} \left\{ e^{\phi/2} \left(F^M_P F^{NP} - \frac{1}{4}G^{MN}F^2 \right) + \frac{1}{2\kappa^2} \partial^M \phi \partial^N \phi - G^{MN} \left[\frac{1}{4\kappa^2} (\partial\phi)^2 + \mathcal{V}(\phi) \right] \right\} - T\kappa^2 \mathcal{B}^{MN}, \quad (2.7)$$

$$D_N \left(e^{\phi/2} F^{NM} \right) = 0, \quad (2.8)$$

$$\frac{1}{2\kappa^2} D^2 \phi = \frac{\partial \mathcal{V}}{\partial \phi}(\phi) + \frac{1}{8} e^{\phi/2} F^2, \quad (2.9)$$

$$\frac{1}{\sqrt{-g}} \partial_\alpha (\sqrt{-g} G_{MN} \partial^\alpha Y^N) = \frac{1}{2} \partial_M G_{NP} \partial Y^N \cdot \partial Y^P, \quad (2.10)$$

where we have fixed $H_{MNP} = 0$, since our interest shall be in backgrounds that enjoy 4D Poincaré invariance. Moreover, in Eq. (2.9) and (2.10) we have introduced the notation $D^2 \phi \equiv D_M D^M \phi$, where D_M is the covariant derivative, and $\partial Y^M \cdot \partial Y^N \equiv \partial_\alpha Y^M \partial^\alpha Y^N$. Recall also that we have suppressed the index k on Y_k^M , which labels each of the branes. The last term in (2.7) represents the brane contribution to the Einstein equations, where \mathcal{B}^{MN} is defined by

$$\mathcal{B}^{MN}(X) \equiv \frac{1}{2} \int d^4x \sqrt{g/G} \delta(X - Y(x)) \partial Y^M \cdot \partial Y^N; \quad (2.11)$$

we note that the bulk quantity G in (2.11) is computed at the position of the brane ($G = G(Y)$) because of the presence of the D -dimensional delta function $\delta(X - Y(x))$. Furthermore, since Eqs. (2.10) come from the variation of the brane action with respect to Y^M , there the bulk fields G_{MN} and $\partial_M G_{NP}$ are computed at the brane position ($G_{MN} = G_{MN}(Y)$ and $\partial_M G_{NP} = \partial_M G_{NP}(Y)$).

In the present paper we will focus mainly on the following ansatz solution to (2.7)-(2.10):

$$Y^\mu = x^\mu, \quad (2.12)$$

$$Y^{\underline{m}} = \text{constant}, \quad (2.13)$$

$$ds^2 = e^{A(\rho)} \eta_{\mu\nu} dx^\mu dx^\nu + d\rho^2 + e^{B(\rho)} K_{mn}(y) dy^m dy^n, \quad (2.14)$$

$$\mathcal{A} = \mathcal{A}_m(\rho, y) dy^m, \quad (2.15)$$

$$\phi = \phi(\rho), \quad (2.16)$$

$$H_{MNP} = 0, \quad (2.17)$$

where $\mu = 0, 1, 2, 3$, $m = 5, \dots, 4 + D_2$, $\underline{m} = (\rho, m)$ (we have $D = 5 + D_2$) and y^m and K_{mn} are respectively the coordinate and the metric on the D_2 -dimensional space. Eq. (2.12) is not really an assumption because we can always use the 4D general coordinate invariance on the branes to set (2.12). Eq. (2.13) is instead a non trivial assumption. Moreover, in Eqs (2.14)-(2.17) we are assuming that the bulk field background has a 4D Poincaré invariance and that the functions A , B and ϕ depend only on the coordinate ρ . We will also assume \mathcal{A} to lie in the Cartan subalgebra of $\text{Lie}(\mathcal{G})$.

One of the simplest models that can be described by this set up is the Randall-Sundrum (RS) model [1], where we have $D = 5$, $\phi = 0$ and $\mathcal{A} = 0$ and the internal space is S^1/Z_2 with two branes on the fixed points of Z_2 , say at $\rho = 0$ and $\rho = \pi r_c$. The explicit form of the solution is given by

$$A = -2k|\rho|, \quad Y_1^\rho = 0, \quad Y_2^\rho = \pi r_c, \quad (2.18)$$

where k is a positive constant. The object $|\rho|$ in (2.18) is equal to the absolute value of ρ in the region $-\pi r_c < \rho < \pi r_c$ and its value anywhere else is obtained by periodicity. In order for (2.18) to be a solution one needs $T_1 = -T_2 = 12k/\kappa^2$ and $\Lambda = -12k^2/\kappa^2$. In Section 5, we shall use this very well-known solution to check the result given in Section 4.

However, in this paper our main interest lies in the analysis of a class of solutions found by Gibbons, Güven and Pope (GGP) [4] to the 6D supergravity: the general set of warped solutions with 4D Poincaré symmetry, and axial symmetry in the transverse dimensions. Here we give only a subset of this general class, namely that which contains singularities no worse than conical and therefore can be sourced by brane terms of the form (2.6).

To give the explicit expression of the conical-GGP solutions, it turns out to be useful to introduce the following radial coordinate [6]

$$u(\rho) \equiv \int_0^\rho d\rho' e^{-A(\rho')/2}, \quad (2.19)$$

whose range is $0 \leq u \leq \bar{u} \equiv \pi r_0/2$. In this frame the metric reads

$$ds^2 = e^{A(u)} (\eta_{\mu\nu} dx^\mu dx^\nu + du^2) + e^{B(u)} \frac{r_0^2}{4} d\varphi^2. \quad (2.20)$$

The explicit conical-GGP solutions¹⁰ are then the following particular case of the ansatz (2.12)-(2.17) [4]:

$$\begin{aligned} e^A &= e^{\phi/2} = \sqrt{\frac{f_1}{f_0}}, & e^B &= 4\alpha^2 e^A \frac{\cot^2(u/r_0)}{f_1^2}, \\ \mathcal{A} &= -\frac{4\alpha}{q\kappa f_1} Q d\varphi, \end{aligned} \quad (2.21)$$

¹⁰The coordinate u is related to the coordinate r in [4] by $r = r_0 \cot(u/r_0)$.

where q and α are generic real numbers and Q is a generator of a $U(1)$ subgroup of a simple factor of \mathcal{G} , satisfying $\text{Tr}(Q^2) = 1$. Also,

$$f_0 \equiv 1 + \cot^2\left(\frac{u}{r_0}\right), \quad f_1 \equiv 1 + \frac{r_0^2}{r_1^2} \cot^2\left(\frac{u}{r_0}\right), \quad (2.22)$$

with $r_0^2 \equiv \kappa^2/(2g_1^2)$ and $r_1^2 \equiv 8/q^2$.

This solution is supported by two branes located at $u = 0$ and $u = \bar{u}$. Indeed, as $u \rightarrow 0$ or $u \rightarrow \bar{u}$, the metric tends to that of a cone, with respective deficit angles

$$\delta = 2\pi \left(1 - |\alpha| \frac{r_1^2}{r_0^2}\right) \quad \text{and} \quad \bar{\delta} = 2\pi (1 - |\alpha|), \quad (2.23)$$

and corresponding delta-function behaviours in the Ricci scalar. We will take $\alpha \geq 0$ without loss of generality. The tensions of the two branes T and \bar{T} are related to the deficit angle as follows [22]:

$$T = 2\delta/\kappa^2 \quad \text{and} \quad \bar{T} = 2\bar{\delta}/\kappa^2. \quad (2.24)$$

Unlike the RS solution, here the warp factor e^A is smooth on the brane positions $u = 0$ and $u = \bar{u}$. In particular we have

$$e^A \xrightarrow{u \rightarrow 0, \bar{u}} \text{constant} \neq 0, \quad \partial_u e^A \xrightarrow{u \rightarrow 0, \bar{u}} 0. \quad (2.25)$$

By using (2.25), (2.12) and (2.13), it is also easy to check that the conical-GGP configuration satisfies the Y -equations (2.10) in addition to the bulk EOMs (2.7)-(2.9).

The expression for the gauge field background in Eq. (2.21) is well-defined in the limit $u \rightarrow 0$, but not as $u \rightarrow \bar{u}$. We should therefore use a different patch to describe the $u = \bar{u}$ brane, and this must be related to the patch including the $u = 0$ brane by a single-valued gauge transformation. This leads to a Dirac quantization condition, which for a field interacting with \mathcal{A} through a charge e gives

$$-e \frac{4\alpha\bar{g}}{\kappa q} = -e \alpha \frac{r_1}{r_0} \frac{\bar{g}}{g_1} = N, \quad (2.26)$$

where N is an integer that is called monopole number and \bar{g} is the gauge coupling constant corresponding to the background gauge field. For example, if \mathcal{A} lies in $U(1)_R$, then $\bar{g} = g_1$. The charge e can be computed once we have selected the background gauge group, since it is an eigenvalue of the generator Q . Also, note that the internal space corresponding to Solutions (2.21) has an S^2 topology (its Euler number equals 2).

Finally, we observe that one can obtain the unwarped ‘‘rugbyball’’ compactification [21] simply by setting $r_0 = r_1$. In this case the metric is

$$ds^2 = \eta_{\mu\nu} dx^\mu dx^\nu + \frac{r_0^2}{4} (d\theta^2 + \alpha^2 \sin^2 \theta d\varphi^2), \quad (2.27)$$

where $\theta \equiv 2u/r_0$, and the background value of the dilaton is zero; therefore this is a solution also to the non-supersymmetric 6D EYMA model. For $\alpha < 1$ the deficit angle is positive. The geometry is also well-defined when $\alpha > 1$ and the deficit angle is negative; we name these spaces ‘‘saddle-spheres’’ (see [9] for a detailed discussion on their properties). Moreover, we can smoothly retrieve the sphere compactification (with radius $r_0/2$) by taking $\alpha = 1$ in addition to $r_0 = r_1$.

3 General Perturbations

The main purpose of this paper is to study the linear perturbations in the above models. We therefore perturb the fields in (2.2) as follows:

$$\begin{aligned} G_{MN} &\rightarrow G_{MN} + h_{MN}, & \mathcal{A}_M &\rightarrow \mathcal{A}_M + V_M, & \phi &\rightarrow \phi + \tau, \\ B_{MN} &\rightarrow B_{MN} + b_{MN}, & Y^M &\rightarrow Y^M + \xi^M. \end{aligned} \quad (3.28)$$

The first terms in the right hand sides of (3.28) represent the background quantities of the corresponding fields. In fact, it is useful to introduce another 2-form field V_{MN} in order to describe the fluctuations of the Kalb-Ramond field. This can be done as follows. Since H_{MNP} appears only quadratically in (2.3), and $H_{MNP} = 0$ at the background level due to 4D Poincaré invariance, the linear approximation (which corresponds to the bilinear level in the action) involves only the linear perturbation of H_{MNP} , that we denote with¹¹ $H_{MNP}^{(1)}$,

$$H_{MNP}^{(1)} = [d(b_2 - \mathcal{A} \wedge V) + 2F \wedge V]_{MNP}, \quad (3.29)$$

where we have used the notation of p-forms and b_2 is the fluctuation in the Kalb-Ramond 2-form, \mathcal{A} and F the background values of the gauge field and its field strength respectively and V the perturbation of the gauge field. We now introduce the 2-form V_2 as follows:

$$V_2 \equiv \kappa(b_2 - \mathcal{A} \wedge V), \quad (3.30)$$

whose components will be denoted by V_{MN} . $H_{MNP}^{(1)}$ can now be expressed in terms of V_2 and V :

$$H_{MNP}^{(1)} = \left(\frac{1}{\kappa} dV_2 + 2\gamma F \wedge V \right)_{MNP}, \quad (3.31)$$

where we have introduced a new parameter γ ; for $\gamma = 1$ we recover the structure of $H_{MNP}^{(1)}$ required by the 6D supergravity, whereas for $\gamma = 0$ the fluctuations of V_{MN} are completely decoupled (at the linear level) from the rest. This will allow us to treat simultaneously the 6D supergravity and the EYMA models.

Finally, we note that the fields $\xi^M(x)$ describe the fluctuations of the brane positions, and as such they are 4D fields.

3.1 Bilinear action

Here we provide the linearized theory which corresponds to the bilinear approximation in the action. The bilinear action has been computed by considering the variation of $S_B + S_b$ under (3.28) and by keeping only terms up to the quadratic order¹². We split it into different contributions as follows:

$$\begin{aligned} &S(h, h) + S(V, V) + S(h, V) + S(\tau, \tau) + S(h, \tau) + S(V, \tau) \\ &+ S(V_2, V_2) + S(V, V_2) + S(\xi, \xi) + S(h, \xi), \end{aligned} \quad (3.32)$$

¹¹Since the background $H_{MNP} = 0$, and the background monopole, \mathcal{A} , lies in the Cartan subalgebra, we see that the exterior derivative acting on the background Kalb-Ramond potential B_2 must be zero. Also, $\mathcal{A} \wedge \mathcal{A} = 0$.

¹²The EOMs (2.7)-(2.10) guarantee that the linear terms vanish.

where $S(h, h)$ is the bilinear action that depends only on the fluctuations h_{MN} , $S(h, V)$ represents the mixing term between h_{MN} and V_M and so on. We have $S(h, V_2) = S(\tau, V_2) = 0$ as a consequence of our background ansatz, for which $H_{MNP} = 0$. We give here the explicit expressions for the bilinear action that depend only on the bulk fields; the dynamics of the ξ^M fields, are explicitly given in Appendix A. We find:

$$\begin{aligned}
S(h, h) = \int d^D X \sqrt{-G} \left\{ \frac{1}{2\kappa^2} \left[\left(h^{MN}{}_{;M} - \frac{1}{2} h^{;N} \right)^2 - \frac{1}{2} h^{NP}{}_{;M} h_{NP}{}^{;M} + \frac{1}{4} h^{;M} h_{;M} - \frac{1}{2} R_1 h^2 \right] \right. \\
- \frac{1}{2} h_{PM} h^P{}_N \left(\frac{1}{2} e^{\phi/2} F^{MR} F^N{}_R + \frac{1}{4\kappa^2} \partial^M \phi \partial^N \phi \right) \\
- \frac{1}{2} h^{MN} h^{PR} \left(\frac{1}{\kappa^2} R_{PMNR} - \frac{1}{2} e^{\phi/2} F_{PM} F_{NR} \right) \\
\left. - \frac{T}{2} \left[\mathcal{B}^{MN} (h_{PM} h^P{}_N - h h_{MN}) + \frac{1}{2} \mathcal{B}^{MNP R} h_{MN} h_{PR} \right] \right\}, \quad (3.33)
\end{aligned}$$

where the semicolon denotes the (background) gravitational covariant derivative, $h \equiv G^{MN} h_{MN}$, $R_{MN}{}^P{}_R$ is the Riemann tensor for the background metric and we have defined

$$\frac{2}{\kappa^2} R_1 \equiv \frac{1}{\kappa^2} R - \frac{1}{4} e^{\phi/2} F^2 - \frac{1}{4\kappa^2} (\partial\phi)^2 - \mathcal{V}(\phi) \quad (3.34)$$

and

$$\begin{aligned}
\mathcal{B}^{MNP R} \equiv \int d^4 x \sqrt{g/G} \delta(X - Y(x)) \left[\frac{1}{2} (\partial Y^M \cdot \partial Y^N) \partial Y^P \cdot \partial Y^R \right. \\
\left. - (\partial Y^M \cdot \partial Y^P) \partial Y^N \cdot \partial Y^R \right]. \quad (3.35)
\end{aligned}$$

The term proportional to $\mathcal{B}^{MNP R}$ in the last line of (3.33) is the contribution to $S(h, h)$ coming from the brane action S_b , whereas the term proportional to \mathcal{B}^{MN} comes from the EOMs (2.7), which we have used to write $S(h, h)$ in the form (3.33). Moreover,

$$\begin{aligned}
S(V, V) = \int d^D X \sqrt{-G} \left[-\frac{1}{2} e^{\phi/2} (D_M V_N D^M V^N - D_M V_N D^N V^M) \right. \\
\left. - \frac{\kappa^2}{12} \gamma^2 e^\phi (F_{[MN} V_{P]}) (F^{[MN} V^{P]}) - \frac{1}{2} \bar{g} e^{\phi/2} F^{MN} V_M \times V_N \right], \quad (3.36)
\end{aligned}$$

$$S(h, V) = - \int d^D X \sqrt{-G} e^{\phi/2} (D^M V^N - D^N V^M) \left(\frac{1}{4} h F_{MN} + h_{PN} F^P{}_M \right), \quad (3.37)$$

$$S(\tau, \tau) = - \int d^D X \sqrt{-G} \left[\frac{1}{4\kappa^2} (\partial\tau)^2 + \frac{1}{2} \frac{\partial^2 \mathcal{V}}{\partial \phi^2} \tau^2 + \frac{1}{32} e^{\phi/2} F^2 \tau^2 \right], \quad (3.38)$$

$$\begin{aligned}
S(h, \tau) = \int d^D X \sqrt{-G} \left[\frac{1}{2\kappa^2} \partial_M \tau \partial_N \phi \left(h^{MN} - \frac{1}{2} G^{MN} h \right) - \frac{1}{2} \frac{\partial \mathcal{V}}{\partial \phi} h \tau \right. \\
\left. + \frac{1}{4} e^{\phi/2} \left(F^{MP} F^N{}_P - \frac{1}{4} F^2 G^{MN} \right) \tau h_{MN} \right], \quad (3.39)
\end{aligned}$$

$$S(V, \tau) = \int d^D X \sqrt{-G} \left[-\frac{1}{4} e^{\phi/2} F_{MN} (D^M V^N - D^N V^M) \tau \right], \quad (3.40)$$

$$S(V_2, V_2) = -\frac{1}{48} \int d^D X \sqrt{-G} e^\phi V_{[NP;M]} V^{[NP;M]}, \quad (3.41)$$

$$S(V, V_2) = -\frac{\kappa}{12} \gamma \int d^D X \sqrt{-G} e^\phi V_{[NP;M]} F^{[MN} V^P], \quad (3.42)$$

where

$$F_{[MN} V_P] \equiv F_{MN} V_P + 2 \text{ cyclic perms}, \quad V_{[NP;M]} \equiv V_{NP;M} + 2 \text{ cyclic perms}.$$

We would like to remind the reader of the assumptions we have made to derive (3.33) and (3.36)-(3.42) (and (A.110)-(A.111) given in Appendix A):

- If the Kalb-Ramond field and the term $H_{MNP} H^{MNP}$ in (2.3) is not included, then the only assumption we made is that the background satisfies the EOMs (2.7)-(2.10).
- If the Kalb-Ramond field and the term $H_{MNP} H^{MNP}$ in (2.3) is instead included, we also assumed $D = 6$ and the background gauge field \mathcal{A} to lie in the Cartan subalgebra.

We observe that if we want to focus on the D -dimensional EYMA system we can restrict ourselves to the terms $S(h, h)$, $S(V, V)$ (for $\gamma = 0$), $S(h, V)$ and the ξ -dependent terms given in Appendix A. Instead, if we want to consider the 6D supergravity, we should put $\gamma = 1$, $\mathcal{V}(\phi) = 8 g_1^2 e^{-\phi/2} / \kappa^4$ and also take into account the terms (3.38)-(3.42). Finally, we note that our results reduce to those of Ref. [12] which studies a general non-supersymmetric class of thick brane models, once we take $T = 0$, $\gamma = 0$ and we neglect the fluctuations V_{MN} .

3.2 Local symmetries

As a consequence of the local symmetries of the complete model, the linearized theory also possesses a number of local symmetries:

$$\delta h_{MN} = -\eta_{N;M} - \eta_{M;N}, \quad (3.43)$$

$$\delta V_M = -\eta^L F_{LM} - D_M \chi, \quad (3.44)$$

$$\delta \tau = -\eta^M \partial_M \phi, \quad (3.45)$$

$$\delta V_{MN} = 2\gamma \kappa \chi F_{MN} + \lambda_{N;M} - \lambda_{M;N}, \quad (3.46)$$

$$\delta \xi^M = \eta^M(Y) - \zeta^\alpha \partial_\alpha Y^M. \quad (3.47)$$

Eqs. (3.43), (3.44) and (3.45) represent the effect of the local symmetries (descending from the D -dimensional coordinate transformation invariance and gauge symmetry) on the metric, the gauge field and the dilaton fluctuations (see *e.g.* Ref. [12]). The bulk functions η and χ are the gauge functions associated with the D -dimensional coordinate invariance and gauge symmetry.

Eq. (3.46) represents instead a local symmetry acting on V_{MN} , which descends from both the gauge symmetry and the Kalb-Ramond symmetry¹³. For this reason χ and λ_M are independent (bulk) gauge functions. Let us explicitly check (3.46). To do so, it is enough to verify the invariance of the 3-form (3.31) under (3.44) and (3.46). We have

$$\delta H^{(1)} = \frac{1}{\kappa} d(\delta V_2) + 2\gamma F \wedge \delta V = 2\gamma d(\chi F) + 2\gamma F \wedge (-\eta \cdot F - D\chi), \quad (3.48)$$

where we have used $d^2\lambda = 0$ and $\eta \cdot F$ represents the 1-form with components $\eta^M F_{MN}$. Now, by using the 4D Poincaré invariance of the background and $D = 6$, which we always assume in the presence of the Kalb-Ramond field, we have $F \wedge (\eta \cdot F) = 0$ and $F \wedge \mathcal{A} = 0$; also, by remembering that \mathcal{A} is assumed to lie in the Cartan subalgebra, we have $dF = 0$. These equations are sufficient to conclude $\delta H^{(1)} = 0$.

Finally, Eq. (3.47) represents the local transformation of the perturbation of the brane position, descending from the D -dimensional coordinate invariance and the 4D brane coordinate transformation invariance (respectively the first and the second term on the right hand side of (3.47)); the latter invariance is associated to ζ^α (a function of x^α), which represents another independent gauge function.

4 Perturbations in the Light Cone Static Gauge

Having derived the general bilinear action, we now have to choose a gauge in order to study the physical spectrum. In this section we will discuss our gauge choice and give the corresponding bilinear action.

4.1 Gauge fixing

We have two types of local symmetries: the bulk local symmetries (which include the D -dimensional coordinate transformation invariance, the gauge symmetry and the Kalb-Ramond symmetry) and the 4D coordinate transformation invariance on the brane. Let us start with the first group.

A very convenient gauge choice for the bulk local symmetry is the light cone gauge, as it ensures that the dynamics of sectors with different spin decouple at the bilinear level¹⁴. Another advantage of the light cone gauge is that it does not involve gauge artifacts such as Faddeev-Popov ghosts, but contains only the physical spectrum [23, 24, 25]. To define this gauge, let us introduce $x^{(\pm)} \equiv (x^3 \pm x^0) / \sqrt{2}$ and $A^{(\pm)} \equiv (A^3 \pm A^0) / \sqrt{2}$, for a general vector A^M . Then the light cone gauge is defined by

$$V_{(-)} = 0, \quad h_{(-)M} = 0, \quad V_{(-)M} = 0, \quad \forall M. \quad (4.49)$$

It can be proved that, after imposing (4.49), the (+) components of the different fields (*i.e.* $V_{(+)}$, $h_{(+)M}$ and $V_{(+)M}$) are not independent, but can be expressed in terms of the

¹³By Kalb-Ramond symmetry we mean the local invariance under $B_2 \rightarrow B_2 + d\Lambda$ of the action, where Λ is a general 1-form.

¹⁴This has been observed in other studies, for example [24, 25, 26, 12].

other components by means of constraint equations [24, 25, 12]. We therefore end up with the following independent bulk fields: h_{ij} , $h_{i\bar{m}}$, V_i , $V_{i\bar{m}}$, $h_{\bar{m}\bar{n}}$, $V_{\bar{m}}$, V_{ij} , $V_{\bar{m}\bar{n}}$ and τ , where $i, j, \dots = 1, 2$. In particular the $h_{(++)}$ field equation simply leads to the constraint

$$h = 0, \quad (4.50)$$

which brings a considerable amount of simplification.

Concerning the 4D coordinate transformation invariance, we instead impose the condition [15]

$$\xi^\mu = 0. \quad (4.51)$$

We will refer to (4.51) as to the static gauge. We observe that the light cone gauge and the static gauge are compatible because, once we fix the light cone gauge by choosing η_M , χ and λ_M in a suitable way, we still have the freedom to perform the local transformations generated by ζ^α . The static gauge is also free from Faddeev-Popov unphysical ghosts [15]. We observe that (4.51) does not remove completely the brane position fields ξ^M , but we are left with their components along the extra dimensions $\xi^{\bar{m}}$. We will refer to them as branons. Even if the branons represent physical degrees of freedom, it can happen that they can be consistently truncated *e.g.* by imposing an orbifold symmetry, as in the RS models or in the conical-GGP compactification [9]. In the following we will confirm that the spin-0 fields $\xi^{\bar{m}}$ do not have any mixing with the spin-2 and spin-1 sectors in the light cone gauge.

4.2 Bilinear action in the light cone static gauge

Here we provide the bilinear action in the light cone static gauge, that we have computed by imposing the gauge conditions (4.49) and (4.51) on the general bilinear action and by using the constraint equations for the (+) components. In this section we assume the form given in (2.12)-(2.17) for the background solution, and give the part of the action that is independent of the branons. Those involving the branons are given in Appendix B.

The results that are presented here reduce to those for the non-supersymmetric model present in¹⁵ [12] once we take $T = 0$, $\gamma = 0$ and we neglect the fluctuations V_{MN} ; they also correctly reduce (for $T = 0$ and $\gamma = 1$) to the results of [27], where the linear perturbations of the sphere-monopole solution to the 6D supersymmetric model are analyzed.

4.2.1 Spin-2 action

The spin-2 action $S^{(2)}$ only contains the field $\tilde{h}_{ij} \equiv h_{ij} - \frac{1}{2}G_{ij}h_k^k$ and has the following simple expression in terms of $\tilde{h}_i^j = G^{jk}\tilde{h}_{ik}$:

$$S^{(2)}(h, h) = -\frac{1}{4\kappa^2} \int d^D X \sqrt{-G} \partial_M \tilde{h}_i^j \partial^M \tilde{h}_j^i. \quad (4.52)$$

We observe that (4.52) has exactly the same form as in [12] even if we have included the brane terms. Therefore, the brane sources do not explicitly contribute to the spin-2 dynamics. We shall use (4.52) to derive the 4D gravitational spectrum for the solutions described in Subsection 2.3.

¹⁵We do, however, correct some typos in that reference.

4.2.2 Spin-1 action

The spin-1 action $S^{(1)}$ involves $h_{\underline{im}}$, V_i and $V_{\underline{im}}$. We have the following explicit expressions.

$$\begin{aligned}
S^{(1)}(h, h) = & \int d^D X \sqrt{-G} \left[-\frac{1}{2\kappa^2} (\partial_\mu h_{\underline{im}} \partial^\mu h^{\underline{im}} + \partial_\rho h_{\underline{im}} \partial_\rho h^{\underline{im}} + h_{\underline{im};n} h^{\underline{im};n}) \right. \\
& - \frac{1}{4\kappa^2} h_{im} h^{im} \left(A'^2 + \frac{B'^2}{2} \right) - \frac{1}{4\kappa^2} h_{\rho i} h_\rho{}^i (D_2 A' B' - A'^2) \\
& - \frac{1}{2} h_{\underline{im}} h^i{}_{\underline{n}} \left(\frac{1}{2} e^{\phi/2} F_{\underline{l}}^m F^{\underline{n}l} + \frac{1}{4\kappa^2} \partial^{\underline{m}} \phi \partial^{\underline{n}} \phi \right) \\
& \left. + \frac{1}{\kappa^2} A' h_\rho{}^i h_{mi}{}^{;m} - \frac{T}{4} \sqrt{g/G} \delta(X_c - Y_c) h_{\underline{m}i} h^{\underline{m}i} \right], \tag{4.53}
\end{aligned}$$

where $' \equiv \partial_\rho$. The last term in (4.53) is the brane contribution. We have introduced the notation X_c and Y_c for the internal components of the coordinate and the brane position respectively, where the label c stands for the codimension of the brane. The other non vanishing terms are the following.

$$\begin{aligned}
S^{(1)}(V, V) = & \int d^D X \sqrt{-G} e^{\phi/2} \left[-\frac{1}{2} (\partial_\mu V_i \partial^\mu V^i + e^{-A} \partial_\rho V_i \partial_\rho V_i + D_m V_i D^m V^i) \right. \\
& \left. - \frac{\kappa^2}{4} \gamma^2 e^{\phi/2} (F_{\underline{mn}} V_i) F^{\underline{mn}} V^i \right], \tag{4.54}
\end{aligned}$$

$$S^{(1)}(h, V) = \int d^D X \sqrt{-G} e^{\phi/2} \left(-D_{\underline{m}} V_i h_{\underline{l}}{}^i F^{\underline{l}m} - \frac{1}{2} A' V_i h^{\underline{l}i} F_{\underline{l}\rho} \right), \tag{4.55}$$

$$\begin{aligned}
S^{(1)}(V_2, V_2) = & -\frac{1}{8} \int d^D X \sqrt{-G} e^\phi \left\{ e^{-A} [\partial_\mu V_{\underline{im}} \partial^\mu V_i{}^{\underline{m}} \right. \\
& + G^{\underline{ml}} G^{\underline{nh}} (\partial_{\underline{m}} V_{\underline{ni}} \partial_{\underline{l}} V_{hi} - \partial_{\underline{m}} V_{\underline{ni}} \partial_{\underline{h}} V_{li})] \\
& - e^{-4A-2\phi} \left(e^{\phi+3A/2} V_{\underline{m}i}{}^{\underline{m}} \right)_{;\underline{m}} \left(e^{\phi+3A/2} V_{\underline{n}i}{}^{\underline{n}} \right)_{;\underline{n}} \\
& \left. - 2e^{-2A} V_{\underline{mi}} \partial^{\underline{m}} \left[e^{-\phi-A/2} \left(e^{\phi+3A/2} V_{\underline{n}i}{}^{\underline{n}} \right)_{;\underline{n}} \right] \right\}, \tag{4.56}
\end{aligned}$$

$$S^{(1)}(V, V_2) = -\frac{\kappa}{2} \gamma \int d^D X \sqrt{-G} e^\phi \left(-\frac{1}{2} A' V_{im} V^i F_\rho{}^m + V_{\underline{ni};\underline{m}} F^{\underline{mn}} V^i \right). \tag{4.57}$$

The term $S^{(1)}(h, V_2)$ vanishes as a consequence of $H_{MNP} = 0$ (at the background level). We observe that the term $S^{(1)}(V, V)$ reduces, as it should, to the corresponding action in Ref. [6] in the case in which V_i is orthogonal to the background gauge field. Finally, we note that the brane tension enters explicitly only in the term $S^{(1)}(h, h)$.

4.2.3 Spin-0 action and singularities due to backreacting, fluctuating branes

The last and most complicated part is the spin-0 action, which involves¹⁶ $h_{\underline{mn}}$, $h_i{}^i$, $V_{\underline{m}}$, τ , V_{ij} , $V_{\underline{mn}}$ and $\xi^{\underline{m}}$. We observe that, in the light cone static gauge, the fields $\xi^{\underline{m}}$ indeed

¹⁶Note that $h_{\underline{mn}}$ and $h_i{}^i$ are not independent as Eq. (4.50) implies $h_i{}^i + h_{\underline{m}}{}^{\underline{m}} = 0$.

only appear here. In other words they are in general completely decoupled from the spin-2 and spin-1 fields. Since it is quite complicated, we give the explicit expression of the spin-0 action in Appendix B.

Having completed the bilinear action, we should make some observations regarding its consistency, in particular given the presence of infinitesimally thin dynamical sources. Indeed, as to be expected, if we include the gravitational backreaction of the branes ($T \rightarrow 0$) then there are singular contributions to the dynamics of both the bulk gravitational fluctuations and the branons.

First, concerning the bulk gravitational fluctuations, we encounter localized contributions to the mass terms in both the spin-1 (see Eq. (4.53)) and spin-0 (see Eq. (B.112)) sectors. These contributions involve the behaviour of background and perturbed fields at the background positions of the branes. They are well-defined in the codimension one RS scenario, where the metric is well-defined everywhere including at the brane positions (although its derivatives are not). They do not appear to be well-defined in the codimension two (or higher) case, where the internal metric is actually singular at the brane positions due to the backreaction of the branes. However, as we shall see in subsequent sections, these terms do not obstruct our derivation of the 4D particle spectra arising from bulk modes in 6D.

Meanwhile, the linearized dynamics for the branons of a backreacting brane would be more problematic. For example, in (B.124), since the action is evaluated at the background position of the branes, the kinetic term for the branons is not well-defined in the codimension two case, because of the conical defect in G_{mn} . Such a singularity was discussed in [13], where it was argued that within the domain of validity of the effective field theory, the curvature singularity could be discarded. Moreover, the mass term for the branons takes the form¹⁷ of a $\delta(0)$. These singularities are not present in the RS model, as there the branons are projected out with an orbifolding¹⁸. Indeed we should reiterate here that we apply our analysis to codimension one branes only on orbifold fixed points (to avoid the appearance of Gibbons-Hawking boundary terms), and so without branon degrees of freedom.

In order to perform a complete analysis of the spin-0 action in codimension two (or higher) models, taking into account both the backreaction of the brane and its dynamical fluctuations, it seems necessary to resolve the thin structure of the brane. Otherwise we can assume a brane tension much smaller than the 6D fundamental scale, so that its backreaction is negligible. Or we can assume a high brane tension so that the brane is very heavy and rigid and does not oscillate. Or else we can assume an additional orbifold symmetry under which the branons are projected out – an example of such a symmetry has been provided in Ref. [9] and is discussed in¹⁹ Appendix C. In these cases, we can avoid the singular branon

¹⁷ $\delta(0)$ singularities due to the localization of fields on a boundary have been discussed in a different context (5D SYM theory on S^1/Z_2) in [28].

¹⁸Indeed, in the RS literature, the radion has been studied in depth [29], but the branons are absent. Although the radion can also be seen as a brane bending in the case of RS, since the branes are at the boundaries of the internal space, one should not confuse the radion with the branon. The radion is a deformation of the bulk metric, whereas the branon is a deformation of the brane itself within the bulk manifold. As a check of our formalism, we will find the radion mode in Subsection 5.4.

¹⁹By using the explicit expression for the mixing terms between branons and bulk fields given in Appendix

action.

5 6D (and 5D) Braneworlds

In the second part of this paper, we apply the results of the previous sections to derive the 4D particle spectra in specific setups. Our main interest is in the warped (and unwarped) axi-symmetric braneworld compactifications of 6D supergravity, but along the way we shall also discuss the rugbyball compactifications in the non-supersymmetric 6D EYMA theory, as well as the 5D Randall-Sundrum models. We discuss in order the spin-2, spin-1 and spin-0 fluctuations.

5.1 Gravitational fluctuations

The simplest application of our results is the analysis of the spin-2 sector. As we have discussed, this sector completely decouples from the rest. The \tilde{h}_i^j fields contain only the maximal helicity components of a spin-2 multiplet; one should look for the remaining components in the spin-1 and spin-0 actions. However, by virtue of 4D Poincaré invariance, the lower helicity components must have the same spectrum [12]. We can therefore focus on Eq. (4.52) to study the spin-2 fluctuations.

In order to analyze this sector we deduce the EOMs from Eq. (4.52):

$$\partial_M \left(\sqrt{-G} \partial^M \tilde{h}_i^j \right) = 0 \quad \forall i, j. \quad (5.58)$$

In deriving this equation we have required as usual that the boundary terms which emerge in the integration by parts vanish, that is

$$\int d^D X \partial_M \left(\sqrt{-G} \delta \tilde{h}_i^j \partial^M \tilde{h}_i^j \right) = 0, \quad (5.59)$$

where $\delta \tilde{h}_i^j$ is the variation of the field \tilde{h}_i^j , which is performed to apply the minimal action principle. Since we assume standard boundary conditions on the 4D boundary, (5.59) reduces to [30, 6]

$$\int d^{D_2+1} X \partial_{\underline{m}} \left(\sqrt{-G} \delta \tilde{h}_i^j \partial^{\underline{m}} \tilde{h}_i^j \right) = 0. \quad (5.60)$$

We now perform a KK decomposition of the fields as follows:

$$\tilde{h}_i^j(X) = \sum_{\mathbf{k}} \tilde{h}_i^{(\mathbf{k})j}(x) f_{\mathbf{k}}(\rho, y), \quad (5.61)$$

where \mathbf{k} represents a collective KK number. By taking $\tilde{h}_i^{(\mathbf{k})j}(x)$ to be an eigenfunction of $\eta^{\mu\nu} \partial_\mu \partial_\nu$, that is $\eta^{\mu\nu} \partial_\mu \partial_\nu \tilde{h}_i^{(\mathbf{k})j}(x) = M_{\mathbf{k}}^2 \tilde{h}_i^{(\mathbf{k})j}(x)$, the EOMs (5.58) become

$$-\frac{1}{\sqrt{-G}} e^A \partial_{\underline{m}} \left(\sqrt{-G} \partial^{\underline{m}} f_{\mathbf{k}} \right) = M_{\mathbf{k}}^2 f_{\mathbf{k}} \quad (5.62)$$

B, it is easy to confirm that symmetry consistently truncates the branons.

and the corresponding boundary conditions (5.60) read (we recall that $\delta\tilde{h}_i^j$ and \tilde{h}_i^j are independent fields)

$$\int d^{D_2+1} X \partial_{\underline{m}} \left(\sqrt{-G} f_{\mathbf{k}'} \partial^{\underline{m}} f_{\mathbf{k}} \right) = 0. \quad \forall \mathbf{k}, \mathbf{k}'. \quad (5.63)$$

Condition (5.63) ensures that the operator acting on $f_{\mathbf{k}}$ in the left hand side of (5.62) is a Hermitian operator [30, 6]; we will therefore refer to (5.63) as the hermiticity condition (HC). In addition to the HC we will also require the wave functions $f_{\mathbf{k}}$ to be normalizable, that is

$$\int d^{D_2+1} X \sqrt{-G} e^{-A} f_{\mathbf{k}}^2 < \infty.$$

This normalizability condition (NC) is equivalent to the finiteness of the kinetic energy of the modes $\tilde{h}_i^{(\mathbf{k})j}(x)$. We observe that there is always a constant massless ($M_{\mathbf{k}}^2 = 0$) solution to (5.62), satisfying the HC (5.63). This solution corresponds to a 4D graviton provided that the NC is satisfied, that is $\int d^{D_2+1} X \sqrt{-G} e^{-A} < \infty$.

5.1.1 Randall-Sundrum

In the special case $D = 5$, and therefore in particular for the RS background (2.18), the EOM (5.62) has the form

$$-e^{-A} \partial_{\rho} (e^{2A} \partial_{\rho} f_{\mathbf{k}}) = M_{\mathbf{k}}^2 f_{\mathbf{k}}. \quad (5.64)$$

Here we do not want to analyze the latter equation as this has been done in the original RS works, but we observe, as a check of our spin-2 action, that (5.64) has exactly the same form as in [2].

5.1.2 6D Brane Worlds

We now move to the conical-GGP solutions to 6D supergravity given in Eqs. (2.21)-(2.22). Since our internal space is topologically S^2 , we require \tilde{h}_i^j to be periodic functions of φ :

$$\tilde{h}_i^j(X) = \sum_{\mathbf{n}, \mathbf{m}} \tilde{h}_{i\mathbf{n}\mathbf{m}}^j(x) f_{\mathbf{n}\mathbf{m}}(\rho) e^{i\mathbf{m}\varphi}, \quad (5.65)$$

where \mathbf{m} is a generic integer and \mathbf{n} is an extra KK number that emerges as we have a number of compact dimensions greater than one. Also we observe that Eq. (5.62) with the HC and NC is formally identical²⁰ to the corresponding problem for 4D gauge fields addressed in Ref. [6]. Therefore, here we only give the result. The wave functions can be expressed in a more compact way by introducing

$$\psi \equiv e^{(3A+B)/4} f, \quad (5.66)$$

²⁰In [6] there is the extra parameter N_V , which is equal to zero here. To check that the two problems are identical it is useful to remember $A = \phi/2$, which is true for the conical-GGP solutions. Also, take care that ϕ in reference [6] is half ϕ here.




Figure 1: Graviton Wave Function Profiles: $\mathbf{n} = 0, 1, 20$ modes plotted for angular momentum numbers $\mathbf{m} = -1, 0$. The parameters are chosen to be $(r_0, \omega, \bar{\omega}) = (1, 1/4, 1)$, corresponding to a single negative tension brane at $u = 0$. Also the normalization constant is set such that $\int du |\psi|^2 = 1$. The number of intersections with the u -axis equals n , according to quantum mechanics. Notice that the $(\mathbf{m}, \mathbf{n}) = (0, 0)$ mode is massless.

where we have suppressed \mathbf{n} and \mathbf{m} . The explicit expression for ψ is

$$\psi \propto z^\epsilon (1-z)^\beta F(a, b, c, z), \quad (5.67)$$

where $z \equiv \cos^2(u/r_0)$, F is Gauss's hypergeometric function and

$$\begin{aligned} \epsilon &\equiv \frac{1}{4} (1 + 2|\mathbf{m}|\bar{\omega}), \quad \beta \equiv \frac{1}{4} (1 + 2\mathbf{m}\omega), \quad c \equiv 1 + |\mathbf{m}|\bar{\omega}, \\ a &\equiv \frac{1}{2} + \frac{\mathbf{m}}{2}\omega + \frac{|\mathbf{m}|}{2}\bar{\omega} + \frac{1}{2}\sqrt{r_0^2 M^2 + 1 + \mathbf{m}^2 (\omega - \bar{\omega})^2}, \\ b &\equiv \frac{1}{2} + \frac{\mathbf{m}}{2}\omega + \frac{|\mathbf{m}|}{2}\bar{\omega} - \frac{1}{2}\sqrt{r_0^2 M^2 + 1 + \mathbf{m}^2 (\omega - \bar{\omega})^2}, \end{aligned} \quad (5.68)$$

with

$$\omega \equiv (1 - \delta/2\pi)^{-1}, \quad \bar{\omega} \equiv (1 - \bar{\delta}/2\pi)^{-1}. \quad (5.69)$$

Moreover the explicit form of the mass spectrum is given by

$$M^2 = \frac{4}{r_0^2} \left[\mathbf{n}(\mathbf{n} + 1) + \left(\frac{1}{2} + \mathbf{n} \right) |\mathbf{m}| (\omega + \bar{\omega}) + \mathbf{m}^2 \omega \bar{\omega} \right] \geq 0, \quad (5.70)$$

where $\mathbf{n} = 0, 1, 2, 3, \dots$ [6]. So we have obtained the exact and complete spectrum (wave functions and masses) for the spin-2 fluctuations of the conical-GGP solutions. We observe that Eq. (5.70) tells us there is a massless normalizable solution (for $\mathbf{n} = \mathbf{m} = 0$), which corresponds to the 4D graviton. This solution is separated from the first KK excitation by a finite mass gap, which is of order $1/r_0$ (if $\omega \sim \bar{\omega} \sim 1$). We plot some representative wave function profiles in Figure 1. As discussed in [6] the asymptotic behaviour close to the branes is universal for each KK tower, and it does not appear possible to separate the infinite number of heavy modes from the light ones by using their respective wave function profiles.

Here we also observe that Eq. (5.58) is independent of γ and the dilaton potential \mathcal{V} . This implies that the spin-2 spectrum of the non supersymmetric and supersymmetric

models are the same (provided the backgrounds are the same). Indeed, the rugbyball configuration (that is $\omega = \bar{\omega}$) leads to the same spin-2 spectrum in the EYMA model and in the 6D supergravity.

Finally, as a check, we can consider the S^2 limit ($\omega, \bar{\omega} \rightarrow 1$), whose mass spectrum is well-known. Our spectrum (5.70) reduces to

$$\frac{r_0^2}{4} M^2 = l(l+1), \quad \text{multiplicity} = 2l+1, \quad (5.71)$$

where $l = 0, 1, 2, 3, \dots$. Since $r_0/2$ represents the radius of S^2 in the sphere limit, this is exactly the result that one finds by using the spherical harmonic expansion [31, 32] from the beginning.

5.2 Vector fluctuations

Here we analyze the vector fluctuations, in particular their wavefunction expansions and mass spectra. In the following subsection we shall study the implications of these results for the structure of the 4D gauge group in the 6D models of interest. The physical 4D vector field spectrum can be extracted from the spin-1 action given in Subsection 4.2.2. However, some of the perturbations in that action are simply the helicity- (± 1) components of massive gravitons and therefore should not be interpreted as independent vector fields.

5.2.1 Randall-Sundrum

To illustrate the previous point we first notice that our spin-1 action leads to the well-known result that there are no physical 4D vector fields in the RS model (unless one introduces bulk gauge fields). Indeed, in that case the only field appearing in the spin-1 action is $h_{i\rho}$, whose action is simply

$$S^{(1)}(h, h) \propto \int d^5 X \sqrt{-G} \left(\partial_\mu h_{i\rho} \partial^\mu h^i{}_\rho + \partial_\rho h_{i\rho} \partial_\rho h^i{}_\rho - \frac{A'^2}{2} h_{i\rho} h^i{}_\rho - \frac{3}{2} A'' h_{i\rho} h^i{}_\rho \right), \quad (5.72)$$

where we used the property

$$A'' + T\kappa^2 \delta(\rho - Y)/3 = 0 \quad (5.73)$$

that follows from the form of the warp factor, Eq. (2.18), in the RS model. Therefore, once (5.73) is used the problem assumes the same form as in Ref. [12], where it is shown that the 4D spectrum from $h_{i\rho}$ exactly reproduces the graviton one with the zero mode removed. By counting the degrees of freedom, it follows that there are no physical vector fields.

5.2.2 6D Brane Worlds

Let us begin by considering what we might expect from the symmetries of the problem – with some benefit of hindsight from the authors. In the limit where the brane tensions go to zero, the smooth sphere-monopole compactification is recovered. In this case, standard

KK theory tells us that there are three massless KK gauge bosons²¹, which manifest the $SU(2)$ isometries of the sphere in the 4D theory [31]. Clearly, any branes break the spherical symmetry in the internal dimensions. For the solutions of present interest an axial isometry survives, and therefore we can expect the 4D theory to enjoy a $U(1)$ KK gauge symmetry²².

At the same time, for the case of an unwarped “saddle-sphere” with the special deficit angles (for $\alpha = 1$ we recover the sphere)

$$\delta = -2\pi, -4\pi, \dots \quad \text{or} \quad \alpha = \frac{1}{\omega} = 2, 3, \dots \quad (5.74)$$

the metric (2.27) – defined everywhere but at the branes – has three single-valued Killing vectors, which obey the Lie algebra of $SU(2)$:

$$K^+ = e^{i\alpha\varphi} \left(\frac{\partial}{\partial\theta} + i \cot\theta \frac{1}{\alpha} \frac{\partial}{\partial\varphi} \right), \quad K^- = e^{-i\alpha\varphi} \left(-\frac{\partial}{\partial\theta} + i \cot\theta \frac{1}{\alpha} \frac{\partial}{\partial\varphi} \right), \quad K^0 = -\frac{i}{\alpha} \frac{\partial}{\partial\varphi}. \quad (5.75)$$

Only one of these Killing vectors, K^0 , implies a genuine continuous isometry, since K^\pm cannot be globally integrated to an isometry²³. In other words, we have an *infinitesimal* $SU(2)$ isometry for the special saddle-spheres, compared to a genuine $SU(2)$ isometry for the sphere. As we will show, this turns out to be sufficient to ensure three massless 4D vectors amongst the KK spectra. From the point of view of the full 4D theory, however, we will argue that these massless fields arise accidentally and that their masslessness is not protected by any symmetry.

Rugbyball Harmonics Let us now see how the above story plays out in detail. Our focus shall then be on the unwarped rugbyballs and saddle-spheres, Eq. (2.27), and indeed all previous results have indicated that warping does not lead to any qualitative changes in the physics (see the spin-2 results in the present paper, as well as Refs.[6]-[11]). We shall thus proceed by finding a set of “rugbyball harmonics” and their mass spectra, in analogy to the spherical harmonics (and, more generally, the so-called Wigner functions) used in the smooth sphere compactification [31].

We first observe that the vector field fluctuations V_i , which are orthogonal to the gauge field background ($V_i F_{mn} = 0$), do not mix with the other perturbations h_{im} and V_{im} . These fluctuations have been already studied in Ref. [6] where the complete KK towers are provided and it is shown that there are as many 4D gauge fields as fluctuations V_i with vanishing monopole charge ($V_i \times F_{mn} = 0$), as expected from group theory. Here we therefore consider only the case when V_i is parallel to the background monopole.

²¹There may be additional massless gauge fields descending from any unbroken higher dimensional gauge symmetries.

²²For the analysis of a very similar model, in which the $SU(2)$ KK gauge symmetry of the sphere is broken down to $U(1)$ by *smooth* axisymmetric deformations, see Ref. [33].

²³To avoid the need of differential geometric results for singular spaces, we can consider removing the brane singularities and taking instead a smooth non-compact manifold, for which $0 < \theta < \pi$. Killing vector fields are the generators of the *infinitesimal isometries* of a manifold, whereas an isometry is a global aspect of the geometry. Whilst for smooth compact manifolds the Killing vectors are always globally integrable to an isometry, for non-compact manifolds this may not always be the case.

Now, from Subsection 4.2.2 it follows that the spin-1 action for unwarped solutions has the following form.

$$\begin{aligned}
S^{(1)}(h, h) &= -\frac{1}{2\kappa^2} \int d^6 X \sqrt{-G} \left(\partial_\mu h_{mi} \partial^\mu h^{mi} + h_{mi;n} h^{mi;n} + \frac{R}{2} h_{mi} h^{mi} \right), \\
S^{(1)}(V, V) &= -\frac{1}{2} \int d^6 X \sqrt{-G} \left[\partial_\mu V_i \partial^\mu V^i + \partial_m V_i \partial^m V^i + \frac{\kappa^2}{2} \gamma^2 F^2 V_i V^i \right], \\
S^{(1)}(V_2, V_2) &= -\frac{1}{8} \int d^6 X \sqrt{-G} \left(\partial_\mu V_{mi} \partial^\mu V^{mi} + V_{mi;n} V^{mi;n} + \frac{R}{2} V_{mi} V^{mi} \right), \\
S^{(1)}(h, V) + S^{(1)}(V, V_2) &= \int d^6 X \sqrt{-G} \left(-\partial_m V_i h_{ni} F^{nm} - \frac{\kappa}{2} \gamma \partial_m V^i V_{ni} F^{nm} \right), \quad (5.76)
\end{aligned}$$

where we have used $V_{i;m} = \partial_m V_i$ because the background solution is unwarped and V_i is uncharged under the background monopole. Also m, n, \dots here run over θ and φ . To derive the last term in $S^{(1)}(h, h)$ we have used the Einstein equations:

$$\frac{2}{\kappa^2} \sqrt{-G} R_{mn} = \sqrt{-G} F_{ml} F_n{}^l + G_{mn} T \delta(X_2 - Y_2), \quad (5.77)$$

which allow us to rewrite the brane contribution in the last term of (4.53) as a combination of the Ricci tensor and the field strength. Also we have used that in two dimensions $R_{mn} = G_{mn} R/2$ and the Maxwell equations $F^{mn}{}_{;m} = 0$. The EOMs descending from (5.76) are the following.

$$\left(\partial^2 + D^2 - \frac{R}{2} \right) h_{mi} - \kappa^2 F_m{}^l \partial_l V_i = 0, \quad (5.78)$$

$$\left(\partial^2 + D^2 - \frac{R}{2} \right) V_{mi} - 2\kappa\gamma F_m{}^l \partial_l V_i = 0, \quad (5.79)$$

$$\left(\partial^2 + D^2 - \frac{\kappa^2}{2} \gamma^2 F^2 \right) V_i + F^{nm} h_{ni;m} + \frac{\kappa}{2} \gamma F^{nm} V_{ni;m} = 0, \quad (5.80)$$

where $\partial^2 \equiv \partial_\mu \partial^\mu$ and $D^2 \equiv D_m D^m$. As for the spin-2 case above, the EOMs come with a set of boundary conditions, which we refer to as Hermiticity Conditions (HCs) [30, 6]:

$$\int d^6 X \sqrt{-G} (\delta h_{mi} h^{mi;n})_{;n} = 0, \quad \int d^6 X \sqrt{-G} (\delta V_{mi} V^{mi;n})_{;n} = 0, \quad (5.81)$$

$$\int d^6 X \sqrt{-G} (\delta V^i h_{li} F^{lm})_{;m} = 0, \quad \int d^6 X \sqrt{-G} (\delta V^i V_{li} F^{lm})_{;m} = 0, \quad (5.82)$$

$$\int d^6 X \sqrt{-G} (\delta V_i \partial^m V^i)_{;m} = 0. \quad (5.83)$$

We will additionally impose the usual Normalizability Conditions (NCs).

We can immediately observe that there is a simple solution to Eqs. (5.78)-(5.80), with $h_{\theta i} = h_{\varphi i} = 0$, $V_{\theta i} = V_{\varphi i} = 0$ and V_i independent of the extra dimensions. Its squared mass is given by

$$M^2 = \frac{\kappa^2}{2} \gamma^2 F^2 = \frac{8}{r_0^2} \gamma^2, \quad (5.84)$$

where we explicitly used the rugbyball solution. We see that the monopole $U(1)$ is a gauge symmetry in the EYMA model ($\gamma = 0$), whereas it is broken in 6D supergravity ($\gamma = 1$), like for the smooth sphere-monopole solution [16].

We now want to find the general solution to Eqs. (5.78)-(5.80) subject to the HCs and NCs. System (5.78)-(5.80) is a rather complicated set of coupled differential equations, but the case at hand can be elegantly solved by using the harmonic expansion of the scalar Laplacian; let us now describe this technique. It is easy to solve the eigenvalue problem for $-D^2$ acting on the 2D scalars (in fact, for $A = 0$ and applying the diagonal HC and the NC, the system is identical to that for the helicity-2 field above). The eigenfunctions are then given by (5.65-5.68) with $\omega = \bar{\omega}$, and the eigenvalues can be written as:

$$\mu_{\mathbf{n}\mathbf{m}}^2 \equiv \frac{4}{r_0^2} (\mathbf{n} + |\mathbf{m}|\omega) (\mathbf{n} + |\mathbf{m}|\omega + 1) \geq 0, \quad (5.85)$$

where $\mathbf{n} = 0, 1, 2, 3, \dots$ and \mathbf{m} run over all the integers. This is the generalization to the rugbyball of the scalar spherical harmonics. They form a complete basis for the 2D scalar fields V_i .

We next proceed by determining a complete basis for the 2D vector fluctuations $\{h_{\theta i}, h_{\varphi i}\}$ and $\{V_{\theta i}, V_{\varphi i}\}$. We focus only on $\{h_{\theta i}, h_{\varphi i}\}$ as the analysis for $\{V_{\theta i}, V_{\varphi i}\}$ is identical. A way to determine such a basis is to look at the eigenvalue problem for the operator $-D^2 + R/2$ appearing in Eq. (5.78), because the diagonal HC (5.81) guarantees that this operator is Hermitian over the space of functions where $\{h_{\theta i}, h_{\varphi i}\}$ lives, and therefore has a complete basis of eigenfunctions. Again, this system is easy to solve, this time using the results²⁴ of Ref. [9]. We therefore just summarise the results. The eigenvalue problem for $-D^2 + R/2$ on 2D vectors generically mixes the $h_{\theta i}$ and $h_{\varphi i}$ components, but reduces to a diagonal form, at least in the rugbyball case, by introducing²⁵

$$h_{\pm i} \equiv \frac{1}{\sqrt{r_0}} \left(e^{B/4} h_{\theta i} \pm i e^{-B/4} h_{\varphi i} \right). \quad (5.86)$$

Eq. (5.86) defines a new basis for tensors on the 2D internal space, and we remind the reader that for the rugbyball $e^B = \alpha^2 \sin^2 \theta$. The squared mass problem can then be transformed into a pair of decoupled Schrödinger equations, which can be solved. Note that the singularities of the spin-1 action discussed in Subsection 4.2.3 appear in the Schrödinger problems as two singular points in the effective potentials (one for each brane), which do not obstruct the determination of the spectrum [6, 9]. The $h_{\pm i}$ fields can be KK expanded as follows

$$h_{\pm i}(X) = \sum_{\mathbf{n}, \mathbf{m}} h_{\pm i \mathbf{n}\mathbf{m}}(x) f_{\mathbf{n}\mathbf{m}}^{\pm}(\theta) e^{i\mathbf{m}\varphi}, \quad (5.87)$$

and, in the case

$$\mathbf{m} = 0 \quad \text{or} \quad |\mathbf{m}| \geq 1/\omega, \quad (5.88)$$

²⁴In Ref. [9] a more general problem has been solved, which reduces to the present one in the unwarped case $A = 0$.

²⁵The \pm appearing in (5.86) and throughout this section should not be confused with the (\pm) used to defined the light-cone gauge in Subsection 4.1, for this reason the latter are written inside brackets.

both the KK tower associated with f^+ and f^- turn out to be exactly that in (5.85), where $\mathbf{n} = 0, 1, 2, 3, \dots$ and \mathbf{m} run over all the integers, but with the constraint $\{\mathbf{n}, \mathbf{m}\} \neq \{0, 0\}$. Condition (5.88) is satisfied by every $|\mathbf{m}|$ for non-negative tensions and it is satisfied by some (but not all) $|\mathbf{m}|$ for negative tensions. This, however, will be enough to show that when the tensions assume the values in (5.74), the KK spectra include extra massless spin-1 fields. In the following we denote (5.88) with $0 \not\prec |\mathbf{m}| \omega \not\prec 1$.

So we have found that, for modes satisfying (5.88), the spectrum of $-D^2 + R/2$ on 2D vectors is made up of two identical copies of the spectrum of $-D^2$ on 2D scalars but with zero mode removed. This suggests that we may be able to express the eigenfunctions of $-D^2 + R/2$ on 2D vectors in terms of eigenfunctions of $-D^2$ on 2D scalars. Indeed, if we consider a solution V to the eigenvalue problem of $-D^2$ with eigenvalue μ^2 , then it is easy to show that $\partial_m V$ is an eigenfunction of $-D^2 + R/2$ with the same eigenvalue. In the case (5.88), this implies that we can write

$$\partial_{\pm} (f_{\mathbf{n}\mathbf{m}}(\theta) e^{i\mathbf{m}\varphi}) = c_{\mathbf{n}\mathbf{m}} f_{\mathbf{n}\mathbf{m}}^{\pm}(\theta) e^{i\mathbf{m}\varphi}, \quad (5.89)$$

where we have used the basis defined in Eq.(5.86) for ∂_{\pm} , and moreover $c_{\mathbf{n}\mathbf{m}}$ are normalization constants which, having chosen a convenient normalization for the wave functions, can be fixed to be $c_{\mathbf{n}\mathbf{m}} = \mu_{\mathbf{n}\mathbf{m}}/\sqrt{2}$. This is the analogue of the derivative relation that exists between Wigner functions for fields of different spin on the sphere (see eq. (3.17) of Ref. [31]). So, remarkably, we can construct the complete harmonic expansion for²⁶ V_i , $h_{\pm i}$ and $V_{\pm i}$ by using the solution to the eigenvalue problem for the scalar Laplacian. Moreover, it is easy to check that, having applied the diagonal HCs to derive the complete basis for $h_{\pm i}$ and $V_{\pm i}$, the mixed HCs (5.82) are automatically satisfied.

Having derived the harmonic expansions one more observation is necessary. It turns out that the factor F_+^+ which appears in the mixing terms $S^{(1)}(h, V)$ and $S^{(1)}(V_2, V)$ is constant for the rugbyball ($F_+^+ = 2\sqrt{2}i/(r_0\kappa)$). Putting everything together, we are then able to transform the differential eigenvalue problem for the squared mass operator into an algebraic problem that can be solved. In particular, after integrating out the extra dimensions, Action (5.76) assumes the following form in terms of the KK modes.

$$\begin{aligned} & S^{(1)}(h, h) + S^{(1)}(V_2, V_2) + S^{(1)}(h, V) + S^{(1)}(V, V_2) + S^{(1)}(V, V) \\ = & \int d^4x \sum_{\mathbf{n}, \mathbf{m}} \left\{ \frac{1}{2\kappa^2} (h_{+i \mathbf{n}\mathbf{m}})^* (\partial^2 - \mu_{\mathbf{n}\mathbf{m}}^2) h_{+i \mathbf{n}\mathbf{m}} + \frac{1}{2\kappa^2} (h_{-i \mathbf{n}\mathbf{m}})^* (\partial^2 - \mu_{\mathbf{n}\mathbf{m}}^2) h_{-i \mathbf{n}\mathbf{m}} \right. \\ & + \frac{1}{8} (V_{+i \mathbf{n}\mathbf{m}})^* (\partial^2 - \mu_{\mathbf{n}\mathbf{m}}^2) V_{+i \mathbf{n}\mathbf{m}} + \frac{1}{8} (V_{-i \mathbf{n}\mathbf{m}})^* (\partial^2 - \mu_{\mathbf{n}\mathbf{m}}^2) V_{-i \mathbf{n}\mathbf{m}} \\ & - \frac{2\mu_{\mathbf{n}\mathbf{m}} i}{r_0\kappa} [(h_{+i \mathbf{n}\mathbf{m}})^* V_{i \mathbf{n}\mathbf{m}} - (h_{-i \mathbf{n}\mathbf{m}})^* V_{i \mathbf{n}\mathbf{m}}] \\ & + \frac{\kappa}{2} \gamma ((V_{+i \mathbf{n}\mathbf{m}})^* V_{i \mathbf{n}\mathbf{m}} - (V_{-i \mathbf{n}\mathbf{m}})^* V_{i \mathbf{n}\mathbf{m}}) \Big] \\ & \left. + \frac{1}{2} (V_{i \mathbf{n}\mathbf{m}})^* \left(\partial^2 - \mu_{\mathbf{n}\mathbf{m}}^2 - \frac{8\gamma^2}{r_0^2} \right) V_{i \mathbf{n}\mathbf{m}} \right\}, \quad (5.90) \end{aligned}$$

²⁶It is easy to show that all we have stated about the harmonic expansion for h_{mi} holds for V_{mi} as well.

		Squared mass	Multiplicity
6D Einstein-Yang-Mills-Λ		0	1
		$\mu_{\mathbf{n}\mathbf{m}}^2 + \frac{2\sqrt{2}}{r_0}\mu_{\mathbf{n}\mathbf{m}}$	1
		$\mu_{\mathbf{n}\mathbf{m}}^2 - \frac{2\sqrt{2}}{r_0}\mu_{\mathbf{n}\mathbf{m}}$	1

		Squared mass	Multiplicity
6D Supergravity		$\frac{8}{r_0^2}$	1
		$\mu_{\mathbf{n}\mathbf{m}}^2$	2
		$\frac{4}{r_0^2} \left(1 + \frac{r_0^2}{4}\mu_{\mathbf{n}\mathbf{m}}^2 + \sqrt{1 + r_0^2\mu_{\mathbf{n}\mathbf{m}}^2} \right)$	1
		$\frac{4}{r_0^2} \left(1 + \frac{r_0^2}{4}\mu_{\mathbf{n}\mathbf{m}}^2 - \sqrt{1 + r_0^2\mu_{\mathbf{n}\mathbf{m}}^2} \right)$	1

Table 1: Squared mass KK towers of physical spin-1 perturbations around the rugbyball solution to the 6D EYMA model and 6D supergravity, for modes satisfying (5.88). $\mu_{\mathbf{n}\mathbf{m}}^2$ is defined in (5.85), but here the KK numbers \mathbf{n}, \mathbf{m} run over $\mathbf{n} = 0, 1, 2, 3, \dots$, $\mathbf{m} = 0, \pm 1, \pm 2, \pm 3, \dots$, with the constraint $\{\mathbf{n}, \mathbf{m}\} \neq \{0, 0\}$.

where the sum over \mathbf{n} and \mathbf{m} is performed over $\mathbf{n} = 0, 1, 2, 3, \dots$ and $\mathbf{m} = 0, \pm 1, \pm 2, \pm 3, \dots$, but with the condition $h_{\pm i 0,0} = 0$ and $V_{\pm i 0,0} = 0$. Also, as a consequence of the reality conditions $h_{+i}(X) = h_{-i}^*(X)$, $V_{+i}(X) = V_{-i}^*(X)$ and $V_i(X) = V_i^*(X)$, we have the relations $h_{+i \mathbf{n}\mathbf{m}}(x) = h_{-i \mathbf{n} - \mathbf{m}}^*(x)$, $V_{+i \mathbf{n}\mathbf{m}}(x) = V_{-i \mathbf{n} - \mathbf{m}}^*(x)$ and $V_{i \mathbf{n}\mathbf{m}}(x) = V_{i \mathbf{n} - \mathbf{m}}^*(x)$.

In this way, the squared mass operator has finally been transformed into an algebraic matrix with constant entries and we can find its eigenvalues exactly.

6D EYMA model To address the spin-1 fluctuations in the 6D EYMA model, we set $\gamma = 0$ and remove the Kalb-Ramond perturbations ($V_{\pm i} = 0$) in the above 4D bilinear action, Eq. (5.90). By diagonalizing the corresponding mass-matrix, we find that the explicit helicity-(± 1) towers are as follows

$$M_{\mathbf{n}\mathbf{m}}^2 = \mu_{\mathbf{n}\mathbf{m}}^2 \geq 0, \quad (5.91)$$

with $\mathbf{n} = 0, 1, 2, 3, \dots$ and $\mathbf{m} = 0, \pm 1, \pm 2, \pm 3, \dots$, but $0 \not\prec |\mathbf{m}| \omega \not\prec 1$ and

$$M_{\mathbf{n}\mathbf{m}}^2 = \mu_{\mathbf{n}\mathbf{m}}^2 \pm \frac{2\sqrt{2}}{r_0}\mu_{\mathbf{n}\mathbf{m}} \geq 0, \quad (5.92)$$

with $\mathbf{n} = 0, 1, 2, 3, \dots$ and $\mathbf{m} = 0, \pm 1, \pm 2, \pm 3, \dots$, but $0 \not\prec |\mathbf{m}| \omega \not\prec 1$ and $\{\mathbf{n}, \mathbf{m}\} \neq \{0, 0\}$. Neither tachyons nor ghosts are found. The $\{\mathbf{n}, \mathbf{m}\} = \{0, 0\}$ mode in (5.91) is the massless

gauge field associated with the 6D monopole $U(1)$, which we have previously discussed in (5.84). The remaining modes in (5.91) are instead the helicity- (± 1) components of massive gravitons; we observe that the massive part of the KK tower (5.70) is exactly reproduced by (5.91), according to 4D Poincaré invariance. The KK towers in (5.92) correspond instead to physical spin-1 fields. The complete set of masses for physical spin-1 fields is given in Table 1.

By analyzing those towers, one easily finds that there are physical massless 4D spin-1 fields (in addition to (5.84)) if and only if $\mu_{\mathbf{n}\mathbf{m}}^2 = 8/r_0^2$, which can be restated as

$$\{\mathbf{n}, \mathbf{m}\omega\} = \{1, 0\} \quad \text{or} \quad \{\mathbf{n}, \mathbf{m}\omega\} = \{0, \pm 1\}. \quad (5.93)$$

Therefore there is at least one massless spin-1 field for any value of the tension (this corresponds to $\{\mathbf{n}, \mathbf{m}\} = \{1, 0\}$). In the sphere case ($\omega = 1$) we have three ways to satisfy this condition, that is

$$\{\mathbf{n}, \mathbf{m}\} = \{1, 0\}, \{0, \pm 1\},$$

which correspond to the three gauge fields of $SU(2)_{KK}$, whereas for positive tension rugbyballs and generic saddle-spheres there is only one choice:

$$\{\mathbf{n}, \mathbf{m}\} = \{1, 0\},$$

corresponding to the KK gauge group $U(1)_{KK}$. However, for the special saddle-spheres for which (5.74) holds, *the number of massless vector fields is enhanced from one plus one to one plus three!*

We shall discuss in detail the physical significance of these modes in the following subsection.

6D supergravity We conclude this subsection by providing the helicity- (± 1) masses for the 6D supergravity (set $\gamma = 1$ and keep the Kalb-Ramond fluctuations in Eq. (5.90)). Diagonalizing the corresponding mass-matrix, we find:

$$M^2 = \frac{8}{r_0^2} > 0, \quad (5.94)$$

which is the vector field associated with the monopole $U(1)$,

$$M_{\mathbf{n}\mathbf{m}}^2 = \mu_{\mathbf{n}\mathbf{m}}^2 \geq 0, \quad \text{with multiplicity } 3 \quad (5.95)$$

and

$$M_{\mathbf{n}\mathbf{m}}^2 = \frac{4}{r_0^2} \left(1 + \frac{r_0^2}{4} \mu_{\mathbf{n}\mathbf{m}}^2 \pm \sqrt{1 + r_0^2 \mu_{\mathbf{n}\mathbf{m}}^2} \right) \geq 0 \quad (5.96)$$

where $\mathbf{n} = 0, 1, 2, 3, \dots$, $\mathbf{m} = 0, \pm 1, \pm 2, \pm 3, \dots$, $\{\mathbf{n}, \mathbf{m}\} \neq \{0, 0\}$ and $0 \not\prec |\mathbf{m}|\omega \not\prec 1$. The masses in (5.94), two towers out of three in (5.95) and the towers in (5.96) correspond to physical spin-1 fields, whereas one of the towers in (5.95) are the helicity- (± 1) components of massive gravitons. The complete set of masses for the physical spin-1 fields is summarized in Table 1. We note that neither tachyons nor ghosts are found.

Regarding the massless vector fields the situation is similar to the 6D EYMA model. It is easy to see that the condition for masslessness is again (5.93) and, therefore, again we have a single KK massless gauge boson for positive tensions and generic negative tensions; instead, for negative tensions of the form (5.74), *the number of massless vector fields is enhanced from one to three!*

As an effective check of the results presented in this subsection, we have also derived the aforementioned spectrum in the sphere case ($\omega = 1$) by expanding the bulk fields over the Wigner functions as in [31] and obtained exactly the sphere limit of our towers.

5.3 Massless vectors and 4D gauge symmetries

In the previous subsection we observed *three* massless 4D vector fields amongst the KK spectra for the 6D models²⁷ on both the sphere and the special saddle-spheres (5.74). We shall now address the physical significance of these modes. One of them, the one with axial quantum number $\mathbf{m} = 0$, should provide the gauge boson for the $U(1)_{KK}$ gauge symmetry, descending from the axial isometry of the internal space. The other two massless vectors, having $\mathbf{m} \neq 0$, are charged under this axial symmetry and so we may expect the three vectors to fit into a non-Abelian structure, like $SU(2)$. For the sphere, this is indeed the case, and the three massless vectors compose the gauge fields of an $SU(2)$ gauge symmetry in the 4D theory. What happens for the special saddle-spheres, where there is no $SU(2)$ isometry in the background? Let's take $\alpha = 2, 3, \dots$, so we consider the special saddle-spheres (we also allow for the smooth sphere with $\alpha = 1$).

5.3.1 Why there are three massless vector modes

Let us begin by understanding why three massless vector modes appear in the spectrum, despite the fact that any branes clearly break the $SU(2)$ isometries of the sphere.

Above, we found that the massless vector fields arise as a linear combination of $h_{\mu}^m{}_{\mathbf{m}\mathbf{n}}(x)$ and $V_{\mu}^m{}_{\mathbf{m}\mathbf{n}}(x)$ (and $V_{\mu}^m{}_{\mathbf{m}\mathbf{n}}(x)$ for 6D supergravity), once we have integrated out the extra dimensions. In detail, if one takes the squared mass matrix defined implicitly by the 4D bilinear action in (5.90) in *e.g.* the EYMA case, one finds that the mass eigenstates are ($\{\mathbf{n}, \mathbf{m}\} \neq (0, 0)$):

$$\begin{aligned} A_{i\ \mathbf{n}\mathbf{m}} &= \frac{i}{2}h_{+i\ \mathbf{n}\mathbf{m}} - \frac{i}{2}h_{-i\ \mathbf{n}\mathbf{m}} + \frac{1}{\sqrt{2}}V_{i\ \mathbf{n}\mathbf{m}}, \\ U_{i\ \mathbf{n}\mathbf{m}} &= -\frac{i}{2}h_{+i\ \mathbf{n}\mathbf{m}} + \frac{i}{2}h_{-i\ \mathbf{n}\mathbf{m}} + \frac{1}{\sqrt{2}}V_{i\ \mathbf{n}\mathbf{m}}, \\ W_{i\ \mathbf{n}\mathbf{m}} &= \frac{1}{\sqrt{2}}h_{+i\ \mathbf{n}\mathbf{m}} + \frac{1}{\sqrt{2}}h_{-i\ \mathbf{n}\mathbf{m}}, \end{aligned} \tag{5.97}$$

corresponding respectively to $M_{\mathbf{n}\mathbf{m}}^2 = \mu_{\mathbf{n}\mathbf{m}}^2 - (2\sqrt{2}/r_0)\mu_{\mathbf{n}\mathbf{m}}$, $M_{\mathbf{n}\mathbf{m}}^2 = \mu_{\mathbf{n}\mathbf{m}}^2 + (2\sqrt{2}/r_0)\mu_{\mathbf{n}\mathbf{m}}$ and $M_{\mathbf{n}\mathbf{m}}^2 = \mu_{\mathbf{n}\mathbf{m}}^2$ in (5.92) and (5.91). Recall that the massless modes emerge from the $A_{i\ \mathbf{n}\mathbf{m}}$ tower, when $\{\mathbf{n}, \mathbf{m}\} = \{1, 0\}, \{0, \pm 1\}$.

²⁷In the EYMA model there is also a massless vector field descending from the higher dimensional $U(1)$ gauge field, which forms a 4D $U(1)$ gauge field.

We can write, then, the harmonic expansion of $h_\mu^m(X)$ as:

$$h_\mu^m(X) = \sum_{I=-,0,+} A_\mu^I(x) K^{Im}(\theta, \varphi) + \text{massive modes}. \quad (5.98)$$

Using the expansion (5.86,5.87), the explicit form for the wave functions (5.89) and the rearrangement in terms of the mass eigenstates (5.97), it is straightforward to show that $K^{Im}(\theta, \varphi)$ indeed correspond to the Killing vectors (5.75) on the special saddle-sphere, where $I = 0$ corresponds to $\{\mathbf{n}, \mathbf{m}\omega\} = \{1, 0\}$ and $I = \pm$ to $\{\mathbf{n}, \mathbf{m}\omega\} = \{0, \pm 1\}$. This is just as in the traditional KK reduction scheme.

In this way, we confirm that the presence of *infinitesimal isometries* on the internal space, which are generated by Killing vector fields, is sufficient for the appearance of massless vector modes – even if they cannot be integrated to genuine isometries.

5.3.2 The absence of enhanced gauge symmetries in the full 4D theory

We now ask whether or not these massless vector modes behave as gauge fields of an $SU(2)$ gauge symmetry. The linearized 4D theory cannot probe any non-Abelian structure, and so to understand the gauge invariance of the full 4D theory, we must go beyond linear order. To this end, we consider a simple extension of the EYMA model, where we add a single complex, massless, neutral scalar field which has an action:

$$S_\Phi = - \int d^6 X \sqrt{-G} \partial_M \Phi^* \partial^M \Phi \quad (5.99)$$

and which assumes a trivial VEV in the saddle-sphere background. It is easy to see that the linearized equation of motion for $Z := \delta\Phi$ gives rise to the rugbyball scalar harmonics (see above Eq. (5.85)):

$$Z(X) = \sum_{\mathbf{m}, \mathbf{n}} z_{\mathbf{m}\mathbf{n}}(x) f_{\mathbf{m}\mathbf{n}}(\theta) e^{i\mathbf{m}\varphi} \quad (5.100)$$

with the corresponding masses (5.85):

$$M^2 = \frac{4}{r_0^2} l(l+1) \quad \text{where} \quad l = \mathbf{n} + |\mathbf{m}|\omega. \quad (5.101)$$

The multiplicity of a given mass is given by $2l+1$ when l is integer or half-odd integer; otherwise it is given by $2([l] + \frac{1}{2}) + 1$, where $[l]$ denotes the integer part of l . We also note that for l integer (which corresponds also to $\mathbf{m}\omega$ integer), the wavefunction $f_{\mathbf{m}\mathbf{n}}(\theta)$ is an Associated Legendre function, just as for the spherical harmonics. The modes with l non-integer are instead additional harmonics, which generically have no corresponding states amongst the spherical harmonics nor indeed any of the Wigner functions.

Now let us ask how the 4D fields $z_{\mathbf{m}\mathbf{n}}(x)$ couple to the massless vector fields, and in particular if they do in a gauge-invariant way. At trilinear level, this coupling descends only from the term:

$$S(Z^*, h_{\mu m}, Z) = - \int d^6 X \sqrt{-G} \partial^\mu Z^* h_\mu^m \partial_m Z \quad (5.102)$$

and its complex conjugate. The above trilinear coupling can now be reexpressed in terms of the 4D fields, and isolating the contributions involving the massless vectors, Eq. (5.98), we find:

$$S(z^*, A_\mu^I, z) = - \int d^4x \sqrt{-g_4} \partial^\mu z^* A_\mu^I z' \int d\theta d\varphi \frac{r_0^2}{4} \alpha \sin \theta f e^{-im\varphi} K^{Im} \partial_m \left(f' e^{im'\varphi} \right), \quad (5.103)$$

where we have suppressed the KK indices $\{\mathbf{n}, \mathbf{m}\}$ and $\{\mathbf{n}', \mathbf{m}'\}$ on z, f and z', f' respectively. Performing the integral over the internal dimensions:

$$g_I = \int d\theta d\varphi \frac{r_0^2}{4} \alpha \sin \theta f e^{-im\varphi} K^{Im} \partial_m \left(f' e^{im'\varphi} \right), \quad (5.104)$$

we see that the wavefunction overlap (5.104) gives the 4D coupling between $z(x)$ and $z'(x)$ via a massless vector field, $A_\mu^I(x)$:

$$- g_I \int d^4x \sqrt{-g_4} \partial^\mu z^* A_\mu^I z'. \quad (5.105)$$

Observe that if the full 4D theory were to respect an $SU(2)$ gauge symmetry whose gauge fields are the three massless vector modes, then z and z' would belong to $SU(2)$ multiplets of the same dimension and (5.105) would correspond to the trilinear terms in the gauge invariant combination $-D_\mu z^{a*} D^\mu z'^a$, where $D_\mu z^a = \partial_\mu z^a + A_\mu^I T^I{}_b^a z^b$, the indices a, b run over $a, b = 1, \dots, r$ and r is the size of the multiplet. For the classic sphere, $\alpha = 1$, this is of course the case, and the wave function overlaps in (5.104) are zero unless z and z' belong to the same $SU(2)$ multiplet, thanks to the properties of the spherical harmonics. We shall now see that such a structure does not hold for the special saddle-spheres.

To this purpose, let us consider the rugbyball harmonics with $0 < \mathbf{m}\omega < 1$. The pattern that emerges for the overlaps (5.104) once both the integrals over $d\theta$ and $d\phi$ are performed²⁸, is that a mode, f , with $0 < \mathbf{m}\omega < 1$ and \mathbf{n} even (*respectively* odd) has a non-zero overlap with the modes, f' , for which $\mathbf{m}'\omega = \mathbf{m}\omega \mp 1$ and all \mathbf{n}' odd (*respectively* even). It can then easily be seen that this prevents the realization of an $SU(2)$ gauge symmetry. Take for instance the set of modes $\{z\}$ with some mass-squared $l(l+1)$ in which $0 < |\mathbf{m}|\omega < 1$ and $\mathbf{n} = 0$. This mass comes only with degeneracy 2, corresponding to KK numbers $\{0, \pm\mathbf{m}\}$. Therefore, if there exists an $SU(2)$ gauge symmetry, then the modes in $\{z\}$ fall either into an $SU(2)$ doublet or two singlets. The overlap (5.104) between the modes $\{0, \mathbf{m}\}$ and $\{0, -\mathbf{m}\}$ is zero, and the subsequent vanishing of the coupling in (5.105) tells us that $\{z\}$ cannot form a doublet. On the other hand, the modes $\{0, \pm\mathbf{m}\}$ do have a non-zero overlap with $\{\mathbf{n} \text{ odd}, \pm\mathbf{m} - 1/\omega\}$ and $\{\mathbf{n} \text{ odd}, \pm\mathbf{m} + 1/\omega\}$, and so the two modes in $\{z\}$ each have a trilinear coupling (5.105) with towers of z' and the massless vectors fields. Thus, they cannot be singlets. In this way we can conclude that there does not exist an $SU(2)$ gauge symmetry corresponding to the massless vector fields.

We would like to draw one more insight into the absence of $SU(2)$ gauge symmetry for the full 4D theory. The Killing vectors (5.75) can be considered as generators of an

²⁸Whilst we have not checked this result for all ω, \mathbf{m} and \mathbf{n} the pattern is quite convincing.

$SU(2)$ algebra, and the mass-squared operator for the saddle-sphere scalars, $-D^2$, can be understood as the Casimir Operator for the algebra: $-\frac{r_0^2}{4}D^2 = \frac{1}{2}(K^+K^- + K^-K^+) + (K^0)^2$. The saddle-sphere scalar harmonics form a basis for the Hilbert space of functions on which the Hermitian operator, $-D^2$ (plus boundary conditions), acts. However, the $SU(2)$ ladder operators, K^\pm , do not act within this Hilbert space: the action of K^\pm on the harmonics $f_{\mathbf{m}\mathbf{n}}(\theta)e^{i\mathbf{m}\varphi}$ with $0 < |\mathbf{m}|\omega < 1$ gives back functions which do not obey the NC and HC boundary conditions. Again, we see that the saddle-sphere harmonics do not furnish well-defined representations of the $SU(2)_{KK}$ generated by the Killing vectors, and it is precisely the modes with $0 < |\mathbf{m}|\omega < 1$ that are the problematic ones²⁹.

As we have implied above, the absence of an $SU(2)$ KK gauge symmetry in the 4D theory can be understood in the 6D picture as being due to the absence of a genuine $SU(2)$ isometry in the internal dimensions.

5.3.3 The emergence of enhanced gauge symmetries at low energies

Finally, notice that although the modes with $0 < |\mathbf{m}|\omega < 1$ do not belong to well-defined $SU(2)$ representations, the massless wave functions that we have found are equivalent to those present in the sphere case (up to an integer constant multiplying φ) and do furnish well-defined $SU(2)$ representations³⁰. This holds also for the massless spin-2 and spin-1 fields above, as well as the massless spin-0 fields discussed below³¹. Therefore, the classical low energy 4D effective theory that results from truncating the massive modes does enjoy an $SU(2)$ KK gauge invariance *to all orders in perturbation theory* – despite the absence of a genuine $SU(2)$ isometry in the extra dimensions. Indeed, this low energy 4D theory does not distinguish between a compactification on a smooth sphere or a special saddle-sphere!

Moreover, we can argue that the above truncation to the massless sector is a consistent one³², at least for the bosonic theory that we have studied whose field content is identical to that of 6D supergravity. Then, if we remove the branes and replace the singular space with a smooth non-compact manifold, the local geometry is the same for the sphere everywhere and the KK ansatz for the special saddle-sphere is essentially identical to that of the smooth sphere. Meanwhile, the sphere reduction of 6D supergravity was shown to be a consistent one in Ref. [35], thanks to a remarkable conspiracy between properties of the 2-sphere and

²⁹Notice that this range of \mathbf{m} is empty for the special saddle-sphere with $\omega = \frac{1}{2}$ if we impose the Z_2 orbifold projection discussed in Appendix C. In this case, then, all the KK modes are in well-defined representations of $SU(2)$ (corresponding to the Wigner functions), and we can expect an $SU(2)$ gauge invariance in the full 4D theory, at least if we remove the branes and discuss a smooth non-compact manifold. This is not surprising, since – outside the branes – the Z_2 orbifolding effectively cancels out the $\delta = -2\pi$ deficit angle, and we return to the standard sphere case.

³⁰This is a consequence of the fact that in our mass-squared's, M^2 , as well as in our $f_{\mathbf{m}\mathbf{n}}(\theta)$ wavefunctions, \mathbf{m} and ω enter only through the combination $\mathbf{m}\omega$. This is obvious for the masses, and for the wave functions it can be seen from Eq. (5.68), after setting $\bar{\omega} = \omega$ to recover the spectra for the rugbyball. Furthermore, the massless modes all have integer $\mathbf{m}\omega$.

³¹We should caution that, although there are no symmetries that suggest them to be massless, our harmonic analysis has not included the modes with $0 < |\mathbf{m}|\omega < 1$ in the spin-1 sector, nor the modes with $0 < |\mathbf{m}|\omega < 1$ and $1 < |\mathbf{m}|\omega < 2$ in the spin-0 sector.

³²Mathematical consistency may of course not be necessary, if the truncation is consistent up to some energy scale.

the structure of supergravity.

5.4 Massless scalars

Finally, we turn to the spectra of 4D spin-0 fields, which are governed by the action given in Appendix B. In Appendix C we give the complete spectra for unwarped braneworld compactifications in 6D supergravity. Here, our focus shall be on the massless scalars featured in the low energy 4D effective theory. Again we shall first review the RS model and then examine the 6D braneworld models.

5.4.1 Randall-Sundrum

In the RS model of Ref. [1], the massless scalar sector involves one normalizable mode (the radion), which becomes non normalizable in the decompactification limit $r_c \rightarrow \infty$ [29]. Let us find this mode in our formalism. We can of course restrict our attention to the spin-0 action $S^{(0)}(h, h)$ as in [1] only gravity is introduced and the branons are consistently projected out by the S^1/Z_2 orbifold conditions. Therefore, we only have to deal with the perturbation $h_{\rho\rho}$, because (4.50) implies $h_i^i = -h_{\rho\rho}$. It is easy to derive the EOM for $h_{\rho\rho}$:

$$-\frac{1}{\sqrt{-G}}\partial_M\left(\sqrt{-G}\partial^M h_{\rho\rho}\right)+\left[-A'^2+\frac{1}{3}T\kappa^2\delta(\rho-Y)\right]h_{\rho\rho}=0, \quad (5.106)$$

where $T\kappa^2\delta(\rho-Y)\equiv T_1\kappa^2\delta(\rho)+T_2\kappa^2\delta(\rho-\pi r_c)$. We now perform a KK decomposition $h_{\rho\rho}(x,\rho)=\sum_{\mathbf{n}}h_{\rho\rho}(x)f_{\mathbf{n}}(\rho)$ and focus on the massless case ($\eta^{\mu\nu}\partial_\mu\partial_\nu=0$); we obtain the simple equation

$$\psi''=0, \quad (5.107)$$

where we have defined $\psi\equiv e^A f$ and used property (5.73). The only solution to (5.107) satisfying the S^1/Z_2 orbifold conditions is ψ constant, which corresponds to

$$f\propto e^{-A}. \quad (5.108)$$

Mode (5.108) is the wave function of the radion. By inserting this mode in the kinetic term of $h_{\rho\rho}$ in (B.112) one easily finds that it is normalizable for any finite r_c , but becomes non normalizable in the limit $r_c\rightarrow\infty$.

5.4.2 6D Brane Worlds

After this non-trivial check of our formalism we now turn to the conical-GGP solutions of 6D supergravity. The stability of the GGP solutions has been investigated in [7] and [9], where no tachyons emerged unless non-Abelian gauge groups are considered. Indeed, in the presence of non-Abelian gauge groups, an instability may arise in the sector described by the action $S^{(0)}(V, V)$, with $V_{\underline{m}}$ orthogonal to the background monopole³³ [9]. We observe that, even in the absence of non-Abelian gauge groups, the stability of the GGP solutions is

³³This instability is also present in the sphere-monopole solution [34], which is a particular case of the GGP solutions.

marginal, in the sense that there are necessarily massless scalars in the physical spectrum. These massless particles are manifestations of two symmetries in the model. One is the following invariance of the EOMs: $G_{MN} \rightarrow w G_{MN}$ and $e^{\phi/2} \rightarrow w e^{\phi/2}$, where w is a real number. Note that this is only a classical symmetry because the action rescales as $S_B \rightarrow w^2 S_B$, so we do not expect the corresponding scalar to remain massless once quantum corrections are included. The other is the Kalb-Ramond symmetry, which acts as $B_2 \rightarrow B_2 + d\Lambda$, where Λ is a general 1-form field. The actual presence of the zero mode corresponding to the former symmetry has been shown in Refs. [11, 7].

Here, by using our bilinear action, we can easily figure out where the other massless scalar is. This emerges as the lightest 4D mode of the field V_{ij} , whose bilinear action is simply (see Eq. (B.118))

$$-\frac{1}{16} \int d^6 X \sqrt{-G} e^{\phi-2A} \partial_M V_{ij} \partial^M V_{ij}. \quad (5.109)$$

This action is equivalent to the spin-2 action (4.52) in the case of the conical-GGP solutions, which satisfy $A = \phi/2$. The wave functions and mass spectrum coming from V_{ij} are therefore identical to the one presented in Subsection 5.1. For $\mathbf{n} = \mathbf{m} = 0$ we obtain the massless scalar field associated to the Kalb-Ramond symmetry. In the spherical limit this corresponds to the $l = 0$ mode in (5.71) [27].

6 Summary of Results

Before concluding, let us provide an overview of our results.

- We have derived the linearized dynamics, Eqs. (4.52)-(4.57) and Appendix B, for the physical perturbations about general backgrounds in a general class of field theories. In particular, we take Einstein-Yang Mills (EYM) theory in D spacetime dimensions, with a bulk dilaton or cosmological constant (Λ), and a number of dynamical 3-branes. Moreover, for $D = 6$ we include a dilaton and 2-form potential. Therefore, 6D chiral supergravity, D -dimensional EYMA theory and the 5D Randall-Sundrum models all fall within our analysis. The backgrounds considered respect 4D Poincaré invariance, but may be warped in a radial transverse coordinate.
- Taking the Randall-Sundrum models as an illustrative example within our formalism, we retrieve the well-known dynamics for spin-2 fluctuations and identify the massless scalar (the radion), which is normalizable in the two brane model and becomes non-normalizable in the one brane model.
- For the 6D EYMA model, we consider the unwarped “rugbyball-monopole” compactifications, sourced by two 3-branes of equal tension. When the tensions are zero, we recover the sphere-monopole compactification, and when the tensions are negative we refer to the 2D geometry as a “saddle-sphere”. By deriving a set of “rugbyball harmonics”, we are able to obtain analytic KK spectra; *i.e.* we discuss how to find physical 4D spin-2, spin-1 and – consistently truncating branons – spin-0 fields and

	spin-2	spin-1	spin-0
Rugby-ball $\delta \geq 0$	<i>all modes</i>	<i>all modes</i>	<i>all modes</i>
Saddle-sphere $\delta = -2\pi$	<i>all modes</i>	<i>all modes</i>	<i>all modes</i>
Generic Saddle-sphere	<i>all modes</i>	$\mathbf{m} = 0; \mathbf{m} \geq 1/\omega$	$ \mathbf{m} = 0, 1/\omega; \mathbf{m} \geq 2/\omega$
Warped Models	<i>all modes</i>	–	–

Table 2: The sectors covered in the present paper for Braneworld Compactifications in 6D Supergravity. In order to address the spin-0 sector, we projected out the branons with an orbifolding. We also here impose the orbifolding for all sectors in the presence of negative tension branes.

their masses. We present the full spin-2 spectrum and the spin-1 spectra for axial momentum number $0 \not\prec |\mathbf{m}|\omega \not\prec 1$.

- For the 6D supergravity, the backgrounds of interest are the warped, axially symmetry braneworld (“conical-GGP”) solutions, which have unwarped limits to the rugbyballs and saddle-spheres, and to the sphere. Our focus is on the bosonic “Salam-Sezgin” sector (from the gravity-tensor supermultiplet and the $U(1)$ gauge multiplet in which the background monopole lies), since the remaining bosonic sectors have been treated elsewhere. We obtain the complete spin-2 spectrum. For the spin-1 and spin-0 sectors, we restrict to the unwarped backgrounds, and employ the rugbyball harmonics to find the spectra. The sectors covered by our analysis³⁴ are summarized in detail in Table 2.

Our main physical results for the 6D braneworlds are as follows.

- The spin-2 spectrum includes the massless 4D graviton separated from the rest of the KK tower by a mass gap, and the mass gap is indeed observed in all sectors. For rugbyballs with positive deficit angles and for generic saddle-spheres, the spin-1 sector contains a massless KK gauge boson due to the $U(1)$ isometry in the background (in addition to any massless 4D gauge bosons descending from unbroken 6D gauge symmetries). For the special saddle-spheres with deficit angles $\delta = -2\pi, -4\pi, \dots$, there is a qualitative difference. Here, there are three Killing vectors, which are well-defined everywhere outside the branes and obey an $SU(2)$ Lie algebra. Although only one of them integrates to a genuine isometry, the number of massless KK vectors fields is consequently enhanced to three. Meanwhile, in the spin-0 sector for supergravity, we identify the two massless scalar fields expected in all cases from the classical scaling symmetry and the Kalb-Ramond symmetry.
- The spin-2 and spin-1 spectra are all well-behaved despite the presence of codimension-two dynamical brane sources, which induce singularities in the bulk geometry. To make progress in the spin-0 sector, we had to discard the branon modes (*e.g.* by placing the branes at orbifold fixed points).

³⁴We also find the spectrum in the sphere case as a check.

- The spectra analysed – which incorporates *all* modes for rugbyballs sourced by positive tension branes – do not harbour any instabilities; neither tachyons nor ghosts.
- To understand the significance of the three massless 4D vector fields that appear for the special saddle-spheres, we go beyond bilinear order. We find that in the full 4D theory, they do *not* represent gauge fields of an $SU(2)$ gauge symmetry. This is due to the presence of KK modes that are not in well-defined $SU(2)$ representations. The classical masslessness of the vector fields is thus not protected by any symmetry, which is in accordance with the absence of a genuine $SU(2)$ isometry in the background.
- In the massless sector, however, all modes fall into well-defined $SU(2)$ representations. Therefore, the low energy 4D effective theory obtained by truncating to the massless sector does seem to enjoy a classical $SU(2)$ KK gauge symmetry, despite the absence of a background $SU(2)$ isometry! Indeed, this low energy effective theory does not distinguish between compactifications on the sphere and the special saddle-spheres.

7 Conclusions

In this paper, we have provided the dynamics of the physical fluctuations in a wide class of models, which incorporate the bosonic fields generically present in bulk supergravity theories – gravity, non-Abelian gauge fields, the dilaton and two-form potential – as well as dynamical 3-branes. Our final equations ((4.52)-(4.57) and those in Appendix B), which can be considered as a generalization of the analysis in [12], provide the starting point to construct a 4D effective field theory emerging from various higher dimensional models, with compactified extra dimensions and/or branes.

We next proceeded with that objective to study the behaviour of braneworlds solutions in six dimensions, taking as representative the rugbyball compactifications of Einstein-Yang Mills theory with a cosmological constant (EYMA) and certain axi-symmetric warped compactifications to 6D minimal gauged supergravity; the so-called conical-GGP solutions. We have obtained the complete KK spectrum for the 4D spin-2 sector in the conical-GGP solutions, which is a step towards understanding the behaviour of gravity in codimension two braneworld models, as for example studied in [36]. The spin-1 and spin-0 sectors present large systems of coupled differential equations to be solved (five-by-five for the spin-1 fluctuations, eight-by-eight for the spin-0 fluctuations after truncating the branons), and we are able to do so in the unwarped cases by developing “rugbyball harmonics”, in analogy to the spherical harmonics. Along the way, we also recovered some familiar features of the 5D Randall-Sundrum models. Our main results are summarized in the previous section.

Previous studies have revealed that codimension-two braneworld compactifications can evade the traditional KK lore in several ways. For instance, in [6] it was found that the KK mass-gap can be decoupled from the size of the extra dimensions in the presence of negative tension branes, in principle allowing not only gravity but also the SM to propagate in large extra dimensions. This phenomenon can also be observed here. We can also now suggest the following. The power-law warping present in the 6D braneworlds studied here does not change qualitatively the physics. Moreover, models with only positive tension

codimension-two branes also have qualitatively the same behaviour as traditional KK compactifications. Meanwhile, the introduction of negative tension codimension-two branes can lead to surprising dynamics.

As yet another example of how the physics of braneworlds in 6D can counter intuition, we have found – for special saddle-sphere compactifications with deficit angles $\delta = -2\pi, -4\pi, \dots$ – three massless vector fields thanks to the presence of three $SU(2)$ Killing vectors in the internal manifold that are well defined everywhere outside the branes. Thus we see that infinitesimal isometries are sufficient to imply massless vector fields, even if they cannot be integrated to genuine isometries. All the massless modes in the models studied here fall into well-defined representations of the $SU(2)$, although there are massive KK modes which do not. In this way we see that the massless vectors provide the gauge fields of an enhanced $SU(2)$ KK gauge symmetry in the classical, low energy, 4D effective theory obtained by truncating to the massless sector, despite the absence of an $SU(2)$ isometry in the background! Apparently, the low energy theory does not distinguish between a compactification on the special saddle-spheres and the smooth sphere.

At the same time, as we approach the energy of the KK mass gap and incorporate the non-zero modes, we see that the $SU(2)$ KK gauge symmetry is broken explicitly to $U(1)$. This is because only the $U(1)$ is a genuine global continuous isometry of the internal manifold. The masslessness of the extra massless vector fields is thus not protected by any symmetry, and should not survive quantum corrections. Meanwhile, reaching energies far above the KK mass-gap, the full 6D symmetries will be restored as usual. The pattern of symmetry breaking and emergence that we have found within our classical approximation, as different energy scales are probed, is thus a novel one.

In the model whose field content and structure corresponds to the bosonic part of 6D supergravity, the low-energy theory describes the graviton, the three vectors in the adjoint of $SU(2)$ and two massless scalars that are $SU(2)$ singlets. Whether the above properties are shared with fermionic modes is not known and their behaviour, though of interest, lies beyond the scope of the present paper. Meanwhile, we argued that we expect the zero-mode truncation to be a consistent one, at least in the aforementioned model once we remove the brane sources and study a non-compact smooth manifold. We thus note that this bosonic model is in principle a complete one, sufficient to demonstrate the unconventional dynamics that we have observed. It would certainly be interesting to check the consistency also in the presence of branes.

This work concludes our study of the bosonic perturbations in the axi-symmetric brane-world solutions to 6D supergravity. We may now turn to the fermionic sector.

Acknowledgments. It is a pleasure to thank Cliff Burgess, Claudia de Rham, Stephan Mohrdieck, Nicola Pagani, Riccardo Rattazzi, Michele Redi, Mikhail Shaposhnikov and Andrea Wulzer for valuable discussions. S.L.P. is supported by the Deutsche Forschungsgemeinschaft under the Collaborative Research Center 676 and by the European Union 6th framework program MRTN-CT-503359 “Quest for Unification”. A.S. has been supported by the Tomalla Foundation and by CICYT-FEDER-FPA2008-01430. S.L.P. and A.S. thank the High Energy Theory Group at ICTP for hospitality at various stages of this work.

Appendix

A General ξ -Dependent Bilinear Action

Here we give the explicit expression for the bilinear action that depends on the fluctuations of the brane positions ξ^M , before any gauge fixing, that is the last two terms in (3.32). These terms have been computed by varying the brane action (2.6) with respect to (3.28) and by keeping only terms up to the quadratic order. Their explicit expression is the following

$$\begin{aligned}
S(\xi, \xi) = & -\frac{T}{2} \int d^4x \sqrt{-g} [G_{MN} \partial \xi^M \cdot \partial \xi^N \\
& + \frac{1}{2} \xi^P \xi^R \partial_P \partial_R G_{MN} \partial Y^M \cdot \partial Y^N + 2 \xi^P \partial_P G_{MN} \partial \xi^M \cdot \partial Y^N \\
& + \frac{1}{2} \xi^P \partial_P G_{MN} \xi^R \partial_R G_{SQ} \left(\frac{1}{2} \partial Y^M \cdot \partial Y^N \partial Y^S \cdot \partial Y^Q - \partial Y^M \cdot \partial Y^S \partial Y^N \cdot \partial Y^Q \right) \\
& + G_{MN} G_{PR} (\partial \xi^M \cdot \partial Y^N \partial \xi^P \cdot \partial Y^R - 2 \partial \xi^M \cdot \partial \xi^P \partial Y^N \cdot \partial Y^R) \\
& + \xi^P \partial_P G_{MN} G_{RS} (\partial Y^M \cdot \partial Y^N \partial \xi^R \cdot \partial Y^S - 2 \partial Y^M \cdot \partial \xi^R \partial Y^N \cdot \partial Y^S)], \quad (\text{A.110})
\end{aligned}$$

and

$$\begin{aligned}
S(h, \xi) = & -\frac{T}{2} \int d^4x \sqrt{-g} [\xi^P \partial_P h_{MN} \partial Y^M \cdot \partial Y^N + 2 h_{MN} \partial \xi^M \cdot \partial Y^N \\
& + h_{MN} \xi^P \partial_P G_{RS} \left(\frac{1}{2} \partial Y^M \cdot \partial Y^N \partial Y^R \cdot \partial Y^S - \partial Y^M \cdot \partial Y^R \partial Y^N \cdot \partial Y^S \right) \\
& + h_{MN} G_{PR} (\partial Y^M \cdot \partial Y^N \partial \xi^P \cdot \partial Y^R - 2 \partial Y^M \cdot \partial \xi^P \partial Y^N \cdot \partial Y^R)]. \quad (\text{A.111})
\end{aligned}$$

The bulk quantities in (A.110) and (A.111), that is the background metric G_{MN} and the fluctuation h_{MN} , are computed in the background brane position. This is because (A.110) and (A.111) come from the variation of the brane action (2.6) where the bulk fields are computed in the brane position.

B Spin-0 Bilinear Action in the Light Cone Static Gauge

Here we provide the spin-0 action in the light cone static gauge defined by (4.49) and (4.51). This is the only part where the branons $\xi^{\underline{m}}$ appear.

Let us start with the spin-0 action that only depends on the bulk fields. The non vanishing terms are the following:

$$\begin{aligned}
S^{(0)}(h, h) = & -\frac{1}{4\kappa^2} \int d^D X \sqrt{-G} \left[\partial_\mu h_{\underline{mn}} \partial^\mu h^{\underline{mn}} + \partial_\rho h_{\underline{mn}} \partial_\rho h^{\underline{mn}} + h_{\underline{mn};l} h^{\underline{mn};l} \right. \\
& + h_{\rho\rho}^2 (D_2 A' B' + 2A'') + 2(A'' + A'^2) h_{\rho\rho} h_i^i \\
& \left. \left(D_2 A' B' + 2A'' - \frac{1}{2} B'^2 - 2B'' \right) h_{\rho m} h_\rho^m - 4A' h_\rho^{\underline{n}} h_{\underline{m}}^i{}^m \right]
\end{aligned}$$

$$\begin{aligned}
& +h_i^i h_m^m A' B' + 2 \left(B'' + \frac{B'^2}{2} \right) h_{\rho\rho} h_m^m + \frac{1}{2} B'^2 h_{mn} h^{mn} + \frac{1}{2} B'^2 (h_m^m)^2 \\
& - 2e^{-B} h_l^m h^n_h \Omega_{mn}{}^{lh} + 2\kappa^2 h_{lm} h^l_n \left(\frac{1}{2} e^{\phi/2} F^m_h F^{nh} + \frac{1}{4\kappa^2} \partial^m \phi \partial^n \phi \right) \\
& + \kappa^2 e^{\phi/2} h^{mn} h^{lh} F_{lm} F_{hn} + \frac{1}{2} \left(\partial_\mu h_i^i \partial^\mu h_j^j + \partial_\rho h_i^i \partial_\rho h_j^j + h_i^i{}_{;m} h_j^j{}^{;m} \right) \\
& + (h_i^i)^2 \left(\frac{1}{2} A'^2 + \frac{T}{2} \kappa^2 \sqrt{g/G} \delta(X_c - Y_c) \right) \Big], \tag{B.112}
\end{aligned}$$

where $\Omega_{mn}{}^l{}_h$ is the Riemann tensor for the metric K_{mn} and X_c and Y_c are defined below Eq. (4.53). We observe that in the last line of (B.112) there is an explicit brane contribution (the tension of the brane T appears explicitly). Moreover,

$$\begin{aligned}
S^{(0)}(V, V) &= -\frac{1}{2} \int d^D X \sqrt{-G} e^{\phi/2} \left[\partial_\mu V_m \partial^\mu V^m + D_m V_n D^m V^n + \left(-2A'^2 + \frac{1}{4} \phi'^2 \right) V_\rho^2 \right. \\
& + (-2A' + \phi') V_\rho D_m V^m + R^{mn} V_m V_n + 2\bar{g} F^{mn} V_m \times V_n \\
& \left. - \frac{1}{2} \phi' V_\rho D_m V^m + \frac{1}{2} \phi' V^m D_m V_\rho + \kappa^2 e^{\phi/2} \left(F_l{}^m V_m \right) F^{ln} V_n \right], \tag{B.113}
\end{aligned}$$

$$\begin{aligned}
S^{(0)}(h, V) &= \int d^D X \sqrt{-G} e^{\phi/2} \left[F^{nm} V_m h_{ln}{}^{;l} + (D_n V_m - D_m V_n) h_l{}^n F^{lm} \right. \\
& \left. - \frac{1}{2} A' F^{\rho m} V_m h_i^i + A' F^{mn} V_n h_{m\rho} \right], \tag{B.114}
\end{aligned}$$

$$S^{(0)}(\tau, \tau) = - \int d^D X \sqrt{-G} \left[\frac{1}{4\kappa^2} \partial_M \tau \partial^M \tau + \frac{1}{2} \left(\frac{\partial^2 \mathcal{V}}{\partial \phi^2} + \frac{1}{16} e^{\phi/2} F^2 + \frac{\phi'^2}{4\kappa^2} \right) \tau^2 \right], \tag{B.115}$$

$$\begin{aligned}
S^{(0)}(h, \tau) &= \int d^D X \sqrt{-G} \left\{ \frac{1}{2\kappa^2} \left[\phi' \tau \left(A' h_{\rho\rho} + h_{m\rho}{}^{;m} - \frac{1}{2} A' h_i^i \right) + h^{m\rho} \partial_m \tau \phi' \right] \right. \\
& \left. + \frac{1}{4} e^{\phi/2} F^{ml} F^n{}_l \tau h_{mn} \right\}, \tag{B.116}
\end{aligned}$$

$$S^{(0)}(V, \tau) = \int d^D X \sqrt{-G} e^{\phi/2} \left[\frac{1}{4} F^{mn} \tau (D_n V_m - D_m V_n) - \frac{1}{2} \phi' F^{\rho m} \tau V_m \right], \tag{B.117}$$

$$\begin{aligned}
S^{(0)}(V_2, V_2) &= -\frac{1}{16} \int d^D X \sqrt{-G} e^\phi \left\{ e^{-2A} \partial_M V_{ij} \partial^M V_{ij} \right. \\
& - 2e^{-4A-2\phi} \left(e^{\phi+2A} V^n{}_m \right)_{;n} \left(e^{\phi+2A} V^{lm} \right)_{;l} \\
& - 4V^{mn} e^{-A} \partial_m \left[e^{-A-\phi} \left(e^{\phi+2A} V^l{}_n \right)_{;l} \right] \\
& \left. + \partial_\mu V_{mn} \partial^\mu V^{mn} + \frac{1}{3} V_{[nl;m]} V^{[nl;m]} + \frac{\kappa^2}{2} \gamma^2 e^{\phi/2} (V_{mn} F^{mn})^2 \right\}, \tag{B.118}
\end{aligned}$$

$$S^{(0)}(V, V_2) = -\frac{\kappa}{4} \gamma \int d^D X \sqrt{-G} e^\phi V_{mn} F^{mn} \left[\left(A' + \frac{1}{2} \phi' \right) V_\rho + D_l V^l \right]. \tag{B.119}$$

We have no mixing of the form $S^{(0)}(h, V_2)$ and $S^{(0)}(\tau, V_2)$ as a consequence of $H_{MNP} = 0$ (at the background level). We have checked that the term $S^{(0)}(V, V)$ reduces, as it should, to

the corresponding action in Ref. [9] in the case in which $V_{\underline{m}}$ is orthogonal to the background gauge field.

Let us consider now the branon-dependent action. This turns out to have the following form³⁵:

$$S^{(0)}(h, \xi) + S^{(0)}(V, \xi) + S^{(0)}(\tau, \xi) + S^{(0)}(\xi, \xi). \quad (\text{B.120})$$

Therefore, the fields $\xi^{\underline{m}}$ in general couple with some bulk fields, but these mixings are confined to the spin-0 action. The explicit expressions for the different pieces are

$$S^{(0)}(h, \xi) = -\frac{T}{2} \int d^4x \sqrt{-g} [2 \xi^{\underline{m}} (A' h_{\rho \underline{m}} + h_{\underline{m} \rho}) + e^{-A} \xi^{\underline{m}} \partial_{\underline{m}} h_{ii}], \quad (\text{B.121})$$

$$S^{(0)}(V, \xi) = T \kappa^2 \int d^4x \sqrt{-g} e^{\phi/2} F_{\underline{m} \underline{n}} V_{\underline{n}} \xi^{\underline{m}}, \quad (\text{B.122})$$

$$S^{(0)}(\tau, \xi) = \frac{T}{2} \int d^4x \sqrt{-g} \xi^{\underline{m}} \partial_{\underline{m}} \phi \tau, \quad (\text{B.123})$$

$$S^{(0)}(\xi, \xi) = -\frac{T}{2} \int d^4x \sqrt{-g} \left[G_{\underline{m} \underline{n}} \partial_{\underline{m}} \xi^{\underline{m}} \partial^{\underline{m}} \xi^{\underline{n}} + \frac{1}{2} \xi^{\underline{m}} \partial_{\underline{m}} g_{\mu \nu} \xi^{\underline{n}} \partial_{\underline{n}} g^{\mu \nu} \left(\frac{1}{2} g^{\mu \nu} g^{\rho \sigma} - g^{\mu \rho} g^{\nu \sigma} \right) \right. \\ \left. + \frac{1}{2} \xi^{\underline{m}} \xi^{\underline{n}} \partial_{\underline{m}} \partial_{\underline{n}} g_{\mu \nu} g^{\mu \nu} + T \kappa^2 \sqrt{g/G} \delta(Y_c - Y_c) \xi^{\underline{m}} \xi_{\underline{m}} \right]. \quad (\text{B.124})$$

We discuss the various singularities that can be observed in the above in Subsection 4.2.3 and below.

C Spin-0 Spectrum for 6D Supergravity Compactification

We finally analyse the (massive) spin-0 fluctuations in 6D braneworlds by using the general spin-0 action given in Appendix B. Here we discard the branons. There are different ways to make this truncation consistently, *e.g.* by introducing an orbifold that projects them out. In Ref. [9] such an orbifold has been defined taking into account the presence of at least two patches in the description of spherical topologies. Here we only use the fact that the orbifold action in the intersection of the two patches is $\varphi \rightarrow \varphi + \pi$. In the absence of the branons the $\delta(0)$ singularities mentioned in Subsection 4.2.3 obviously disappear. We shall see that it is also possible to deal with the other type of singularities mentioned there and extract a finite spectrum.

Here we focus on the unwarped solutions and in particular on the rugbyballs and saddlespheres defined in Subsection 2.3. In this case we will be able to generalize the harmonic analysis developed in Subsection 5.2.2 to the spin-0 sector, which involves 2D tensors as well as 2D vectors and scalars. This technique allows us to transform complicated coupled differential equations into algebraic equations whose solutions can be found exactly. The relevant fluctuations are h_{mn} , V_m , τ , V_{ij} and V_{mn} , where m and n run over θ and φ . We observe that the fluctuations V_m orthogonal to background gauge field decouple to the other fields and have already been analyzed in [9]; therefore here we assume V_m to be parallel to

³⁵The term of the form $S^{(0)}(V_2, \xi)$ vanishes as a consequence of $H_{MNP} = 0$ at the background level, which in turn follows from our background ansatz.

the background gauge field. One should keep in mind that, if the branons are projected out by the above-mentioned orbifold, only the modes with \mathbf{m} even survive (in the Fourier expansion over $e^{i\mathbf{m}\varphi}$). The spin-0 action in the light cone gauge assumes the following form:

$$\begin{aligned}
S^{(0)}(h, h) &= -\frac{1}{4\kappa^2} \int d^6 X \sqrt{-G} \left[\partial_\mu h_{mn} \partial^\mu h^{mn} + h_{mn;l} h^{mn;l} \right. \\
&\quad - 2h^m_l h^n_h R_{mn}{}^{lh} + \kappa^2 h_{lm} h^l_n F^m_h F^{nh} + \kappa^2 h^{mn} h^{lh} F_{lm} F_{hn} \\
&\quad \left. - \frac{1}{2} h_m{}^m (\partial^2 + D^2) h_n{}^n + \frac{T}{2} \kappa^2 (h_m{}^m)^2 \sqrt{g/G} \delta(X_2 - Y_2) \right], \\
S^{(0)}(V, V) &= -\frac{1}{2} \int d^6 X \sqrt{-G} [\partial_\mu V_m \partial^\mu V^m + V_{m;n} V^{m;n} \\
&\quad + \frac{1}{2} R V_m V^m + \frac{\kappa^2}{2} F^2 V_m V^m], \\
S^{(0)}(\tau, \tau) &= \frac{1}{4\kappa^2} \int d^6 X \sqrt{-G} \left[\tau \left(\partial^2 + D^2 - \frac{4}{r_0^2} \right) \tau \right], \\
S^{(0)}(V_2, V_2) &= \int d^6 X \sqrt{-G} \left[\frac{1}{16} V_{ij} (\partial^2 + D^2) V_{ij} + \frac{1}{16} V_{mn} (\partial^2 + D^2) V^{mn} \right. \\
&\quad \left. - \frac{\kappa^2 \gamma^2}{32} (V_{mn} F^{mn})^2 \right], \\
S^{(0)}(h, V) &= \int d^6 X \sqrt{-G} \left[-V_{n;m} h_l{}^n F^{lm} \right], \\
S^{(0)}(h, \tau) &= \frac{1}{8} \int d^6 X \sqrt{-G} F^2 \tau h_m{}^m, \\
S^{(0)}(V, \tau) &= \int d^6 X \sqrt{-G} \frac{\tau}{4} F^{mn} (V_{m;n} - V_{n;m}), \\
S^{(0)}(V, V_2) &= -\frac{\kappa}{4} \gamma \int d^6 X \sqrt{-G} V_{mn} F^{mn} V_l{}^{:l}, \tag{C.125}
\end{aligned}$$

where we have used the light cone gauge relation $h_i{}^i + h_m{}^m = 0$ in $S^{(0)}(h, h)$ and the property $F^{ml} F_l{}^n = G^{mn} F^2/2$ in $S^{(0)}(V, V)$ and $S^{(0)}(h, \tau)$, which is a consequence of (5.77).

We now want to use a technique similar to that explained in the spin-1 sector, in order to transform the above differential problem into an algebraic one. Note that the method provided in Subsection 5.2.2 can be already applied to perform this transformation in the terms $S^{(0)}(V, V)$, $S^{(0)}(\tau, \tau)$, $S^{(0)}(V_2, V_2)$, $S^{(0)}(V, \tau)$ and $S^{(0)}(V, V_2)$ as they only involve 2D scalars and 2D vectors³⁶. What we have done there is to identify appropriate mass-squared operators from the diagonal part of the bilinear action, which are Hermitian once the HCs are imposed. In this way we were able to define complete sets of 2D scalar and vector harmonics. Then we focused on the cases in which the derivative relation, Eq. (5.89), between scalar and vector harmonics holds. That relation is what allowed us to deal with the derivative couplings between scalars and vectors and transform the spin-1 differential problem into an algebraic one, which could easily be solved.

³⁶The analysis of the V_{mn} -EOMs shows that $V_{mn}/\sqrt{-G}$ is a 2D scalar.

Here we generalize the above procedure to include the 2D tensor fluctuations in h_{mn} . Indeed, h_{mn} can be decomposed into its trace, h_m^m , and traceless part, $\tilde{h}_{mn} \equiv h_{mn} - G_{mn}h_l^l/2$, so that the first entry in (C.125) decomposes into the two terms:

$$\begin{aligned} S^{(0)}(h_m^m, h_m^m) &= \frac{1}{8\kappa^2} \int d^6 X \sqrt{-G} \left[h_m^m \left(\partial^2 + D^2 - \frac{\kappa^2}{4} F^2 \right) h_n^n \right], \\ S^{(0)}(\tilde{h}_{mn}, \tilde{h}_{mn}) &= \frac{1}{4\kappa^2} \int d^6 X \sqrt{-G} \left[\tilde{h}_{mn} (\partial^2 + D^2 - R) \tilde{h}^{mn} \right], \end{aligned} \quad (\text{C.126})$$

where we used the following identities:

$$R_{pmqn} = \frac{R}{2}(G_{pq}G_{mn} - G_{mq}G_{pn}), \quad F_{lm}F_{hn} = \frac{F^2}{2}(G_{lh}G_{mn} - G_{mh}G_{ln}). \quad (\text{C.127})$$

Observe that h_m^m is in fact a 2D scalar field, and we can expand it in terms of the 2D scalar harmonics found in Subsection 5.1.2. The fluctuations \tilde{h}_{mn} are instead genuine 2D tensor fluctuations, and the appropriate mass-squared operator is $-D^2 + R$. Thus, we would like to solve the eigenproblem:

$$(-D^2 + R) \tilde{h}_{mn} = \mu_T^2 \tilde{h}_{mn}, \quad (\text{C.128})$$

with the given NCs and HCs, where μ_T^2 are the corresponding mass-eigenvalues.

In a general basis, Eq. (C.128) is a set of two coupled differential equations (the traceless property removes one out of the three components of a rank two symmetric tensor in two dimensions). However, by writing down Eq. (C.128) in the \pm basis defined in (5.86):

$$(-D^2 + R) h_{\pm\pm} = \mu_T^2 h_{\pm\pm}, \quad (\text{C.129})$$

where we used $h_{\pm\pm} = \tilde{h}_{\pm\pm}$, and by explicitly evaluating $D^2 h_{\pm\pm}$, one finds that the equations for h_{++} and h_{--} are decoupled, like those of the h_{+i} and h_{-i} fields in the spin-1 sector. After a long but straightforward calculation we find

$$-\partial_\theta^2 f^{\pm\pm} + \frac{\dot{B}}{2} \partial_\theta f^{\pm\pm} + \left(\mathbf{m}^2 e^{-B} \pm 2\mathbf{m} \dot{B} e^{-B/2} + \frac{\dot{B}^2}{2} - \frac{\ddot{B}}{2} \right) f^{\pm\pm} = \frac{r_0^2}{4} \mu_T^2 f^{\pm\pm}, \quad (\text{C.130})$$

where a dot represents a derivative with respect to θ and $f^{\pm\pm}$ is the wave function of $h_{\pm\pm}$, defined by a KK expansion

$$h_{\pm\pm}(X) = \sum_{\mathbf{n}, \mathbf{m}} h_{\pm\pm \mathbf{n}\mathbf{m}}(x) f_{\mathbf{n}\mathbf{m}}^{\pm\pm}(\theta) e^{i\mathbf{m}\varphi}, \quad \text{with } \mathbf{m} = \text{generic integer} \quad (\text{C.131})$$

and in (C.130) the KK numbers \mathbf{n} and \mathbf{m} are understood. The eigenvalues μ_T^2 can be found by using the technique discussed in Ref. [6]: one can put the equations into the hypergeometric form, consider the general solution to the hypergeometric equation and then impose the HCs and NCs. We find

- For $|\mathbf{m}| \omega \geq 2$

$$\mu_T^2 = \frac{4}{r_0^2} [(\mathbf{n} + |\mathbf{m}| \omega)(\mathbf{n} + |\mathbf{m}| \omega + 1) - 2] \quad (\text{C.132})$$

- For $-2 < \mathbf{m}\omega < 2$

$$\mu_T^2 = \frac{4}{r_0^2} [(\mathbf{n} + 2)(\mathbf{n} + 3) - 2] \quad (\text{C.133})$$

where $\mathbf{n} = 0, 1, 2, 3, \dots$. In this way we have found a complete set of 2D tensor harmonics.

We now remember that, in the spin-1 sector analysed in Subsection 5.2.2, one can generate the 2D vector harmonics by acting with derivatives over the 2D scalar harmonics (see Eq. (5.89) and the discussion right above). We can imagine that something similar happens here and the 2D tensor harmonics (C.129) can be generated by acting with derivatives over 2D vector harmonics. This is indeed the case and in order to see it let us consider the 2D vector harmonics for V_m :

$$\left(-D^2 + \frac{R}{2}\right) V_m = \mu_V^2 V_m, \quad (\text{C.134})$$

where μ_V^2 are the 2D vector mass-eigenvalues. From now on we shall assume Condition (5.88), so that $\mu_V^2 = \mu^2$, with μ^2 the 2D scalar mass-eigenvalues given in (5.85). After some manipulation it is easy to show that if V_m satisfies the previous equation then we also have

$$-D^2 \tilde{V}_{m;n} + R(\tilde{V}_{m;n} + \tilde{V}_{n;m}) + \frac{1}{2}(R_{;m}V_n + R_{;n}V_m - G_{mn}R_{;l}V^l) = \mu^2 \tilde{V}_{m;n}, \quad (\text{C.135})$$

where $\tilde{V}_{m;n} \equiv V_{m;n} - G_{mn}V_l{}^{;l}/2$. This equation is valid for any unwarped compactification, but in the rugbyball case it can be simplified. Although the Ricci scalar is not constant everywhere like in the sphere limit as it contains delta-functions, these additional delta function terms can be discarded in Eq. (C.135) because they are dominated by stronger singularities³⁷, which emerge from $D^2 \tilde{V}_{m;n}$. This allows us to write (C.135) as follows:

$$-D^2 \tilde{V}_{m;n} + R_s(\tilde{V}_{m;n} + \tilde{V}_{n;m}) = \mu^2 \tilde{V}_{m;n}, \quad (\text{C.136})$$

where R_s is the Ricci scalar of the sphere ($R_s = 8/r_0^2$), or, in the \pm basis,

$$(-D^2 + 2R_s) V_{\pm;\pm} = \mu^2 V_{\pm;\pm}, \quad (\text{C.137})$$

where we used $V_{\pm;\pm} = \tilde{V}_{\pm;\pm}$. Now, comparing the eigenproblems for $h_{\pm\pm}$ and $V_{\pm;\pm}$, Eqs. (C.129) and (C.137), we see that their eigenfunctions will belong to the same orthogonal set provided that:

$$\mu_T^2 = \mu^2 - R_s = \mu^2 - 8/r_0^2. \quad (\text{C.138})$$

By comparing the 2D vector mass-eigenvalues, μ^2 given in (5.85), with the 2D tensor eigenvalues, μ_T^2 given in Eqs. (C.132) and (C.133), we find that Condition (C.138) is indeed true in the following cases:

- For $\mathbf{m} = 0$ or $|\mathbf{m}| \geq 2/\omega$, which we denote by $0 \not\leq |\mathbf{m}|\omega \not\leq 2$, with the constraint $\{\mathbf{n}, \mathbf{m}\} \neq \{0, 0\}, \{1, 0\}$.
- For $|\mathbf{m}|\omega = 1$, with the constraint $\mathbf{n} \neq 0$.

³⁷This is a quite generic property of rugbyball compactifications [6, 9].

- The sphere case ($\omega = 1$), with the constraint $\{\mathbf{n}, \mathbf{m}\} \neq \{0, 0\}, \{1, 0\}, \{0, \pm 1\}$. This result is in agreement with that obtained by using the Wigner functions [31].

When (C.138) is true a derivative relation between the 2D tensor and 2D vector wave functions holds:

$$D_{\pm}(f_{\mathbf{nm}}^{\pm}(\theta)e^{i\mathbf{m}\varphi}) = c_{T\mathbf{nm}}f_{\mathbf{nm}}^{\pm\pm}(\theta)e^{i\mathbf{m}\varphi}, \quad (\text{C.139})$$

where $c_{T\mathbf{nm}}$ are normalization constants which, having chosen a convenient normalization for the wave functions, can be fixed to be $c_{T\mathbf{nm}} = \sqrt{\mu_{T\mathbf{nm}}^2}/\sqrt{2}$.

It remains to expand the 6D fields in the action (C.125) into their harmonics on the rugbyball and integrate over the extra dimensions. Thanks to the derivative relations (5.89, C.139), and $F^2 = \text{const}$, the mass-squared operator reduces to an algebraic matrix with constant entries, which can easily be diagonalized. We note that the mass-matrix turns out to be well-defined despite the singularities mentioned in Subsection 4.2.3.

We end with the resulting spectrum for spin-0 fields (which can be trusted when Eq. (C.138) holds). For definiteness we focus here on the 6D supergravity setup, but there are no problems in deriving the squared masses in the EYMA case as well. We split the spectrum according to the values of $l \equiv \mathbf{n} + |\mathbf{m}|$:

- For $l = 0$

$$\frac{r_0^2}{4}M^2 = 0, 0, 2, [2]$$

- For $l = 1$

$$\frac{r_0^2}{4}M^2 = 2, 6, [2], [2], [2], [6]$$

- For $l > 1$

$$\begin{aligned} M^2 &= \mu_{\mathbf{nm}}^2 \quad \text{with multiplicity } 1[+3] \\ M^2 &= \frac{4}{r_0^2} \left(1 + \frac{r_0^2}{4}\mu_{\mathbf{nm}}^2 - \sqrt{1 + r_0^2\mu_{\mathbf{nm}}^2} \right) \quad \text{with multiplicity } 1[+1] \\ M^2 &= \frac{4}{r_0^2} \left(1 + \frac{r_0^2}{4}\mu_{\mathbf{nm}}^2 + \sqrt{1 + r_0^2\mu_{\mathbf{nm}}^2} \right) \quad \text{with multiplicity } 1[+1] \end{aligned}$$

where the square brackets denote helicity-0 components of higher spin fields and the remaining modes are physical spin-0 fields. We observe that there are neither ghosts nor tachyons and we recover the two massless fields discussed in Subsection 5.4.2.

As an effective check of the above spectrum we observe that it correctly reduces, when $\omega = 1$, to the sphere result obtained by directly expanding the bulk fields over the Wigner functions [27].

References

- [1] L. Randall and R. Sundrum, “A large mass hierarchy from a small extra dimension,” *Phys. Rev. Lett.* **83** (1999) 3370 [arXiv:hep-ph/9905221].
- [2] L. Randall and R. Sundrum, “An alternative to compactification,” *Phys. Rev. Lett.* **83** (1999) 4690 [arXiv:hep-th/9906064].
- [3] H. Nishino and E. Sezgin, “Matter And Gauge Couplings Of N=2 Supergravity In Six-Dimensions,” *Phys. Lett. B* **144** (1984) 187.
- [4] G. W. Gibbons, R. Guven and C. N. Pope, “3-branes and uniqueness of the Salam-Sezgin vacuum,” *Phys. Lett. B* **595** (2004) 498 [arXiv:hep-th/0307238].
- [5] Y. Aghababaie *et al.*, “Warped brane worlds in six dimensional supergravity,” *JHEP* **0309** (2003) 037 [arXiv:hep-th/0308064].
- [6] S. L. Parameswaran, S. Randjbar-Daemi and A. Salvio, “Gauge fields, fermions and mass gaps in 6D brane worlds,” *Nucl. Phys. B* **767** (2007) 54 [arXiv:hep-th/0608074].
- [7] C. P. Burgess, C. de Rham, D. Hoover, D. Mason and A. J. Tolley, “Kicking the rugby ball: Perturbations of 6D gauged chiral supergravity,” *JCAP* **0702** (2007) 009 [arXiv:hep-th/0610078].
- [8] H. M. Lee and A. Papazoglou, “Gravitino in six-dimensional warped supergravity,” *Nucl. Phys. B* **792** (2008) 166 [arXiv:0705.1157 [hep-th]].
- [9] S. L. Parameswaran, S. Randjbar-Daemi and A. Salvio, “Stability and Negative Tensions in 6D Brane Worlds,” *JHEP* **0801** (2008) 051 [arXiv:0706.1893 [hep-th]].
- [10] C. P. Burgess, S. L. Parameswaran and I. Zavala, arXiv:0812.3902 [hep-th].
- [11] H. M. Lee and A. Papazoglou, “Scalar mode analysis of the warped Salam-Sezgin model,” *Nucl. Phys. B* **747** (2006) 294 [Erratum-ibid. *B* **765** (2007) 200] [arXiv:hep-th/0602208].
- [12] S. Randjbar-Daemi and M. Shaposhnikov, “A formalism to analyze the spectrum of brane world scenarios,” *Nucl. Phys. B* **645** (2002) 188 [arXiv:hep-th/0206016].
- [13] R. Sundrum, “Compactification for a three-brane universe,” *Phys. Rev. D* **59** (1999) 085010 [arXiv:hep-ph/9807348].
- [14] *See for e.g.:* M. Bando, T. Kugo, T. Noguchi and K. Yoshioka, “Brane fluctuation and suppression of Kaluza-Klein mode couplings,” *Phys. Rev. Lett.* **83** (1999) 3601 [arXiv:hep-ph/9906549]; N. Arkani-Hamed, S. Dimopoulos and G. R. Dvali, “Phenomenology, astrophysics and cosmology of theories with sub-millimeter dimensions and TeV scale quantum gravity,” *Phys. Rev. D* **59** (1999) 086004 [arXiv:hep-ph/9807344]; A. Dobado and A. L. Maroto, “The dynamics of the Goldstone bosons on the brane,” *Nucl. Phys. B* **592** (2001) 203 [arXiv:hep-ph/0007100];

- H. Murayama and J. D. Wells, “Graviton emission from a soft brane,” *Phys. Rev. D* **65** (2002) 056011 [arXiv:hep-ph/0109004].
- [15] R. Sundrum, “Effective field theory for a three-brane universe,” *Phys. Rev. D* **59** (1999) 085009 [arXiv:hep-ph/9805471].
- [16] S. Randjbar-Daemi, A. Salam, E. Sezgin and J. A. Strathdee, “An Anomaly Free Model In Six-Dimensions,” *Phys. Lett. B* **151** (1985) 351.
- [17] S. D. Avramis, A. Kehagias and S. Randjbar-Daemi, “A new anomaly-free gauged supergravity in six dimensions,” *JHEP* **0505** (2005) 057 [arXiv:hep-th/0504033]; S. D. Avramis and A. Kehagias, “A systematic search for anomaly-free supergravities in six dimensions,” *JHEP* **0510** (2005) 052 [arXiv:hep-th/0508172]; R. Suzuki and Y. Tachikawa, “More anomaly-free models of six-dimensional gauged supergravity,” *J. Math. Phys.* **47**, 062302 (2006) [arXiv:hep-th/0512019].
- [18] H. Nishino and E. Sezgin, “The Complete N=2, D = 6 Supergravity With Matter And Yang-Mills Couplings,” *Nucl. Phys. B* **278** (1986) 353.
- [19] G. W. Gibbons and S. W. Hawking, “Action Integrals And Partition Functions In Quantum Gravity,” *Phys. Rev. D* **15** (1977) 2752; H. A. Chamblin and H. S. Reall, “Dynamic dilatonic domain walls,” *Nucl. Phys. B* **562** (1999) 133 [arXiv:hep-th/9903225].
- [20] Z. Lalak and R. Matyszkiewicz, “Boundary terms in brane worlds,” *JHEP* **0111**, 027 (2001) [arXiv:hep-th/0110141].
- [21] S. M. Carroll and M. M. Guica, “Sidestepping the cosmological constant with football-shaped extra dimensions,” arXiv:hep-th/0302067; I. Navarro, “Codimension two compactifications and the cosmological constant problem,” *JCAP* **0309**, 004 (2003) [arXiv:hep-th/0302129].
- [22] R. Sundrum, “Compactification for a three-brane universe,” *Phys. Rev. D* **59** (1999) 085010 [arXiv:hep-ph/9807348]; J. W. Chen, M. A. Luty and E. Ponton, “A critical cosmological constant from millimeter extra dimensions,” *JHEP* **0009** (2000) 012 [arXiv:hep-th/0003067].
- [23] M. Kaku, “Ghost-Free Formulation of Quantum Gravity in the Light-Cone Gauge,” *Nucl. Phys. B* **91** (1983) 99.
- [24] C. A. Lutken and C. R. Ordonez, “Vacuum Energy Of Eleven-Dimensional Supergravity,” *Class. Quant. Grav.* **4** (1987) 1543.
- [25] S. Randjbar-Daemi, A. Salam and J. A. Strathdee, “Towards A Selfconsistent Computation Of Vacuum Energy In Eleven-Dimensional Supergravity,” *Nuovo Cim. B* **84** (1984) 167.
- [26] S. Randjbar-Daemi and M. H. Sarmadi, “Graviton Induced Compactification In The Light Cone Gauge,” *Phys. Lett. B* **151** (1985) 343.

- [27] *For explicit computations in the spherical case, see the Appendix C of the PhD thesis:* A. Salvio, “Aspects of physics with two extra dimensions,” arXiv:hep-th/0701020.
- [28] E. A. Mirabelli and M. E. Peskin, “Transmission of supersymmetry breaking from a 4-dimensional boundary,” Phys. Rev. D **58**, 065002 (1998) [arXiv:hep-th/9712214]; S. Ichinose and A. Murayama, “The delta(0) singularity in the warped Mirabelli-Peskin model,” arXiv:hep-th/0606167.
- [29] *See for e.g.:* W. D. Goldberger and M. B. Wise, “Bulk fields in the Randall-Sundrum compactification scenario,” Phys. Rev. D **60**, 107505 (1999) [arXiv:hep-ph/9907218]; C. Csaki, M. Graesser, L. Randall and J. Terning, “Cosmology of brane models with radion stabilization,” Phys. Rev. D **62** (2000) 045015 [arXiv:hep-ph/9911406]. C. Charmousis, R. Gregory and V. A. Rubakov, “Wave function of the radion in a brane world,” Phys. Rev. D **62** (2000) 067505 [arXiv:hep-th/9912160].
- [30] H. Nicolai and C. Wetterich, “On The Spectrum Of Kaluza-Klein Theories With Noncompact Internal Spaces,” Phys. Lett. B **150** (1985) 347; G. W. Gibbons and D. L. Wiltshire, “Space-Time As A Membrane In Higher Dimensions,” Nucl. Phys. B **287**, 717 (1987) [arXiv:hep-th/0109093]; A. Kehagias, “On non-compact compactifications with brane worlds,” arXiv:hep-th/9911134.
- [31] S. Randjbar-Daemi, A. Salam and J. A. Strathdee, “Spontaneous Compactification In Six-Dimensional Einstein-Maxwell Theory,” Nucl. Phys. B **214** (1983) 491.
- [32] A. Salam and E. Sezgin, “Chiral Compactification On Minkowski $\times S^{*2}$ Of $N=2$ Einstein-Maxwell Supergravity In Six-Dimensions,” Phys. Lett. B **147** (1984) 47.
- [33] S. Randjbar-Daemi, A. Salvio and M. Shaposhnikov, “On the decoupling of heavy modes in KK theories,” Nucl. Phys. B **741** (2006) 236 [arXiv:hep-th/0601066].
- [34] S. Randjbar-Daemi, A. Salam and J. A. Strathdee, “Instability Of Higher Dimensional Yang-Mills Systems,” Phys. Lett. B **124** (1983) 345 [Erratum-ibid. B **144** (1984) 455]; G. R. Dvali, S. Randjbar-Daemi and R. Tabbash, “The origin of spontaneous symmetry breaking in theories with large extra dimensions,” Phys. Rev. D **65** (2002) 064021 [arXiv:hep-ph/0102307].
- [35] G. W. Gibbons and C. N. Pope, Nucl. Phys. B **697** (2004) 225 [arXiv:hep-th/0307052].
- [36] M. L. Graesser, J. E. Kile and P. Wang, “Gravitational perturbations of a six dimensional self-tuning model,” Phys. Rev. D **70**, 024008 (2004) [arXiv:hep-th/0403074]; M. Peloso, L. Sorbo and G. Tasinato, “Standard 4d gravity on a brane in six dimensional flux compactifications,” Phys. Rev. D **73**, 104025 (2006) [arXiv:hep-th/0603026].