

DESY 18-121

KW 18-008

MBnumerics: Numerical integration of Mellin-Barnes integrals in physical regions

Johann Usovitsch*

*Hamilton Mathematics Institute, Trinity College Dublin,
College Green, Dublin 2, Ireland*
E-mail: usovitsj@maths.tcd.ie

Ievgen Dubovyk

II. Institut für Theoretische Physik, Universität Hamburg, 22761 Hamburg, Germany
DESY Deutsches Elektronen-Synchrotron, 15738 Zeuthen, Germany
E-mail: e.a.dubovyk@gmail.com

Tord Riemann

University of Silesia, 40-007 Katowice, Poland
DESY Deutsches Elektronen-Synchrotron, 15738 Zeuthen, Germany
E-mail: tordriemann@gmail.com

We introduce techniques to treat numerically Mellin-Barnes integrals in physical regions, which arise in the need of the computation of Feynman integrals for the electroweak two-loop corrections to the pseudo observables at the Z-boson resonance.

Loops and Legs in Quantum Field Theory (LL2018)

29 April 2018 - 04 May 2018

St. Goar, Germany

*Speaker.

1. Introduction

Our starting point will be the loop-momenta integral representation of a scalar Feynman integral:

$$G_L = \int \prod_{j=1}^L \frac{d^D k_j}{i\pi^{D/2}} \frac{1}{P_1^{v_1} \dots P_N^{v_N}}. \quad (1.1)$$

The functions $P_i^{v_i}$ in the denominator are expressed in terms of the L loop-momenta k_l which are not fixed through momentum conservation at each vertex and the E linearly independent external momenta p_e :

$$P_i = \left(\sum_{l=1}^L a_{il} k_l + \sum_{e=1}^E b_{ie} p_e \right)^2 - m_i^2 + i\delta, \quad a_{il}, b_{ie} \in \{-1, 0, 1\}, \quad (1.2)$$

where the m_i denote the masses of the corresponding virtual particles. The $i\delta$ is the Feynman prescription. In the most general case the P_i are a linear combination of N linearly independent scalar products depending on the loop-momenta k_l . The propagator exponents v_i are complex variables if not stated otherwise. Within dimensional regularization, $D = 4 - 2\varepsilon$ denotes the dimension of space-time. As usual $D \neq 4$ is used to regularize infrared or ultraviolet divergences.

Before evaluating these integrals one often applies the Feynman trick:

$$\frac{(-1)^{\mathbf{v}}}{\prod_{j=1}^N (-P_j)^{v_j}} = \frac{(-1)^{\mathbf{v}} \Gamma(\mathbf{v}) \left(\prod_{j=1}^N \tilde{n}_j \right) \delta(1 - \sum_{j=1}^{N_G} x_j)}{(-k_l^\mu M_{ll'} k_{l'\mu} + 2k_l^\mu Q_{l\mu} + J - i\delta)^{\mathbf{v}}}, \quad \mathbf{v} = \sum_{j=1}^N v_j, \quad (1.3)$$

where

$$M_{ll'} = \sum_{j=1}^N a_{jl} a_{j l'} x_j \quad (1.4)$$

is an $L \times L$ symmetric matrix,

$$Q_l^\nu = - \sum_{j=1}^N x_j a_{jl} \sum_{e=1}^E b_{je} p_e^\nu \quad (1.5)$$

is a vector with L components and

$$J = - \sum_{j=1}^N x_j \left(\sum_{e=1}^E b_{je} p_e^\mu \sum_{e'=1}^E p_{e'}^\nu b_{j e'} g_{\mu\nu} - m_j^2 \right), \quad (1.6)$$

where x_j are the Feynman parameters introduced with the Feynman trick. The set of Feynman parameters $\{x_1, \dots, x_{N_G}\}$ corresponds to the set of functions $\{P_1, \dots, P_{N_G}\}$ with positive $\{v_1, \dots, v_{N_G}\}$ in Eq.(1.1). The metric tensor is $g_{\mu\nu} = \text{diag}(1, -1, \dots, -1)$. The \tilde{n}_j is defined as:

$$\tilde{n}_j \phi(\vec{x}) = \begin{cases} \int_{\{x_j \geq 0\}} \frac{dx_j x_j^{v_j-1}}{\Gamma(v_j)} \phi(\vec{x}), & v_j \neq -m, \\ (-1)^{v_j} \phi^{(-v_j)}(0, x_{i \neq j}), & v_j = -m, \end{cases} \quad m \in \mathbb{N}_0, \quad (1.7)$$

where $\phi^{(-v_j)}(0, x_{i \neq j})$ means to take $(-v_j)$ derivative in x_j and then set x_j to zero.

The Feynman integral can now be written in the Feynman parameter integral representation:

$$G_L = (-1)^{\nu} \Gamma(\nu - LD/2) \left(\prod_{j=1}^N \tilde{n}_j \right) \delta\left(1 - \sum_{j=1}^{N_G} x_j\right) \frac{\mathcal{U}(x)^{\nu - (L+1)D/2}}{\mathcal{F}(x)^{\nu - LD/2}}, \quad (1.8)$$

where

$$\mathcal{U}(x) = \det M, \quad (1.9)$$

$$\mathcal{F}(x) = \mathcal{U}(x) (Q_i^{\mu} M_{\mu\nu}^{-1} Q_{\nu}^{\mu} + J - i\delta). \quad (1.10)$$

From these definitions it follows that the functions $\mathcal{F}(x)$ and $\mathcal{U}(x)$ are homogeneous in the Feynman parameters x_i . The function $\mathcal{U}(x)$ is of degree L and the function $\mathcal{F}(x)$ is of degree $L+1$. The functions $\mathcal{U}(x)$ and $\mathcal{F}(x)$ are also known as Symanzik polynomials.

1.1 Mellin-Barnes integral

Feynman integrals may be infrared and ultraviolet divergent. To treat these integrals in a consistent and automated way two methods are known: The Mellin-Barnes integral approach [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11] and the sector decomposition approach [12, 13, 14, 15, 16, 17].

To derive a Mellin-Barnes integral one will use either the loop-by-loop approach [7] or the global approach [9]. Both techniques apply in their core to the $\mathcal{F}(x)$ and $\mathcal{U}(x)$ functions in Eq. (1.8) the Mellin-Barnes integral master formula:

$$\frac{1}{(a+b)^{\nu}} = \int_{-i\infty}^{i\infty} \frac{dz}{2\pi i} \frac{a^z b^{-z-\nu} \Gamma(-z) \Gamma(\nu+z)}{\Gamma(\nu)}, \quad |\arg a - \arg b| < \pi, \quad (1.11)$$

until the integrations over the Feynman parameters can be all carried out in terms of Euler's Beta-functions:

$$B(\xi, \chi) = \int_0^{\infty} \frac{x^{\xi-1}}{(1+x)^{\xi+\chi}} dx = \frac{\Gamma(\xi) \Gamma(\chi)}{\Gamma(\xi+\chi)}, \quad \Re \xi > 0, \Re \chi > 0. \quad (1.12)$$

These steps lead to Mellin-Barnes integrands depending on a ratio of Euler's Gamma-functions Γ depending on the integration variables z_i and some kinematics raised to the powers of z_i .

As an example we study the Feynman integral in Fig. 1, whose loop-momenta integral representation contains one nontrivial numerator $k_1 p_1$:

$$I_{0h0w14r} = \int \frac{d^D k_1}{i\pi^{D/2}} \frac{d^D k_2}{i\pi^{D/2}} \frac{k_1 p_1}{k_1^2 ((k_1 - k_2)^2 - M_Z^2) k_2^2 ((k_2 + p_1)^2 - M_Z^2) (k_1 + p_1 + p_2)^2}, \quad (1.13)$$

and its Mellin-Barnes integral representation is:

$$I_{0h0w14r} = \int_{-\frac{1}{3}-i\infty}^{-\frac{1}{3}+i\infty} \frac{dz_1}{2\pi i} \int_{-\frac{1}{3}-i\infty}^{-\frac{1}{3}+i\infty} \frac{dz_2}{2\pi i} \frac{\Gamma(-z_1) \Gamma(-z_2) \Gamma(z_2+1) \Gamma(-\varepsilon-z_1) \Gamma(2\varepsilon+z_1+1) \left(-\frac{M_Z^2}{s}\right)^{z_1}}{2\Gamma(1-z_2) \Gamma(-3\varepsilon-z_1+2) \Gamma(-2\varepsilon-2z_1-z_2)} \times (-s)^{-2\varepsilon} \Gamma(-2\varepsilon-z_1-z_2)^2 \Gamma(-\varepsilon-z_1-z_2) \Gamma(\varepsilon+z_1+z_2+1), \quad (1.14)$$

where $\varepsilon = (4-D)/2$. A straight integration contour parallel to the imaginary axis is chosen, such




Fig. 1: Two-loop vertex Feynman integral with two internal massive lines and the kinematics are $p_{1,2}^2 = 0$ and $2p_1 p_2 = s$. The Z -boson mass M_Z indicates massive propagators.




Fig. 2: The black dots are the poles of the integrand in Eq. (2.1) in the z_1 complex plane. The dashed line is the integration contour parallel to the imaginary axis.

that all poles are well separated, see Fig. 2 for $z_{1,2} = -1/3 + it_{1,2}$, $t_{1,2} \in [-\infty, \infty]$. The expansion around $\varepsilon = 0$ in Eq. (1.14) leads to a finite contribution in the lowest order:

$$I_{0h0w14r} = - \int_{-\frac{1}{3}-i\infty}^{-\frac{1}{3}+i\infty} \frac{dz_1}{2\pi i} \int_{-\frac{1}{3}-i\infty}^{-\frac{1}{3}+i\infty} \frac{dz_2}{2\pi i} \frac{\Gamma(-z_1)^2 \Gamma(z_1+1) \Gamma(-z_2) \Gamma(z_2+1) \left(-\frac{M_Z^2}{s}\right)^{z_1} \Gamma(-z_1-z_2)^3}{2\Gamma(2-z_1) \Gamma(1-z_2) \Gamma(-2z_1-z_2)} \times \Gamma(z_1+z_2+1) + \mathcal{O}(\varepsilon). \quad (1.15)$$

1.2 Minkowskian kinematics

Whether we derive the Mellin-Barnes integrals with the loop-by-loop or the global approach, we face problems in the numerical treatment of these integrals in Minkowskian kinematics. To illustrate this we apply the well known Stirling approximation formula

$$\Gamma(z) \underset{|z| \rightarrow \infty}{\approx} z^{z-1/2} e^{-z} \sqrt{2\pi}, \quad |\arg z| < \pi, \quad (1.16)$$

to the integrand in Eq. (1.15) and examine the asymptotic behavior for $z_1 = -\frac{1}{3} + it_1$, and $z_2 = -\frac{1}{3} + it_2$, $t_1 \rightarrow -t$ and $t_2 \rightarrow t$:

$$\mathcal{I}_{0h0w14r} \underset{t \rightarrow \infty}{\approx} t^{-2+2x_1+2x_2} \Big|_{x_1=x_2=-1/3}. \quad (1.17)$$

In comparison to the Euclidean kinematics, where the asymptotic behavior is everywhere exponentially damped, we see that for Minkowskian kinematics the asymptotic behavior is polynomial. In the case of a Mellin-Barnes integral this polynomial asymptotic behavior leads to numerous numerical instabilities, some of which are:

- Oscillations are less damped compared to the Euclidean case.
- Integrals may be not absolutely convergent if the asymptotic behavior is worse than $1/t^a$, with $a < 2$.
- At any level of accuracy, we need to evaluate the integrands for bigger values t_i than in the case of Euclidean kinematics.
- In particular, if we are interested in high accuracy results, we have to evaluate the Γ functions for very big arguments and this leads again to numerical instabilities.

2. Techniques to treat Mellin-Barnes integrals in Minkowskian kinematics

We assume that the treatment of one-dimensional Mellin-Barnes integrals is a solved problem by means of contour deformation [18, 19, 20, 10]. We describe techniques which are applied to multi-dimensional Mellin-Barnes integrals. These techniques are automatized in the Mathematica package MBnumerics, which was developed to treat numerically Feynman integrals appearing in the calculation of the electroweak two-loop corrections to the pseudo observables at the Z-boson resonance [21], [22].

2.1 Linear transformation of integration variables

In the case of the Mellin-Barnes integral the linear transformation of integration variables may lead to improvements of the numerical integration. If we apply the variable change $z_2 \rightarrow z_2 - z_1$ to the example integral in Eq. (1.15) we get

$$I_{0h0w14} = - \int_{-\frac{1}{3}-i\infty}^{-\frac{1}{3}+i\infty} \frac{dz_1}{2\pi i} \int_{-\frac{2}{3}-i\infty}^{-\frac{2}{3}+i\infty} \frac{dz_2}{2\pi i} \frac{(-\frac{M_2^2}{s+i\delta})^{z_1} \Gamma(-z_1)^2 \Gamma(1+z_1) \Gamma(z_1-z_2) \Gamma(-z_2)^3}{2\Gamma(2-z_1) \Gamma(-z_1-z_2) \Gamma(1+z_1-z_2)} \times \Gamma(1+z_2) \Gamma(1-z_1+z_2). \quad (2.1)$$

After this simple change of variable the asymptotic behavior of the Mellin-Barnes integrand has been changed. If we apply again the Stirling formula in Eq. (1.16) to the integrand in Eq. (2.1), and study the asymptotic behavior for $z_1 = -\frac{1}{3} + it_1$, $z_2 = -\frac{2}{3} + it_2$, $t_1 \rightarrow -t$ and $t_2 \rightarrow 0$, we find

$$\mathcal{I}_{0h0w14} \underset{t \rightarrow \infty}{\approx} t^{-2+2x_2} \Big|_{x_2=-2/3}, \quad (2.2)$$

i.e. the polynomial asymptotic behavior depends only on x_2 . Linear integration variable transformations give a possibility for a nontrivial cross check of the numerical evaluation of the Mellin-Barnes integrals, since the integrands have different asymptotic behavior before and after the linear transformation.

2.2 Integrand mappings

An obvious improvement is the application of the cotangent mapping $t = \frac{1}{\tan(-\pi d)}$, which maps the integration boundaries from $t \in [-\infty, \infty]$ to $d \in [0, 1]$. We apply this mapping to a polynomial function, which gives

$$\frac{1}{t^a} = \tan(-\pi d)^a, \quad (2.3)$$

and the Jacobian is:

$$\frac{\pi}{\sin(\pi d)^2}, \quad (2.4)$$

where the limits of the integrand at the boundaries of the new integration domain are:

$$\lim_{d \rightarrow 0, d \rightarrow 1} \frac{\pi \tan(-\pi d)^a}{\sin(\pi d)^2} = \begin{cases} \frac{1}{0}, & a < 2, \\ \pi, & a = 2, \\ 0, & a > 2. \end{cases} \quad (2.5)$$

Compared to the cotangent mapping, a logarithmic mapping, as it is advocated in the program MB.m [5], always leads to infinities at the new integration boundaries, which would lead to numerical instabilities.

Since we use the cotangent mapping it is mandatory to transform the integrand as follows:

$$\prod_i \Gamma_i \rightarrow \exp\left(\sum_i \log \Gamma_i\right), \quad (2.6)$$

where the key idea is that the $\log \Gamma(z_i)$ functions grow slower than the $\Gamma(z_i)$ functions for large values of $|z_i|$.

2.3 Shifts

If we shift the Mellin-Barnes integration variables according to

$$z_i = x_i + it_i + n_i, \quad n_i \in \mathbb{R}, \quad (2.7)$$

the asymptotic behavior of a given Mellin-Barnes integrand may depend explicitly on the shifts n_i :

$$\mathcal{J}_{0h0w14} \underset{t \rightarrow \infty}{\approx} t^{-2+2x_2+2n_2} \Big|_{x_2=-2/3}. \quad (2.8)$$

It is then possible to improve the polynomial asymptotic behavior by tuning the shifts n_i . If, by changing the values of n_i , the contour crosses some poles of the Mellin-Barnes integrand, we have to collect their residues.

The shifts may also be used as a method to evaluate Mellin-Barnes integrals in Minkowskian regions due to one more observation: the integral along a shifted contour may be numerically smaller by orders of magnitude compared to the original integral.

For example, the original integral in Eq. (2.1), with $M_Z/\sqrt{s} = 1 - i\delta$, evaluated along the contour C_1 , see Fig. 3, gives $0.3923828588857 + 0.7456388536613i$. We chose $\delta = 10^{-16}$. The shifted




Fig. 3: The original contour C_1 is shifted by $n_1 = -2$ to a contour C_2 . The third contour C_3 encircles the poles to correct the shift.

integral with $n_1 = -2$, evaluated along the contour C_2 gives -0.00974965823202 . In addition the following equation holds:

$$\int \frac{dz_2}{2\pi i} \int_{C_1} \frac{dz_1}{2\pi i} \mathcal{I}_{0h0w14} = \int \frac{dz_2}{2\pi i} \int_{C_2} \frac{dz_1}{2\pi i} \mathcal{I}_{0h0w14} + \overbrace{\int \frac{dz_2}{2\pi i} \left(\sum_{z_0} \text{Res}_{z_0} \mathcal{I}_{0h0w14} \right)}^{\text{1 dim integrals}}. \quad (2.9)$$

We have added to the result of the shifted integral the contributions from the residues of the three poles enclosed by the contour C_3 . Upon integrating them over z_2 , their sum is $0.402132517117807 + 0.745638853661318i$. In general, shifting the contour of an n -fold Mellin-Barnes integral will yield residue terms, which will be $(n - 1)$ -fold Mellin-Barnes integrals and hence simpler to evaluate.

3. Nontrivial example




Fig. 4: This Feynman integral depends non-trivially on the scales s , M_W and m_t .

Method	Numerics	
MB	$0.0602664865576999 \varepsilon^{-2}$	
SD - 90 Mio	$0.06026648655 \varepsilon^{-2}$	
MB	$(-0.0315124890$	$+0.1893327514i) \varepsilon^{-1}$
SD - 90 Mio	$(-0.031512481$	$+0.189332716i) \varepsilon^{-1}$
MB	$(-0.22823186755$	$-0.08824794573i) + \mathcal{O}(\varepsilon)$
SD - 90 Mio	$(-0.2282265$	$-0.0882459i) + \mathcal{O}(\varepsilon)$

Tab. 1: The numbers labeled MB are evaluated with MBnumerics. The numbers labelled with SD are evaluated with SecDec v.3 [23].

The Feynman integral shown in Fig. 4,

$$\int \frac{d^D k_1}{i\pi^{D/2}} \frac{d^D k_2}{i\pi^{D/2}} \frac{\exp(2\varepsilon\gamma_E) M_Z^{2+2\varepsilon} (k_2 p_2)}{(k_1)^2 ((k_1 - k_2)^2 - m_t^2) ((k_2)^2 - M_W^2) ((k_1 + p_1)^2) ((k_2 + p_1)^2 - m_t^2) (k_1 + p_1 + p_2)^2}, \quad (3.1)$$

depends in a nontrivial way on the scalar product $k_2 p_2$ in the numerator. We evaluate this integral with MBnumerics, which implements the method of the shifts. The results are collected in Tab. 1. The scales are fixed to: $s = M_Z^2 + i\delta$, $M_Z = 91.1876$ GeV, $M_W = 80.385$ GeV and $m_t = 173.2$ GeV. In this example the Mellin-Barnes integral representation is at most a three-dimensional integral. With the sector decomposition approach the Feynman integral (3.1) is five-dimensional. If one can find a Mellin-Barnes integral representation whose integration dimension is smaller than or equal to that of the sector decomposition representation, the method of shifts turns out to be very successful to compute numerically Feynman integrals in Minkowskian regions.

Acknowledgments

We would like to thank Ayres Freitas and Janusz Gluza for fruitful discussions. We enjoyed the opportunity to complete together with them the calculation of the electroweak two-loop corrections to the Z-boson resonance physics. The work of I.D. is supported by a research grant of Deutscher Akademischer Austauschdienst (DAAD) and is supported partly by the Polish National Science Centre (NCN) under the Grant Agreement 2017/25/B/ST2/01987. The work of T.R. is supported in part by an Alexander von Humboldt Polish Honorary Research Fellowship. This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme under grant agreement No 647356 (CutLoops). This work is supported by Graduiertenkolleg 1504 "Masse, Spektrum, Symmetrie" of Deutsche Forschungsgemeinschaft (DFG).

References

- [1] V. A. Smirnov, *Analytical result for dimensionally regularized massless on shell double box*, *Phys. Lett.* **B460** (1999) 397 [[hep-ph/9905323](#)].
- [2] J. B. Tausk, *Nonplanar massless two loop Feynman diagrams with four on-shell legs*, *Phys. Lett.* **B469** (1999) 225 [[hep-ph/9909506](#)].

- [3] G. Heinrich and V. A. Smirnov, *Analytical evaluation of dimensionally regularized massive on-shell double boxes*, *Phys. Lett.* **B598** (2004) 55 [hep-ph/0406053].
- [4] M. Czakon, J. Gluza and T. Riemann, *Master integrals for massive two-loop bhabha scattering in QED*, *Phys. Rev.* **D71** (2005) 073009 [hep-ph/0412164].
- [5] M. Czakon, *Automatized analytic continuation of Mellin-Barnes integrals*, *Comput. Phys. Commun.* **175** (2006) 559 [hep-ph/0511200].
- [6] C. Anastasiou and A. Daleo, *Numerical evaluation of loop integrals*, *JHEP* **10** (2006) 031 [hep-ph/0511176].
- [7] J. Gluza, K. Kajda and T. Riemann, *AMBRE: A Mathematica package for the construction of Mellin-Barnes representations for Feynman integrals*, *Comput. Phys. Commun.* **177** (2007) 879 [0704.2423].
- [8] J. Gluza, K. Kajda, T. Riemann and V. Yundin, *Numerical Evaluation of Tensor Feynman Integrals in Euclidean Kinematics*, *Eur. Phys. J.* **C71** (2011) 1516 [1010.1667].
- [9] I. Dubovyk, J. Gluza and T. Riemann, *Non-planar Feynman diagrams and Mellin-Barnes representations with AMBRE 3.0*, *J. Phys. Conf. Ser.* **608** (2015) 012070.
- [10] I. Dubovyk, J. Gluza, T. Riemann and J. Usovitsch, *Numerical integration of massive two-loop Mellin-Barnes integrals in Minkowskian regions*, *PoS* **LL2016** (2016) 034 [1607.07538].
- [11] M. Prausa, *Mellin-Barnes meets Method of Brackets: a novel approach to Mellin-Barnes representations of Feynman integrals*, *Eur. Phys. J.* **C77** (2017) 594 [1706.09852].
- [12] K. Hepp, *Proof of the Bogolyubov-Parasiuk theorem on renormalization*, *Commun. Math. Phys.* **2** (1966) 301.
- [13] T. Binoth and G. Heinrich, *An automatized algorithm to compute infrared divergent multiloop integrals*, *Nucl. Phys.* **B585** (2000) 741 [hep-ph/0004013].
- [14] T. Binoth and G. Heinrich, *Numerical evaluation of multiloop integrals by sector decomposition*, *Nucl. Phys.* **B680** (2004) 375 [hep-ph/0305234].
- [15] T. Binoth and G. Heinrich, *Numerical evaluation of phase space integrals by sector decomposition*, *Nucl. Phys.* **B693** (2004) 134 [hep-ph/0402265].
- [16] A. Denner and S. Pozzorini, *An Algorithm for the high-energy expansion of multi-loop diagrams to next-to-leading logarithmic accuracy*, *Nucl. Phys.* **B717** (2005) 48 [hep-ph/0408068].
- [17] G. Heinrich, *Sector Decomposition*, *Int. J. Mod. Phys.* **A23** (2008) 1457 [0803.4177].
- [18] A. Freitas and Yi-Cheng Huang, *On the Numerical Evaluation of Loop Integrals With Mellin-Barnes Representations*, *JHEP* **04** (2010) 074 [1001.3243].
- [19] J. Gluza, T. Jelinski and D. A. Kosower, *Efficient Evaluation of Massive Mellin-Barnes Integrals*, *Phys. Rev.* **D95** (2017) 076016 [1609.09111].
- [20] Z. Peng, *Topics in $N = 4$ Yang-Mills theory*, Ph.D. thesis, IPHT, Saclay, 2012. Submitted on 14 Jun 2013. HAL Id: tel-00834200. HAL is a multi-disciplinary open access archive. <https://tel.archives-ouvertes.fr/tel-00834200>.
- [21] I. Dubovyk, A. Freitas, J. Gluza, T. Riemann and J. Usovitsch, *The two-loop electroweak bosonic corrections to $\sin^2 \theta_{\text{eff}}^b$* , *Phys. Lett.* **B762** (2016) 184 [1607.08375].

-
- [22] I. Dubovyk, A. Freitas, J. Gluza, T. Riemann and J. Usovitsch, *Complete electroweak two-loop corrections to Z boson production and decay*, *Phys. Lett.* **B783** (2018) 86 [1804.10236].
- [23] S. Borowka, G. Heinrich, S. P. Jones, M. Kerner, J. Schlenk and T. Zirke, *SecDec-3.0: numerical evaluation of multi-scale integrals beyond one loop*, *Comput. Phys. Commun.* **196** (2015) 470 [1502.06595].