

Thraxions: Ultralight Throat Axions

Arthur Hebecker,¹ Sascha Leonhardt,¹ Jakob Moritz,² Alexander Westphal²

¹ *Institute for Theoretical Physics, University of Heidelberg, Philosophenweg 19,
69120 Heidelberg, Germany*

² *Deutsches Elektronen-Synchrotron, DESY, Notkestraße 85,
22607 Hamburg, Germany*

(a.hebecker, s.leonhardt @thphys.uni-heidelberg.de
jakob.moritz, alexander.westphal @desy.de)

May 3, 2019

Abstract

We argue that a new type of extremely light axion is generically present in the type IIB part of the string theory landscape. Its mass is suppressed by the third power of the warp factor of a strongly warped region (Klebanov-Strassler throat), suggesting the name *thraxion*. Our observation is based on the generic presence of several throats sharing the same 2-cycle. This cycle shrinks to zero volume at the end of each throat. It is hence trivial in homology and the corresponding C_2 axion is massive. However, the mass is warping-suppressed since, if one were to cut off the strongly warped regions, a proper 2-cycle would re-emerge. Since the kinetic term of the axion is dominated in the UV, an even stronger, quadratic mass suppression results. Moreover, if the axion is excited, the angular modes of the throats backreact. This gives our effective C_2 axion a finite monodromy and flattens its potential even further. Eventually, the mass turns out to scale as the third power of the warp factor. We briefly discuss possible implications for phenomenology and potential violations of the Weak Gravity Conjecture for axions. Moreover we identify a mechanism for generating super-Planckian axionic field ranges which we call *drifting monodromies*. However, in the examples we consider, the potential oscillates on sub-Planckian distances in field space, preventing us from building a natural inflation model on the basis of this idea.

Contents

1	Introduction	3
2	Backreacted Potential of the Thraxion from 10d	6
2.1	Geometric and Flux-Background	6
2.1.1	Geometric Features of Generic CYs	6
2.1.2	Moduli Stabilization by Fluxes	9
2.2	Local Backreaction in the Throat	10
2.3	The CY Breaking Potential	12
2.4	Discussion of Results	15
2.5	The B_2 -axion	17
3	Four-Dimensional SUGRA Completion	19
3.1	Counting Moduli Through the Conifold Transition	19
3.2	The Thraxion Superpotential	20
3.2.1	The GVW Superpotential of a Multi Conifold System	20
3.2.2	The Thraxion as a Stabilizer Field	22
3.2.3	The Double Throat: $n = 2, m = 1$	25
3.2.4	The General Multi Throat	27
3.3	Comments on the b -Axion	28
4	The Axion Potential and the Gauge/Gravity Correspondence	29
5	Applications	31
5.1	Thraxions on the Quintic: Drifting Monodromy	31
5.2	A Clash with the Weak Gravity Conjecture	34
5.3	The Spectrum of Effective Instantons	35
5.4	Axion Phenomenology	36
5.5	Uplifting	37
6	Conclusion and Discussion	40
A	Deformed Conifold for General Complex Structure Modulus	43
B	The Axion Potential in the Local Throat	46

C	Derivation and Solution of the 5d Equations of Motion	48
C.1	The 5d Action for φ	48
C.2	Schrödinger Equations and Exact Solutions for Free Fields	49
D	Axion Decay Constant and Potential Parameters	52
E	Background on Multi Conifolds	55
	References	58

1 Introduction

Axion-like particles (axions for short) have become a main player in beyond-the-standard-model (BSM) physics in general [1, 2] and in string phenomenology in particular [3–7]. They are relevant e.g. in the context of the strong CP problem, inflation and as dark matter candidates. Furthermore, in the recent debate surrounding the Landscape/Swampland program [8, 9] and the Weak Gravity Conjecture (WGC) [10], axions have occupied a prominent place [11–19]. This is also related to the question whether or with which parameters axion monodromy [20, 21] can be realized in consistent quantum gravity settings [22–27].

In this paper, we present a novel type of ultralight axion which, as we argue, is generically present in the type IIB part of the landscape, building on a proposal made in [28]. Its extreme lightness, both in absolute terms and in relation to its decay constant (i.e. compared to the scale $M_{\text{P}}^4 \exp(-M_{\text{P}}/f)$ of generic non-perturbative potentials) lets it stand out among the many other stringy axions. It is surprising that, to the best of our knowledge, this very generic axionic degree of freedom was missed at the time when the string axiverse was being intensely studied.

Before turning to the details, we want to explain our central and, in our opinion, rather surprising, parametric results: Consider type IIB Calabi-Yau orientifold or F-theory models stabilized by fluxes and non-perturbative effects [29–31]. It is generally accepted that Klebanov-Strassler (KS) throats [32] with warp factor $w_{\text{IR}} \ll 1$ will be present in an order-one fraction of such models [33–35]. This warp factor can easily be *exponentially* small, such that it is justified to focus for the moment only on the dependence on w_{IR} . In other words, let us for now set $R_{\text{CY}} \sim M_{\text{string}}^{-1}$ and $N_{\text{flux}} \sim \mathcal{O}(1)$, thus ignoring all parameters except for the warp factor. Naively, the lightest states are then the glueballs (or warped-throat KK modes) with mass $\sim w_{\text{IR}}$ (in Planck units). By contrast, we claim

that an ultralight axion with mass $\sim w_{\text{IR}}^3$ is frequently present in such settings. To be more precise, this happens at least in all cases where the fluxes stabilize the complex structure moduli near a conifold transition point in moduli space.

Moreover, our axion has a decay constant $f \sim \mathcal{O}(1)$ in the simplest models¹, which can be enhanced by products of flux numbers to super-Planckian values in more general settings, and an effective potential which can be much smaller than the naive expectation $V \sim \exp(-1/f) \cos(\phi/f)$ (again in Planck units). Clearly, this has potentially many interesting applications, from the WGC for axions to inflation and uplifting.

The paper is organized as follows. We start with the background solution in Sect. 2.1. We consider a Calabi-Yau with a conifold point in complex structure moduli space at which multiple three-cycles degenerate simultaneously. We explain why this is a generic feature of Calabi-Yaus. Concentrating on the case of two degenerate three-cycles, we introduce separate deformation parameters z_i with phases $\varphi_i = \arg z_i$, $i = 1, 2$, for the two deformed conifold regions. Crucially, the two conifolds, specifically the S^3 -cycles describing the apices, are related in homology. As a result, the Calabi-Yau condition ensures that only one complex structure modulus, $z = z_1 = z_2$, is present. Deformations with $z_1 \neq z_2$ are massive. We then introduce fluxes stabilizing the complex structure modulus z near the conifold point $|z| \ll 1$. The resulting geometry is illustrated in Fig. 1. One can see that the so-called \mathcal{B} -cycle is an S^3 which can be thought of as a family of S^2 's. This S^2 family reaches into both throats such that the S^2 's collapse at the apices. The corresponding dual \mathcal{A} -cycle is an S^3 over every point of the double throat in Fig. 1.

In Sect. 2.2, we introduce the axion² $c \sim \int_{S^2} C_2$, called *thraxion* from now on. An excursion of the thraxion generates non-zero opposite values of the RR-field strength F_3 at the ends of the two throats. Local backreaction of the resulting energy density then deforms the two throats independently: While the phase φ_1 of the local deformation parameter of one throat is displaced by fluxes, the phase φ_2 of the other throat is displaced in the opposite direction by anti-fluxes. This breaks the constraint $\varphi_1 = \varphi_2$ coming from the CY condition and the homology relation between the two throats. In Sect. 2.3, we calculate the potential induced by non-vanishing 10d Ricci curvature that stabilizes the two deformation

¹Note that despite the fact that strongly warped throats are needed to generate a small scalar potential for the axion, the decay constant is not suppressed by warping effects. This is because its internal field-profile is *not* localized at the bottom of the throats, in contrast to some examples that have appeared in the literature [36, 37].

²This may seem like a misnomer since the shift symmetry is completely broken: The 10d gauge invariance $C_2 \sim C_2 + d\Lambda_1$ does not imply a 4d shift symmetry because the two-sphere is trivial in homology $S^2 = \partial\Sigma_3$, $\int_{S^2} d\Lambda_1 = \int_{\Sigma_3} d^2\Lambda_1 = 0$. However, from the unwarped UV-perspective the two-sphere is non-trivial and the field is a proper axion, with a monodromy created only by the warped-down IR region.




Figure 1: An illustration of the setup of the double throat including the phases φ_i and the axion c . The phases φ_i describe physical rotations of each throat. We have not drawn the S^3 over every point of the double throat.

parameters against each other. After integrating out heavy degrees of freedom, the result is an effective potential for the thraxion with the properties described above and discussed in Sect. 2.4.

Sect. 3 rederives the effective axion potential from a proposed generalization of the Gukov-Vafa-Witten (GVW) superpotential [38] that includes the axion. We thereby reproduce the results in 4d supergravity language, and identify the *saxion* partner of the axion. In Sect. 3.2.4 we generalize our results to general multi throat systems where one or more ultra-light thraxions appear.

We discuss the consistency of our results with the holographic dictionary in the KS context [39, 32] in Sect. 4 by matching the enhancement of the decay constant of our axion $\int C_2$ with gaugino condensation on the gauge theory side. Applications and implications of these results are the content of Sect. 5. We consider as an explicit example the quintic three-fold stabilized near a conifold transition point. We study the scalar potential for judicious choices of flux quanta. Interestingly, the overall monodromy enhancement is given by the least common multiple of all the flux quanta which can easily become parametrically super-Planckian. However, the presence of sub-Planckian modulations generically prevents successful slow-roll inflation. The underlying idea is that a large monodromy is generated by unsynchronized phases (of monodromies of individual throats) drifting away from one another. We call this mechanism *drifting monodromies*. This mechanism can also be thought of as the well-known beat phenomenon in acoustics, in which the interference of harmonics with slightly different small wavelengths leads to

large wavelength oscillations. For related alternative possibilities of generating large decay constants see [40, 41, 16, 42–48]³. We also describe a clash with the WGC: The effective Euclidean instanton action determined from the scale of the effective potential violates the axionic WGC ($S \lesssim qM_{\text{P}}/f_{\text{eff}}$) parametrically, but instead a slightly weaker inequality holds ($S \lesssim (qM_{\text{P}}/f_{\text{eff}})^2$). After commenting on the relevance of our findings to light axion phenomenology, we finally consider interesting possibilities for uplifting to de Sitter vacua. We draw our conclusions in Sect. 6.

2 Backreacted Potential of the Thraxion from 10d

2.1 Geometric and Flux-Background

2.1.1 Geometric Features of Generic CYs

First we will explain the basic geometric requirements for our discussion to apply. We will explain why we expect them to be *generically* met.

Let us consider compactifications of type IIB string theory on a Calabi-Yau (CY) threefold, which leads to an effective $\mathcal{N} = 2$ supergravity theory in four dimensions. There is a moduli space of vacua parametrized by the $h^{2,1}$ complex structure moduli and $h^{1,1}$ Kähler moduli. There are special points in complex structure moduli space called conifold points where the CY develops conical singularities, and one or more three-cycles degenerate to zero volume [49, 50]. We will consider a CY near such a conifold point, where *multiple* three-cycles degenerate.

To understand in what sense this is generic, we consider cases in which it is also possible to *resolve* conifold singularities while preserving the CY condition. This does not restore the degenerate three-cycles to finite size, but rather produces non-trivial two-cycles. In going through this so-called *conifold transition*, a new CY threefold with Hodge numbers $\tilde{h}^{1,1} > h^{1,1}$ and $\tilde{h}^{2,1} < h^{2,1}$ is produced [51]. Whenever two CY threefolds are connected via a conifold transition, at the conifold transition point two or more three-cycles \mathcal{A}_i that are related in homology shrink to zero size. This is a consequence of demanding Kählerity on the resolved side of the transition [52]. It is widely believed that a *generic* CY threefold is

³Note in particular the following two papers: The work of [45] is closely related to ours in making use of the conifold complex structure modulus z to create super-Planckian decay constants, while on the technical level the approach is very different. Ref. [46] defines the 5d axion $\int B_2$ on the KT background, analogously to our thraxion. There, the geometric backreaction via the 5d breathing mode allows for monodromy-induced super-Planckian field ranges to be explored in an anisotropic and inhomogeneous 5d spacetime.

related to other CY threefolds via conifold transitions [53,51]. While research on this is still on-going, there are large classes of CY's for which this has been shown [54–59]⁴. Therefore, a generic CY has loci in complex structure moduli space where multiple three-cycles \mathcal{A}_i degenerate together. We expand on this in Sect. 3. Being related in homology, the number of homology classes is smaller than the number of collapsing three-cycles. For now we focus on the case of precisely two cycles $\mathcal{A}_{1,2}$ that degenerate. From the above it immediately follows that they are related in homology $[\mathcal{A}] \equiv [\mathcal{A}_1] = [\mathcal{A}_2]$. There is a symplectic dual three-cycle \mathcal{B} connecting the two singular points. We will call this system a *double conifold*. Its complex structure will be denoted by z and the double conifold singularity develops in the limit $|z| \rightarrow 0$.

We introduce the fields z_1 and z_2 as illustrated in Fig. 1. These fields may be thought of as ‘local complex structure deformations’ $z_i = \int_{\mathcal{A}_i} \Omega$, with the holomorphic three-form Ω of the CY, and describe independent *local* deformations of the manifold near one of the two apices. Thus, in the vicinity of either apex of the double conifold we want to describe the manifold by embedding it into \mathbb{C}^4 via

$$w_1^2 + w_2^2 + w_3^2 + w_4^2 = z_i, \quad w \in \mathbb{C}^4. \quad (1)$$

While the homology relation $[\mathcal{A}_1] = [\mathcal{A}_2]$ enforces $z = z_1 = z_2$ on complex structure moduli space⁵, we will also consider deformations of the manifold such that $z_1 \neq z_2$, i.e. deformations away from complex structure moduli space⁶ (i.e. $d\Omega \neq 0$). It is important to note that a deformation of the local phases φ_i of z_i becomes an isometry far away from the tip of either conifold: The cross-section of either deformed conifold at a given radial coordinate r possesses a spontaneously broken $U(1)_R$ isometry⁷ realized as a rotation of the local deformation parameter z_i [61], see Fig. 2. In the limit of large radial coordinates, $r^3/|z_i| \rightarrow \infty$, the deformed conifold becomes indistinguishable from the singular conifold. Therefore, the symmetry remains unbroken in this limit. We make this explicit in App. A.

To complete the discussion of the geometric setting, we note that one may go beyond

⁴Note that this does not mean that every conifold singularity (or even a generic one) is also such a transition point. For example, the mirror quintic threefold at vanishing complex structure has a single shrunken three-cycle. Hence there is no resolved CY geometry [52].

⁵This is because the difference $\mathcal{A}_1 - \mathcal{A}_2$ is the boundary of a 4-chain \mathcal{C} . Therefore, one has $z_1 - z_2 = \int_{\mathcal{A}_1} \Omega - \int_{\mathcal{A}_2} \Omega = \int_{\partial\mathcal{C}} \Omega = \int_{\mathcal{C}} d\Omega$. On complex structure moduli space one has $d\Omega = 0$, and hence $z_1 = z_2$.

⁶For similar considerations with deformations away from Kähler moduli space, i.e. $dJ \neq 0$, see ref. [60].

⁷The index R is due to this symmetry group being the R-symmetry of the dual supersymmetric gauge theory. While this is of no importance to us, we keep it for notational consistency.




Figure 2: The double conifold with asymptotic $U(1)_R$ symmetric regions.

the simplest case with exactly two collapsing three-cycles. Such multi conifold situations are analyzed in Sect. 3. Furthermore, for reasons of tadpole cancellation we are interested in CY threefolds which are orientifolded such that O3/O7-planes arise. This projection should leave the conifold transition intact and preserve the key ingredient of a \mathcal{B} -cycle reaching down into several conifold regions. In the double throat case, this is realized if two originally present pairs of throats are mapped to each other by the orientifold projection, see Fig. 3. This is completely analogous to the widely-discussed double throat system of the oldest axion-monodromy models, see e.g. [21, 62] (just simpler, since we need no 2-cycle for the NS5 brane and can hence use standard KS throats). More generally, F-theory solutions with the analogous geometric properties can be considered. Here the tadpole cancellation relies on the fourfold Euler number and no orientifolding is required. Either way, we do not expect that the orientifolding condition or the fourfold embedding endangers the generality of our setting.




Figure 3: A sketch of the the orientifold projection σ . It maps the two originally independent double throat cycles \mathcal{B} and \mathcal{B}' onto one another.

2.1.2 Moduli Stabilization by Fluxes

As was shown in [29], generic choices of three-form flux quanta stabilize the axio-dilaton as well as all the complex structure moduli. Whenever the flux quanta $K \equiv -\frac{1}{(2\pi)^2\alpha'} \int_{\mathcal{B}} H_3$ and $M \equiv \frac{1}{(2\pi)^2\alpha'} \int_{\mathcal{A}_1} F_3 = \frac{1}{(2\pi)^2\alpha'} \int_{\mathcal{A}_2} F_3$ satisfy⁸ $K \gg g_s M$ (where g_s is the string coupling), our complex structure modulus $z = |z| e^{i\varphi}$ is stabilized near the conifold point

$$|z| \propto \exp\left(-2\pi \frac{K}{g_s M}\right) \ll 1. \quad (2)$$

Separating the equation $D_z W = 0$ into real and imaginary parts (see Sect. 3 of [29]), we also find that the phase is stabilized. Its value is set by the RR-3-form flux $Q = \frac{1}{(2\pi)^2\alpha'} \int_{\mathcal{B}} F_3$

$$\varphi = 2\pi \frac{Q}{M}. \quad (3)$$

Locally, we can always set φ to 0 by an appropriate redefinition of the angle. Conversely, without loss of generality, we will choose $Q = 0$.

Moreover, backreaction of fluxes leads to the formation of warped throats (or Klebanov-Strassler throats). Within these, the metric is well approximated by the Klebanov-Tseytlin solution⁹ [39]

$$ds^2 = w(r)^2 \eta_{\mu\nu} dx^\mu dx^\nu + w(r)^{-2} (dr^2 + r^2 ds_{T^{1,1}}^2), \quad w(r)^2 \sim \frac{r^2}{g_s M \alpha'} \log(r/r_{\text{IR}})^{-\frac{1}{2}}, \quad (4)$$

with radial coordinate r and warp factor $w(r)$. The radial coordinate is cut off in the IR by the Klebanov-Strassler region and in the UV by the gluing into the bulk CY. One has $w_{\text{IR}} \equiv w(r_{\text{IR}}) \sim r_{\text{IR}}/r_{\text{UV}} \sim |z|^{1/3}$ giving rise to an exponential hierarchy à la Randall-Sundrum [64, 32, 29]. As we have explained, in the vicinity of a conifold transition point a *double throat* (or even *multi throat*) forms¹⁰, see Fig. 1.

The three-cycle \mathcal{B} can be thought of as an S^2 fibered over the radial direction of the

⁸It has recently been claimed that achieving this hierarchy may be impossible [63]. We think that the assumptions of [63] are too restrictive (as is pointed out by the authors themselves). In particular, O7-planes with curvature contribute to the D3 tadpole and huge Euler numbers are achievable in F-theory. We expect that this will allow to make the tadpole large enough for our purposes.

⁹Near the bottom of the throat, it has to be replaced by the full Klebanov-Strassler solution [32].

¹⁰It may not seem obvious that the units of NS-flux on the \mathcal{B} -cycle are split democratically so that *each* conifold region is replaced by a warped throat. In fact we will see that there is a light dynamical field that controls this relative distribution (see Sect. 3). In the vacuum however this field is stabilized such that fluxes are indeed distributed democratically.

conifold [50]. The S^2 collapses at the two *tips* of the deformed conifolds. As discussed in [28], there exists a 4d mode $c(x)$ on the double throat background that can be thought of as the integral of the Ramond-Ramond (RR) two form C_2 over the S^2 as measured far away from the tips of the double throat. A non-trivial field excursion leads to the creation of regions with flux on the two respective ends of the cycle \mathcal{B} , and hence a red-shifted potential $V(c) = \frac{1}{2}m^2c^2 + \dots$, with $m^2 \sim w_{\text{IR}}^4$. Backreaction of the geometry was neglected in [28].

In the rest of this section, we will establish the following points:

- The fields z_1 and z_2 of the two respective throats adjust to the flux/anti-flux pair in such a way that within the two throats supersymmetry is restored locally.
- This adjustment of z_1 and z_2 takes us away from the complex structure moduli space, which is characterized by $z_1 = z_2$ (cf. Fig. 4). For $z_1 \neq z_2$, the CY condition is broken and a scalar potential is generated. This potential is of the order $|z|^2 \sim w_{\text{IR}}^6$ and receives its dominant contributions from the bulk CY.
- The backreacted scalar potential is periodic in c with periodicity $2\pi M$. Hence, the naive 2π periodicity of the c -axion is enhanced by a *finite* factor M . While this does *not* allow for a super-Planckian effective axion decay constant $f \gg M_{\text{P}}$, approximately Planckian values are possible (see however Sect. 5 for a way to generate large axion periodicities).




Figure 4: Illustration of the z_1 - z_2 deformation space. The complex structure moduli space is the subspace $z_1 = z_2$. We only consider deformations away from $z_1 = z_2$ outside the conifold point.

2.2 Local Backreaction in the Throat

We start by discussing how a single throat reacts locally to a finite field excursion c . Since the outcome will be that the throat almost perfectly adjusts to produce a locally

supersymmetric configuration, we are entitled to use the 4d description in terms of the GVW superpotential [38, 29] for two KS throats. As far as (say) the first local throat is concerned, a non-vanishing field excursion c cannot be distinguished from additional flux $P \equiv c/2\pi$ on the local portion of the \mathcal{B} -cycle¹¹, see Fig. 5,

$$c = \frac{1}{2\pi\alpha'} \int_{S^2} C_2 = \frac{1}{2\pi\alpha'} \int_{\frac{1}{2}\mathcal{B}} dC_2 = (2\pi)P. \quad (5)$$

Considering a single throat with complex structure modulus $z_1 \equiv |z_1|e^{i\varphi_1}$, the arguments




Figure 5: Fluxes induced by non-vanishing c are localized at the tips of the throats.

of GKP [29] show that there are SUSY configurations for

$$\varphi_1 = 2\pi \frac{P}{M} = c/M, \quad (6)$$

compare (3). Hence, the throat can locally relax the SUSY breaking induced by the extra RR-flux by adjusting the phase of the deformation parameter z_1 . However, the second throat sees the field excursion c as the flux $-P = -c/2\pi$ on the \mathcal{B} -cycle for which there exists a locally supersymmetric configuration with $\varphi_2 = -c/M$.

Since there is no additional flux that would lead to $|z_1| \neq |z_2|$, we keep $|z| = |z_1| = |z_2|$ fixed at the stabilized value (3) for what follows. These two modes decouple from the discussion at hand.

We can encode the discussion above in a 4d EFT potential. To quadratic order, the

¹¹We will use the term flux also for the non-quantized integral $\int F_3$ over some region.

discrepancy between the local fluxes and local deformations induces a potential

$$V_{\text{flux}}(c, \varphi_1, \varphi_2) = \frac{1}{2}\mu^4(M\varphi_1 - c)^2 + \frac{1}{2}\mu^4(M\varphi_2 + c)^2, \quad (7)$$

We have $\mu \sim w_{\text{IR}}$ since the potential is generated locally near the tip of the throats.

The fact that only the combinations $M\varphi_i \pm c$ appear in the scalar potential can be derived also via ten-dimensional considerations. As shown in App. B, in the local throats, the combined transformation $\varphi_{1,2} \rightarrow \varphi_{1,2} \pm \delta$, $c \rightarrow c + M\delta$ is a diffeomorphism acting on the KS solution. Hence, only the invariant combinations $M\varphi_{1,2} \mp c$ can appear in the scalar potentials that are generated locally at the bottom of the throats.

The potential derived so far possesses a flat direction which we parametrize by c . This flat direction is given by

$$\varphi_1 = -\varphi_2 = c/M. \quad (8)$$

2.3 The CY Breaking Potential

In the preceding section we have argued that the individual throats react to the field excursion c by adjusting their local deformation parameters z_1 and z_2 , more specifically their phases φ_1 and φ_2 respectively. Since the corresponding CY has only one complex structure modulus $z \equiv z_1 \equiv z_2$, the mode z_1/z_2 or rather $\varphi_1 - \varphi_2$ must be massive already before fluxes are turned on. This eliminates the remaining flat direction in the potential.

We parametrize the part of the scalar potential that is due to the breaking of the CY condition as

$$V_{\text{CY-breaking}} = \Lambda^4(1 - \cos(\varphi_1 - \varphi_2)), \quad (9)$$

with a yet undetermined scale Λ . In writing this we have assumed the following:

- (a) The potential is a function of the difference $\varphi_1 - \varphi_2$ only.
- (b) It satisfies $V_{\text{CY-breaking}}(\varphi_1 - \varphi_2) = V_{\text{CY-breaking}}(\varphi_1 - \varphi_2 + 2\pi)$.
- (c) The lowest harmonic dominates.

Condition a) must hold because only the local fluxes of the throats stabilize $\varphi_{1,2}$ individually and, without the flux potential, the complex structure modulus $\varphi = (\varphi_1 + \varphi_2)/2$ should be a flat direction. We expect condition b) to hold because we see no reason for a monodromy. Condition c) is a rather unimportant assumption that we make for ease of exposition.

We combine (7) and (9) and integrate out $\varphi_{1,2}$ under the assumption $\Lambda^4 \ll \mu^4$ (to be justified below). This corresponds to imposing (8). The effective potential takes the form

$$V_{\text{eff}}(c) = \Lambda^4 (1 - \cos(2c/M) + \mathcal{O}(\Lambda^4/\mu^4)) . \quad (10)$$

The height of this potential can be estimated using the 10d solution. To do so, we need to develop a clear picture of how field profiles and 10d geometry change if we excite c . Recall that c is originally defined by a particular ‘Wilson line’ VEV of C_2 in the UV of the two throats (as well as in the piece of the CY connecting them). Turning on this VEV and focussing on one throat only, we observe a backreaction of the throat geometry which maintains SUSY and corresponds to the motion along a flat direction in 4d field space. This is independently true for the second throat, which backreacts in the opposite way: $\varphi_1 = -\varphi_2 = c/M$.

Now, the crucial point is that these rotations, defined by the IR parameters $\varphi_{1,2}$, must by continuity be accompanied by a corresponding r -dependent rotational profile along the whole throat. We encode this in a five-dimensional field $\phi(x^\mu, r)$ that interpolates between $\varphi_1 = c/M$ and $\varphi_2 = -c/M$ at the respective ends of the throats. This is illustrated in Fig. 6, which also displays the expected symmetry: The solution should be antisymmetric under the exchange of the two throats.¹²




Figure 6: The expected profile of the 10d/5d mode along the radial direction.

For computational simplicity, we model the transition region between the throats by a single point, $r = r_{\text{UV}}$ ¹³. In doing so, we ignore effects of the unwarped CY region

¹²By a slight abuse of notation we stick with the familiar variable r , although according to our figure this variable must now be growing as one goes down the second throat.

¹³In fact, the exact UV geometry and UV fluxes are irrelevant as long as we do not consider perturbative

(accepting an $\mathcal{O}(1)$ error). The field ϕ must be zero at this point for symmetry reasons. This symmetry also ensures that we may limit our attention to one of the two throats when computing the energy density associated with an excursion of c .

The key point is that, after these preliminaries, we are actually able to estimate this energy. It is given by the gradient energy of ϕ , which accounts precisely for the clash between the opposite rotations of φ_1 and φ_2 . The relevant action for $\phi = \phi(x^\mu, r)$ is obtained by dimensionally reducing the 10d Ricci scalar to quadratic order on the warped conifold background (see App. C):

$$\begin{aligned} S[\phi] &= \frac{M_{10d}^8}{2} \int d^4x \int_{r_{\text{IR}}}^{r_{\text{UV}}} dr \sqrt{-g_{5d}} r^5 w(r)^{-5} \epsilon(r)^2 \left(-\frac{1}{2} g_{5d}^{MN} \partial_M \phi \partial_N \phi \right) \\ &= \frac{M_{10d}^8}{2} |z|^2 \int d^4x \int_{r_{\text{IR}}}^{r_{\text{UV}}} \frac{dr}{r} \left(-\frac{1}{2} (\partial_r \phi)^2 - \frac{1}{2} w(r)^{-4} (\partial_\mu \phi)^2 \right). \end{aligned} \quad (11)$$

Considering the metric (4), this form of the 5d action is easily understood. The metric naturally splits into a 5d part g_{5d} in the external and radial direction and an angular part $\propto g_{T^{1,1}}$. The latter contributes the r -dependent terms $\sqrt{g_{T^{1,1}}} \propto r^5 w(r)^{-5}$ to the metric determinant. The function $\epsilon(r)$ encodes the degree of $U(1)_R$ symmetry breaking, compare Fig. 2. Since a field excursion ϕ is obtained by acting with a $U(1)_R$ transformation, any terms in the action that contain the field must be multiplied by the factor $\epsilon(r)^2$. This symmetry breaking is due to the deformation near the tip and takes the form $\epsilon(r) \sim |z|/r^3$ far from the tip of the throat, see App. A.

We now apply the static approximation (i.e. disregard the $(\partial_\mu \phi)^2$ in (11)), derive the equation of motion and solve it for the boundary conditions $\phi(x^\mu, r_{\text{UV}}) = 0$ and $\phi(x^\mu, r_{\text{IR}}) = \varphi_1 = c/M$. This gives

$$\phi(x^\mu, r) = c/M \frac{r_{\text{UV}}^2 - r^2}{r_{\text{UV}}^2 - r_{\text{IR}}^2} \quad (12)$$

which, inserting back in (11), leads to a 4d potential $V \sim |z|^2 c^2$. This is a result at quadratic order in c but, comparing to (10), this is sufficient to infer that $\Lambda^4 \sim |z|^2$. Finally inserting the stabilized value $|z| \propto w_{\text{IR}}^3$ [29], we arrive at $\Lambda^4 \sim w_{\text{IR}}^6$. Our assumption $\mu^4 \gg \Lambda^4$ is now *a posteriori* justified. It is also apparent that the effective mass of our ultra-light field is $m_c \sim w_{\text{IR}}^3$.

Let us justify the use of the static approximation above. Really, we should have

and non-perturbative corrections [65].

supplemented (11) by the kinetic term $S_{\text{kin}}[c] \sim \int d^4x (\partial_\mu c)^2$, imposed the constraint $\phi(x^\mu, r_{\text{UV}}) = c(x^\mu)/M$, and determined the mass of the lowest-lying KK mode of the resulting 5d action. However, it is intuitively clear that the UV-dominated kinetic term of c is much more important than the warped-down 4d-gradient term $(\partial_\mu \phi)^2$ in (11). Thus, c is the most inert part of the system and it is an excellent approximation to assume that the ϕ profile extremizes just the 5d-gradient-part of the action. To make this quantitative, one may substitute c on the r.h. side of (12) with the plane wave $c = \exp(ikx)$ (with $k^2 = -m_c^2$) and check that the resulting $(\partial_\mu \phi)^2$ contribution from (11) is negligible compared to $S_{\text{kin}}[c]$.¹⁴

More details are given in App. C.

2.4 Discussion of Results

The information we have gathered can be summarized in an effective Lagrangian¹⁵,

$$\begin{aligned} \mathcal{L} = & -\frac{1}{2}f_\varphi^2(\partial\varphi_1)^2 - \frac{1}{2}f_\varphi^2(\partial\varphi_2)^2 - \frac{1}{2}f_c^2(\partial c)^2 \\ & - \frac{1}{2}\mu^4(M\varphi_1 - c)^2 - \frac{1}{2}\mu^4(M\varphi_2 + c)^2 - \Lambda^4(1 - \cos(\varphi_1 - \varphi_2)), \end{aligned} \quad (13)$$

with coefficients

$$\begin{aligned} f_\varphi^2 & \sim \log(w_{\text{IR}}^{-1})^{-3/2} w_{\text{IR}}^2 M_{\text{P}}^2, & f_c^2 & \approx \frac{2}{9M^2} \log(w_{\text{IR}}^{-1})^{-1} M_{\text{P}}^2, \\ \Lambda^4 & \sim \frac{g_s^2}{(g_s M)^4} \log(w_{\text{IR}}^{-1})^{-7/2} w_{\text{IR}}^6 M_{\text{P}}^4, & \mu^4 & \sim \frac{g_s^4}{(g_s M)^6} \log(w_{\text{IR}}^{-1})^{-3} w_{\text{IR}}^4 M_{\text{P}}^4. \end{aligned} \quad (14)$$

The above expressions are valid for the special case of the bulk CY having a single characteristic length-scale, $R_{\text{CY}}^6 \sim \text{Vol}(\text{CY})$, and where the throats marginally fit into the bulk, i.e. $R_{\text{CY}}^4 \sim R_{\text{throat}}^4 \sim g_s M K \alpha'^2$. For the general case see App. D. We briefly pause to explain what is meant by the requirement that the throats fit into the bulk marginally: On the one hand the bulk CY has an overall size R_{CY} that is set by a combination of the Kähler moduli. On the other hand the throats can really be thought of as objects of a characteristic physical size R_{throat} embedded into the bulk CY. This size R_{throat}^4 is well

¹⁴Note that there is no contribution from $S_{\text{pot}}[c] \sim \int d^4x dr (\partial_r c)^2$. When exciting c in the UV, the local deformation parameter adjusts as explained above. The profile $c(r)$ is now stabilized in turn, for energies $\Lambda < E < \mu$, to the radially constant value of the phase of the deformation parameter $\pm M\varphi_i$. We expand on this in App. B.

¹⁵For ease of exposition we have written down a diagonal kinetic matrix. This is not quite the case but is not relevant for our discussion. See App. B for details.

known to be set by the local D3 brane charge stored in the fluxes of the throat [32, 29], independently of the size of the bulk CY. For this to be a geometrically consistent configuration, we should require $R_{\text{CY}} > R_{\text{throat}}$. Taking $R_{\text{CY}} \sim R_{\text{throat}}$ is the case where the throats fit into the bulk CY only *marginally*.¹⁶

Far below the scale $w_{\text{IR}} M_{\text{P}}$ we may integrate out $\varphi_{1,2}$, to obtain the effective Lagrangian

$$\mathcal{L}' = -\frac{1}{2} f_c^2 (\partial c)^2 - \Lambda^4 (1 - \cos(2c/M)). \quad (15)$$

We would like to highlight the following points,

- Our simplification $R_{\text{throat}} = R_{S^2} = R_{\text{CY}}$ gives the largest possible value for the decay constant $f_{\text{eff}} = M f_c$; any hierarchy $R_{\text{throat}} < R_{S^2} < R_{\text{CY}}$ suppresses its value. Taking into account logarithmic corrections, the maximal periodicity one can achieve is $\mathcal{O}(M_{\text{P}}/\sqrt{\log w_{\text{IR}}^{-1}})$ (see App. D). A large hierarchy $w_{\text{IR}} \ll 1$ suppresses the periodicity only very mildly. By taking $g_s M$ and g_s^{-1} to be large, the 10d perturbative expansion becomes better controlled without affecting the axion periodicity. In this sense, our axion can be made approximately Planckian.
- The mass of the axion is $\mathcal{O}(w_{\text{IR}}^3)$ which is parametrically smaller than both the warped Kaluza-Klein scale ($\mathcal{O}(w_{\text{IR}})$), and the estimate of [28], where backreaction of the local geometry was not taken into account ($\mathcal{O}(w_{\text{IR}}^2)$). The mass-spectrum is essentially *gapped*.
- As pointed out before, the scale of the effective potential is set by the $U(1)_R$ breaking induced by the deformation of the conifold as measured in the UV, $\epsilon^2(r_{\text{UV}}) \propto |z|^2$. Strictly speaking this is *not* a warp factor suppression, although for moderate CY volumes $|z|$ and w_{IR}^3 are of the same order¹⁷.

The following caveats should be noted: The effective Lagrangians (13) and (15) are incomplete: We have worked in the regime of classical type IIB solutions so at least the universal Kähler modulus T is not yet stabilized. Moreover, we have not included the b -axion that complexifies c . Finally, there is no parametric separation between the mass scale of the complex structures and the warped Kaluza-Klein scale. Hence, the Lagrangian (13) does not define a useful effective field theory in the Wilsonian sense. Equation (15) however

¹⁶For a recent discussion on this see [66].

¹⁷One might for instance be tempted to consider the large volume limit where warping becomes negligible. In this case the scale of the potential would still be given by $|z|^2 \ll 1$.

does give rise to a Wilsonian effective Lagrangian once it is completed by the b -axion and the Kähler modulus T .

2.5 The B_2 -axion

In the preceding sections we have focused on the ultralight c -axion that can be thought of as the integral of the RR two-form C_2 over a sphere between the two throats. Similarly, we can define a b -axion by integrating the NS two-form B_2 instead,

$$b \equiv \frac{1}{2\pi\alpha'} \int_{S^2} B_2. \quad (16)$$

By the same arguments as before (see (5)) a non-vanishing field excursion induces a pair of H_3 flux/anti-flux on the portions of the \mathcal{B} -cycle that reach down into the two throats. Now, in the vacuum the \mathcal{B} -cycle is already filled with quantized H_3 -flux,

$$K \equiv K_1 + K_2 \equiv \frac{1}{(2\pi)^2\alpha'} \left(\int_{\mathcal{B}_1} H_3 + \int_{\mathcal{B}_2} H_3 \right). \quad (17)$$

Here \mathcal{B}_1 and \mathcal{B}_2 are the three-chains that reach into the respective throats and are bounded by the sphere between the throats, so that $\mathcal{B} = \mathcal{B}_1 + \mathcal{B}_2$. Clearly the continuous field excursion of the b -axion does not change the quantized flux integer K . However, by Stokes' theorem it *does* change the relative flux distribution,

$$K_1 \longrightarrow K_1 + \frac{b}{2\pi}, \quad K_2 \longrightarrow K_2 - \frac{b}{2\pi}. \quad (18)$$

By definition, K_1 and K_2 are the (non-quantized) H_3 fluxes that reside in the respective throats. Again, treating the local throat deformation parameters $z_{1,2}$ as independent it is clear that the throats can restore supersymmetry by an appropriate adjustment [29]:

$$|z_{1,2}| \sim \exp\left(-2\pi \frac{K_{1,2}}{g_s M}\right) \longrightarrow \exp\left(-2\pi \frac{K_{1,2} \pm b/2\pi}{g_s M}\right). \quad (19)$$

Thus, the discussion of the previous section applies also to the b -axion if one replaces the phases of the local deformation parameters by $\log |z_i|$. In other words, while the c -axion rotates the throats against each other, the b -axion makes one throat longer and the other shorter (see Fig. 7).

Expanding on the above, we can now comment on the interesting difference between the




Figure 7: The physical effect of a field excursion of the b -axion in the double throat system. One throat becomes shorter, whereas the other becomes longer.

5d and 4d perspective on excitations of $\int_{S^2} B_2$ in the local throat. This will in particular facilitate comparison with the related discussion in [46].

We start by noting that, instead of the 4d field b defined by an integral inbetween the two throats, one may also consider the 5d field $b(r) \sim \int_{S^2} B_2$ within either of the throats. Here r is the radial location of the relevant S^2 .¹⁸ Away from the tip of the throat and from the bulk CY, where the 5d language can be used, one has a continuum of solutions to the 10d supergravity equations [39]. In particular, there is a continuum of solutions for $b(r)$, parametrized by z via the boundary condition $b(r = |z|^{1/3}) = 0$. Two such solutions are plotted in Fig. 8. The relevant 5d equation for a static solution is, symbolically,

$$[\partial_r^2 - f(r)\partial_r - m^2(r)] b(r) = 0, \quad (20)$$

where $m(r)$ is the 5d mass. One immediately sees that the non-trivial (non-constant) profile of $b(r)$ is enforced by the non-zero $m^2(r)$. This non-zero potential comes from the CS term $\int F_5 \wedge B_2 \wedge F_3$. One might be concerned that such a potential clashes with the presence of an ultralight (in the present approximation massless) 4d mode of b . But such concerns are unfounded: Indeed, the 4d flat direction parameterized by b is also present in the 5d description, in spite of the non-zero 5d potential. It corresponds to a change of the whole $b(r)$ -profile within the available continuum of solutions accompanied by a change of boundary condition, e.g. $z \rightarrow z'$, cf. Fig. 8. The variable z is, of course, not accessible to a local observer in 5d at some fixed position r_* .

Finally, when gluing together throats to create a double throat one expects a small potential for the b -axion due to the misalignment of the *magnitudes* of the two deformation

¹⁸In order to avoid introducing additional symbols, by a slight abuse of notation, we will denote by $b(r)$ the 5d mode while b without radial argument denotes the 4d mode.




Figure 8: Two solutions $b(r) \sim \int B_2$ to the supergravity equations of motion in the Klebanov-Tseytlin throat. The 4d flat direction of the potential corresponds to a change of solution.

parameters, compare the discussion of Sect. 2.3 in the C_2 -case. We will make a quantitative statement in the next section.

3 Four-Dimensional SUGRA Completion

So far we have discussed how the c -axion backreacts on the phases of the local deformation parameters of the throats. In this section we propose a completion of the model in the language of 4d supergravity. The C_2 -axion pairs with the analogous B_2 -axion into a complex field $\mathcal{G} = c - \tau b$. The b -axion backreacts on the *magnitudes* of the deformation parameters in a way that is analogous to the backreaction of the c -axion on their phases.

3.1 Counting Moduli Through the Conifold Transition

Throughout Sect. 2 we have focused on the case of two S^3 -cycles related in homology, i.e. $[\mathcal{A}_1] = [\mathcal{A}_2]$. In general we denote by n the number of collapsing three-spheres \mathcal{A}^i , $i = 1, \dots, n$ and by m the number of homology relations between them $\sum_{i=1}^n p_i^I [\mathcal{A}^i] = 0$, $I = 1, \dots, m$.

Before fluxes are turned on and orientifold projections are imposed the physical degrees of freedom assemble into $\mathcal{N} = 2$ multiplets. The $n - m$ complex structure moduli z^i are the scalar components of $n - m$ vector multiplets. The z^i parametrize the Coulomb branch of the gauge theory. Whenever some of the three-cycles shrink to zero size, charged hypermultiplets (*Strominger black holes*) become massless and have to be ‘integrated in’

[67]. These can be thought of as D3-branes that wrap the shrunken cycles. At the origin of the Coulomb branch there are hence n massless charged hypermultiplets and n singular nodes have developed in the CY threefold. There exists an m -dimensional Higgs-branch where the singular nodes are resolved into m (homologically independent) \mathbb{P}^1 's [68]. On this branch, the $n - m$ vector multiplets eat $n - m$ black brane hypermultiplets and become massive. Geometrically speaking this is the resolution of the conifold [50, 51].

In the $\mathcal{N} = 1$ flux compactification that we are considering the tips of the conifolds become strongly red-shifted. Moreover backreaction of fluxes ensures that even at tiny complex structure the S^3 's stay at finite size so that the Strominger black holes play no role. However, since the deformed and the resolved conifold differ only by their strongly red-shifted tip geometries we expect to recover some remnant of the resolved phase of the conifold theory in the light spectrum. As outlined in Sect. 2 we expect the ‘local complex structures’ to decouple from one another so that all the n local deformation parameters z_1, \dots, z_n become equally light. In other words there are m additional light geometric modes. Moreover, on the resolved side of the transition there would be m massless axion modes. Since the obstruction for them to be massless is also localized at the tips of the conifolds where the would-be two-cycles collapse we also expect m complex light axionic modes \mathcal{G}^I . As we will argue in the next section, these modes indeed appear quite naturally in the discussion of the flux superpotential.

3.2 The Thraxion Superpotential

In this section we make a proposal for the 4d supergravity completion of the Lagrangians (13) and (15) for a general number of throats n with m homology relations among the shrinking cycles. Throughout this section we work in units $M_{\text{P}} = 1$.

3.2.1 The GVW Superpotential of a Multi Conifold System

As a starting point we consider the GVW flux superpotential for a multi conifold system. All the necessary ingredients are derived in ref. [69] and summarized in Appendix E. We choose to treat the redundant set of the n complex structure parameters z_i associated with the n vanishing cycles \mathcal{A}^i democratically, and impose the m *CY conditions* via a set of Lagrange multipliers λ_I , $I = 1, \dots, m$. The superpotential reads

$$W(z) = \sum_{i=1}^n \left(M_i \frac{z_i}{2\pi i} \log(z_i) + M_i g^i(z) - \tau K^i z_i \right) + \sum_{I=1}^m \lambda_I P^I + \hat{W}_0(z). \quad (21)$$

The m homology relations among the vanishing cycles $\sum_{i=1}^n p_i^I \mathcal{A}^i = \partial \mathcal{C}^I$, $I = 1, \dots, m$ lead to the following m CY conditions for the $z_i \equiv \int_{\mathcal{A}^i} \Omega$,

$$0 \stackrel{d\Omega=0}{=} \int_{\mathcal{C}^I} d\Omega = \int_{\partial \mathcal{C}^I} \Omega = \sum_{i=1}^n p_i^I \int_{\mathcal{A}^i} \Omega = \sum_{i=1}^n p_i^I z_i \equiv P^I, \quad I = 1, \dots, m. \quad (22)$$

In this language, the m CY conditions $P^I = 0$ are equivalent to the F-term equations of the Lagrange multipliers λ_I , $\partial_{\lambda_I} W \stackrel{!}{=} 0$. For details we refer the reader to App. E¹⁹.

Here, the M_i and K^i are the flux numbers associated to the \mathcal{A} - and \mathcal{B} -cycle of the i -th throat, and the holomorphic function $\hat{W}_0(z)$ denotes contributions to the flux superpotential from other cycles. The $M_i \in \mathbb{Z}$ cannot all be chosen independently but must comply with the m homology conditions

$$\sum_{i=1}^n p_i^I M_i = 0, \quad I = 1, \dots, m. \quad (23)$$

The K^i can be chosen independently but there is an m -fold redundancy in their definition because we may transform $K^i \rightarrow K^i + \sum_I \alpha_I p_i^I$ for any $\alpha \in \mathbb{C}^m$ leaving the superpotential invariant upon imposing the constraint equations²⁰. Furthermore, there are n unknown functions $g^i(z)$ defined on complex structure moduli space that are holomorphic near the origin.

¹⁹Note that we restrict ourselves to regions in complex structure moduli space close to the conifold transition point, where all throats degenerate simultaneously, compare the discussion in Sect. 2.1.1. This might be more restrictive than is needed for our analysis: If the matrix p_i^I is block-diagonal, we can separate the multi throat system into smaller multi throats whose deformations are independent of one another. In this case we can go through a conifold transition by local degeneration of the throats of a smaller system. Even away from the trivial case of multi throats factorizing, one might be able to achieve small thraion masses by ‘freezing’ individual throats with larger deformation z . Given a multi throat with some large z ’s one has to check the thraion potential as proposed in this section for flat directions. We leave a more thorough analysis of this possibility for future work.

²⁰The $n - m$ physical H_3 flux quantization conditions can be stated as $K^a - \sum_{I=1}^m p_a^I K^{n-m+I} \in \mathbb{Z}$, $a = 1, \dots, n - m$. This is because we can always choose the first $n - m$ of the shrinking cycles to correspond to integral basis elements $[\mathcal{A}^1], \dots, [\mathcal{A}^{n-m}]$ in homology. The Lagrange constraints can be stated as $0 = P^I = \sum_{a=1}^{n-m} p_a^I z_a + z_{n-m+I}$, i.e. $z_{n-m+I} = -\sum_{a=1}^{n-m} p_a^I z_a$. In the superpotential the terms that multiply z_1, \dots, z_{n-m} are given by the above combination of K^i and correspond to the integer flux numbers on the cycles $\mathcal{B}_1, \dots, \mathcal{B}_{n-m}$. Alternatively, one may demand the sufficient but not necessary conditions that $K^i \in \mathbb{Z}$ for $i = 1, \dots, n$. In this more restrictive but democratic formulation the i -th throat carries K^i units of flux. We can still reach all possible integer values for flux numbers on the cycles \mathcal{B}_a .

The Kähler potential is given by

$$\begin{aligned}
K_{\text{cs}}(z_i, \bar{z}_i) &= -\log \left(-i \int \Omega \wedge \bar{\Omega} \right) = -\log \left(ig_K(z) - i\overline{g_K(z)} + \sum_{a=1}^{n-m} i\bar{z}_a G^a + c.c. \right) \\
&= -\log \left(ig_K(z) - i\overline{g_K(z)} + \sum_{i=1}^n \left[\frac{|z_i|^2}{2\pi} \log(|z_i|^2) + i\bar{z}_i g^i(z) - iz_i \overline{g^i(z)} \right] \right), \tag{24}
\end{aligned}$$

where the holomorphic function $g_K(z)$ encodes contributions from other cycles. We would like to stress that despite the fact that we have written the unknown functions g^i , g_K and \hat{W}_0 as functions of all the z_i , $i = 1, \dots, n$, knowledge of the periods of the various cycles (and the flux quanta) only determines their behavior *along* complex structure moduli space and not beyond.

3.2.2 The Thraxion as a Stabilizer Field

We are now ready to formulate a proposal for the thraxion superpotential. First, we note the following. By expanding the Lagrange multiplier terms, one may rewrite the superpotential (21) as

$$W(z) = \sum_{i=1}^n \left(M_i \frac{z_i}{2\pi i} \log(z_i) + M_i g^i(z) + \left[-\tau K^i + \sum_{I=1}^m \lambda_I p_i^I \right] z_i \right) + \hat{W}_0(z). \tag{25}$$

One observes immediately that the combinations $\sum_{I=1}^m \lambda_I p_i^I$ can be interpreted as an *additional*, unquantized contribution to the complex three-form flux $G_3 = F_3 - \tau H_3$ on the (local portion $\tilde{\mathcal{B}}^i$ of the) \mathcal{B} -cycle of the i -th conifold. But we know that such a flux is detected by a boundary integral

$$\hat{\mathcal{G}}_i \equiv c_i - \tau b_i \equiv \frac{1}{2\pi\alpha'} \int_{S^2|_{i\text{-th throat}}} (C_2 - \tau B_2) = \frac{1}{2\pi\alpha'} \int_{\tilde{\mathcal{B}}^i} (F_3 - \tau H_3) \tag{26}$$

over the S^2 at the top of the i -th throat. Crucially, the variables $\hat{\mathcal{G}}_i$ define axionic field excursions as measured near the entrance of the i -th throat.

We would like to interpret (a subset of) these as light physical degrees of freedom. This is motivated by the fact that there are m light axions on the other side of the conifold transition that correspond to the integrals of $C_2 - \tau B_2$ over the independent resolution 2-cycles. Indeed, the counting is correct. A consistent axionic field excursion must not induce any overall flux on *any* of the global \mathcal{B} -cycles (see Fig. 9). There are hence $n - m$ *no-*




Figure 9: A cartoon of the \mathcal{B} -cycle of the triple throat, $n = 3$ and $m = 2$. The local c -axion excursions c_1, c_2 and c_3 , $c_i = \text{Re } \hat{\mathcal{G}}_i$, must be chosen so that no overall flux is generated on \mathcal{B} , i.e. $0 \stackrel{!}{=} c_1 + c_2 + c_3$ ²¹ or rather $\sum_i \hat{\mathcal{G}}^i = 0$.

flux conditions, one for each linearly independent \mathcal{B} -cycle, leaving only m physical axions. These can be parametrized as $\hat{\mathcal{G}}^i = \sum_{I=1}^m p_i^I \mathcal{G}_I$ and we are led to the following conjecture:

The Lagrange multipliers λ_I must be promoted to m light axionic degrees of freedom, $\lambda_I \rightarrow \frac{\mathcal{G}_I}{2\pi}$. Moreover, the z^i are promoted to n physically independent degrees of freedom.

The normalization factor 2π is chosen such that locally in the i -th throat a shift of axionic field excursion \mathcal{G}_I by 2π (or $2\pi\tau$) for some I is indistinguishable from an increase of the F_3 -flux (respectively H_3 -flux) on the \mathcal{B} -cycle of the throat by an integer amount p_i^I .

Thus, our proposal for the superpotential is

$$W = \sum_{i=1}^n \left(M_i \frac{z_i}{2\pi i} \log(z_i) + M_i g^i(z) - \tau K^i z_i \right) - \sum_{I=1}^m \frac{\mathcal{G}_I}{2\pi} P^I + \hat{W}_0(z). \quad (27)$$

We find it interesting to note that the axions \mathcal{G}^I now serve as the *stabilizer fields* for the combinations of the local deformation parameters that break the m CY conditions $P^I = 0$, $I = 1, \dots, m$. This form of the dynamical thaxion superpotential is fairly unique in that it preserves the set of *discrete* shift-symmetries

$$z_i \longrightarrow z_i e^{\frac{2\pi i}{M_i} \sum_I \eta_I p_i^I}, \quad \mathcal{G}_I \longrightarrow \mathcal{G}_I + 2\pi \eta_I, \quad \forall \eta \in \mathbb{C}^m : \sum_I \eta_I p_i^I \in M_i \mathbb{Z} \quad \forall i. \quad (28)$$

²¹Compare this to Fig. 5: The ‘no-flux’ condition in the double throat setup amounts to $c_1 = -c_2$. The two axions c_1 and c_2 are actually identified, up to a sign due to different orientation of the two-sphere in the definition. This is why we only had one axion c to begin with in the 10d analysis of Sect. 2.

Our proposal for the Kähler potential is

$$K(\mathcal{G}_I, \bar{\mathcal{G}}_I, T, \bar{T}, z, \bar{z}) = K_1(\mathcal{G}_I - \bar{\mathcal{G}}_I, T + \bar{T}) + K_{\text{cs}}(z, \bar{z}), \quad (29)$$

where K_{cs} is the Kähler potential (24) and K_1 is the Kähler potential of the m axions (and Kähler moduli T) on the other side of the conifold transition as derived in [70]

$$K_1 = K_0 - 3 \log \left(T + \bar{T} - \frac{3i}{4(\tau - \bar{\tau})} \kappa_{1IJ} (\mathcal{G} - \bar{\mathcal{G}})^I (\mathcal{G} - \bar{\mathcal{G}})^J \right), \quad (30)$$

where K_0 contains a constant part and the Kähler potential of the axio-dilaton and κ_{1IJ} are triple intersection numbers.²²

We expect (24) and (27) to hold even when we break the CY condition $P^I \neq 0$ with the important subtlety that the domain of the holomorphic functions g^i, g_K and \hat{W}_0 must be extended *beyond* complex structure moduli space. We find it reasonable to expect that such an extension *exists* although even full knowledge of the CY periods would not determine their behavior away from the moduli space. The detailed form of these functions will be of no importance in what follows. Moreover, we expect that using the potential K_1 gives an excellent approximation because the kinetic term of the axions is dominated by contributions from the UV where the deformation or resolution of the conifold plays but a tiny role. Note, that the behavior of the kinetic terms of the complex structure moduli is dominated by the logarithmic terms in (24). In particular, the functions $g^i(z)$ and $g_K(z)$ contribute to kinetic terms only at sub-leading order.

Since we are interested in small z_i we Taylor-expand

$$\begin{aligned} g^i(z) &= g_0^i + \sum_{j=1}^n g_1^{ij} z_j + \mathcal{O}(z^2), & \hat{W}(z) &= g_{W,0} + \sum_{i=1}^n g_{W,1}^i z_i + \mathcal{O}(z^2), \\ g_K(z) &= g_{K,0} + \sum_{i=1}^n g_{K,1}^i z_i + \mathcal{O}(z^2). \end{aligned} \quad (31)$$

This should really be understood as a Taylor expansion in n *independent* variables z_i and makes our conjectured extension of the domain of these functions beyond the complex structure moduli space manifest.

²²For ease of exposition we have stated the Kähler potential for the case $h_{1,1}^+ = 1$. For $\tau = \text{const.}$ the Kähler potential fulfills the no-scale relation $(\partial_T K_1)(K_1^{-1})^{T\bar{T}}(\partial_{\bar{T}} K_1) = 3$, and $(K_1^{-1})^{g^I \bar{X}^{\bar{j}}} \partial_{\bar{X}^{\bar{j}}} K_1 = 0$ where $X^i = (T, \mathcal{G}^I)$.

We absorb all $\mathcal{O}(z^0)$ terms in the superpotential in the definition $\hat{W}_0 \equiv g_{W,0} + \sum_{i=1}^n M_i g_0^i$. The coefficients in (31) should all be viewed as independent of the flux quanta that thread the cycles of the multi throat system, and only $(g_{W,0}, g_{W,1}^i)$ depend on fluxes on other cycles.

3.2.3 The Double Throat: $n = 2, m = 1$

It is clear that to obtain the effective superpotential for the \mathcal{G} -fields we should integrate out the local deformation parameters. Before we discuss this in full generality it is instructive to first consider the simplest case of the double throat, i.e. $n = 2$ and $m = 1$. There are two \mathcal{A} -cycles \mathcal{A}^1 and \mathcal{A}^2 and we choose the homology relation to be $\mathcal{A}^1 \sim \mathcal{A}^2$. Hence, there are two deformation parameters z_1 and z_2 and one axion \mathcal{G} . For ease of exposition we assume that of all the coefficients defined in (31), only $g_{W,0}$ and $g_{K,0}$ are non-vanishing, in other words, we choose \hat{W}_0 as well as all non-logarithmic terms in the Kähler potential to be constant. In doing so we accept an $\mathcal{O}(1)$ error in all expressions, in particular in the resulting superpotential W_{eff} for \mathcal{G} . This simplifying assumption will be dropped when we generalize the discussion to the multi throat case in Sect. 3.2.4.

We must set $M_1 = M_2 \equiv M$ due to the homology relation between the shrinking cycles, and we choose $K^1 \equiv K/2 \equiv K^2$ which results in flux $K^1 + K^2 = K$ on the \mathcal{B} -cycle. All choices of the pair (K^1, K^2) that satisfy $K^1 + K^2 = K$ are physically equivalent to this choice and can be brought back to the symmetric choice via a linear redefinition of \mathcal{G} (compare the discussion below (23)). The superpotential takes the form

$$W(z_1, z_2, \mathcal{G}) = \sum_{i=1}^2 \left(\frac{z_i}{2\pi i} \log(z_i) M - \frac{1}{2} K \tau z_i \right) - \frac{\mathcal{G}}{2\pi} (z_1 - z_2) + \hat{W}_0 + \mathcal{O}(z_i^2). \quad (32)$$

First, the F-terms F_{z_i} are given by

$$\begin{aligned} D_{z_i} W &= \partial_{z_i} W + (\partial_{z_i} K) W = \frac{\log(z_i) + 1}{2\pi i} M - \frac{K}{2} \tau \mp \frac{\mathcal{G}}{2\pi} + \mathcal{O}(z_i) \\ &= \frac{M}{2\pi i} (\log(z_i/z_0) \mp i\mathcal{G}/M) + \mathcal{O}(z_0), \end{aligned} \quad (33)$$

where

$$z_0 = e^{-1} \exp\left(2\pi i \tau \frac{K/2}{M}\right) + \mathcal{O}\left(e^{-4\pi \frac{K/2}{g_s M}}\right) = \mathcal{O}\left(e^{-2\pi \frac{K/2}{g_s M}}\right). \quad (34)$$

As usual, with $K/2 > g_s M$ one obtains $|z_0| \ll 1$ with universal dependence on the flux numbers. Following Sect. 2 we may integrate out the local deformation parameters, which

yields

$$z_1 = z_0 e^{i\mathcal{G}/M}, \quad z_2 = z_0 e^{-i\mathcal{G}/M}. \quad (35)$$

The effective superpotential for the axion \mathcal{G} reads

$$W_{\text{eff}}(\mathcal{G}) = 2\epsilon (1 - \cos(\mathcal{G}/M)) + W_0 + \mathcal{O}(z_0^2), \quad (36)$$

with $\epsilon \equiv M \frac{z_0}{2\pi i}$, and $W_0 \equiv \hat{W}_0 - \frac{z_0}{\pi i} M$.

This is the expression we were after. Crucially, it is consistent with the results of Sect. 2: Using the Kähler potential for the Kähler moduli and \mathcal{G} -axions (30), one may show that $V(\mathcal{G}, \bar{\mathcal{G}}) \propto |\partial_{\mathcal{G}} W(\mathcal{G})|^2$ ²³. If we restrict to $\mathcal{G} = c \in \mathbb{R}$, we reproduce the periodic potential (10) with the correct scaling $|\epsilon|^2 \sim |z_0|^2 \sim w_{\text{IR}}^6$.

Note that because we have made use of the *unwarped* Kähler potential we do not reproduce the correct mass-scale of the local deformation parameters z_i . Here, this is of no importance because all degrees of freedom that are related to strongly warped regions are integrated out supersymmetrically. In particular, the potential energy induced by a non-vanishing field excursion of the field \mathcal{G} receives its dominant contributions from the bulk CY where warping plays no role. Because in going from weak to strong warping, the solutions of the complex structure F-terms are left invariant [38], and because the z_i are parametrically heavier than \mathcal{G} even when the appropriate red-shift factors are introduced in the scalar potential, this procedure is justified.

We are now ready to expand on the conclusions we have drawn in Sect. 2. First of all, the kinetic term of the full complex field \mathcal{G} lives in the bulk. This implies that the mass² of \mathcal{G} is of order $|z_0|^2 \ll 1$. Since the Kähler potential is independent of $\text{Re}(\mathcal{G})$ a discrete shift-symmetry $\mathcal{G} \rightarrow \mathcal{G} + 2\pi M$ is manifest²⁴, while the IR superpotential breaks the shift-symmetry corresponding to $\text{Im}(\mathcal{G})$ completely.

While in principle the target space distance traversed by $\text{Im}(\mathcal{G})$ can be made large, the scalar potential grows exponentially as a function thereof as is common for saxionic directions in field space. In particular, this direction in field space is of little use for (slow-roll) inflation. There is a *critical* field excursion $|\text{Im}(\mathcal{G})_{\text{crit}}| \lesssim 3M \log(w_{\text{IR}}^{-1})$ beyond which one side of the double throat is entirely pulled up into the bulk CY, $z_1 \sim 1$ or $z_2 \sim 1$. Near this field excursion we no longer know the form of the potential because we work to lowest order in $|z_1|, |z_2|$. Moreover, there is a tower of warped KK-modes with masses that scale

²³Where we use that K_1 is of no-scale type.

²⁴In an exact no-scale background the scalar potential even has periodicity πM . However, any no-scale breaking effects will break it to the periodicity of the superpotential.

as

$$m_n^2 \sim n^2 w_{\text{IR}}^2 \exp(-2|\text{Im}(\mathcal{G})|/3M), \quad (37)$$

where the warp factor $w^2 \sim |z_i|^{2/3}$ now depends on $\text{Im}(\mathcal{G})$. Since these modes have been integrated out, the ratio $\Lambda/m_{\mathcal{G}}$ of the cutoff of the \mathcal{G} -EFT (i.e. the smallest KK-mass) over the mass-scale of \mathcal{G} , comes down as

$$\Lambda/m_{\mathcal{G}} \sim w_{\text{IR}}^{-2} e^{-\frac{4}{3M}|\text{Im}\mathcal{G}|} \leq w_{\text{IR}}^{-2} \exp\left(-\frac{\phi_b}{M_{\text{P}}}\right), \quad (38)$$

at large field excursion, consistent with a *distance conjecture* [9,27,71–76]. Here ϕ_b measures the canonical field distance from the origin along the imaginary \mathcal{G} -axis²⁵.

At strong warping the maximal allowed field excursion is $|\text{Im}(\mathcal{G})_{\text{max}}| \sim \frac{3}{2}M \log(w_{\text{IR}}^{-1})$ before the 4d EFT description breaks down. Near this field excursion, a large fraction of the reservoir of fluxes of one of the throats has been transferred to the other one and the mass scale of the \mathcal{G} -field is of the same order as the warped KK-scale of the longer throat. At this point, contributions to the scalar potential from non-vanishing F -terms $D_{z_i}W$ start to play a significant role, or from a 10d point of view, the potential energy sinks down into the longer throat.

3.2.4 The General Multi Throat

For general m and n that satisfy $n - m > 0$ homology relations, there are n local deformation parameters z_1, \dots, z_n that have to be integrated out. We are left with an effective supergravity theory of m axions \mathcal{G}^I . The computational steps are analogous to the double throat case that was laid out in detail. Hence, we only state the effective axion superpotential

$$W_{\text{eff}}(\mathcal{G}^I) = - \sum_{i=1}^n \epsilon_i e^{i \sum_{I=1}^m p_I^i \mathcal{G}^I / M_i} + \hat{W}_0, \quad (39)$$

and we have defined

$$\epsilon_i \equiv \frac{M_i}{2\pi i} z_{0,i} (1 - 2\pi \tilde{g}_0^i \hat{W}_0 / (a M_i)), \quad \tilde{g}_0^i \equiv g_0^i - \overline{g_{K,1}^i}, \quad a \equiv -2\text{Im}(g_{K,0}), \quad (40)$$

and

$$z_{0,i} = e^{-1} e^{-\frac{2\pi i}{M_i} (\sum_j M_j g_1^{j,i} + g_{W,1}^{i,i} + i \tilde{g}_0^i \hat{W}_0 / a)} \exp\left(2\pi i \frac{K^i \tau}{M_i}\right) + \mathcal{O}\left(e^{-4\pi \frac{K^i}{g_s M_i}}\right). \quad (41)$$

²⁵See App. D for the conversion rule between \mathcal{G} and the canonical distance in field space ϕ_b . At any point in field space, $g_{\mathcal{G}\bar{\mathcal{G}}} < M_{\text{P}}^2/M^2$. Hence, $\phi_b = \int_0^{\text{Im}\mathcal{G}} d\text{Im}\mathcal{G}' \sqrt{g_{\mathcal{G}\bar{\mathcal{G}}}} < \text{Im}\mathcal{G} M_{\text{P}}/M$.

It is important to note that the $z_{0,i}$ as defined above can in general *not* be interpreted as the values of the local deformation in the vacuum. The physical local deformation parameters are given by

$$z_{\text{ph},i} \equiv z_{0,i} \exp \left(i \sum_I p_I^i \mathcal{G}^I / M_i \right), \quad (42)$$

where in the vacuum the \mathcal{G}^I need not vanish in general.

3.3 Comments on the b -Axion

In the above supergravity completion we have ‘complexified’ the c -axion by pairing it with the analogous b -axion. We have outlined the 10d backreaction of the b -axion already in Sect. 2.5. Now that we have addressed the scalar potential of the b -axion quantitatively, in this section we would like to comment on a potential worry and how to resolve it: We recall that the effect of a non-vanishing field excursion of the b -axion is the creation of a pair of fluxes of the NS field strength H_3 . Since both throats are filled up with H_3 -fluxes already in the vacuum one should think about this process more properly as a transfer of H_3 -flux from one throat to the other. Since the magnitudes of the local deformation parameters (and the associated hierarchy) are set by the ratio of local H_3 -flux (on the \mathcal{B} -cycle) and F_3 -flux (on the \mathcal{A} -cycle) it is clear that these will backreact when the H_3 -fluxes are redistributed, see (19).

However, it is also clear that when the local H_3 -fluxes are changed, the circumference of the throat at the UV end is affected strongly. This is because it is set by the total D3-charge that is stored in the throat which is itself proportional to the amount of H_3 -flux [32], compare Fig. 7. Hence, naively one might worry that such considerable change at the UV ends of the throats could lead to a large potential energy. One may convince one-self that this is not the case as follows. Starting from the supersymmetric situation we can redistribute a small amount of fluxes from one throat to the other, so that throat A has δ units of H -flux more than throat B . We can now proceed to convert the extra fluxes into a number of D3-branes by going through the Kachru-Pearson-Verlinde (KPV) transition [77]. From the UV perspective this process is only detected by a change in the throat complex structure which is a tiny perturbation far from the tip of the throat. Now we are back to an even flux distribution with a number of mobile D3-branes. These can be moved out of the throat at no cost in energy so the situation with the mobile branes should be a vacuum again. In other words, the redistribution of fluxes creates an energy density that is *only* due to the misalignment of local deformation parameters and the change of size

of the throats at their UV ends does not generate an extra contribution to the potential. We reiterate that the situation is analogous to the backreaction of the c -axion with the phases of the local deformation parameters replaced by the logarithms of their magnitude.

Finally, note that in the Kähler potential (30) the b -axion appears explicitly, while the approximate c -axion shift symmetry is manifest. One might suspect that the small scale of the b -axion is therefore accidental due to our use of tree level supergravity. This conclusion would be incorrect: The target space manifold with Kähler metric derived from (30) is shift symmetric also in the b -direction [78, 79]. In general we expect both shift symmetries to be preserved to all orders in the perturbative expansion with explicit breaking only due to the superpotential (39). When moreover *non-perturbative* Kähler moduli dependent terms are generated in the superpotential such as the ones considered in [30], we expect the b -axion mass to be lifted to the scale of Kähler moduli stabilization while the c -axion can remain parametrically lighter.

4 The Axion Potential and the Gauge/Gravity Correspondence

We have derived the axion potential via a classical computation within 10d SUGRA, and proposed a 4d SUGRA description that matches it. Since the local throats are believed to have a dual description in terms of KS gauge theories [32], it is useful to give an alternative derivation of our results on the gauge theory side of the correspondence. The KS gauge theory is a $SU(N + M) \times SU(N)$ gauge theory with (classical) global symmetry group $SU(2) \times SU(2) \times U(1)_R$. It contains matter in bi-fundamental representations $(\square, \bar{\square})$ and $(\bar{\square}, \square)$ of the gauge group that transform as doublets under the first respectively second global $SU(2)$ factor. The holomorphic gauge couplings of the two gauge theory factors τ_{YM} and $\tilde{\tau}_{YM}$ have been argued to be set by [61, 80]

$$\tau_{YM} + \tilde{\tau}_{YM} = \tau, \quad \tau_{YM} - \tilde{\tau}_{YM} = -\tau + \mathcal{G}/\pi \pmod{2(m - n\tau)}, \quad (43)$$

with $(m, n) \in \mathbb{Z}^2$. The radial running of the \mathcal{G} -field together with $\tau = \text{const.}$ matches the RG-running of the gauge theory coupling constants. Throughout this section \mathcal{G} takes values in its suitable fundamental domain.

As the KS gauge theory flows to the infrared, it undergoes repeated steps of Seiberg

dualities that reduce the ranks of the gauge groups according to

$$\begin{aligned} SU(N_0 + M) \times SU(N_0) &\longrightarrow SU(N_1 + M) \times SU(N_1) \\ &\longrightarrow \cdots \longrightarrow SU(N_k + M) \times SU(N_k), \end{aligned} \quad (44)$$

with $N_k \equiv N - kM$, $k \in \mathbb{N}$. If we start with $N = KM$, after K steps in the duality cascade the gauge group is $SU(M)$. Since, roughly speaking, it corresponds to the first gauge group factor in $SU(M) \equiv SU(N_K + M) \times SU(N_K)$, its holomorphic scale is given by $\Lambda^{3M} = \mu_{\text{IR}}^{3M} \exp(2\pi i \tau_{\text{YM}}(\mu_{\text{IR}})) = \mu_{\text{IR}}^{3M} \exp(i\mathcal{G})$, where μ_{IR} is the infrared scale of the throat and where we make use of the KS dictionary $\tau_{\text{YM}} \simeq \mathcal{G}/2\pi$. Gaugino condensation leads to an effective ADS-superpotential [81, 82, 32]

$$W_{\text{eff}}(\mathcal{G}) = M\Lambda^3 \sim M\mu_{\text{IR}}^3 \exp(2\pi i \tau_{\text{YM}}/M) \sim M\mu_{\text{UV}}^3 \exp\left(\frac{2\pi i}{M} \left(\tau K + \frac{\mathcal{G}}{2\pi}\right)\right), \quad (45)$$

where we have used that the IR-scale is related to the UV-scale by $\mu_{\text{IR}}^3 = \mu_{\text{UV}}^3 \exp(-2\pi \frac{K}{g_s M})$.

The superpotential that we have proposed on the gravity side of the correspondence (39) indeed takes this form,

$$W \propto \sum_{i=1}^n M_i A_i \exp\left(\frac{2\pi i}{M_i} \left(\tau K^i + \frac{\hat{\mathcal{G}}^i}{2\pi}\right)\right) + \hat{W}_0, \quad (46)$$

with $\hat{\mathcal{G}}^i = \sum_{I=1}^m p_I^i \mathcal{G}^I$, and²⁶

$$A_i \equiv \exp\left(-\frac{2\pi i}{M_i} \left(\sum_{j=1}^n M_j g_1^{ji} + g_{W,1}^i + i\tilde{g}_0^i \hat{W}_0/a\right)\right) + \log\left(1 - \frac{2\pi \tilde{g}_0^i \hat{W}_0}{a M_i}\right). \quad (47)$$

From the gauge theory perspective we should interpret the appearance of the constants g_1^{ji} , $g_{W,1}^i$, \tilde{g}_0^i and \hat{W}_0 as a parameterization of threshold corrections near the UV cutoff²⁷. Indeed, as they are taken to zero the A_i become unity.

It is now obvious that the M -fold extension of the periodicity of the c -axion is related to gaugino condensation in the KS gauge theory²⁸ [39, 32, 85, 80]: As usual there is a $U(1)_R$ symmetry that is broken to \mathbb{Z}_{2M} by gauge theory instantons. Gaugino condensation

²⁶Note that in (39) we have set $M_{\text{P}} = 1$. Therefore we identify $\mu_{\text{UV}} \sim M_{\text{P}}$.

²⁷Of course these are in general functions of all other complex structure moduli that do not control the infrared regions of the throats and are frozen at a high scale.

²⁸Related observations were made in the non-compact flux-less multi-node setting of [83, 84].

spontaneously breaks $\mathbb{Z}_{2M} \rightarrow \mathbb{Z}_2$, so there are M gauge theory vacua. As we transform $c \rightarrow c+2\pi$, we move from one gauge theory vacuum to the next, and the gaugino condensate (which corresponds to the local deformation parameters on the gravity side) picks up a phase $\exp(2\pi i/M)$. This is as in Sect. 2.2 where we learned that the M different vacua are reached by dialing the RR flux quanta on the \mathcal{B} -cycle $Q = 0, \dots, M-1$ (see (3)).

5 Applications

5.1 Thractions on the Quintic: Drifting Monodromy

In this section we will give a concrete example of a string compactification where a light thractions appears. Along the way we identify concrete setups in which parametrically super-Planckian racetrack-type axion periodicities are possible. We choose the CY to be the quintic three-fold which is defined as the vanishing locus of a homogeneous polynomial P of rank five in \mathbb{P}^4 . Following [69], it can be brought to a conifold transition point by choosing the complex structures such that

$$P(X^0, \dots, X^4) = X^3 f_4(X^0, \dots, X^4) + X^4 g_4(X^0, \dots, X^4), \quad (48)$$

where f_4 and g_4 are generic homogeneous rank four polynomials of the projective coordinates $\{X^0, \dots, X^4\}$ of \mathbb{P}^4 . The conditions $P = 0, dP = 0$ are satisfied whenever

$$0 = X^3 = X^4 = f_4(X^0, X^1, X^2, 0, 0) = g_4(X^0, X^1, X^2, 0, 0). \quad (49)$$

Since f_4 and g_4 are chosen to be generic polynomials of rank 4, there exist $4 \cdot 4 = 16$ distinct solutions. These are 16 conifold points. Hence, there are 16 vanishing three-cycles \mathcal{A}^i , $i = 1, \dots, 16$. Because the solution set lies on a \mathbb{P}^2 submanifold of \mathbb{P}^4 , there is precisely one homology relation among them,

$$\sum_{i=1}^{16} [\mathcal{A}^i] = 0. \quad (50)$$

Hence, we have a multi throat system with $n = 16$ and $m = 1$ so there is one light axion.

Let us give two examples that differ by choices of flux numbers. In both examples we set the coefficients A_i defined in (47) to unity. Generically we expect these to be of order

one. Inserting $\mathcal{O}(1)$ factors below does not change the physical outcome.²⁹

Example 1: A simple thraxion potential

$$M_i = (-1)^{i+1}M, \quad K^i = (-1)^{i+1}K, \quad (51)$$

with $K/g_s M \gg 1$. Then we have $\epsilon_i \equiv (-1)^{i+1}\epsilon$, and

$$W_{\text{eff}}(\mathcal{G}) = -16i\epsilon \sin 2\mathcal{G}/M + \hat{W}_0 = 16i\epsilon(1 - \cos 2\mathcal{G}'/M) + W_0, \quad (52)$$

with $W_0 = \hat{W}_0 - 16\epsilon$, $\mathcal{G}' = \mathcal{G} - \pi M/4$, and small $|\epsilon| \propto \exp(-2\pi K/g_s M)$. Up to the numerical pre-factor this is exactly what we found for $n = 2$ and $m = 1$.

Example 2: Drifting Monodromy

We now slightly detune the F_3 fluxes from one another:

$$M_1 = M, \quad M_2 = M + 1, \quad M_3 = -M, \quad M_4 = -(M + 1), \quad (53)$$

and $M_{i+4} = M_i$, with $K^i \equiv \text{sign}(M_i)|K|$, and again $K/g_s M \gg 1$. In this case

$$W_{\text{eff}}(\mathcal{G}) \approx -8iz_0(M \sin(\mathcal{G}/M) + (M + 1) \sin(\mathcal{G}/(M + 1))) + \hat{W}_0, \quad (54)$$

with $z_0 \sim \exp 2\pi i K \tau / M$. Additionally to the previous simplification, we have also neglected order one prefactors that arise from the fact that the ratios K^i/M_i are not all exactly equal. Again, this is of no consequences for our purposes.

The superpotential (54) is a *racetrack*-type superpotential³⁰ for \mathcal{G} . The axion periodicity is now given by $2\pi M(M + 1)$. Crucially, this implies another M -fold extension of the axion field range *on top* of the one already discussed in the simpler examples of the double throat and the first example of this section. Clearly, one may take this even further to periodicities such as $2\pi M \cdots (M + 3)$.³¹ Since we still only have to fulfill the requirement that the throats fit into the bulk CY, this implies the existence of a simple, concrete and explicit mechanism in string theory that can generate huge super-Planckian axion periodicities. In general the full periodicity of the superpotential is given by the least common multiple of the different RR flux numbers M_i . We dub this mechanism of generating a parametrically

²⁹If some coefficients can be tuned parametrically smaller than others, new qualitative features might arise. We leave an investigation of this possibility to future research.

³⁰For a discussion of racetrack superpotentials in connection with the WGC, see [86].

³¹But not further because we have to respect the orientifold action.




Figure 10: The axion potential of example 2 for the case $M = 10$. There are sub-Planckian oscillations within a long super-Planckian envelope.

large axion monodromy *drifting monodromies* since it relies on a frequency drift within a set of several finite-order monodromy effects. This is related but different from the winding idea, where a constraint forces the effective axion on a long trajectory in a multi-axion moduli space [40, 41, 16, 43, 44]. Here, by contrast, one may think of a single fundamental axion extended by several small, finite-order monodromy effects. The result of this can still be large as explained above. The intended outcome, namely to realize an effective large- f axion accepting a short-wavelength oscillatory potential, is of course the same (see in particular the very recent analysis of [48]).

The minima of the potential $V \propto |\partial_{\mathcal{G}} W|^2$ are located along the slice $\text{Im } \mathcal{G} = 0$ where it takes the form

$$\begin{aligned} V(c) &\propto [\cos(c/M) + \cos(c/(M+1))]^2, \quad c \equiv \text{Re } \mathcal{G}, \\ &\propto \cos^2\left(\frac{2M+1}{2M(M+1)}c\right) \cdot \cos^2\left(\frac{1}{2M(M+1)}c\right), \end{aligned} \quad (55)$$

which has $2M + 1$ distinct Minkowski vacua (see Fig. 10).

We note that despite the long $2\pi M(M+1)$ periodicity the scalar potential oscillates on shorter wavelengths of order $2\pi M$. This is essentially due to the rank condition (23) which forces us to introduce flux numbers of both signs.³² We have not shown in general that suppressing such shorter wavelength oscillations in order to produce a smooth super-Planckian axion potential is impossible. At this point we only note that the condition (23) presents a severe obstacle towards this.

These examples also serve to illustrate that by scanning over flux numbers one may obtain a vast number of possible effective superpotentials and axionic potentials.

³²This condition is *global* in the sense that it need not hold in a non-compact CY where gravity is decoupled.

5.2 A Clash with the Weak Gravity Conjecture

In this section we would like to point out that the axion potential we have derived clashes with what has been put forward as the *weak gravity conjecture for axions* [10, 14, 17]. We have computed the axion potential via a classical supergravity calculation. However, one may equally well associate it to non-perturbative effects in the KS gauge theory (namely gaugino condensation), as argued in Sect. 4. As such, (if true) the weak gravity conjecture should apply to our construction.

In its form adapted to axions and instantons the conjecture states that there should exist an instanton with Euclidean action S and axionic charge q such that³³

$$S \leq \mathcal{O}(1) q M_{\text{P}}/f_{\text{eff}}. \quad (56)$$

Such instantons (if they contribute to the superpotential) generically induce terms in the scalar potential of the form

$$V(\phi) \supset e^{-S}(1 - \cos(q\phi/f_{\text{eff}})), \quad (57)$$

where ϕ is the canonically normalized axion. Thus, $2\pi f_{\text{eff}}/q$ is the canonically normalized periodicity of the term in the (super)potential that is generated by a given instanton.

By comparison with the above we may associate an (effective) Euclidean instanton action to each of the leading exponentially suppressed terms in the axion (super)potential³⁴.

$$S_{\text{eff}}^i \approx 3 \log(1/w_{\text{IR}}^i) \approx 2\pi \frac{K^i}{g_s M_i}. \quad (58)$$

As computed in App. D, in the regime where the throats marginally fit into the bulk CY, the periodicities f_{eff}/q^i of the dominant terms in the superpotential associated to each throat $i = 1, \dots, n$ read

$$f_{\text{eff}}/q_i \approx \frac{2}{3} (r_{\text{D3}}^i)^{1/2} \log(1/w_{\text{IR}}^i)^{-1/2} M_{\text{P}} \approx \frac{2}{3} (r_{\text{D3}}^i)^{1/2} \left(\frac{2\pi}{3} \frac{K^i}{g_s M_i} \right)^{-1/2} M_{\text{P}}, \quad (59)$$

where $r_{\text{D3}}^i = M_i K^i / N_{\text{D3}}^{\text{flux}}$ is the fraction of the total D3 brane charge of three-form fluxes

³³Note that $M_{\text{P}}/f_{\text{eff}}$ is the analogue of the gauge coupling for zero-forms (i.e. axions).

³⁴Taking the correspondence with instantons seriously, these are ($\frac{1}{2}$ BPS-)instantons.

which is stored in the i -th throat. Hence,

$$S_{\text{eff}}^i \cdot f_{\text{eff}}/q^i \sim 2(r_{\text{D3}}^i)^{1/2} \sqrt{\log(1/w_{\text{IR}}^i)} M_{\text{P}} \approx 2(r_{\text{D3}}^i)^{1/2} \sqrt{\frac{2\pi}{3} \frac{K^i}{g_s M_i}} M_{\text{P}}. \quad (60)$$

In the regime $w_{\text{IR}} \ll 1$ (i.e. $K^i \gg g_s M_i$) the r.h. side is parametrically larger than $\mathcal{O}(1)$ so the objects that generate the relevant terms in the superpotential do not satisfy a weak gravity conjecture bound.

Of course as is always true in string theory compactifications [87] there *does* exist a tower of instantons that satisfies the weak gravity bound (56) but generates no monodromy.³⁵ It is also apparent that these instantons occupy a sub-lattice of the full charge lattice. This sub-lattice corresponds to all the possible wrapping numbers of a Euclidean D1 string. However, in our setup this sub-lattice can be made parametrically *coarse*.³⁶ Let us illustrate this with a concrete example: We consider a variant of the *drifting monodromies* example given in Sect. 5.1, with flux numbers $M_i \in \{5, 6, 7, 8, -5, -6, -7, -8\}$. The axion decay constant is enhanced by the least common multiple of 5, 6, 7, 8 which is 840. The instantons that satisfy (56) respect the periodicity of the axion *before* monodromy. Thus the possible charges take values in $840\mathbb{Z} \subset \mathbb{Z}$. Clearly, a lattice WGC is parametrically violated, while a sub-lattice WGC [89, 88] (see also [90]) is always satisfied but with parametrically coarse sub-lattice. Note that generically these instantons only give rise to sub-leading corrections to the scalar potential (if they contribute at all), compare Sect. 5.5.

However, we observe that the n effective instantons that *do* give the dominant contribution to the superpotential satisfy a relation

$$S_{\text{eff}}^i \leq \frac{4}{3} r_{\text{D3}}^i (q^i M_{\text{P}}/f_{\text{eff}})^2, \quad \forall i = 1, \dots, n. \quad (61)$$

Together with $S_{\text{eff}}^i > 1$, as is required for controlled expansion in powers of $e^{-S_{\text{eff}}^i}$, this *still* implies the existence of short wavelength harmonics in the superpotential. This motivates a closer look at the full spectrum of our effective instantons to which we now turn.

5.3 The Spectrum of Effective Instantons

We now set aside the spectrum of instantons that satisfy the WGC but instead ask what are the properties of the effective instantons that generate the superpotential. We have

³⁵In our case, these are Euclidean D1 strings wrapping representative S^2 's in the UV, compare Sect. 5.5.

³⁶Hence we seem to realize explicitly the loop hole mentioned in footnote 25 of ref. [88].

written down the most dominant contributions to the superpotential (39) and noted that they satisfy (61). This condition is weaker than (56) because at fixed control parameter $S_{\text{eff}}^i \stackrel{!}{>} 1$ the value of $qM_{\text{P}}/f_{\text{eff}}$ is constrained to be bigger only than $\sqrt{S_{\text{eff}}}$ rather than S_{eff} . Nevertheless, as a consequence the dominant effective instantons give rise to sub-Planckian wavelength oscillations in the scalar potential.

In general it is easy to see that beyond these most dominant effective instantons there exists a whole \mathbb{Z}^n lattice worth of these effective instantons³⁷. These correspond to the higher orders in the $|z_i|$, $i = 1, \dots, n$, that we have neglected in Sect. 3. Therefore, the n dominant effective instantons serve as n basis vectors of the lattice \mathbb{Z}^n and a general effective instanton is labeled by an effective charge vector $\vec{k} \in \mathbb{Z}^n$. The bound (61) for these general effective instantons reads

$$S_{\text{eff}}^{\vec{k}} \leq \frac{4}{3} M_{\text{P}}^2 \sum_{i=1}^n |k^i| r_{\text{D3}}^i \left(\frac{q^i}{f_{\text{eff}}} \right)^2, \quad (62)$$

and the r.h. side defines a (1-)norm on \mathbb{Z}^n .

The effective instantons can also be embedded into the full one-dimensional charge lattice \mathbb{Z} of the preceding section via $\vec{k} \mapsto \sum_{i=1}^n k^i q^i$, although they do not satisfy the WGC bound (56).

Again, this is perhaps best understood using the explicit example given in the preceding section. The dominant instantons have charges

$$q^i \in \left\{ \pm \frac{840}{M_i} \right\}_i = \left\{ \pm \frac{840}{8}, \pm \frac{840}{7}, \pm \frac{840}{6}, \pm \frac{840}{5} \right\} = \{\pm 105, \pm 120, \pm 140, \pm 168\}. \quad (63)$$

They induce harmonics in the superpotential with periodicities f_{eff}/q^i . Due to the relation (61) precisely these combinations are restricted to be sub-Planckian as long as $S_{\text{eff}}^i > 1$. They can be understood to occupy a coarse $105\mathbb{Z} + 120\mathbb{Z} + 140\mathbb{Z} + 168\mathbb{Z}$ sub-lattice of \mathbb{Z} while the Euclidean D1-instantons satisfying the WGC occupy an even coarser sub-lattice $840\mathbb{Z} \subset \mathbb{Z}$.

5.4 Axion Phenomenology

We have identified a string theory axion with remarkable properties. It is parametrically lighter than the tower of states that is usually associated to strongly warped regions

³⁷We consider general combinations of holomorphic and anti-holomorphic instantons.

$m_{\text{tower}} \propto w_{\text{IR}} M_{\text{P}}$. The axion mass can be tuned almost independently of the periodicities of the dominant oscillations in the scalar potential, since we have $m \propto w_{\text{IR}}^3 M_{\text{P}}$, while the oscillation period f_{eff}/q of the scalar potential depends only weakly on the warp factor $f_{\text{eff}}/q \sim M_{\text{P}}/\sqrt{\log(w_{\text{IR}}^{-1})}$. Conversely, the mass scales unusually strongly with the oscillation wavelength,

$$\frac{m^2}{M_{\text{P}}^2} \propto w_{\text{IR}}^6 \approx e^{-2S_{\text{eff}}} \approx \exp\left(-\alpha \left(\frac{qM_{\text{P}}}{f_{\text{eff}}}\right)^2\right), \quad (64)$$

with $\alpha = \mathcal{O}(1)$.

In contrast most other stringy axions usually satisfy the relation [6]

$$\frac{m^2}{M_{\text{P}}^2} \sim \exp\left(-\alpha \frac{qM_{\text{P}}}{f_{\text{eff}}}\right), \quad (65)$$

As such the thraXion assumes a rather special place in the string theory landscape. This is potentially interesting for axion phenomenology. We refer the reader to [6] for a range of phenomenological applications for different axion mass scales.

We have to emphasize that at least in the simplest setups our axion is not a generic inflaton candidate because of the generic presence of dominant sub-Planckian wavelength modes in the scalar potential, *despite* the large monodromy enhancement of the effective axion decay constant.

5.5 Uplifting

We would like to briefly comment on some possible scenarios of uplifting. To actually implement these ideas in concrete models involves the complicated interplay of different effects. We highlight the assumptions needed for the following two scenarios in turn. Both of them are based on the idea of adding an oscillating potential of different wavelength to the known thraXion potentials of the form (10) or (55).

Uplifting requires as a precondition, that a full mechanism of Kähler moduli stabilization is in place. Stabilizing the Kähler moduli by definition breaks the no-scale property of GKP-type flux compactifications. Hence, in a full setup of our multi throat system there are two sources of no-scale breaking – the Calabi-Yau breaking potential of our C_2 -axion(s), and also the scalar potential that stabilizes Kähler moduli. In general we expect these two sources of no-scale breaking to mix non-trivially, and we leave a detailed analysis of this for future research. For the rest of this discussion we now assume that these subtleties get resolved for both KKLT and large volume scenario (LVS)-type setups of Kähler moduli

stabilization.

We now wish to look at situations where c -dependent corrections to the Kähler potential may become relevant. This is certainly the case in the regime $|W_0| \sim 1$, leading us to consider LVS-like moduli stabilization [31]. Potentially interesting corrections may arise from Euclidean D1-brane instantons that wrap members of the family of two-spheres that vanish at the tips of the conifold. Since the cycle is trivial in homology we expect no corrections to the superpotential but at most corrections to the Kähler potential of the form

$$\delta e^{-K_1} \sim \mathcal{C} e^{-S_{\text{DBI}} - i S_{\text{CS}}} + c.c., \quad (66)$$

with $\mathcal{C} = \mathcal{O}(1)$ and DBI and CS actions

$$S_{\text{DBI}} = \frac{1}{g_s} \frac{\text{Vol}(S^2)|_{\text{UV}}}{2\pi\alpha'}, \quad \text{and} \quad S_{\text{CS}} = \frac{1}{2\pi\alpha'} \int_{S^2} \left(\sum_p C_p \right) \wedge e^B|_{2\text{-form}} = \text{Re } \mathcal{G}. \quad (67)$$

Here, we have evaluated the DBI action on a representative sphere in the UV, i.e. in the bulk CY. This is because we expect such a representative to give the dominant contribution: As explained in App. B, there are different two-spheres at a given radial coordinate in the throat that are labeled by a $U(1)$ phase and that all share the same volume. As we scan over this phase, the corresponding integrals of C_2 at a given radial coordinate pass through their fundamental domain. Therefore, integrating over all Euclidean brane instantons on the two-spheres should cancel all contributions due to the oscillatory behavior of the correction (66). This is consistent with the fact that after accounting for backreaction of the phases of the throat deformations the C_2 field excursion cannot be measured in the local throats. In the analysis of Sect. 3, we extended this result to $\text{Re}(\mathcal{G})$, i.e. to $C_2 - C_0 B_2$. In passing towards the UV, our description of the throat breaks down. In particular, we do not expect the different sphere representatives to all share the same volume. Thus, we expect non-vanishing instanton corrections.

Using $\text{Vol}(S^2)|_{\text{UV}} \gtrsim R_{\text{throat}}^2 \propto (g_s M K)^{1/2} \alpha'$, this leads to corrections to the scalar potential of the form

$$\delta V \lesssim e^{-\alpha \sqrt{\frac{KM}{g_s}}} (1 - \cos(\text{Re } \mathcal{G})), \quad (68)$$

with $\alpha = \mathcal{O}(1)$. Assuming that the exponentially small prefactors of the classical warping suppressed potential (10) (or that of example 1 of Sect. 5.1) and the non-perturbative correction terms are of the same order, it is feasible that additional local minima appear in




Figure 11: The axion potential of Fig. 10 with an additive correction $\delta V(c) \propto \text{const.} + \cos(c/f_{\text{eff}})$ that shares the periodicity of the superpotential.

the scalar potential that could in principle lift to meta-stable non-supersymmetric minima, possibly even de Sitter vacua.

The exponential terms are of comparable magnitude when

$$\sqrt{\frac{K}{g_s M}} \gtrsim M. \quad (69)$$

In F-theory models with large Euler characteristic we do not see an immediate obstacle to realizing this.

We may turn this around and add large-wavelength corrections to shorter-wavelength oscillations such as those of the example of *drifting monodromies* given in Sect. 5.1. On the large scale of $f_{\text{eff}} \sim MM_{\text{P}}$ there are several Minkowski vacua of the potential (55), compare Fig. 10. It is conceivable that these are uplifted to de Sitter vacua once further corrections to the potential are taken into account. This might happen automatically when the no-scale properties of the Kähler potential (30) get broken by perturbative or non-perturbative corrections, since we know that the existence of Minkowski vacua strongly depends on the cancellation of different terms in the scalar potential. We expect these scalar potential corrections to follow the periodicity of the superpotential, which is given by the super-Planckian decay constant f_{eff} . An optimistic sketch of this is illustrated in Fig. 11. A different approach might be to consider *drifting monodromies* in which we allow for hierarchies between fluxes M_i of individual throats.

We leave a more thorough investigation of these uplifting ideas for future research.

6 Conclusion and Discussion

In this paper we have shown that a novel type of axion-like particle is present in many flux compactifications of type IIB string theory. While its exponentially small mass is due to the existence of strongly red-shifted regions in the compactification manifold (warped throats), it is parametrically lighter than the red-shifted tower of KK-modes that is usually associated with such throats. We would like to emphasize that for fixed value of the axion decay constant the *thraxion* mass is far smaller than all other stringy axions that we are aware of. Moreover, we are able to find explicit models with even parametrically super-Planckian axion decay constants (but with generically dominant sub-Planckian oscillations in the scalar potential). As such the thraxion assumes a rather special place in the string theory landscape that is potentially interesting for axion phenomenology.

The existence of this type of ultra-light axion is intimately linked to so-called conifold transitions between (topologically) distinct CY manifolds. Light thraxions arise whenever fluxes drive a CY close to the transition point. Moreover they come paired with light scalar degrees of freedom (‘saxions’) which are the relevant light degrees of freedom that control the global stabilization of the multi throat system that arises near such a point in moduli space. At the perturbative level the saxion is as light as the axion while non-perturbative Kähler moduli stabilization effects would generically lift this degeneracy. The extremely low (s)axion mass implies that multi throat systems are surprisingly weakly stabilized.

We now summarize the key steps in the derivation of the (s)axion potential. Throughout most of this paper we have focused on the case of a double throat system. As shown in [28] in such a setting there exists a light axion mode that can be thought of as the integral of the RR two-form C_2 over a family of spheres that degenerate at the infrared ends of the two throats. When holding the geometry fixed, a finite field excursion leads to the formation of a flux/anti-flux pair at the two ends, so an appropriately red-shifted potential energy $V \sim w_{\text{IR}}^4$ is induced where w_{IR} is the infrared warp factor of the throats. One of the key observations made in this paper is that when the throats are long there exist *two* light complex scalars z_1, z_2 that control the infrared geometry of the individual throats. This is despite the fact that only a diagonal combination of the two is an actual complex structure modulus of the CY. A finite field excursion of the axion mode drives z_1 and z_2 away from complex structure moduli space, i.e. to $z_1 \neq z_2$. After this geometrical backreaction is accounted for, locally in each throat supersymmetry is almost perfectly restored. We have determined the scale of the remaining scalar potential by finding the higher dimensional field profile that interpolates the 4d modes z_1 and z_2 between the two

throats. It turns out to be dominated by tiny contributions from the bulk CY where the mismatch between the two throats is detected. The final potential energy scales as w_{IR}^6 which is parametrically smaller than the estimate of [28]. Since the axion kinetic term is dominated by contributions from the bulk geometry, the axion mass is of order w_{IR}^3 which is parametrically smaller than the infrared mass-scale of the local throats. Furthermore, the final axion potential turns out to be periodic, and the periodicity is enhanced by a flux number M via monodromy. In other words, the available amount of axion monodromy after backreaction is finite.³⁸ The low scale of the axion potential implies that throughout the compact axion field space the local throats are essentially frozen supersymmetrically.

We have cross-checked and expanded on the 10d double throat calculation in several ways: First, we have shown that a natural proposal for the extension of the GVW flux superpotential leads to an axion potential that matches precisely our 10d conclusions. Moreover, in this framework it is immediate that the axion is paired with a saxion that can be thought of as the integral of the NS two-form B_2 over the same family of spheres. Together they form the complex scalar component of a chiral multiplet. Furthermore we have extended the discussion to the case of a general multi throat system where n three-cycles degenerate near a conifold transition point, with m homology relations among them. There are n ‘local complex structure moduli’ z_i that are relevant for the discussion of which only $n - m$ linear combinations correspond to complex structure moduli space. In this setup there are m families of two-spheres that can be used to define m (s)axions. Again, a finite field excursion of the (s)axions drives the local deformation parameters z_i away from complex structure moduli space and a small periodic potential remains which is of order w_{IR}^6 . We have checked our conclusions also from the perspective of the Klebanov-Strassler gauge theory [39, 32] that is believed to be the gauge theory dual of the throat solution. The (s)axions set a combination of gauge couplings at the UV-ends of the throats and receive a non-perturbative superpotential from gaugino condensation in the infrared. This superpotential again matches precisely the one we have obtained from *classical* 10d/4d supergravity. As is common in the gauge/gravity duality, a classical effect on one side matches a non-perturbative quantum effect on the other.

Our construction can be used to investigate a vast number of axion-models, by scanning

³⁸We note that the local two-node setup of [91] on the deformed side of the geometric transition resembles our setup in the non-compact flux-less limit. Hence, it is tempting to speculate that fluxed (warped) versions of this, as proposed in [21] and further explicated in [62] with its D5- $\overline{\text{D5}}$ (or NS5- $\overline{\text{NS5}}$) system, will show similar backreaction effects as discussed in our paper, which might significantly affect the axion monodromy potential. It is an open challenge to construct the geometry for the wrapped NS5 - $\overline{\text{NS5}}$ -pair at a level such that this question can be addressed explicitly.

over different conifold transition points of CY threefolds and three-form flux quanta. We have illustrated this by giving two examples based on a well-studied conifold transition point of the quintic threefold. With one of these examples we exhibit a mechanism that may be able to generate parametrically super-Planckian axion periodicities. This happens when several throats carry different flux numbers M_i that each give rise to a finite monodromy enhancement. The overall enhancement is given by the least common multiple of all the flux numbers. In the regime considered, the validity of the effective field theory is *not* undermined by the appearance of a large number of light states below or near the axion mass-scale as is usually expected for large f [87, 89, 88] or as parametrically large geodesics in field space are traversed [9, 27, 71–76]. We call this mechanism to generate large axion periodicities *drifting monodromies*. Nevertheless, we identify a global constraint among the flux numbers that presents a serious obstacle to actually realizing a scalar potential with parametrically sub-dominant sub-Planckian oscillations.

Clearly, the examples we have given form only a tiny subset of possible *thra*xion models. Moreover, these models are a promising playground for testing swampland conjectures such as the weak gravity conjecture for axions. The simplest models already suggest that the simplest form of the conjecture need not hold in general. We have briefly commented on thra-xion-related ideas for uplifting to de Sitter minima. However, whether or not these ideas can be realized in a controlled way is left for future work. This would require a detailed understanding of the interplay between the thra-xions and no-scale breaking effects such as gaugino condensation on seven-branes [30] or perturbative α' corrections [92], which are needed for full moduli stabilization [30, 31].

Acknowledgements We would like to thank Federico Carta, Richard Eager, Markus Dierigl, Thomas Grimm, Lukas Hahn, Craig Lawrie, Evan McDonough and Ander Retolaza for useful discussions and comments. This work is supported by the ERC Consolidator Grant STRINGFLATION under the HORIZON 2020 grant agreement no. 647995.

A Deformed Conifold for General Complex Structure Modulus

The unwarped internal geometry before backreaction by fluxes is that of the deformed conifold [50]. It can be embedded in \mathbb{C}^4 via

$$\sum_i w_i^2 = z, \quad (70)$$

where w_i are complex coordinates and z is some complex parameter. For now, we set $z = |z| \in \mathbb{R}^+$. One can check, that the five-dimensional hypersurface at a given fixed radial coordinate $\rho^2 \equiv \sum |w_i|^2 > |z|$ allows for a transitive $SU(2) \times SU(2)$ action³⁹ with isotropy group $U(1) \subset SU(2) \times SU(2)$. The space can therefore be written as the homogeneous space $\frac{SU(2) \times SU(2)}{U(1)}$. At $\rho^2 = |z|$, the isotropy group is enhanced to a diagonal $SU(2) \subset SU(2) \times SU(2)$, therefore leading to the homogeneous space $\frac{SU(2) \times SU(2)}{SU(2)} \sim \frac{SO(4)}{SO(3)} = S^3$. Given that topologically $\frac{SU(2) \times SU(2)}{U(1)} = S^3 \times S^2$, we arrive at the picture of a family of two-spheres along some radial coordinate fibered over a three-sphere, where the two-sphere vanishes or collapses at the apex, while the three-sphere stays finite everywhere.

Following ref. [94], we may derive the most general $SU(2) \times SU(2)$ -invariant metric, that is also invariant under a \mathbb{Z}_2 exchange of the two $SU(2)$ -factors. It takes the form

$$\begin{aligned} ds^2 = & (B^2 + H^2) \left((g^1)^2 + (g^2)^2 \right) + C^2 \left((g^3)^2 + (g^4)^2 \right) \\ & + D^2 \left(d\rho^2 + (g^5)^2 \right) + 2CH (g^1 g^4 - g^2 g^3). \end{aligned} \quad (71)$$

Here, we used the vielbein

$$g^1 = \frac{e^1 - e^3}{\sqrt{2}}, \quad g^2 = \frac{e^2 - e^4}{\sqrt{2}}, \quad g^3 = \frac{e^1 + e^3}{\sqrt{2}}, \quad g^4 = \frac{e^2 + e^4}{\sqrt{2}}, \quad g^5 = e^5, \quad (72)$$

with

$$\begin{aligned} e^1 &= -\sin \theta_1 d\phi_1, \quad e^2 = d\theta_1, \quad e^3 = \cos \psi \sin \theta_2 d\phi_2 - \sin \psi d\theta_2, \\ e^4 &= \sin \psi \sin \theta_2 d\phi_2 + \cos \psi d\theta_2, \quad e^5 = d\psi + \cos \theta_1 d\phi_1 + \cos \theta_2 d\phi_2. \end{aligned} \quad (73)$$

³⁹Acting as $SO(4)$ on the vector $(w_1, w_2, w_3, w_4)^T$ in the obvious way [93]. It is more convenient to parametrize the embedding via the matrix $T = \frac{1}{\sqrt{2}} \sum w_i \sigma_i$, with Pauli matrices σ_i . General solutions to $\det T = -\frac{|z|}{2}$ and $\rho^2 = \text{Tr}(T^\dagger T)$ then take the form $T = LT_0R$, with a specific solution $T_0(|z|, \rho)$ and $(L, R) \in SU(2) \times SU(2)$. The $SU(2) \times SU(2)$ action is now the obvious one $(L, R) \rightarrow (U_L L, U_R R)$ [94].

The real coefficients B , C , D and H are functions of ρ and $|z|$. We impose the constraints on the coefficients from the embedding (70) and insert the definition of the radial coordinate, as well as demand Ricci-flatness of the metric (71). The resulting CY-metric with $SU(2) \times SU(2)$ -isometry is now best expressed in the coordinate $\tau \geq 0$ defined via $\rho^2 = |z| \cosh \tau$ [94],

$$ds_{\text{dc}}^2 = D_{\text{dc}}^2(\tau) \left(d\tau^2 + (g^5)^2 \right) + B_{\text{dc}}^2(\tau) \left((g^1)^2 + (g^2)^2 \right) + C_{\text{dc}}^2(\tau) \left((g^3)^2 + (g^4)^2 \right), \quad (74)$$

with

$$B_{\text{dc}}^2(\tau) = |z|^{\frac{2}{3}} \frac{K(\tau)}{2} \sinh^2(\tau/2), \quad C_{\text{dc}}^2(\tau) = |z|^{\frac{2}{3}} \frac{K(\tau)}{2} \cosh^2(\tau/2), \quad (75)$$

$$D_{\text{dc}}^2(\tau) = |z|^{\frac{2}{3}} \frac{1}{6K(\tau)^2}, \quad \text{where} \quad K(\tau) \equiv \frac{(\sinh(2\tau) - 2\tau)^{\frac{1}{3}}}{2^{\frac{1}{3}} \sinh(\tau)}, \quad (76)$$

and $H_{\text{dc}} \equiv 0$.

We now generalize this metric to arbitrary values of $z = |z| e^{i\varphi} \in \mathbb{C}$. This is most easily done by considering the action of $e^{i\varphi} \in U(1)_R$ on the coordinates

$$w_i \rightarrow w_i e^{i\varphi/2}. \quad (77)$$

Looking at the embedding (70), this gives the rotation $|z| \rightarrow |z| e^{i\varphi}$ we are after. After some calculation, one can reinterpret the rotation of coordinate w_i as an action on the coefficients B , C , D and H . The new coefficients after applying a $U(1)_R$ rotation to second order in φ read

$$\begin{aligned} B &= B_{\text{dc}} + \varphi^2 \frac{B_{\text{dc}} C_{\text{dc}}^2 - B_{\text{dc}}^2}{C_{\text{dc}} 8C_{\text{dc}}}, \\ C &= C_{\text{dc}} + \varphi^2 \frac{B_{\text{dc}}^2 - C_{\text{dc}}^2}{8C_{\text{dc}}}, \\ H &= \varphi \frac{C_{\text{dc}}^2 - B_{\text{dc}}^2}{2C_{\text{dc}}}. \end{aligned} \quad (78)$$

In the full result (i.e. beyond quadratic order in φ) the periodicity $\varphi \rightarrow \varphi + 2\pi$ is of course preserved.

We now want to determine the metric at large $\rho^2 \gg |z|$. For $\varphi = 0$ one may express it

in the form

$$ds^2 = f_0(r)^2 dr^2 + r^2 \left(\frac{1}{9} f_5(r)^2 (g^5)^2 + \frac{1}{6} \sum_{i=1}^4 f_i(r)^2 (g^i)^2 \right), \quad (79)$$

where we have introduced the asymptotic conifold radial coordinate $r = \sqrt{\frac{3}{2}} \rho^{2/3}$. Defining $\epsilon(r) \equiv \frac{|z|}{r^3}$, at order ϵ^0 one has $f_i \equiv 1$ which corresponds to the metric of the singular conifold. However, also far away from the deformation $\epsilon \ll 1$, we see some remnant of the apex geometry

$$\begin{aligned} f_0^2(r) = f_5^2(r) &= \frac{9}{r^2} D^2(\tau(r)) = 1 + \mathcal{O}(\epsilon^2), \\ f_1^2(r) = f_2^2(r) &= \frac{6}{r^2} B^2(\tau(r)) = 1 + \left(\frac{3}{2}\right)^{3/2} \epsilon + \mathcal{O}(\epsilon^2), \\ f_3^2(r) = f_4^2(r) &= \frac{6}{r^2} C^2(\tau(r)) = 1 - \left(\frac{3}{2}\right)^{3/2} \epsilon + \mathcal{O}(\epsilon^2). \end{aligned} \quad (80)$$

Neglecting numerical $\mathcal{O}(1)$ -factors, we arrive at

$$ds^2 = dr^2 + r^2 \left\{ d\Omega_{T^{1,1}}^2 + \epsilon(r) \left[(g^1)^2 + (g^2)^2 - (g^3)^2 - (g^4)^2 \right] \right\} + \mathcal{O}(\epsilon^2). \quad (81)$$

Using the transformation behaviour of the coefficients of the angular terms (78), it is now straightforward to write down the same asymptotic expansion for non-zero φ

$$\begin{aligned} ds^2 &= dr^2 + r^2 \left\{ d\Omega_{T^{1,1}}^2 + \epsilon(r) d\Omega_5^2(\varphi) \right\}. \\ d\Omega_5^2(\varphi) &\equiv (1 + \varphi^2) \left((g^1)^2 + (g^2)^2 - (g^3)^2 - (g^4)^2 \right) + \varphi (g^1 g^4 - g^2 g^3) \end{aligned} \quad (82)$$

Finally, we consider the full 10d metric found as a solution to 10d SUGRA with M units of F_3 -flux on the three-sphere described above and some H_3 -flux (unquantized in the non-compact setting) on the dual cycle [32]

$$ds^2 = w^2(r) \eta_{\mu\nu} dx^\mu dx^\nu + w^{-2}(r) ds_{dc}^2, \quad (83)$$

where the warp factor goes like

$$w(r)^2 = \frac{r^2}{g_s M \alpha'} \frac{1}{\sqrt{\log\left(\frac{r^3}{|z|}\right)}}, \quad (84)$$

for $\frac{r^3}{|z|} \gg 1$. For arbitrary φ , we will work with the asymptotic metric

$$ds^2 = w^2(r)\eta_{\mu\nu}dx^\mu dx^\nu + w^{-2}(r) [dr^2 + r^2 (d\Omega_{T^{1,1}}^2 + \epsilon(r)d\Omega_5(\varphi)^2)] . \quad (85)$$

B The Axion Potential in the Local Throat

In the main text we have repeatedly made use of the fact that the C_2 - and B_2 -axions c and b can only enter the scalar potential that is generated in the local throat in certain combinations with the ‘local complex structure’ of the throat, namely the real and imaginary part of

$$M \log(z) - i\mathcal{G} , \quad (86)$$

where $\mathcal{G} = c - \tau b$, and z is the ‘local complex structure’. Here, we would like to derive this without using ‘local flux stabilization’ as in Sect. 2.3 but rather rely only on asymptotic properties of the KS/KT solution [39, 32]. For simplicity we will set the RR zero form to zero, i.e. $\tau = ig_s^{-1}$.

We cut off the throat at a radial coordinate r_{UV} and define the b -axion at that value of the radial coordinate. Since the B_2 profile runs along the radial direction [95], changing $b \rightarrow b + \delta b$ can be realized by choosing a *different* UV-cutoff r'_{UV} . Since the absolute value of the complex structure is defined in units of the UV-cutoff it scales as

$$z \rightarrow e^{-\frac{\delta b}{g_s M}} z . \quad (87)$$

Since this is just a coordinate transformation from the perspective of the local KS throat, the combination $g_s M \log(|z|) + b$ cannot appear in the scalar potential that is generated within the throat. It acquires physical meaning only if the throat is cut off at fixed, finite r_{UV} .

Similarly, in the limit $|z|/r^3 \rightarrow 0$ the RR three form takes the form

$$F_3 = 2\pi\alpha' M (g^5 + dc/M) \wedge \omega_\Sigma , \quad (88)$$

where $g^5 = d\psi + \dots$ is given by (72), and ω_Σ is the normalized harmonic 2-form of $T^{1,1}$. The field $c(x)$ transforms like a Goldstone boson under (local) coordinate transformations [80]

$$\psi \rightarrow \psi + 2\omega(x) , \quad c(x) \rightarrow c(x) - 2M\omega(x) . \quad (89)$$

Shifting along this angular direction is an isometry of the asymptotic KS solution (called $U(1)_R$). Near the IR this is not the case precisely because (by definition) the phase of the complex structure also transforms like a Goldstone boson, see App. A

$$\arg z \longrightarrow \arg z - 2\omega. \quad (90)$$

Again, in the local throat the combination $M \arg z + c$ has no physical meaning as it is eaten via the Higgs mechanism. Only when the throat is glued into the CY space at finite radial coordinate r_{UV} does the c -axion gain its independent physical meaning because the $U(1)_R$ symmetry is badly broken by the CY geometry. Putting together the real and imaginary part of $\mathcal{G} = c - \tau b$, we arrive at the conclusion that only the combination (86) is physical when considering a single throat. A second degree of freedom only becomes physical by finiteness of the throat, i.e. by breaking the asymptotic $U(1)_R$ symmetry. This also implies that the kinetic terms for the fields $\varphi_{1,2}$ stated in (13) actually take the form $[\partial(\varphi_{1,2} \pm c/M)]^2$ since they arise from local throat physics. We have disregarded some unimportant off-diagonal terms in the kinetic matrix.

The transformation behavior of $c(x)$ under a $U(1)_R$ can also be calculated directly [96]. For this, we choose some S^2 -submanifold of the cross-section $T^{1,1}$ of the deformed conifold far in the UV and apply the $U(1)_R$ action on the coordinates as in App. A. One can show that the manifold X defined by the $U(1)_R$ -orbit of S^2 -submanifolds is an element of $H^3(T^{1,1})$, and therefore a multiple of the S^3 -cycle of $T^{1,1}$. It turns out that the multiplicity is 2. Denoting a member of the family of two-spheres as $S^2(\omega)$, $\omega \in U(1)_R$, we therefore find by applying Stokes' theorem

$$\int_{S^2(0)} C_2 - \int_{S^2(2\pi)} C_2 = \int_X F_3 = 2 \int_{S^3} F_3 = 2(2\pi)^2 \alpha' M. \quad (91)$$

Using homogeneity of the flux distribution we extrapolate $\int_{S^2(0)} C_2 - \int_{S^2(\omega)} C_2 = 4\pi \alpha' M \omega$. In terms of c defined on (say) $S^2(0)$, we arrive at the transformation law as stated above. Interpreting $\int_{S^2} C_2$ as a generalized Wilson line, we find the usual behavior under deforming the integration path over some region with non-vanishing associated field-strength F_3 . Under $\pi \in U(1)_R$, the family $S^2(\omega)$ sweeps out S^3 completely; accordingly the Wilson line picks up the flux $\int_{S^3} F_3 = (2\pi)^2 \alpha' M$.

C Derivation and Solution of the 5d Equations of Motion

C.1 The 5d Action for φ

In App. A we explained how the phase φ of the complex structure modulus z enters the metric to leading order in $|z|$. In this subsection we now want to explicitly derive how we arrive at the action (11) from this. The 10d metric (85) can be decomposed into the external and radial five-dimensional metric and the internal angular metric

$$\begin{aligned} G_{MN} &= (w^2 \eta_{\mu\nu} \oplus w^{-2} g_{rr}) \oplus (w^{-2} r^2 g_{\Omega, mn}) , \quad m, n = \phi_1, \phi_2, \theta_1, \theta_2, \psi , \\ g_{\Omega, mn} &= g_{T^{1,1}, mn} + \epsilon(r) h_{mn} , \quad g_{T^{1,1}, mn} dy^m dy^n \equiv d\Omega_{T^{1,1}}^2 , \quad h_{mn} dy^m dy^n = d\Omega_5^2(\varphi) , \end{aligned} \quad (92)$$

where $\epsilon(r) = |z|/r^3$ is the $U(1)_R$ -breaking parameter. As explained in Sect. 2.3, we promote φ to a dynamical field $\varphi = \varphi(x^\mu, r)$. While this field extends into multiple throats, we make use of its antisymmetry in order to reduce the problem to a single throat. The corresponding boundary conditions are given below.

We expand the 5d action for φ in ϵ via the expansion of the 10d Ricci scalar about the $U(1)_R$ -preserving metric

$$\begin{aligned} R(G) &= R(G_{U(1)_R}) + \epsilon(r) \nabla_m \nabla_n h^{mn} - \square(\epsilon(r) g_{T^{1,1}}^{mn} h_{mn}) + \mathcal{O}(\epsilon^2) , \\ G_{U(1)_R} &= w^2 \eta \oplus w^{-2} g_{rr} \oplus w^{-2} r^2 g_{T^{1,1}} , \\ G_{5d} &= w^2 \eta \oplus w^{-2} g_{rr} . \end{aligned} \quad (93)$$

We know that for $\varphi = \text{const.}$ we simply describe a deformed conifold with general complex deformation parameter $z = |z| e^{i\varphi}$, which is by construction Ricci-flat. Therefore, we have $R(G) = 0$ in every order in ϵ , such that the only non-vanishing contribution can come from derivative terms of φ

$$\begin{aligned} \square(\epsilon(r) g_{T^{1,1}}^{mn} h_{mn}) &= \epsilon(r) g_{T^{1,1}}^{mn} G_{5d}^{ij} \partial_i \partial_j h_{mn}(\varphi(x^\mu, r)) \\ &= \epsilon(r) g_{T^{1,1}}^{mn} [\partial_\varphi h_{mn} G_{5d}^{ij} \partial_i \partial_j \varphi + \partial_\varphi^2 h_{mn} G_{5d}^{ij} \partial_i \varphi \partial_j \varphi] . \end{aligned} \quad (94)$$

We can write

$$g_{T^{1,1}, mn} dx^m dx^n = \frac{1}{6} \left((g^1)^2 + (g^2)^2 + \left((g^3)^2 + (g^4)^2 \right) \right) + \frac{1}{9} (g^5)^2 . \quad (95)$$

With this and looking at the perturbation h_{mn} above, one finds that to order ϵ , there is no contribution to the 5d action. This is due to the fact that there is no non-zero contraction $g_{T^{1,1}}^{mn} \partial_\varphi h_{mn}$ or $g_{T^{1,1}}^{mn} \partial_\varphi^2 h_{mn}$. There is no term $g^1 g^4 - g^2 g^3$ in $g_{T^{1,1}}$ and the terms coming from contractions of $(g^1)^2 + (g^2)^2$ and $(g^3)^2 + (g^4)^2$ cancel due to the different signs between these terms in $g_{T^{1,1}}$ and h ⁴⁰. The first non-vanishing order is ϵ^2 .

Using $\sqrt{G} \approx \sqrt{G_{U(1)_R}} = w^{-2} r^5$, the following terms may now arise up to order φ^2 after integrating out the angular directions in the 10d action

$$S_{5d} = \frac{1}{g_s^2 \kappa_{10}^2} \int d^4x \int dr r^5 w(r)^{-2} \left(R_{5d} + \Lambda(r) - \frac{1}{2} \epsilon(r)^2 G_{5d}^{ij} \partial_i \varphi \partial_j \varphi \right). \quad (96)$$

Here, we have implemented 5d Lorentz symmetry and used the fact, that there are no linear terms in φ once we go to order ϵ^2 . Importantly there are no non-derivative terms for φ , because $\varphi = \text{const.}$ is a point in complex moduli space. Dropping 5d gravity, we arrive at

$$S_{5d} = \frac{1}{g_s^2 \kappa_{10}^2} \int d^4x \int_{r_{\text{IR}}}^{r_{\text{UV}}} dr \epsilon(r)^2 r^5 \left(-\frac{1}{2} w(r)^{-4} (\partial_\mu \varphi)^2 - \frac{1}{2} (\partial_r \varphi)^2 \right). \quad (97)$$

Explicitly calculating $R(G)$ with $\varphi = \varphi(x^\mu, t)$ and expanding to first order in $|z|$ and second order in ϵ confirms this explicitly.

C.2 Schrödinger Equations and Exact Solutions for Free Fields

Using the previous subsection and the results of [28], we collect the two actions⁴¹ that will give us all relevant parameters. Note that we derive the action for φ for $c = 0$ and that the one for c for $\varphi = 0$. We deal with the form of interaction terms in App. B.

$$\begin{aligned} S_\varphi &= \frac{1}{g_s^2 \alpha'^4} \int d^4x \int \frac{dr}{r} \epsilon(r)^2 r^6 \left(-\frac{1}{2} w(r)^{-4} (\partial_\mu \phi_\varphi)^2 - \frac{1}{2} (\partial_r \phi_\varphi)^2 \right), \\ S_c &= \frac{1}{\alpha'^2} \int d^4x \int \frac{dr}{r} r^2 w(r)^4 \left(-\frac{1}{2} w(r)^{-4} (\partial_\mu \phi_c)^2 - \frac{1}{2} (\partial_r \phi_c)^2 \right), \end{aligned} \quad (98)$$

⁴⁰In other words: The term in $g_{T^{1,1}}$ is symmetric under the exchange of the $SU(2)$'s defining the cone, $(\phi_1, \theta_1) \leftrightarrow (\phi_2, \theta_2)$ in coordinates, while the term in h is antisymmetric.

⁴¹For ease of exposition we work with only two real fields although strictly speaking they are both real components of complex fields (see Sect. 3).

where we use a general notation with fields $\phi_i(x^\mu, r)$, $i = \varphi, c$. The equations of motion using the plane wave ansatz $\phi_i(x^\mu, r) = e^{ip_i x} \chi_i(r)$, with $p_i^2 = -m_i^2$, read

$$\begin{aligned} w(r)^4 \epsilon(r)^{-2} r^{-5} \partial_r (\epsilon(r)^2 r^5 \partial_r \chi_\varphi(r)) &= -m_\varphi^2 \chi_\varphi(r), \\ r^{-1} \partial_r (r w^4 \partial_r \chi_c(r)) &= -m_c^2 \chi_c(r). \end{aligned} \quad (99)$$

We insert the warp-factor (84), dropping logarithmic dependencies, and the $U(1)_R$ -breaking factor to arrive at

$$\begin{aligned} r^5 \partial_r \left(\frac{1}{r} \partial_r \chi_\varphi(r) \right) &= -R^4 m_\varphi^2 \chi_\varphi(r), \\ \frac{1}{r} \partial_r (r^5 \partial_r \chi_c(r)) &= -R^4 m_c^2 \chi_c(r), \end{aligned} \quad (100)$$

where we introduced the abbreviation $R = \sqrt{g_s M \alpha'}$ for the normalization of the warp factor of KT and KS. Using the reparametrization

$$\begin{aligned} x_\varphi &= \frac{R^2 m_\varphi}{r}, & g_\varphi(x_\varphi) &= \frac{1}{r} \chi_\varphi(r(x_\varphi)), \\ x_c &= \frac{R^2 m_c}{r}, & g_c(x_c) &= r^2 \chi_c(r(x_c)), \end{aligned} \quad (101)$$

we arrive at the following form of the equations of motion

$$x_i^2 g_i''(x_i) + x_i g_i'(x_i) + (x_i^2 - \alpha_i^2) g_i(x_i) = 0, \quad (102)$$

where $i = \varphi, c$ and where $\alpha_\varphi = 1$ and $\alpha_c = 2$. This is the standard form of the Bessel equation. Plugging back in our reparametrization, we write down the general solutions to the equations of motion

$$\begin{aligned} \chi_\varphi(r) &= r \left(A J_1 \left(\frac{R^2 m_\varphi}{r} \right) + B Y_1 \left(\frac{R^2 m_\varphi}{r} \right) \right), \\ \chi_c(r) &= \frac{1}{r^2} \left(A J_2 \left(\frac{R^2 m_c}{r} \right) + B Y_2 \left(\frac{R^2 m_c}{r} \right) \right), \end{aligned} \quad (103)$$

with Bessel functions J_{α_i} and Y_{α_i} of first and second kind respectively.

We now have to apply the boundary conditions

$$\begin{aligned} \chi_\varphi(r_{\text{UV}}) &= 0, & \partial_r \chi_\varphi(r_{\text{IR}}) &= 0, \\ \partial_r \chi_c(r_{\text{UV}}) &= 0, & w_{\text{IR}} \partial_r \chi_c(r_{\text{IR}}) &= \frac{\delta}{R} \chi_c(r_{\text{IR}}), \end{aligned} \tag{104}$$

for some $\mathcal{O}(1)$ -number δ depending on how we model the flux distribution in the IR. While the UV boundary conditions are easily implemented to fix the constants A and B up to an overall normalization, the ansatz $m_i \propto \frac{w_{\text{IR}}^2}{R}$ gives consistent solutions to the IR boundary conditions when using the small argument expansions of the Bessel functions

$$\begin{aligned} m_\varphi^2 &= 8 \frac{w_{\text{IR}}^4}{R^2}, \\ m_c^2 &= \frac{8\delta}{4 + \delta} \frac{w_{\text{IR}}^4}{R^2}, \end{aligned} \tag{105}$$

where we now normalized the warp factor $w_{\text{UV}} = 1$. We plot the solutions with $\delta = 1$, $w_{\text{IR}} = e^{-10/3} \sim 10^{-2}$ and m_i as above in Fig. 12.




Figure 12: The lowest lying radial eigenmodes for φ and c for $w_{\text{IR}} \sim 10^{-2}$ plotted in the physical radial coordinate $t = 3R \log(r/r_{\text{IR}})$.

Referring back to the discussion in Sect. 2.3, we plot the expected solutions over the entirety of the double throat in Fig. 13, where we simply mirrored one throat at the UV end with appropriate symmetry, so both $r = r_{\text{IR}}$ and $r = 2r_{\text{UV}} - r_{\text{IR}}$ correspond to IR regions.

We are left with calculating the 4d kinetic terms. From Fig. 12, we would guess (correctly) that φ is an IR mode, while c is a UV mode. We derive this by considering the lowest KK mode $\phi_0(x^\mu, r) = f_0(x^\mu) \chi_0(r)$, where f_0 is now an arbitrary superposition of




Figure 13: The eigenmodes for φ and c on the double throat. Here, $t < 10$ corresponds to the first throat, $t > 10$ corresponds to the second, with $t = 0$ and $t = 10$ corresponding to the respective ends of the throats.

plane waves, to find the kinetic terms induced by the actions (98)

$$\begin{aligned}
 f_\varphi^2 &= \frac{1}{g_s^2 \alpha'^4} \int_{r_{\text{IR}}}^{r_{\text{UV}}} \frac{dr}{r} \epsilon(r)^2 r^6 w(r)^{-4} \chi_\varphi(r)^2 \approx \frac{R^6}{g_s^2 \alpha'^4} w_{\text{IR}}^2, \\
 f_c^2 &= \frac{1}{\alpha'^2} \int_{r_{\text{IR}}}^{r_{\text{UV}}} \frac{dr}{r} r^2 \chi_c(r)^2 \approx \frac{R^2}{\alpha'^2}.
 \end{aligned}
 \tag{106}$$

Here, we have again expanded the solution in $w_{\text{IR}} \ll 1$, and neglected logarithmic corrections. Furthermore we have normalized the internal field profiles such that $\chi_c(r_{\text{UV}}) = \chi_\varphi(r_{\text{IR}}) = 1$.

Comparing with the quadratic terms in c in the potential (7) and the quadratic terms in φ_i in the potential (9), we find the parameters μ and Λ

$$\mu^4 = m_c^2 f_c^2 \propto w_{\text{IR}}^4, \quad \Lambda^4 = m_\varphi^2 f_\varphi^2 \propto w_{\text{IR}}^6.
 \tag{107}$$

D Axion Decay Constant and Potential Parameters

In this appendix we extend the results of the previous App. C by including logarithmic corrections to the first-order approximations and by taking care of the different length scales that may appear in the parameters (14) of the effective potential (10).

We compute the axion decay constant (without accounting for monodromy factors) for the c -axion from the 10d action, paying attention to numerical factors of $\mathcal{O}(1)$. We may dimensionally reduce the $|F_3|^2$ term of the 10d type IIB SUGRA and plug in $C_2 = 2\pi\alpha'c(x)\omega_\Sigma$, where ω_Σ is the associated (quasi-)harmonic form. This leads to

$$\frac{1}{(2\pi)^7 \alpha'^4} \left(-\frac{1}{2} \int *|F_3|^2 \right) = \int d^4x \left(-\frac{1}{2} f^2 (\partial c)^2 \right),
 \tag{108}$$

with

$$f^2 = (2\pi)^2 g_s^2 \frac{\int_{\text{CY}} d^6 y \sqrt{g_6} w^2 \alpha'^2 |\omega_\Sigma|^2}{\int_{\text{CY}} d^6 y \sqrt{g_6} w^2} M_{\text{P}}^2 \equiv (2\pi)^2 g_s^2 \frac{\alpha'^2}{(\text{Vol}(S^2)|_{\text{UV}})^2} M_{\text{P}}^2, \quad (109)$$

where g_s is the string coupling, g_6 is the internal 6d metric, w^2 is the warp factor and M_{P} is the four-dimensional Planck mass. Due to the appearance of the warp factor, the integrals are dominated by UV contributions. Thus, for large values of the overall Kähler modulus $\text{Re } T \gg N_{\text{D3}}$ we have $f^2 \sim g_s (\text{Re } T)^{-1}$ in Planck units⁴², where N_{D3} is the total D3 brane charge stored in fluxes (and mobile D3 branes). This is consistent with the 4d SUGRA of Sect. 3. We expect to be allowed to make the string frame volume as small as $\mathcal{O}((g_s N_{\text{D3}})^{3/2})$. At even smaller volume backreaction of fluxes becomes significant throughout the CY and we lose the notion of an unwarped bulk CY [29]. For simplicity we now take into consideration only the D3 brane charge stored in a single throat, i.e. $N_{\text{D3}} = MK$. In this regime the top of the throat marginally fits into the bulk CY and we expect the ratio of integrals in (109) to be well approximated by the value of $\alpha'^2 |\omega_\Sigma|^2$ at the top of the KS throat. Since the hierarchy is set by $\log w_{\text{IR}}^{-1} \sim \frac{K}{g_s M}$ it follows immediately that f scales as $f \sim \sqrt{\frac{g_s}{MK}} M_{\text{P}} \sim \frac{1}{M} \log(w_{\text{IR}}^{-1})^{-1/2} M_{\text{P}}$.

We can compute this in more detail by using the asymptotic form of the throat metric far away from the deformation, i.e. the KT metric (4)

$$ds_6^2 = w^{-2} (dr^2 + r^2 ds_{T^{1,1}}^2), \quad ds_{T^{1,1}}^2 = \frac{1}{9} (g^5)^2 + \frac{1}{6} \sum_{i=1}^4 (g^i)^2, \quad (110)$$

$$w^2 = \frac{2\sqrt{2}}{9} \frac{r^2}{g_s M \alpha'} \log(r/r_{\text{IR}})^{-\frac{1}{2}}, \quad \omega_\Sigma = \frac{1}{8\pi} (g^1 \wedge g^2 + g^3 \wedge g^4). \quad (111)$$

We see, that the physical radius at the top of the throat is given by

$$R_{\text{throat}}^2 = \frac{9}{2\sqrt{2}} g_s M \alpha' \sqrt{\log r_{\text{UV}}/r_{\text{IR}}} \approx \frac{9}{2\sqrt{2}} g_s M \alpha' \sqrt{\log w_{\text{IR}}^{-1}}, \quad (112)$$

where in the last step we use the normalized warp factor with $w_{\text{UV}} = 1$. Taking into account angular dependencies of $T^{1,1}$, we arrive at $\alpha' |\omega_\Sigma| = (3\pi g_s M)^{-1} \log(w_{\text{IR}}^{-1})^{-\frac{1}{2}}$, and hence

$$f \approx \frac{2}{3M} \log(w_{\text{IR}}^{-1})^{-1/2} M_{\text{P}}. \quad (113)$$

Actually, we are interested in the case where two or more throats each store only part of the total D3 brane charge and also fluxes on other cycles contribute to N_{D3} . It is useful

⁴²Recall that $\text{Re } T$ measures a 4-cycle volume in 10d Einstein frame.

to define as r_{D3}^i the fraction of the total D3 brane charge that is stored in the i -th throat, i.e.

$$r_{\text{D3}}^i \equiv M_i K^i / N_{\text{D3}}, \quad i = 1, \dots, n. \quad (114)$$

Since the decay constant scales as $(\text{Re } T)^{-1/2}$ and the marginal case corresponds to $\text{Re } T \sim N_{\text{D3}} = M_i K^i / r_{\text{D3}}^i$, for any $i = 1, \dots, n$, we should really replace (113) by

$$f \approx \frac{2}{3M_i} (r_{\text{D3}}^i)^{1/2} \log((w_{\text{IR}}^i)^{-1})^{-1/2} M_{\text{P}}, \quad (115)$$

and this expression is correct when evaluated for any $i = 1, \dots, n$.

In the double throat case, we have $r_{\text{D3}}^1 = r_{\text{D3}}^2 \equiv r_{\text{D3}}$ and the effective (monodromy-) potential (10) reads

$$V(c)/M_{\text{P}}^4 \sim w_{\text{IR}}^6 \left(1 - \cos \left(\frac{3}{(r_{\text{D3}})^{1/2}} \sqrt{\log(1/w_{\text{IR}})} \frac{c}{M_{\text{P}}} \right) \right). \quad (116)$$

Even in this marginal case, the decay constant of c is slightly sub-Planckian in regimes of parametric control since then $w_{\text{IR}} \ll 1$, and $r_{\text{D3}} < 1$.

We may also derive the quantities given in (14): Using (98) the kinetic term of φ is given by

$$f_\varphi^2 \sim \frac{(g_s M \alpha')^6}{g_s^2 \alpha'^4} w_{\text{IR}}^2 \sim \frac{(g_s M)^3}{\mathcal{V}} w_{\text{IR}}^2 M_{\text{P}}^2, \quad (117)$$

where we use that the integral for the kinetic term (106) is dominated by the IR, such that we may insert the IR throat radius, $R^2 \sim g_s M \alpha'$, and where we have made use of the fact that $M_{\text{P}}^2 \sim g_s^{-2} \alpha'^{-1} \mathcal{V}$, with the (bulk) CY volume \mathcal{V} as measured in string units. Neglecting geometric back-reaction of the double throat system in the UV, the flux energy density $M^2 \mu^4$ can be understood as a pure IR effect: It is induced by an excursion of $\varphi_{1,2}$ in the IR whilst keeping c fixed (for the factor of M^2 compare (7)). Therefore the resulting mass must be of the order of the warped KK-scale $m_{\text{wKK}}^2 \sim R^{-2} w_{\text{IR}}^2 \sim (g_s M \alpha')^{-1} w_{\text{IR}}^2$

$$\mu^4 \sim \frac{1}{M^2} f_\varphi^2 m_{\text{wKK}}^2 \sim \frac{g_s^4}{\mathcal{V}^2} w_{\text{IR}}^4 M_{\text{P}}^4. \quad (118)$$

Finally, the scale Λ^4 is dominated by contributions from the bulk CY (which we again

assume to have only one length-scale, $R_{\text{CY}}^2 \sim \mathcal{V}^{1/3} \alpha'$), so

$$\begin{aligned} \Lambda^4 &\sim \underbrace{\frac{1}{g_s^2} \frac{1}{\alpha'^4}}_{\sim M_{\text{P},10\text{d}}^8} \underbrace{\mathcal{V} \alpha'^3}_{\sim \int d^6 y \sqrt{g_{\text{CY}}}} \underbrace{\epsilon^2(r_{\text{UV}})}_{\sim \text{UV tail of deformation}} \underbrace{\frac{1}{\mathcal{V}^{1/3} \alpha'}}_{\sim \mathcal{L}_{10\text{d}} \supset (\nabla \varphi)^2 \propto 1/R_{\text{CY}}^2} \\ &\sim \frac{g_s^2}{\mathcal{V}^{4/3}} \epsilon^2(r_{\text{UV}}) M_{\text{P}}^4 \sim \frac{g_s^2}{\mathcal{V}^{4/3}} \log(w_{\text{IR}}^{-1})^{-3/2} w_{\text{IR}}^6 M_{\text{P}}^4. \end{aligned} \quad (119)$$

Here, we have made use of the fact that $\epsilon^2(r_{\text{UV}}) \sim (r_{\text{IR}}/r_{\text{UV}})^6 \sim w_{\text{IR}}^6 \log(w_{\text{IR}}^{-1})^{-3/2}$. The scale Λ^4 determines the axion potential and can be compared with the SUGRA result. Using the Kähler potential (30) one easily shows that the parametric dependencies come out correctly.

Going to the limit where the throats marginally fit into the bulk CY means taking $\mathcal{V} \sim (g_s N_{\text{D3}})^{3/2} \sim (r_{\text{D3}})^{3/2} (g_s M)^3 \log(w_{\text{IR}}^{-1})^{3/2}$. In this limit,

$$\begin{aligned} \Lambda^4/M_{\text{P}}^4 &\sim (r_{\text{D3}})^{-2} \frac{g_s^2}{(g_s M)^4} \log(w_{\text{IR}}^{-1})^{-7/2} w_{\text{IR}}^6, \quad \mu^4/M_{\text{P}}^4 \sim (r_{\text{D3}})^{-3} \frac{g_s^4}{(g_s M)^6} \log(w_{\text{IR}}^{-1})^{-3} w_{\text{IR}}^4, \\ f_\varphi^2/M_{\text{P}}^2 &\sim (r_{\text{D3}})^{-3/2} \log(w_{\text{IR}}^{-1})^{-3/2} w_{\text{IR}}^2. \end{aligned} \quad (120)$$

For the double throat case considered in Sect. 2, we may assume a configuration where almost all the D3 brane charge resides in the two throats, and therefore $r_{\text{D3}} \approx 1/2$.

E Background on Multi Conifolds

In this appendix we discuss preliminaries that are important for Sect. 3. We follow mainly Chapter 8 of [69].

For a general CY threefold M we can choose $2h^{2,1}+2$ three-cycles $\mathcal{A}^a, \mathcal{B}_a, a = 1, \dots, h^{2,1}+1$ as a symplectic basis of $H^3(M)$, i.e.

$$\int_{\mathcal{A}^b} \alpha^a = \int_M \alpha^a \wedge \beta_b = \delta_b^a, \quad \int_{\mathcal{B}_b} \beta_a = \int_M \beta_a \wedge \alpha^b = -\delta_a^b, \quad (121)$$

where α^a and β_a are the harmonic three-forms that are Poincaré dual to \mathcal{B}_a , respectively \mathcal{A}^a . One may define the periods

$$z_a = \int_{\mathcal{A}^a} \Omega, \quad G^a = \int_{\mathcal{B}_a} \Omega, \quad (122)$$

where Ω is the holomorphic three-form. The z_a form a set of projective coordinates on complex structure moduli space and the G^a are functions of them. We are interested in what happens when n cycles γ^i with m homology relations among them shrink at a conifold point in moduli space. The Picard-Lefschetz formula states that upon encircling a conifold point in moduli space, a three-cycle δ undergoes the monodromy [97, 98, 68]

$$\delta \longrightarrow \delta + \sum_{i=1}^n (\delta \cap \gamma^i) \gamma^i. \quad (123)$$

Knowing this monodromy transformation is enough to determine that

$$\int_{\delta} \Omega = \frac{1}{2\pi i} \sum_{i=1}^n (\delta \cap \gamma^i) \int_{\gamma^i} \Omega \log\left(\int_{\gamma^i} \Omega\right) + \text{single-valued}. \quad (124)$$

We may choose $n - m$ of the degenerating cycles as part of the basis $\mathcal{A}^i = \gamma^i$ for $i = 1, \dots, n - m$, while the remaining m are integer linear combinations $\gamma^i = \sum_{a=1}^{n-m} c_a^i \mathcal{A}^a$ for $i = n - m + 1, \dots, n$. By applying (123) to the cycles \mathcal{B}_a we arrive at

$$G^a = \int_{\mathcal{B}_a} \Omega = \frac{1}{2\pi i} z_a \log(z_a) + \frac{1}{2\pi i} \sum_{i=n-m+1}^n c_a^i z_i \log(z_i) + g^a(z), \quad a = 1, \dots, n - m, \quad (125)$$

where $g^a(z)$ are $n - m$ holomorphic functions. We have defined $z_i \equiv \sum_{a=1}^{n-m} c_a^i z_a$ for $i = n - m + 1, \dots, n$, i.e. $z_i \equiv \int_{\gamma^i} \Omega$ when applying (123)⁴³. At frozen values of z^a , $a = n - m + 1, \dots, h^{2,1} + 1$ the periods associated to other cycles, $G^a = \int_{\mathcal{B}_a} \Omega$, with $a = n - m + 1, \dots, h^{2,1} + 1$, are holomorphic in the complex structures that parametrize the multi conifold deformations, i.e. in the z^i , with $i = 1, \dots, n - m$. In what follows we denote by z^a only the multi conifold deformation parameters.

We may now evaluate the GVW superpotential $W = \int_M G_3 \wedge \Omega$ where we choose flux quanta M_a and K^a according to $G_3 = -\sum_{a=1}^{n-m} (M_a \alpha^a - \tau K^a \beta_a)$. Using that

$$\int_M \alpha^a \wedge \Omega = -\int_{\mathcal{B}_a} \Omega = -G^a, \quad \int_M \beta_a \wedge \Omega = -\int_{\mathcal{A}^a} \Omega = -z_a, \quad (126)$$

⁴³When using a local expression for the holomorphic three-form Ω in the vicinity of smoothed conical singularity described by (1) one can calculate $\int_{\gamma^i} \Omega = z_i$ [85]. This identifies the z_i defined here with the local deformation parameter of the i -th throat.

one obtains

$$\begin{aligned}
W(z_a) &= \sum_{a=1}^{n-m} \frac{M_a}{2\pi i} z_a \log(z_a) + \sum_{i=n-m+1}^n \frac{M_i}{2\pi i} z_i \log(z_i) \\
&\quad + \sum_{a=1}^{n-m} M_a g^a(z) - \tau \sum_{a=1}^{n-m} K^a z_a + \hat{W}_0(z_a),
\end{aligned} \tag{127}$$

where we have defined $M_i \equiv \sum_{a=1}^{n-m} c_a^i M_a$, and the holomorphic function $\hat{W}_0(z_a)$ parametrizes the contributions from fluxes on other cycles. We may use the z_a and z_i with $i = n - m + 1, \dots, n$ on the same footing by interpreting our definition of the z_i as m constraint equations

$$0 = P^I \equiv \sum_{i=1}^n p_i^I z_i \equiv z_{n-m+I} - \sum_{a=1}^{n-m} c_a^{n-m+I} z_a, \quad I = 1, \dots, m. \tag{128}$$

Here, the $m \times n$ matrix p_i^I is implicitly defined as

$$p_i^I = \begin{cases} -c_i^{n-m+I}, & i = 1, \dots, n-m, \\ \delta_i^{n-m+I}, & i = n-m+1, \dots, n. \end{cases} \tag{129}$$

We may now write the superpotential as

$$W(z_i) = \sum_{i=1}^n \left(M_i \frac{z_i}{2\pi i} \log(z_i) + M_i g^i(z) - \tau K^i z_i \right) + \sum_{I=1}^m \lambda_I P^I + \hat{W}_0(z_i), \tag{130}$$

with m Lagrange multipliers λ_I . The homology relation is now enforced via the F-term of the fields λ_I , compare [85] where Lagrange multipliers in the superpotential are also used to impose additional constraints on chiral superfields. In doing so, we have defined g^i to be zero for $i > n - m$.

The F_3 -flux on γ^i is given by M_i . By the definition of M_i for $i = n - m + 1, \dots, n$, the flux numbers automatically fulfill $\sum_{i=1}^n p_i^I M_i = 0$ for all I . In democratic terms, the n flux numbers M_i must be chosen in compliance with the m homology constraints $\sum_{i=1}^n p_i^I M_i = 0$. The H_3 -flux on \mathcal{B}_a is given by $K^a + \sum_{I=1}^m c_a^{n-m+I} K^{n-m+I}$, as this is the coefficient appearing in front of z_a . In other words the $n - m$ flux quantization conditions read $K^a + \sum_{I=1}^m c_a^{n-m+I} K^{n-m+I} \in \mathbb{Z}$. Note that we may transform $K^i \rightarrow K^i + \sum_I \alpha_I p_i^I$ for any $\alpha \in \mathbb{C}^m$ because the superpotential is left invariant upon imposing the constraint

equations, that is to say, we can undo such a transformation by also shifting the Lagrange multipliers $\lambda^I \rightarrow \lambda^I + \tau\alpha^I$. Of course, the flux quantization conditions are invariant under these shifts. Finally, the Kähler potential is given by

$$K_{\text{cs}}(z_i, \bar{z}_i) = -\log\left(-i \int \Omega \wedge \bar{\Omega}\right) = -\log\left(ig_K(z) - i\overline{g_K}(z) + \sum_{a=1}^{n-m} i\bar{z}_a G^a(z) + c.c.\right) \quad (131)$$

$$= -\log\left(ig_K(z) - i\overline{g_K}(z) + \sum_{i=1}^n \left[\frac{|z_i|^2}{2\pi} \log(|z_i|^2) + i\bar{z}_i g^i(z) - iz_i \overline{g^i}(z)\right]\right), \quad (132)$$

where $g_K = \sum_{a=n-m+1}^{h^{2,1}+1} \bar{z}_a G^a(z)$ encodes contributions from other cycles. It is holomorphic in z_a , $a = 1, \dots, n - m$. Note that despite the democratic formulation, the Kähler and superpotential are strictly defined only along complex structure moduli space, where $P^I = 0$. As explained in Sect. 3 we propose to extend the domain of these functions to the deformation space parametrized by *all* z_i by introducing general Taylor expansions of $g^i(z_i)$, $g_K(z_i)$ and $\hat{W}_0(z_i)$ in (31).

References

- [1] P. Sikivie, *Axion Cosmology, Lect. Notes Phys.* **741** (2008) 19 [[astro-ph/0610440](#)].
- [2] J. Jaeckel and A. Ringwald, *The Low-Energy Frontier of Particle Physics, Ann. Rev. Nucl. Part. Sci.* **60** (2010) 405 [[1002.0329](#)].
- [3] J. P. Conlon, *The QCD axion and moduli stabilisation, JHEP* **05** (2006) 078 [[hep-th/0602233](#)].
- [4] P. Svrcek and E. Witten, *Axions In String Theory, JHEP* **06** (2006) 051 [[hep-th/0605206](#)].
- [5] T. W. Grimm, *Axion inflation in type II string theory, Phys. Rev.* **D77** (2008) 126007 [[0710.3883](#)].
- [6] A. Arvanitaki, S. Dimopoulos, S. Dubovsky, N. Kaloper and J. March-Russell, *String Axiverse, Phys. Rev.* **D81** (2010) 123530 [[0905.4720](#)].
- [7] M. Cicoli, M. Goodsell and A. Ringwald, *The type IIB string axiverse and its low-energy phenomenology, JHEP* **10** (2012) 146 [[1206.0819](#)].

- [8] C. Vafa, *The String landscape and the swampland*, hep-th/0509212.
- [9] H. Ooguri and C. Vafa, *On the Geometry of the String Landscape and the Swampland*, *Nucl. Phys.* **B766** (2007) 21 [hep-th/0605264].
- [10] N. Arkani-Hamed, L. Motl, A. Nicolis and C. Vafa, *The String landscape, black holes and gravity as the weakest force*, *JHEP* **06** (2007) 060 [hep-th/0601001].
- [11] A. de la Fuente, P. Saraswat and R. Sundrum, *Natural Inflation and Quantum Gravity*, *Phys. Rev. Lett.* **114** (2015) 151303 [1412.3457].
- [12] T. Rudelius, *Constraints on Axion Inflation from the Weak Gravity Conjecture*, *JCAP* **1509** (2015) 020 [1503.00795].
- [13] M. Montero, A. M. Uranga and I. Valenzuela, *Transplanckian axions!?*, *JHEP* **08** (2015) 032 [1503.03886].
- [14] J. Brown, W. Cottrell, G. Shiu and P. Soler, *Fencing in the Swampland: Quantum Gravity Constraints on Large Field Inflation*, *JHEP* **10** (2015) 023 [1503.04783].
- [15] T. C. Bachlechner, C. Long and L. McAllister, *Planckian Axions and the Weak Gravity Conjecture*, *JHEP* **01** (2016) 091 [1503.07853].
- [16] A. Hebecker, P. Mangat, F. Rompineve and L. T. Witkowski, *Winding out of the Swamp: Evading the Weak Gravity Conjecture with F-term Winding Inflation?*, *Phys. Lett.* **B748** (2015) 455 [1503.07912].
- [17] B. Heidenreich, M. Reece and T. Rudelius, *Weak Gravity Strongly Constrains Large-Field Axion Inflation*, *JHEP* **12** (2015) 108 [1506.03447].
- [18] D. Junghans, *Large-Field Inflation with Multiple Axions and the Weak Gravity Conjecture*, *JHEP* **02** (2016) 128 [1504.03566].
- [19] K. Kooner, S. Parameswaran and I. Zavala, *Warping the Weak Gravity Conjecture*, *Phys. Lett.* **B759** (2016) 402 [1509.07049].
- [20] E. Silverstein and A. Westphal, *Monodromy in the CMB: Gravity Waves and String Inflation*, *Phys. Rev.* **D78** (2008) 106003 [0803.3085].
- [21] L. McAllister, E. Silverstein and A. Westphal, *Gravity Waves and Linear Inflation from Axion Monodromy*, *Phys. Rev.* **D82** (2010) 046003 [0808.0706].

- [22] F. Marchesano, G. Shiu and A. M. Uranga, *F-term Axion Monodromy Inflation*, *JHEP* **09** (2014) 184 [1404.3040].
- [23] R. Blumenhagen and E. Plauschinn, *Towards Universal Axion Inflation and Reheating in String Theory*, *Phys. Lett.* **B736** (2014) 482 [1404.3542].
- [24] A. Hebecker, S. C. Kraus and L. T. Witkowski, *D7-Brane Chaotic Inflation*, *Phys. Lett.* **B737** (2014) 16 [1404.3711].
- [25] L. E. Ibanez, M. Montero, A. Uranga and I. Valenzuela, *Relaxion Monodromy and the Weak Gravity Conjecture*, *JHEP* **04** (2016) 020 [1512.00025].
- [26] A. Hebecker, F. Rompineve and A. Westphal, *Axion Monodromy and the Weak Gravity Conjecture*, *JHEP* **04** (2016) 157 [1512.03768].
- [27] D. Klaewer and E. Palti, *Super-Planckian Spatial Field Variations and Quantum Gravity*, *JHEP* **01** (2017) 088 [1610.00010].
- [28] A. Hebecker, J. Moritz, A. Westphal and L. T. Witkowski, *Towards Axion Monodromy Inflation with Warped KK-Modes*, *Phys. Lett.* **B754** (2016) 328 [1512.04463].
- [29] S. B. Giddings, S. Kachru and J. Polchinski, *Hierarchies from fluxes in string compactifications*, *Phys. Rev.* **D66** (2002) 106006 [hep-th/0105097].
- [30] S. Kachru, R. Kallosh, A. D. Linde and S. P. Trivedi, *De Sitter vacua in string theory*, *Phys. Rev.* **D68** (2003) 046005 [hep-th/0301240].
- [31] V. Balasubramanian, P. Berglund, J. P. Conlon and F. Quevedo, *Systematics of moduli stabilisation in Calabi-Yau flux compactifications*, *JHEP* **03** (2005) 007 [hep-th/0502058].
- [32] I. R. Klebanov and M. J. Strassler, *Supergravity and a confining gauge theory: Duality cascades and chi SB resolution of naked singularities*, *JHEP* **08** (2000) 052 [hep-th/0007191].
- [33] S. Ashok and M. R. Douglas, *Counting flux vacua*, *JHEP* **01** (2004) 060 [hep-th/0307049].

- [34] F. Denef and M. R. Douglas, *Distributions of flux vacua*, *JHEP* **05** (2004) 072 [hep-th/0404116].
- [35] A. Hebecker and J. March-Russell, *The Ubiquitous throat*, *Nucl. Phys.* **B781** (2007) 99 [hep-th/0607120].
- [36] K. Dasgupta, H. Firouzjahi and R. Gwyn, *On The Warped Heterotic Axion*, *JHEP* **06** (2008) 056 [0803.3828].
- [37] S. Franco, D. Galloni, A. Retolaza and A. Uranga, *On axion monodromy inflation in warped throats*, *JHEP* **02** (2015) 086 [1405.7044].
- [38] S. Gukov, C. Vafa and E. Witten, *CFT's from Calabi-Yau four folds*, *Nucl. Phys.* **B584** (2000) 69 [hep-th/9906070].
- [39] I. R. Klebanov and A. A. Tseytlin, *Gravity duals of supersymmetric $SU(N) \times SU(N+M)$ gauge theories*, *Nucl. Phys.* **B578** (2000) 123 [hep-th/0002159].
- [40] J. E. Kim, H. P. Nilles and M. Peloso, *Completing natural inflation*, *JCAP* **0501** (2005) 005 [hep-ph/0409138].
- [41] M. Berg, E. Pajer and S. Sjors, *Dante's Inferno*, *Phys. Rev.* **D81** (2010) 103535 [0912.1341].
- [42] E. Palti, *On Natural Inflation and Moduli Stabilisation in String Theory*, *JHEP* **10** (2015) 188 [1508.00009].
- [43] K. Choi and S. H. Im, *Realizing the relaxion from multiple axions and its UV completion with high scale supersymmetry*, *JHEP* **01** (2016) 149 [1511.00132].
- [44] D. E. Kaplan and R. Rattazzi, *Large field excursions and approximate discrete symmetries from a clockwork axion*, *Phys. Rev.* **D93** (2016) 085007 [1511.01827].
- [45] R. Blumenhagen, D. Herschmann and F. Wolf, *String Moduli Stabilization at the Conifold*, *JHEP* **08** (2016) 110 [1605.06299].
- [46] G. Buratti, J. Calderón and A. M. Uranga, *Transplanckian Axion Monodromy !?*, 1812.05016.

- [47] A. Hebecker, D. Junghans and A. Schachner, *The Higgs and other Thorns in the side of the Swampland*, talk given by A. Hebecker at ‘Vistas over the Swampland Program’, Madrid, Spain, Sept. 19-21, 2018.
- [48] A. Hebecker, D. Junghans and A. Schachner, *Large Field Ranges from Aligned and Misaligned Winding*, 1812.05626.
- [49] P. Candelas, P. S. Green and T. Hubsch, *Rolling Among Calabi-Yau Vacua*, *Nucl. Phys.* **B330** (1990) 49.
- [50] P. Candelas and X. C. de la Ossa, *Comments on Conifolds*, *Nucl. Phys.* **B342** (1990) 246.
- [51] P. Candelas, P. S. Green and T. Hubsch, *Finite Distances Between Distinct Calabi-yau Vacua: (Other Worlds Are Just Around the Corner)*, *Phys. Rev. Lett.* **62** (1989) 1956.
- [52] P. Candelas, X. C. De La Ossa, P. S. Green and L. Parkes, *A Pair of Calabi-Yau manifolds as an exactly soluble superconformal theory*, *Nucl. Phys.* **B359** (1991) 21.
- [53] M. Reid, *The Moduli Space of 3-Folds with $K = 0$ may Nevertheless be Irreducible.*, *Mathematische Annalen* **278** (1987) 329.
- [54] T.-m. Chiang, B. R. Greene, M. Gross and Y. Kanter, *Black hole condensation and the web of Calabi-Yau manifolds*, *Nucl. Phys. Proc. Suppl.* **46** (1996) 82 [hep-th/9511204].
- [55] A. C. Avram, M. Kreuzer, M. Mandelberg and H. Skarke, *The Web of Calabi-Yau hypersurfaces in toric varieties*, *Nucl. Phys.* **B505** (1997) 625 [hep-th/9703003].
- [56] V. V. Batyrev, *Dual polyhedra and mirror symmetry for Calabi-Yau hypersurfaces in toric varieties*, *J. Alg. Geom.* **3** (1994) 493 [alg-geom/9310003].
- [57] V. Batyrev and M. Kreuzer, *Constructing new Calabi-Yau 3-folds and their mirrors via conifold transitions*, *Adv. Theor. Math. Phys.* **14** (2010) 879 [0802.3376].
- [58] M. Lynker and R. Schimmrigk, *Conifold transitions and mirror symmetries*, *Nucl. Phys.* **B484** (1997) 562 [hep-th/9511058].

- [59] V. V. Batyrev, I. Ciocan-Fontanine, B. Kim and D. van Straten, *Conifold transitions and mirror symmetry for Calabi-Yau complete intersections in Grassmannians*, *Nucl. Phys.* **B514** (1998) 640.
- [60] T. W. Grimm and A. Klemm, *$U(1)$ Mediation of Flux Supersymmetry Breaking*, *JHEP* **10** (2008) 077 [0805.3361].
- [61] I. R. Klebanov and E. Witten, *Superconformal field theory on three-branes at a Calabi-Yau singularity*, *Nucl. Phys.* **B536** (1998) 199 [hep-th/9807080].
- [62] A. Retolaza, A. M. Uranga and A. Westphal, *Bifid Throats for Axion Monodromy Inflation*, *JHEP* **07** (2015) 099 [1504.02103].
- [63] I. Bena, E. Dudas, M. Graña and S. Lüster, *Uplifting Runaways*, 1809.06861.
- [64] L. Randall and R. Sundrum, *A Large mass hierarchy from a small extra dimension*, *Phys. Rev. Lett.* **83** (1999) 3370 [hep-ph/9905221].
- [65] D. Baumann, A. Dymarsky, S. Kachru, I. R. Klebanov and L. McAllister, *$D3$ -brane Potentials from Fluxes in AdS/CFT*, *JHEP* **06** (2010) 072 [1001.5028].
- [66] F. Carta, J. Moritz and A. Westphal, *Gaugino condensation and small uplifts in KKL T* , 1902.01412.
- [67] A. Strominger, *Massless black holes and conifolds in string theory*, *Nucl. Phys.* **B451** (1995) 96 [hep-th/9504090].
- [68] B. R. Greene, D. R. Morrison and A. Strominger, *Black hole condensation and the unification of string vacua*, *Nucl. Phys.* **B451** (1995) 109 [hep-th/9504145].
- [69] B. R. Greene, *String theory on Calabi-Yau manifolds*, in *Fields, strings and duality. Proceedings, Summer School, Theoretical Advanced Study Institute in Elementary Particle Physics, TASI'96, Boulder, USA, June 2-28, 1996*, pp. 543–726, 1996, hep-th/9702155.
- [70] T. W. Grimm and J. Louis, *The Effective action of $N = 1$ Calabi-Yau orientifolds*, *Nucl. Phys.* **B699** (2004) 387 [hep-th/0403067].
- [71] R. Blumenhagen, I. Valenzuela and F. Wolf, *The Swampland Conjecture and F -term Axion Monodromy Inflation*, *JHEP* **07** (2017) 145 [1703.05776].

- [72] T. W. Grimm, E. Palti and I. Valenzuela, *Infinite Distances in Field Space and Massless Towers of States*, *JHEP* **08** (2018) 143 [1802.08264].
- [73] B. Heidenreich, M. Reece and T. Rudelius, *Emergence of Weak Coupling at Large Distance in Quantum Gravity*, *Phys. Rev. Lett.* **121** (2018) 051601 [1802.08698].
- [74] S.-J. Lee, W. Lerche and T. Weigand, *Tensionless Strings and the Weak Gravity Conjecture*, *JHEP* **10** (2018) 164 [1808.05958].
- [75] S.-J. Lee, W. Lerche and T. Weigand, *A Stringy Test of the Scalar Weak Gravity Conjecture*, *Nucl. Phys.* **B938** (2019) 321 [1810.05169].
- [76] T. W. Grimm, C. Li and E. Palti, *Infinite Distance Networks in Field Space and Charge Orbits*, 1811.02571.
- [77] S. Kachru, J. Pearson and H. L. Verlinde, *Brane / flux annihilation and the string dual of a nonsupersymmetric field theory*, *JHEP* **06** (2002) 021 [hep-th/0112197].
- [78] I. Antoniadis, J.-P. Derendinger, P. M. Petropoulos and K. Siampos, *Heisenberg symmetry and hypermultiplet manifolds*, *Nucl. Phys.* **B905** (2016) 293 [1512.06418].
- [79] P. Corvilain, T. W. Grimm and D. Regalado, *Shift-symmetries and gauge coupling functions in orientifolds and F-theory*, *JHEP* **05** (2017) 059 [1607.03897].
- [80] C. P. Herzog, I. R. Klebanov and P. Ouyang, *D-branes on the conifold and $N=1$ gauge / gravity dualities*, in *Progress in string, field and particle theory: Proceedings, NATO Advanced Study Institute, EC Summer School, Cargese, France, June 25-July 11, 2002*, pp. 189–223, 2002, hep-th/0205100.
- [81] G. Veneziano and S. Yankielowicz, *An Effective Lagrangian for the Pure $N=1$ Supersymmetric Yang-Mills Theory*, *Phys. Lett.* **113B** (1982) 231.
- [82] I. Affleck, M. Dine and N. Seiberg, *Dynamical Supersymmetry Breaking in Supersymmetric QCD*, *Nucl. Phys.* **B241** (1984) 493.
- [83] J. J. Heckman, J. Seo and C. Vafa, *Phase Structure of a Brane/Anti-Brane System at Large N* , *JHEP* **07** (2007) 073 [hep-th/0702077].
- [84] J. J. Heckman and C. Vafa, *Geometrically Induced Phase Transitions at Large N* , *JHEP* **04** (2008) 052 [0707.4011].

- [85] F. Cachazo, K. A. Intriligator and C. Vafa, *A Large N duality via a geometric transition*, *Nucl. Phys.* **B603** (2001) 3 [[hep-th/0103067](#)].
- [86] J. Moritz and T. Van Riet, *Racing through the swampland: de Sitter uplift vs weak gravity*, *JHEP* **09** (2018) 099 [[1805.00944](#)].
- [87] T. Banks, M. Dine, P. J. Fox and E. Gorbatov, *On the possibility of large axion decay constants*, *JCAP* **0306** (2003) 001 [[hep-th/0303252](#)].
- [88] B. Heidenreich, M. Reece and T. Rudelius, *Evidence for a sublattice weak gravity conjecture*, *JHEP* **08** (2017) 025 [[1606.08437](#)].
- [89] B. Heidenreich, M. Reece and T. Rudelius, *Sharpening the Weak Gravity Conjecture with Dimensional Reduction*, *JHEP* **02** (2016) 140 [[1509.06374](#)].
- [90] S. Andriolo, D. Junghans, T. Noumi and G. Shiu, *A Tower Weak Gravity Conjecture from Infrared Consistency*, *Fortsch. Phys.* **66** (2018) 1800020 [[1802.04287](#)].
- [91] M. Aganagic, C. Beem, J. Seo and C. Vafa, *Geometrically Induced Metastability and Holography*, *Nucl. Phys.* **B789** (2008) 382 [[hep-th/0610249](#)].
- [92] K. Becker, M. Becker, M. Haack and J. Louis, *Supersymmetry breaking and alpha-prime corrections to flux induced potentials*, *JHEP* **06** (2002) 060 [[hep-th/0204254](#)].
- [93] C. P. Herzog, I. R. Klebanov and P. Ouyang, *Remarks on the warped deformed conifold*, in *Modern Trends in String Theory: 2nd Lisbon School on g Theory Superstrings Lisbon, Portugal, July 13-17, 2001*, 2001, [hep-th/0108101](#).
- [94] R. Minasian and D. Tsimpis, *On the geometry of nontrivially embedded branes*, *Nucl. Phys.* **B572** (2000) 499 [[hep-th/9911042](#)].
- [95] I. R. Klebanov and N. A. Nekrasov, *Gravity duals of fractional branes and logarithmic RG flow*, *Nucl. Phys.* **B574** (2000) 263 [[hep-th/9911096](#)].
- [96] S. Leonhardt, *Axion Monodromy in the Klebanov-Strassler Background*, Master's thesis, Heidelberg University, 2017.
- [97] E. Picard and G. Simart, *Théorie des fonctions algébriques de deux variables indépendantes*, no. Bd. 1 in PCMI collection. Gauthier-Villars, 1897.

[98] S. Lefschetz, *L'analysis situs et la géométrie algébrique*. Gauthier-Villars, Paris, 1950.