

Open charm production at high energies and the quark Reggeization hypothesis

B.A. Kniehl* and A.V. Shipilova†

*II. Institut für Theoretische Physik, Universität Hamburg,
Luruper Chaussee 149, 22761 Hamburg, Germany*

V.A. Saleev‡

Samara State University, Academic Pavlov Street 1, 443011 Samara, Russia

Abstract


We study open charm production at high energies in the framework of the quasi-multi-Regge-kinematics approach applying the quark-Reggeization hypothesis implemented with Reggeon-Reggeon-particle and Reggeon-particle-particle effective vertices. Adopting the Kimber-Martin-Ryskin unintegrated quark and gluon distribution functions of the proton and photon, we thus nicely describe the proton structure function $F_{2,c}$ measured at DESY HERA as well as the transverse-momentum distributions of D mesons created by photoproduction at HERA and by hadroproduction at the Fermilab Tevatron.

PACS numbers: 12.39.St, 12.40.Nn, 13.60.Hb, 13.85.Ni

*Electronic address: kniehl@desy.de

†Electronic address: alexshipilova@ssu.samara.ru

‡Electronic address: saleev@ssu.samara.ru



I. INTRODUCTION

The study of open charm production in high-energy lepton-hadron and hadron-hadron collisions is considered as a test of the general applicability of perturbative quantum chromodynamics (QCD) and also provides information on the parton distribution functions (PDFs) of protons and photons. The present analysis is to explore our potential to access a new dynamical regime, namely the high-energy Regge limit, which is characterized by the condition $\sqrt{S} \gg \mu \gg \Lambda_{\text{QCD}}$, where \sqrt{S} is the total collision energy in the center-of-mass (CM) reference frame, Λ_{QCD} is the asymptotic scale parameter of QCD, and μ is the typical energy scale of the hard interaction. In the processes of heavy-quark production, one has $\mu \geq m$, where m is the heavy-quark mass. In this high-energy regime, the contribution from partonic subprocesses involving t -channel parton (quark or gluon) exchanges to the production cross section can become dominant. Thus, the transverse momenta of the incoming partons and their off-shell properties can no longer be neglected, and we deal with *Reggeized* t -channel partons.

The quasi-multi-Regge-kinematics (QMRK) approach [1, 2] is particularly appropriate for this kind of high-energy phenomenology. It is based on an effective quantum field theory implemented with the non-Abelian gauge-invariant action, as suggested a few years ago [3]. Our previous analyses of charmonium and bottomonium production at the Fermilab Tevatron [4] demonstrated the advantages of the high-energy factorization scheme over the collinear parton model as far as the description of experimental data is concerned. These observations were substantiated for B -meson production at the Tevatron in Ref. [5], where the experimental data were again well described using the Fadin-Lipatov effective Reggeon-Reggeon-gluon vertex [2]. In Ref. [6], where the effective photon-Reggeon-quark vertex was obtained and for the first time, the hypothesis of quark Reggeization was successfully used to describe experimental data on single prompt-photon production and on the proton structure functions F_2 and F_L .

The CDF Collaboration measured the differential cross sections $d\sigma/dp_T$ for the inclusive production of D^0 , D^\pm , $D^{*\pm}$, and D_s^\pm mesons in $p\bar{p}$ collisions in run II at the Tevatron as functions of transverse momentum ($p_T = |\vec{p}_T|$) in the central rapidity (y) region [7]. These measurements were compared with theoretical predictions obtained at next-to-leading order (NLO) in the collinear parton model of QCD [8, 9] taking into account quark and hadron

mass effects, and it was found that the latter improve the description of the data.

The differential cross sections $d\sigma/dp_T$ and $d\sigma/dy$ for inclusive $D^{*\pm}$ and D_s^\pm photoproduction measured by the H1 [10] and ZEUS [11, 12] collaborations at the DESY HERA Collider were compared with NLO predictions in the collinear parton model. For $D^{*\pm}$ mesons, this was done in three approaches: the zero-mass variable-flavor-number scheme (ZM-VFNS) [13, 14], the fixed-flavor-number scheme (FFNS) [15], and the general-mass variable-flavor-number scheme (GM-VFNS) [16]. The experimental results were found to generally lie above the NLO expectations. For D_s^\pm mesons, the calculations were performed in the FFNS [15] and in the model suggested by Berezhnoy, Kiselev, and Likhoded [17].

In this paper, we study D -meson production under HERA and Tevatron experimental conditions as well as the charm structure function $F_{2,c}$ of the proton for the first time in the framework of the QMRK approach [1, 2] complemented with the quark-Reggeization hypothesis. This paper is organized as follows. In Sec. II, we present the basic formalism of our calculations and briefly recall the QMRK approach in connection with the quark-Reggeization hypothesis. In Sec. III, we consider the charm structure function $F_{2,c}$ of the proton and compare our results with experimental data. In Secs. IV and V, we describe D -meson production via c -quark fragmentation at HERA and the Tevatron, respectively. In Sec. VI, we summarize our conclusions.

II. BASIC FORMALISM

In the phenomenology of the strong interactions at high energies, it is necessary to describe the QCD evolution of the PDFs of the colliding particles (hadrons or photons) starting from some scale μ_0 which controls a non-perturbative regime up to the typical scale μ of the hard-scattering processes, which is typically of the order of the transverse mass $M_T = \sqrt{M^2 + p_T^2}$ of the produced particle (or hadron jet) with (invariant) mass M and transverse momentum \vec{p}_T . In the region of very high energies, which corresponds to the so-called Regge limit, the typical ratio $x = \mu/\sqrt{S}$ becomes very small, $x \ll 1$. This leads to large logarithmic contributions of the type $[\alpha_s \ln(1/x)]^n$, where α_s is the strong-coupling constant, which are conveniently resummed in the Balitsky-Fadin-Kuraev-Lipatov [18] formalism by the evolution of unintegrated gluon and quark distribution functions $\Phi_{g,q}^{p,\gamma}(x, q_T^2, \mu^2)$, where x and \vec{q}_T are the longitudinal-momentum fraction and transverse momentum of the

Reggeized parton w.r.t. the parent particle, respectively. Correspondingly, in the QMRK approach [1, 2], the initial-state t -channel gluons and quarks are considered as Reggeons, or Reggeized gluons (R) and quarks (Q). They carry finite transverse momenta \vec{q}_T with respect to the hadron or photon beam from which they stem and are off mass shell.

The advantages of the QMRK approach in comparison with the conventional k_T -factorization scheme [19] include: firstly, it uses gauge-invariant amplitudes and is based on a factorization hypothesis that is proven in the leading logarithmic approximation; secondly, it carries over to non-leading orders in the strong-coupling constant, as recently proven [20]. The Reggeization of amplitudes provides the opportunity to efficiently take into account large radiative corrections to processes in the Regge limit beyond what is included in the collinear approximation, which is of great practical importance.

Recently, the Feynman rules for the induced and some important effective vertices of the effective theory based on the non-Abelian gauge-invariant action [3] have been derived in Ref. [21]. However, these rules only refer to processes with Reggeized gluons in the initial state. As for t -channel quark-exchange processes, such rules are still unknown, so that it is necessary to construct effective vertices involving Reggeized quarks using QMRK approach prescriptions in each application from first principles. Of course, a certain set of Reggeon-Reggeon-Particle effective vertices are known, for example for the transitions $RR \rightarrow g$ [22], $Q\bar{Q} \rightarrow g$ [23], and $RQ \rightarrow q$ [24]. The effective $\gamma^*Q \rightarrow q$ vertex, which describes the production of a quark in the collision of a virtual photon with a Reggeized quark, has been recently obtained in Ref. [6].

In our numerical calculations below, we adopt the prescription proposed by Kimber, Martin, Ryskin, and Watt [25] to obtain unintegrated gluon and quark distribution functions for the proton from the conventional integrated ones, as implemented in Watt's code [26]. To obtain the analogous unintegrated functions for the photon, we modify Watt's code [26]. As input for this procedure, we use the Martin-Roberts-Stirling-Thorne [27] proton and the Glück-Reya-Vogt [28] photon PDFs.

III. CHARM STRUCTURE FUNCTION $F_{2,c}$ OF THE PROTON

On the experimental side, the charm structure function $F_{2,c}$ of the proton was measured by H1 [29] and ZEUS [30] in deep inelastic scattering (DIS) of electrons and positrons on

protons at HERA. In this section, we consider this quantity in the framework of the QMRK approach endowed with the quark-Reggeization hypothesis. We thus need the partonic cross section for the production of a c quark in the collision of a virtual photon and a Reggeized charm quark. The relevant vertex was found in Ref. [6] and reads:

$$C_{\gamma Q}^q = -ee_q \left[\frac{q_1^2}{q_1^2 + q_2^2} \gamma^\mu - \frac{2k^\mu}{q_1^2 + q_2^2} \not{q}_2 + \frac{2x_2 q_2^2 P_2^\mu}{(q_1^2 + q_2^2)^2} \not{q}_2 \right], \quad (1)$$

where the four-momenta of the virtual photon, the proton, the Reggeized charm quark, and the outgoing charm quark are denoted as q_1 , P_2 , $q_2 = x_2 P_2 + q_{2T}$, and $k = q_1 + q_2$, respectively. We concentrate on photons with large virtuality $Q^2 = -q_1^2 \gg m_c^2$, so that the massless approximation for describing DIS structure functions is appropriate [6]. We then obtain the following master formula for $F_{2,c}$:

$$F_{2,c}(x_B, Q^2) = 2e_c^2 \int_0^{Q^2} dt_2 \Phi_c^p(x_2, t_2, \mu^2) \frac{Q^2(Q^4 + 6Q^2 t_2 + 2t_2^2)}{(Q^2 + t_2)^3}, \quad (2)$$

where $e_c = 2/3$ is the fractional electric charge of the c quark and $x_2 = x_B(Q^2 + t_2)/Q^2$, with x_B being the Bjorken variable. For definiteness, we choose the factorization scale to be $\mu^2 = Q^2$.

In Fig. 1, we compare the x_B distributions of $F_{2,c}$ for various values of Q^2 with the H1 [29] and ZEUS [30] data. We find good agreement for all values of Q^2 , except for the highest one, $Q^2 = 500 \text{ GeV}^2$, where our prediction is about 50% below the data. This disagreement shows the importance of higher-order corrections at large values of Q^2 , which are beyond the scope of our present study.

IV. D-MESON PRODUCTION AT HERA

On the experimental side, ZEUS measured the p_T distributions of $D^{*\pm}$ [11] and D_s^\pm [12] mesons with rapidity [34] $|y| \leq 1.5$ inclusively produced in photoproduction at HERA I, with proton energy $E_p = 820 \text{ GeV}$ and lepton energy $E_e = 27.5 \text{ GeV}$ in the laboratory frame, in the ranges $2 \leq p_T \leq 12 \text{ GeV}$ and $3 \leq p_T \leq 12 \text{ GeV}$, respectively. In this section, we compare this data with our QMRK predictions. At leading order (LO), we need to consider only three $2 \rightarrow 1$ partonic subprocesses, namely $C_p \gamma \rightarrow c$ for direct photoproduction and $C_p R_\gamma \rightarrow c$ and $R_p C_\gamma \rightarrow c$ for resolved photoproduction, where the subscript indicates the mother particle.

Exploiting the factorization theorem, the p_T distribution of direct photoproduction takes the form

$$p_T^3 \frac{d\sigma}{dp_T} = 2\pi \int dy \int dz x_\gamma f_{\gamma/e}(x_\gamma) z^2 D_{c \rightarrow D}(z, \mu^2) \times \Phi_c^p(x_1, t_1, \mu^2) \overline{|M(C_p \gamma \rightarrow c)|^2}, \quad (3)$$

where

$$x_1 = \frac{p_T e^y}{2zE_p}, \quad x_\gamma = \frac{p_T e^{-y}}{2zE_e}, \quad t_1 = k_T^2, \quad \vec{k}_T = \frac{\vec{p}_T}{z}, \quad (4)$$

with \vec{k}_T being the transverse momentum of the produced c quark. We evaluate the quasi-real-photon flux $f_{\gamma/e}$ in Weizsäcker-Williams approximation using

$$f(x_\gamma) = \frac{\alpha}{2\pi} \left[\frac{1 + (1 - x_\gamma)^2}{x_\gamma} \ln \frac{Q_{\max}^2}{Q_{\min}^2} + 2m_e^2 x_\gamma \left(\frac{1}{Q_{\min}^2} - \frac{1}{Q_{\max}^2} \right) \right], \quad (5)$$

where α is Sommerfeld's fine-structure constant, m_e is the electron mass, $Q_{\min}^2 = m_e^2 x_\gamma^2 / (1 - x_\gamma)$, and Q_{\max}^2 is determined by the experimental setup, with $Q_{\max}^2 = 1 \text{ GeV}^2$ in our case [11, 12]. As for the $c \rightarrow D$ fragmentation function (FF) $D_{c \rightarrow D}$, we adopt the non-perturbative $D^{*\pm}$ and D_s^\pm sets determined in the ZM-VFNS with initial evolution scale $\mu_0 = m_c$ [31] from fits to OPAL data from CERN LEP1. We choose the renormalization and initial- and final-state factorization scales to be $\mu = \sqrt{m_D^2 + p_T^2}$, where m_D is the D -meson mass. Using the Reggeized-quark-photon effective vertex from Ref. [6], the square of the hard-scattering amplitude is found to be

$$\overline{|M(C\gamma \rightarrow c)|^2} = 8\pi\alpha e_c^2 k_T^2. \quad (6)$$

It is understood that also the contribution from the charge-conjugate partonic subprocess is to be included in Eq. (3).

In the case of resolved photoproduction via the partonic subprocess $C_p R_\gamma \rightarrow c$, the factorization formula reads:

$$p_T^3 \frac{d\sigma}{dp_T} = \int dy \int dz \int dx_\gamma \int dt_2 \int d\phi_2 f_{\gamma/e}(x_\gamma) z^2 D_{c \rightarrow D}(z, \mu^2) \times \Phi_c^p(x_1, t_1, \mu^2) \Phi_g^\gamma(x_2, t_2, \mu^2) \overline{|M(C_p R_\gamma \rightarrow c)|^2}, \quad (7)$$

where

$$x_1 = \frac{p_T e^y}{2zE_p}, \quad x_2 = \frac{p_T e^{-y}}{2x_\gamma z E_e}, \quad t_1 = t_2 - 2k_T \sqrt{t_2} \cos \phi_2 + k_T^2, \quad t_2 = q_{2T}^2, \quad \vec{k}_T = \frac{\vec{p}_T}{z}, \quad (8)$$

with ϕ_2 being the angle enclosed between \vec{p}_T and \vec{q}_{2T} . Using the Reggeized-quark–Reggeized-gluon effective vertex from Ref. [24], we have

$$\overline{|M(CR \rightarrow c)|^2} = \frac{2}{3}\pi\alpha_s(\mu^2)k_T^2. \quad (9)$$

Again, the charge-conjugate partonic subprocess is to be included in Eq. (7). Resolved photoproduction via the partonic subprocess $R_p C_\gamma \rightarrow c$ is treated very similarly.

In Figs. 2(a) and (b), our results for $D^{*\pm}$ and D_s^\pm mesons, respectively, are broken down to the $C_p \gamma \rightarrow c$, $C_p R_\gamma \rightarrow c$, and $R_p C_\gamma \rightarrow c$ contributions and are compared with the ZEUS data [11, 12]. We find that the theoretical predictions are dominated by direct photoproduction and agree rather well with the experimental data over the whole p_T range considered.

V. D-MESON PRODUCTION AT THE TEVATRON

CDF [7] measured the p_T distributions of D^0 , D^\pm , $D^{*\pm}$, and D_s^\pm mesons with rapidity $|y| \leq 1$ inclusively produced in hadroproduction in run II at the Tevatron, with $\sqrt{S} = 1.96$ TeV. To LO in the QMRK approach, the factorization formula for the $C_p R_{\bar{p}} \rightarrow c$ channel reads:

$$\begin{aligned} p_T^3 \frac{d\sigma}{dp_T} &= \int dy \int dz \int dt_1 \int d\phi_1 z^2 D_{c \rightarrow D}(z, \mu^2) \\ &\times \Phi_c^p(x_1, t_1, \mu^2) \Phi_g^{\bar{p}}(x_2, t_2, \mu^2) \overline{|M(C_p R_{\bar{p}} \rightarrow c)|^2}, \end{aligned} \quad (10)$$

where $\overline{|M(C_p R_{\bar{p}} \rightarrow c)|^2}$ is given by Eq. (9),

$$x_1 = \frac{p_T e^y}{z\sqrt{S}}, \quad x_2 = \frac{p_T e^{-y}}{z\sqrt{S}}, \quad t_2 = t_1 - 2\frac{p_T}{z}\sqrt{t_1} \cos \phi_2 + \frac{p_T^2}{z^2}. \quad (11)$$

The result for the $R_p C_{\bar{p}} \rightarrow c$ channel is similar and has to be included in Eq. (10) together with those from the charge-conjugate partonic subprocesses.

In Figs. 3(a)–(d), our results for D^0 , D^\pm , $D^{*\pm}$, and D_s^\pm mesons, respectively, are compared with the CDF data [7]. We find that the theoretical predictions generally agree rather well with the experimental data, except perhaps for the slope. In fact, the predictions exhibit a slight tendency to undershoot the data at small values of p_T and to overshoot them at large values of p_T . However, we have to bear in mind that these are just LO predictions, so that there is room for improvement by including higher orders.

In the framework of the collinear parton model, comparisons with the experimental data of Ref. [7] were performed beyond LO, namely in the fixed-order-next-to-leading-logarithm (FONLL) scheme [8] and at NLO in the GM-VFNS [9, 32]. The FONLL predictions systematically undershoot the CDF data [7]. The GM-VFNS predictions of Ref. [9], which are evaluated with FFs determined in the ZM-VFNS [31], describe these data within their errors, but are still somewhat on the low side. The degree of agreement is further improved [32] by evaluating the GM-VFNS predictions of Ref. [9] using FFs extracted [33] from a global fit to B - and Z -factory data of e^+e^- annihilation in the very same scheme.

VI. CONCLUSIONS

In this paper, we explored the usefulness of the quark-Reggeization hypothesis in the framework of the QMRK approach by studying several observables of inclusive charm production at LO, namely the charm structure function $F_{2,c}$ of the proton measured at HERA [29, 30] as well as the one-particle-inclusive cross sections of $D^{*\pm}$ and D_s^\pm photoproduction in ep collisions at HERA [11, 12] and of D^0 , D^\pm , $D^{*\pm}$, and D_s^\pm hadroproduction in $p\bar{p}$ collisions at the Tevatron [7]. In all three cases, we found satisfactory agreement between our default predictions and the experimental data, which is quite encouraging in view of the simplicity of our LO expressions for the partonic cross sections. By contrast, in the collinear parton model of QCD, the inclusion of NLO corrections is necessary to achieve such a degree of agreement. We thus recover the notion that the QMRK approach is a powerful tool for the theoretical description of QCD processes in the high-energy limit and automatically accommodates an important class of corrections that lie beyond the reach of the collinear parton model at LO [4].

Acknowledgments

We thank L. N. Lipatov and O. V. Teryaev for useful discussions. A. V. S. is grateful to the International Center of Fundamental Physics in Moscow and to the Dynastiya Foundation for financial support. This work was supported in part by the German Federal Ministry for Education and Research BMBF through Grant No. 05 HT6GUA and by the German

- [1] V. S. Fadin and L. N. Lipatov, Nucl. Phys. **B406**, 259 (1993).
- [2] V. S. Fadin and L. N. Lipatov, Nucl. Phys. **B477**, 767 (1996) [arXiv:hep-ph/9602287].
- [3] L. N. Lipatov, Nucl. Phys. **B452**, 369 (1995) [arXiv:hep-ph/9502308].
- [4] V. A. Saleev and D. V. Vasin, Phys. Rev. D **68**, 114013 (2003) [arXiv:hep-ph/0304114]; Phys. Atom. Nucl. **68**, 94 (2005) [Yad. Fiz. **68**, 95 (2005)]; Phys. Part. Nucl. **38**, 635 (2007); B. A. Kniehl, D. V. Vasin, and V. A. Saleev, Phys. Rev. D **73**, 074022 (2006) [arXiv:hep-ph/0602179]; B. A. Kniehl, V. A. Saleev, and D. V. Vasin, *ibid.* **74**, 014024 (2006) [arXiv:hep-ph/0607254].
- [5] Ph. Hägler, R. Kirschner, A. Schäfer, L. Szymanowski, and O. V. Teryaev, Phys. Rev. D **62**, 071502(R) (2000) [arXiv:hep-ph/0002077].
- [6] V. A. Saleev, Phys. Rev. D **78**, 034033 (2008) [arXiv:0807.1587 [hep-ph]].
- [7] D. E. Acosta *et al.* [CDF Collaboration], Phys. Rev. Lett. **91**, 241804 (2003) [arXiv:hep-ex/0307080].
- [8] M. Cacciari and P. Nason, JHEP **0309**, 006 (2003) [arXiv:hep-ph/0306212].
- [9] B. A. Kniehl, G. Kramer, I. Schienbein, and H. Spiesberger, Phys. Rev. D **71**, 014018 (2005) [arXiv:hep-ph/0410289]; Eur. Phys. J. C **41**, 199 (2005) [arXiv:hep-ph/0502194]; Phys. Rev. Lett. **96**, 012001 (2006) [arXiv:hep-ph/0508129]; Phys. Rev. D **77**, 014011 (2008) [arXiv:0705.4392 [hep-ph]].
- [10] S. Aid *et al.* [H1 Collaboration], Nucl. Phys. **B472**, 32 (1996) [arXiv:hep-ex/9604005]; C. Adloff *et al.* [H1 Collaboration], *ibid.* **B545**, 21 (1999) [arXiv:hep-ex/9812023].
- [11] J. Breitweg *et al.* [ZEUS Collaboration], Eur. Phys. J. C **6**, 67 (1999) [arXiv:hep-ex/9807008].
- [12] J. Breitweg *et al.* [ZEUS Collaboration], Phys. Lett. B **481**, 213 (2000) [arXiv:hep-ex/0003018].
- [13] J. Binnewies, B. A. Kniehl, and G. Kramer, Z. Phys. C **76**, 677 (1997) [arXiv:hep-ph/9702408]; B. A. Kniehl, G. Kramer, and M. Spira, *ibid.* **76**, 689 (1997) [arXiv:hep-ph/9610267]; J. Binnewies, B. A. Kniehl, and G. Kramer, Phys. Rev. D **58**, 014014 (1998) [arXiv:hep-ph/9712482].
- [14] M. Cacciari, M. Greco, S. Rolli, and A. Tanzini, Phys. Rev. D **55**, 2736 (1997) [arXiv:hep-ph/9608213]; M. Cacciari and M. Greco, *ibid.* **55**, 7134 (1997) [arXiv:hep-ph/9702389].

- [15] S. Frixione, M. L. Mangano, P. Nason, and G. Ridolfi, Phys. Lett. B **348**, 633 (1995) [arXiv:hep-ph/9412348]; S. Frixione, P. Nason, and G. Ridolfi, Nucl. Phys. **B454**, 3 (1995) [arXiv:hep-ph/9506226].
- [16] G. Kramer and H. Spiesberger, Eur. Phys. J. C **22**, 289 (2001) [arXiv:hep-ph/0109167]; **28**, 495 (2003) [arXiv:hep-ph/0302081]; **38**, 309 (2004) [arXiv:hep-ph/0311062].
- [17] A. V. Berezhnoy, V. V. Kiselev, and A. K. Likhoded, Phys. Atom. Nucl. **63**, 1595 (2000) [Yad. Fiz. **63**, 1682 (2000)] [arXiv:hep-ph/9905555].
- [18] E. A. Kuraev, L. N. Lipatov, and V. S. Fadin, Sov. Phys. JETP **44**, 443 (1976) [Zh. Eksp. Teor. Fiz. **71**, 840 (1976)]; I. I. Balitsky and L. N. Lipatov, Sov. J. Nucl. Phys. **28**, 822 (1978) [Yad. Fiz. **28**, 1597 (1978)].
- [19] L. V. Gribov, E. M. Levin, and M. G. Ryskin, Phys. Rept. **100**, 1 (1983); J. C. Collins and R. K. Ellis, Nucl. Phys. **B360**, 3 (1991); S. Catani, M. Ciafaloni, and F. Hautmann, *ibid.* **B366**, 135 (1991).
- [20] V. S. Fadin, M. I. Kotsky, and L. N. Lipatov, Phys. Lett. B **415**, 97 (1997); D. Ostrovsky, Phys. Rev. D **62**, 054028 (2000) [arXiv:hep-ph/9912258]; V. S. Fadin, M. G. Kozlov, and A. V. Reznichenko, Phys. Atom. Nucl. **67**, 359 (2004) [Yad. Fiz. **67**, 377 (2004)] [arXiv:hep-ph/0302224]; J. Bartels, A. Sabio Vera, and F. Schwennsen, JHEP **0611**, 051 (2006) [arXiv:hep-ph/0608154].
- [21] E. N. Antonov, L. N. Lipatov, E. A. Kuraev, and I. O. Cherednikov, Nucl. Phys. **B721**, 111 (2005) [arXiv:hep-ph/0411185].
- [22] V. S. Fadin and L. N. Lipatov, JETP Lett. **49**, 352 (1989) [Pis'ma Zh. Eksp. Teor. Fiz. **49**, 311 (1989)]; Sov. J. Nucl. Phys. **50**, 712 (1989) [Yad. Fiz. **50**, 1141 (1989)].
- [23] V. S. Fadin and V. E. Sherman, JETP Lett. **23**, 548 (1976) [Pis'ma Zh. Eksp. Teor. Fiz. **23**, 599 (1976)]; Sov. Phys. JETP **45**, 861 (1977) [Zh. Eksp. Teor. Fiz. **72**, 1640 (1977)].
- [24] A. V. Bogdan and V. S. Fadin, Nucl. Phys. **B740**, 36 (2006) [arXiv:hep-ph/0601117].
- [25] M. A. Kimber, A. D. Martin, and M. G. Ryskin, Phys. Rev. D **63**, 114027 (2001) [arXiv:hep-ph/0101348]; G. Watt, A. D. Martin, and M. G. Ryskin, Eur. Phys. J. C **31**, 73 (2003) [arXiv:hep-ph/0306169]; Phys. Rev. D **70**, 014012 (2004); **70**, 079902(E) (2004) [arXiv:hep-ph/0309096].
- [26] G. Watt, URL: <http://www.hep.ucl.ac.uk/~watt/>.
- [27] A. D. Martin, R. G. Roberts, W. J. Stirling, and R. S. Thorne, Phys. Lett. B **531**, 216 (2002)

- [arXiv:hep-ph/0201127].
- [28] M. Glück, E. Reya, and A. Vogt, Phys. Rev. D **46**, 1973 (1992).
- [29] C. Adloff *et al.* [H1 Collaboration], Phys. Lett. B **528**, 199 (2002) [arXiv:hep-ex/0108039].
- [30] S. Chekanov *et al.* [ZEUS Collaboration], Phys. Rev. D **69**, 012004 (2004) [arXiv:hep-ex/0308068].
- [31] J. Binnewies, B. A. Kniehl, and G. Kramer, Phys. Rev. D **58**, 014014 (1998) [arXiv:hep-ph/9712482]; B. A. Kniehl and G. Kramer, *ibid.* **71**, 094013 (2005) [arXiv:hep-ph/0504058]; **74**, 037502 (2006) [arXiv:hep-ph/0607306].
- [32] B. A. Kniehl, in *Proceedings of the XVI International Workshop on Deep-Inelastic Scattering and Related Subjects (DIS 2008)*, London, England, 2008, edited by R. Devenish and J. Ferrando (Science Wise Publishing, 2008), <http://dx.doi.org/10.3360/dis.2008.195> [arXiv:0807.2215 [hep-ph]].
- [33] T. Kneesch, B. A. Kniehl, G. Kramer, and I. Schienbein, Nucl. Phys. **B799**, 34 (2008) [arXiv:0712.0481 [hep-ph]].
- [34] Since we neglect finite quark and hadron mass effects, pseudorapidity and rapidity coincide.




FIG. 1: $F_{2,c}(x_B, Q^2)$ as a function of x_B at (a) $Q^2 = 25$, (b) 30, (c) 45, (d) 60, (e) 130, and (f) 500 GeV^2 . The H1 [29] (open circles) and ZEUS [30] (filled circles) are compared with LO predictions from the QMRK approach with the quark-Reggeization hypothesis.




FIG. 2: p_T distributions of inclusive (a) $D^{*\pm}$ and (b) D_s^\pm photoproduction for $\sqrt{S} = 300$ GeV and $|y| \leq 1.5$. The ZEUS data from (a) Ref. [11] and (b) Ref. [12] are compared with LO predictions from the QMRK approach with the quark-Reggeization hypothesis.




FIG. 3: p_T distributions of inclusive (a) D^0 , (b) D^\pm , (c) $D^{*\pm}$, and (d) D_s^\pm hadroproduction for $\sqrt{S} = 1.96$ TeV and $|y| \leq 1$. The CDF data from Ref. [7] are compared with LO predictions from the QMRK approach with the quark-Reggeization hypothesis.