

Vacuum Birefringence as a Probe of Planck Scale Noncommutativity

Steven A. Abel^a, Joerg Jaeckel^b, Valentin V. Khoze^a and Andreas Ringwald^b

^a*Centre for Particle Theory, Durham University, Durham, DH1 3LE, UK*


^b*Deutsches Elektronen-Synchrotron DESY, Notkestrasse 85, D-22607 Hamburg, Germany*

s.a.abel@durham.ac.uk, jjaeckel@mail.desy.de, valya.khoze@durham.ac.uk,
andreas.ringwald@desy.de

Abstract

Because of ultraviolet/infrared (UV/IR) mixing, the low energy physics of noncommutative gauge theories in the Moyal-Weyl approach seems to depend crucially on the details of the ultraviolet completion. However, motivated by recent string theory analyses, we argue that their phenomenology with a very general class of UV completions can be accurately modelled by a cutoff close to the Planck scale. In the infrared the theory tends continuously to the commutative gauge theory. If the photon contains contributions from a trace- $U(1)$, we would observe vacuum birefringence, i.e. a polarisation dependent propagation speed, as a residual effect of the noncommutativity. Constraints on this effect require the noncommutativity scale to be close to the Planck scale.

arXiv:hep-ph/0607188 v1 17 Jul 2006



1 Introduction

Field theories on noncommuting spacetime,

$$[x^\mu, x^\nu] = i\theta^{\mu\nu} , \quad (1.1)$$

receive a great deal of attention, not least because they arise naturally in a particular, Seiberg-Witten, limit [1] of string theories, see [2–4] for reviews. The corresponding effective field theories can be derived from their commutative cousins by simply replacing products of fields in the Lagrangian by Weyl-Moyal star products¹,

$$(\phi * \varphi)(x) \equiv \phi(x) e^{\frac{i}{2}\theta^{\mu\nu}\overleftarrow{\partial}_\mu\overrightarrow{\partial}_\nu} \varphi(x) . \quad (1.2)$$

The parameter $\theta^{\mu\nu}$ then appears in the vertices of perturbation theory, and, since it has dimensions of $mass^{-2}$, defines a second mass scale in the theory (besides the string scale, M_s), the so-called noncommutativity scale, M_{NC} . A natural question to ask is *what is the allowed range of M_{NC} ?*

In this paper we shall consider the above question in as general a manner as possible using UV cutoffs to mimic the effects of the UV complete theory. As we go along we will compare our results with the string realisation of noncommutative field theory, namely strings in background magnetic (B) fields, which provide a nice, divergence-free framework in which to examine noncommutativity. Despite this obvious attraction of strings, most of the phenomenology depends on very general properties of any UV completion (for example that they should be divergence free, continuous and so on) and we will see that they are well modelled by UV cutoffs.

Before proceeding, let us try to make our question a little more precise. We can estimate the *possible* range of M_{NC} by invoking the notion of naturalness. Considering the specific example of string theory for a moment, pure noncommutative field theory is realised as a special limit of open strings in a background $B^{\mu\nu}$ field, in which closed string (i.e. gravitational) modes are decoupled, leaving only open string interactions. There is no potential for B which as far as the string theory is concerned is just a rather mild background, so in principle $\theta^{\mu\nu}$ could be anything. Nevertheless it seems reasonable to suppose that, if nonperturbative string physics fixes the value of B to be nonzero, it does so with vacuum expectation values (VEVs) of order one in string units.² A natural scale for θ would in that case be $\theta \sim M_s^{-2}$. Depending on the scenario in question that still leaves open a huge possible range: $M_{\text{P}}^{-2} < \theta < M_{\text{W}}^{-2}$, with the Planck scale M_{P} and the weak scale M_{W} , the latter arising, for example, in large extra dimension scenarios. What about other possible UV completions? One role of any UV completion would almost certainly be

¹In the following, we will not consider a more indirect alternative approach to noncommutativity which uses the Seiberg-Witten map. For comments on the relation between the two approaches see [5–7]).

²Note that $M_{\text{NC}} \gg M_{\text{P}}$ does not imply large VEVs for the B fields. In general, since $\theta \sim \frac{1}{\text{const}+B} B \frac{1}{\text{const}-B}$ (Lorentz indices suppressed) vanishing B fields imply vanishing θ and therefore $M_{\text{NC}} \rightarrow \infty$.

to describe quantum gravity. As $\theta^{\mu\nu}$ is intimately involved in the properties of spacetime, a mild assumption is that its “natural” value would inevitably be determined by the only mass scale in quantum gravity. Then one would assume that typically $\theta \sim M_{\text{P}}^{-2}$. As the string theory example shows, the natural range of θ could be beefed-up by, for example, large volumes of extra compact dimensions, but it is difficult to see how much smaller but nonzero values could arise very easily. If there is noncommutativity, therefore, it is natural that $\theta > M_{\text{P}}^{-2}$, or equivalently, $M_{\text{NC}} < M_{\text{P}}$.

So our slightly refined question is, *can noncommutativity at energy scales as high as M_{P} lead to observable effects?* Surprisingly, the answer is yes. As we shall see in this paper, current observations and experiments already severely restrict the range of allowed noncommutativity scales. The reason for this lies in two interesting properties of noncommutative field theories that need to be taken into account in the construction of a viable noncommutative standard model extension [8; 9]. First, there are strong constraints on both the dynamics and the field content. Only $U(N)$ gauge groups with matter fields in fundamental, bifundamental and adjoint representations are allowed [10–14] (for $U(1)$ gauge groups charges are restricted to $\pm 1, 0$ [15]). Second, as we will detail below, universality does not hold and ultraviolet/infrared (UV/IR) mixing occurs [16–19].

In four *continuous* dimensions (i.e. without any quantum gravity, high energy cutoff or UV completion), noncommutative models of this type seem to conflict badly with experiment, as outlined in Ref. [6]. Either there are superfluous massless degrees of freedom or a nonvanishing (and Lorentz symmetry violating) mass term for the photon. Since neither is observed this presents a challenge for any attempt to construct a realistic extension of the Standard Model based on a noncommutative space time.

However, the result of Ref. [6] was based on the assumption that the gauge fields live on a *continuous* four dimensional space time. In particular, it assumed that the four dimensional noncommutative gauge theory is valid up to arbitrarily large momentum scales. But if noncommutative gauge theories are realised as low energy effective field theories of some underlying theory such as string theory, this assumption almost certainly requires modification. It is likely that noncommutative field theory gets spectacularly modified at energy scales approaching M_s . One possible avenue to explore then is the possible effects of “stringy features” such as additional compactified space dimensions. These make the theory effectively higher dimensional at large momentum scales which can be thought of as an intermediate stage towards the string theory. Thanks to UV/IR mixing, the effects of extra dimensions can be transmitted to the IR in the trace- $U(1)$ photon [20]. Such effects can be analysed in a field theoretic framework, and one may search for helpful properties such as the amelioration of constraints on the noncommutativity scale due to, for example, power law decoupling of the trace- $U(1)$ photon in the IR [20].

Despite the obvious attraction of the field-theoretical approach in [6; 20], the drawback is that it is unable to describe effects arising from physics above M_s . This regime is described by the UV completion of the theory, whatever that may be. Normally of course we would not have to worry about such a thing because of universality: the influence

of physics above a cutoff Λ_{UV} on the physics at a momentum scale k is suppressed by powers of k/Λ_{UV} . If universality holds, a modification at very high momentum scales *cannot* modify the physics at much smaller momentum scales. However, although ordinary renormalizable commutative theories fall into this category, noncommutative theories do not, because of UV/IR mixing, [16; 17] and [18; 19].

The phenomenon of UV/IR mixing can be understood from a simple argument. To account for the effects of noncommutativity, we are instructed to replace ordinary products in our field theory by Weyl-Moyal star products (1.2). This results in factors of $\exp(i\tilde{k}\cdot p)$ in the (non-planar) loop integrals [21], where $\tilde{k}^\mu = \theta^{\mu\nu}k_\nu$. Consider a typical loop integral with massless particles in the loop,

$$\int \frac{d^4p}{(2\pi)^4} \frac{1}{p^2(p+k)^2} \exp(i\tilde{k}\cdot p). \quad (1.3)$$

The oscillating phase regularises the integral for large values of momentum p , and the integral is dominated by regions where $\tilde{k}\cdot p \sim 1$, or $|p| \sim M_{\text{NC}}^2/|k|$. This value of $|p|$ is large when the external momentum k is small. The large momenta in the loop $p \sim M_{\text{NC}}^2/k$ indeed influence the physics at small external momentum k . Now consider the effect of heavy particles of mass M in the loop. When $|k| \gg M_{\text{NC}}^2/|p|$, the loop integral is killed when $|p| \ll M$, and (broadly speaking) we may neglect the contribution of heavy particles. But when $|k| \ll M_{\text{NC}}^2/|p|$ the phase is irrelevant and the integral receives contributions from large values of p , $|p| > M$. In other words as we lower our external momentum k , we access ever heavier modes in the loop.

In general, therefore, a modification of the noncommutative theory above a UV scale Λ_{UV} will indeed influence physics below an infrared scale $\Lambda_{\text{IR}} \sim M_{\text{NC}}^2/\Lambda_{\text{UV}}$, as we will see in detail in Sect. 2. The problematic mass term for the photon is an effect of this UV/IR mixing. Hence, it seems plausible that this problem can be treated with a UV modification of the theory. As we already stated, our aim here is to determine some general phenomenological features of noncommutative models and test them against experimental constraints. At first sight this looks like a hopeless task, since constraints corresponding to the lowest energy scales (for example photon masses) are influenced by the highest mass modes in the loop integrals. It looks as though sooner or later we will run up against the UV completion of the theory, at which point all hopes of generality will be lost. However, guided by recent work in Ref. [22], we can determine some generic properties for a large class of theories (cf. Sect. 4). Indeed with fairly mild assumptions (which are true for string theory), the phenomenological effects of the UV completion, such as for example the restoration of normal Wilsonian behaviour in the deep IR, are well modelled by a simple UV cutoff.

As we have already mentioned, there is an important difference between the set-up we use in the present paper and that of Refs. [6; 8; 20] in the way we interpret the underlying noncommutative gauge theory from the perspective of standard particle physics at low energies. The UV/IR mixing effects illustrated by the integral (1.3) occur only in the trace- $U(1)$ factors of the $U(N)$ gauge group(s); the $SU(N)$ degrees of freedom are free from the

UV/IR mixing. The results of the present paper show, that, in presence of a fundamental cutoff Λ_{UV} , the mixing of those U(1) gauge fields (affected by the UV/IR mixing) with the photon does not cause severe problems such as generating a polarisation dependent photon mass. This differs from the approach in Refs. [6; 8; 20], where $\Lambda_{\text{UV}} = \infty$. As will be explained in the next Section, through the UV/IR mixing, an ultimate UV cutoff induces an infrared scale $\Lambda_{\text{IR}} \sim M_{\text{NC}}^2/\Lambda_{\text{UV}}$. At energy scales below Λ_{IR} , physics of all degrees of freedom is governed by an ordinary commutative low-energy effective theory and the effects of the UV/IR mixing are very small. We now place the Standard Model at energies below Λ_{IR} . This is different from the set-up in Refs. [6; 8; 20] which in the language of this paper amounts to $\Lambda_{\text{UV}} = \infty$ and $\Lambda_{\text{IR}} = 0$, thus implying that the Standard Model was embedded into a noncommutative theory at energies E_{SM} in the opposite region: $\Lambda_{\text{IR}} < E_{\text{SM}} < M_{\text{NC}} < \Lambda_{\text{UV}}$.

We will see in Sect. 3 that the problem of the unwanted mass term for the trace-U(1) photon caused by the UV/IR mixing [6] softens considerably at energies below Λ_{IR} . Instead of a mass term one gets, at low momentum scales, vacuum birefringence, i.e. a polarisation dependent propagation speed. If M_{NC} is close enough to the cutoff scale $\Lambda_{\text{UV}} \sim M_{\text{P}}$, this vacuum birefringence can be pushed beyond the current experimental limits. Thereby, a window opens for M_{NC} where noncommutativity is still allowed. As experimental and observational sensitivity is likely to improve in the near future, this provides an interesting probe for scales M_{NC} very close to the Planck scale.

In the following we will concentrate on the case of a pure U(1) noncommutative gauge theory. A pure U(1) gauge theory behaves qualitatively like the trace-U(1) factors of U(N) theories and captures all essential features of the UV/IR mixing.³

The paper is organized as follows. In Sect. 2, we discuss the essential features of UV/IR mixing and the running gauge coupling in the presence of an ultimate UV cutoff. In the following Sect. 3, we demonstrate how this leads to vacuum birefringence. We discuss experimental and observational bounds on this effect and the resulting constraints on the scale of noncommutativity M_{NC} . The validity of using a fundamental UV cutoff to simulate the UV completion is outlined in Sect. 4, where we make a comparison with string theory, and use it as evidence in support of our claim that the phenomenology outlined here is very generic. Finally, we conclude in Sect. 5.

2 UV/IR mixing in presence of a finite UV cutoff

In noncommutative gauge theories, Lorentz symmetry is explicitly broken since the matrix θ on the right hand side of (1.1) is a constant matrix to be specified in a fixed reference frame. This allows an additional transverse (gauge invariant) structure that might be

³We recall that the noncommutative U(1) is an interacting theory, which is asymptotically free in the UV. Its commutative counterpart is of course a free theory.

j=	scalar	Weyl fermion	gauge boson	ghost
α_j	-1	$\frac{1}{2}$	$-\frac{1}{2}$	1
C_j	0	$\frac{1}{2}$	2	0
d_j	1	2	4	1

Table 1: Coefficients appearing in the evaluation of the loop diagrams.

present in the polarisation tensor⁴,

$$\Pi_{\mu\nu} = \Pi_1(k^2, \tilde{k}^2) (k^2 g_{\mu\nu} - k_\mu k_\nu) + \Pi_2(k^2, \tilde{k}^2) \frac{\tilde{k}_\mu \tilde{k}_\nu}{\tilde{k}^2} \quad \text{with} \quad \tilde{k}^\mu = \theta^{\mu\nu} k_\nu. \quad (2.1)$$

The Π_1 part multiplies the ordinary transverse structure and is related to the gauge running coupling via [18]

$$\frac{1}{g^2(k, \tilde{k})} = \frac{1}{g_0^2} + \Pi_1(k, \tilde{k}). \quad (2.2)$$

Π_2 is a new Lorentz symmetry violating structure [16; 17]. In theories with exact supersymmetry (SUSY) it is absent [17; 18]. Its size is therefore related to the SUSY breaking scale [5].

Performing a one loop calculation for the polarisation tensor one obtains [5; 18],

$$\Pi_{\mu\nu}(k) = \Pi_{\mu\nu}(k, l=0) - \Pi_{\mu\nu}(k, l=\tilde{k}), \quad (2.3)$$

with

$$\begin{aligned} \Pi_{\mu\nu}(k, l) = 2 \sum_j \alpha_j \int \frac{d^4 q}{(2\pi)^4} \left\{ d(j) \left[\frac{(2q+k)_\mu (2q+k)_\nu}{(q^2+m_j^2)((q+k)^2+m_j^2)} - \frac{2\delta_{\mu\nu}}{q^2+m_j^2} \right] \right. \\ \left. + 4C(j) \frac{k^2 \delta_{\mu\nu} - k_\mu k_\nu}{(q^2+m_j^2)((q+k)^2+m_j^2)} \right\} \exp(iq \cdot l), \end{aligned} \quad (2.4)$$

where the coefficients α_j , $d(j)$ and $C(j)$ are given in Tab. 1.

As we already stated in the introduction we want to model the UV finiteness of an underlying theory by cutting off all fluctuations above a UV scale Λ_{UV} . One suitable way to do this is by introducing a factor of $\exp(-\frac{1}{\Lambda_{UV}^2 t^2})$ in the integral over the Schwinger

⁴Here, and in the following we will concentrate on the case of a noncommutative U(1) gauge group. The generalisation to U(N) gauge groups is straightforward. All statements remain valid, when applied to the trace-U(1) part of the gauge group. The SU(N) part is unaffected by noncommutativity, independent of the presence of a cutoff.

time t . One obtains (s. [5]),

$$\begin{aligned}
\Pi_{\mu\nu}(k) &= \frac{1}{4\pi^2} (k^2 \delta_{\mu\nu} - k_\mu k_\nu) \\
&\quad \times \sum_j \alpha_j \int_0^1 dx [4C(j) - (1-2x)^2 d(j)] \left[K_0 \left(\frac{\sqrt{A_j}}{\Lambda_{\text{UV}}} \right) - K_0 \left(\frac{\sqrt{A_j}}{\Lambda_{\text{eff}}} \right) \right] \\
&\quad + \frac{1}{(4\pi)^2} \tilde{k}_\mu \tilde{k}_\nu \Lambda_{\text{eff}}^2 \sum_j \alpha_j d(j) \int_0^1 dx A_j K_2 \left(\frac{\sqrt{A_j}}{\Lambda_{\text{eff}}} \right) \\
&\quad + \delta_{\mu\nu} [\text{gauge non-invariant term}],
\end{aligned} \tag{2.5}$$

where

$$A_j = m_j^2 + x(1-x)k^2 \tag{2.6}$$

and

$$\frac{1}{\Lambda_{\text{eff}}^2} = \frac{1}{\Lambda_{\text{UV}}^2} + \tilde{k}^2. \tag{2.7}$$

We will neglect the gauge non-invariant terms in the following. They can be treated and eliminated by using modified Ward-Takahashi identities [23–25].

The employed regularisation cuts off the modes $p \gtrsim \Lambda_{\text{UV}}$ in the loop integral in a smooth way. Of course there are lots of different possibilities to do this. Since universality does not hold, different regularisations will, in principle, lead to different results. However, as long as we leave the qualitative feature “all momenta $p \gtrsim \Lambda_{\text{UV}}$ are cut off” holds, we expect that the qualitative results we obtain remain true. For some details on other choices for the implementation of the cutoff, see Appendix A.

Let us first concentrate on Π_1 , i.e. the running gauge coupling, and for the moment eliminate Π_2 by considering a theory with unbroken supersymmetry.

In Fig. 1, we plot the running gauge coupling for various values of the cutoff Λ_{UV} . As expected the running stops at the UV scale Λ_{UV} . In an ordinary commutative theory we would expect no further changes. Here, however, we observe that the running stops, again, at an infrared scale $\Lambda_{\text{IR}} \sim M_{\text{NC}}^2/\Lambda_{\text{UV}}$. The running for $k < \Lambda_{\text{IR}}$ vanishes up to threshold effects and is therefore essentially the same as that of a pure commutative U(1) gauge theory (recalling that the β function of a pure commutative U(1) gauge theory vanishes). It is easy to check that a similar picture holds also for a more general matter content. Stated differently, only in the range $\Lambda_{\text{IR}} < k < \Lambda_{\text{UV}}$ do we observe a truly noncommutative behavior of the running gauge coupling. Outside this range the behavior is strongly affected by the presence of the UV cutoff.

So far we have rather sloppily been using the scale M_{NC}^2 . Let us now give a more precise definition,

$$|\tilde{k}| = M_{\text{NC}}^{-2} |k|, \tag{2.8}$$




Figure 1: Running gauge coupling for a massless supersymmetric pure U(1) gauge theory. The red, blue and black lines (bottom to top) are for $\Lambda_{\text{UV}} = 1000 M_{\text{NC}}, 10^5 M_{\text{NC}}, \infty M_{\text{NC}}$, respectively. We have fixed the maximal gauge coupling to be $g_{\text{max}}^2 = 4$. One can clearly see that for finite values of the cutoff the running stops at $\sim \Lambda_{\text{UV}}$ in the UV and at $\Lambda_{\text{IR}} \sim M_{\text{NC}}^2/\Lambda_{\text{UV}}$ in the IR.

where M_{NC} is the noncommutativity mass-scale. Heuristically, $M_{\text{NC}}^{-2} \sim |\theta|$ but it may depend on the direction. E.g., for $\theta^{\mu\nu}$ in the canonical basis,

$$\theta^{\mu\nu} = \begin{pmatrix} 0 & \theta_1 & 0 & 0 \\ -\theta_1 & 0 & 0 & 0 \\ 0 & 0 & 0 & \theta_2 \\ 0 & 0 & -\theta_2 & 0 \end{pmatrix}, \quad (2.9)$$

only when $\theta_1 \simeq \theta_2$ does one have $M_{\text{NC}}^{-2} = |\theta|$. Otherwise the scale M_{NC} depends on k_μ ,

$$M_{\text{NC}}^{-2} = \frac{|\theta^{\mu\nu} k_\nu|}{|k|} = |\theta_2| \sqrt{1 + \frac{\theta_1^2 - \theta_2^2 k_0^2 + k_1^2}{\theta_2^2 k^2}}. \quad (2.10)$$

If for example one of the $\theta_i = 0$ one can have a situation where $M_{\text{NC}} \rightarrow \infty$. In general the truly noncommutative region $\Lambda_{\text{IR}} < k < \Lambda_{\text{UV}}$ will depend on the direction in momentum space. This is depicted in Fig. 2. While we have truly noncommutative behavior inside we have cutoff dominated nearly commutative behavior outside this region.

The crucial question is now in which region we perform experiments. If the noncommutativity scale is low and the cutoff sufficiently high, say $M_{\text{NC}} \sim \text{few TeV}$, $\Lambda_{\text{UV}} \sim 10^{18} \text{ GeV}$ we would live in the fully noncommutative region (shaded area in Fig. 2). However, this has already been excluded for a four dimensional theory [6]. If we consider high scales for M_{NC} , say $M_{\text{NC}} \sim (10^{-3} - 1)M_{\text{P}}$, we find (using $\Lambda_{\text{UV}} = M_{\text{P}}$)

$$\Lambda_{\text{IR}} \sim (10^{-6} - 1)M_{\text{P}} \gg k_{\text{max}} \sim 1 \text{ TeV}, \quad (2.11)$$




Figure 2: Depiction of the regions where the model behaves like a fully noncommutative theory (grey shaded area) and where the theory behaves more or less like the commutative theory (white). The green (left) line gives Λ_{IR} , blue (middle) M_{NC} and red (right) Λ_{UV} . In the left panel we have used $|\tilde{k}| = |k|\theta_0 \cos(\alpha)$, $\theta_0 = 10^{10}/\Lambda_{\text{UV}}^2$. In the right panel we use $\alpha = \arctan(10^\xi - 1)$ to show that the lines for Λ_{IR} and Λ_{UV} intersect. The different grey shades also depict the deviation of $1/g^2$ from the UV value $1/g_0^2$ (lighter colors less deviation). This shows that in the IR and in the commutative directions (α close to $\pi/2$) the coupling is given by the UV value. For a more general matter content the coupling will attain (up to threshold corrections) the same value as a purely commutative theory.

which is much bigger than the highest momentum transfer k_{max} that has been reached in experiments so far. Therefore, with sufficiently high scale noncommutativity we expect to live in the nearly commutative region (white areas in Fig. 2) where the gauge coupling behaves like that of a commutative theory. Nevertheless, we will see in the next section that even in this region Π_2 leads to some possibly observable remnant effects. One final remark concerning Fig. 2 is that the deep IR and the far UV are continuously connected. In this sense the UV and the deep IR are all part of the “UV phase”.

3 Vacuum birefringence - a remnant effect from high scale noncommutativity

Let us now turn to the Π_2 part of the polarisation tensor. It, too, is affected by the presence of a finite UV cutoff. From Eq. (2.5) we can easily see that it vanishes for a supersymmetric theory since

$$\sum_j \alpha_j d(j) = 0 \quad (3.1)$$

for supersymmetric theories. When supersymmetry is softly broken⁵ we can easily derive the following approximate expressions (for some additional details see Appendix A),

$$\begin{aligned}\Pi_2 &= D\Delta M_{\text{SUSY}}^2, & \text{for } \frac{M_{\text{NC}}^2}{\Lambda_{\text{UV}}} \ll k \ll \Delta M_{\text{SUSY}}, \\ \Pi_2 &= D'\Delta M_{\text{SUSY}}^2\Lambda_{\text{UV}}^2\tilde{k}^2, & \text{for } k \ll \frac{M_{\text{NC}}^2}{\Lambda_{\text{UV}}}, \quad m_j^2 \ll \Lambda_{\text{UV}}^2,\end{aligned}\tag{3.2}$$

where D, D' are known constants and

$$\Delta M_{\text{SUSY}}^2 = \frac{1}{2} \sum_b M_b^2 - \sum_f M_f^2\tag{3.3}$$

is the (super-)trace of the mass matrix. Following the arguments given in [6] we can now solve the equations of motion for the photon,

$$\Pi^{\mu\nu}(k)A_\nu(k) = 0.\tag{3.4}$$

For concreteness, we now specify the noncommutativity,

$$\theta^{13} = -\theta^{31} = \theta := \frac{1}{M_{\text{NC}}^2},\tag{3.5}$$

and all other components of $\theta^{\mu\nu}$ vanishing (in the 3-direction, this use of M_{NC} coincides with our direction dependent definition (2.8)). The photon flies in the three direction,

$$k^\mu = (k^0, 0, 0, k^3).\tag{3.6}$$

Due to gauge invariance, only the two transverse polarisations are physical. They have the polarisation vectors

$$A_1^\mu = (0, 1, 0, 0), \quad A_2^\mu = (0, 0, 1, 0).\tag{3.7}$$

Inserting into Eq. (3.4) we find

$$\begin{aligned}(\Pi_1 k^2 - \Pi_2)A_1^\mu &= 0, \\ \Pi_1 k^2 A_2^\mu &= 0.\end{aligned}\tag{3.8}$$

The photon polarized along A_2^μ obviously behaves like an ordinary massless photon. However, in the A_1^μ direction we observe new and interesting effects. To study these in more detail let us now insert the approximate expressions (3.2). For the A_1^μ polarisation we obtain the following dispersion relations,

$$k^2 - D\frac{\Delta M_{\text{SUSY}}^2}{\Pi_1} = 0, \quad \text{for } \Lambda_{\text{IR}} = \frac{M_{\text{NC}}^2}{\Lambda_{\text{UV}}} \ll k \ll \Delta M_{\text{SUSY}},\tag{3.9}$$

$$k^2 + D'\frac{1}{\Pi_1}\frac{\Delta M_{\text{SUSY}}^2\Lambda_{\text{UV}}^2}{M_{\text{NC}}^4}(k^3)^2 = 0, \quad \text{for } k \ll \frac{M_{\text{NC}}^2}{\Lambda_{\text{UV}}} = \Lambda_{\text{IR}}.\tag{3.10}$$

⁵Meaning that numbers of bosonic and fermionic degrees of freedom of the theory still match.

Equation (3.9) yields a Lorentz symmetry violating mass term of the order of ΔM_{SUSY}^2 that was already discussed in detail in [6]. Without cutoff, i.e. in the limit $\Lambda_{\text{UV}} \rightarrow \infty$, this mass term persists down to $k \rightarrow 0$, thereby excluding any chance that this can be the photon observed in nature. In presence of the cutoff, Eq. (3.9) is only applicable for $k \gg \Lambda_{\text{IR}}$. Masslessness of the photon is well tested up to at least 1 GeV. Using $M_{\text{P}} \sim \Lambda_{\text{UV}} = 10^{18}$ GeV, this gives us a conservative lower bound of $M_{\text{NC}} > 10^9$ GeV. Nevertheless, this opens a rather large window of opportunity compared to the $\Lambda_{\text{UV}} \rightarrow \infty$ case where all $M_{\text{NC}} < M_{\text{P}}$ are excluded.

For small photon momentum, Eq. (3.10) applies (recall from our discussion at the end of Sect. 2 that we actually expect to live in this limit). To understand (3.10) better, let us restore the light speed c in our equations and use $k^0 = \omega$ for the frequency of the wave,

$$\omega^2 - c^2 \left(\frac{1}{1 + \Delta n} \right)^2 (k^3)^2 = 0, \quad (3.11)$$

with

$$\begin{aligned} \Delta n &\approx \frac{D'}{2} \frac{1}{\Pi_1} \frac{\Delta M_{\text{SUSY}}^2 \Lambda_{\text{UV}}^2}{M_{\text{NC}}^4} \\ &= 10^{-34} \left(\frac{D'/2\Pi_1}{10^{-4}} \right) \left(\frac{\Delta M_{\text{SUSY}}}{10^3 \text{ GeV}} \right)^2 \left(\frac{\Lambda_{\text{UV}}}{10^{18} \text{ GeV}} \right)^2 \left(\frac{M_{\text{NC}}}{10^{18} \text{ GeV}} \right)^{-4} \ll 1. \end{aligned} \quad (3.12)$$

Here, we have combined the regularisation dependent loop factor $D' = \mathcal{O}(1/4\pi^2)$ and the field content dependent factor $\Pi_1 = \mathcal{O}(10 - 100)$ to parameterise the model dependence.

From Eq. (3.11) we can see that the photon A_1^μ propagates with a speed⁶ $\approx c(1 - \Delta n)$. Since the A_1^μ photon propagates with c we observe birefringence, i.e. different polarisations propagate with different speed.

Although Δn seems to be quite small we should compare this to the current experimental sensitivity. In Ref. [26], a study of all possible dimension four Lorentz violating operators in electrodynamics was conducted and constraints derived. The most general dimensions four Lagrangian which is gauge and CPT invariant, but violates Lorentz symmetry is,

$$\mathcal{L}_{\text{general}} = -\frac{1}{4} F^{\mu\nu} F_{\mu\nu} - \frac{1}{4} (k_F)_{\mu\nu\alpha\beta} F^{\mu\nu} F^{\alpha\beta}. \quad (3.13)$$

Comparing the propagator derived from Eq. (3.13) with Eq. (2.1) we find

$$(k_F)_{\mu\nu\alpha\beta} = \frac{D'}{2} \frac{1}{\Pi_1} \Delta M_{\text{SUSY}}^2 \Lambda_{\text{UV}}^2 \theta_{\mu\nu} \theta_{\alpha\beta}. \quad (3.14)$$

In Ref. [26], the coefficients of k_F have been constrained using various methods. For laboratory measurements, their estimate translates to

$$|\Delta n_{\text{lab}}| \lesssim 10^{-14} - 10^{-10}, \quad (3.15)$$

⁶Note that $\Delta n < 0$ is not inconsistent. Since Lorentz symmetry is explicitly broken, propagation with speeds $> c$ is, in principle, possible.




Figure 3: Bounds on the scale of noncommutativity M_{NC} in a four dimensional non-commutative theory with an ultimate UV cutoff Λ_{UV} . The red area is excluded by the requirement that $M_{\text{NC}} < \Lambda_{\text{UV}}$. The other curves show lower limits on M_{NC} derived via Eq. (3.12) from bounds on vacuum birefringence. The grey band corresponds to estimates from lab measurements. The blue (lower) and black (upper) thick solid curves originate from observations at astrophysical and cosmological distances, respectively. The thick dashed line gives the most recent constraint from polarisation measurements of gamma ray bursts [28]. We used $D'/2\Pi_1 = 10^{-4}$.

depending on the pattern of the noncommutativity. Astrophysical observations already provide a much tighter bound of

$$|\Delta n_{\text{astro}}| \lesssim 10^{-16}, \quad (3.16)$$

while the strongest constraints come from observations of objects at cosmological distances (see also [27; 28]),

$$|\Delta n_{\text{cosmo}}| \lesssim 10^{-37} - 10^{-32}. \quad (3.17)$$

In Fig. 3, we show the lower limits on M_{NC} originating from these experimental and observational upper limits on the birefringence of the vacuum.

4 Cutoffs as a mimic of UV physics

After having found that an ultimate UV cutoff leads to interesting physics in noncommutative gauge theories, let us now discuss why such a cutoff provides a good approximation to the effect of UV completion. As we stated in the Introduction, our evidence for this comes from the understanding one gains from the string theory theory realisation of non-commutativity.

In order to appreciate what happens in string theory, consider what would happen in a more general field theory containing a tower of massive modes of mass m_i . In Euclidean space, the typical one loop diagrams would have a sum over the modes as follows

$$I(\theta, k, \Lambda_{\text{UV}}) = \sum_i \int \frac{d^4 p}{(2\pi)^4} \frac{1}{(p^2 + m_i^2)((p+k)^2 + m_i^2)} \exp(i\tilde{k} \cdot p). \quad (4.1)$$

Again we will go to the Schwinger parameterisation. Using the identity

$$\frac{1}{A_1 A_2} = \int_0^1 dx \int_0^\infty dt t e^{-t(xA_1 + (1-x)A_2)}, \quad (4.2)$$

we may recast (4.1) as

$$I(\theta, k, m_i) = \int_0^\infty dt t e^{-\frac{\tilde{k}^2}{4t}} \sum_i \int \frac{d^4 p}{(2\pi)^4} \int_0^1 dx e^{-t(p^2 + k^2 x(1-x) + m_i^2)}, \quad (4.3)$$

where we used $k \cdot \tilde{k} = 0$. Very heuristically, the way string theory works as a finite UV completion is to arrange the masses m_i so that the integrand resums into functions with modular properties that render the integral finite. The additional modes that are required to do this have masses of order the string scale, i.e. the typical masses of the lowest lying extra modes is order M_s ; we call them UV modes⁷.

More generally, we can consider the class of theories where the UV completion yields a one-loop contribution of the form

$$I(\theta, k, \Lambda_{\text{UV}}) = \int_0^\infty dT T f\left(\frac{\tilde{k}^2 \Lambda_{\text{UV}}^2}{4T}\right) Z\left(T, \frac{k}{\Lambda_{\text{UV}}}\right), \quad (4.4)$$

where we have rescaled to a dimensionless Schwinger parameter, $t = T/\Lambda_{\text{UV}}^2$. The function f contains all the effects of noncommutativity, whereas Z would also be present in a commutative theory. Since the commutative theory should be finite, too, Z implements the UV finiteness of the integral. This property is typically provided by the sum over an appropriate spectrum of massive modes as indicated by the sum in Eq. (4.3). It is this connection which imbues Λ_{UV} with a physical meaning as the scale at which the UV completion modifies the integrand. Roughly, it corresponds to the typical mass of the lightest UV modes m_i . How this works in a string theoretical setting with a non-zero B -field has been shown in Ref. [22]. There, the role of Λ_{UV} is played by $1/\sqrt{\alpha'}$, with the string tension α' . Moreover, in that case the form of the function f is indeed given by

$$f = e^{-\frac{\tilde{k}^2 \Lambda_{\text{UV}}^2}{4T}}, \quad (4.5)$$

⁷Actually the question of finiteness in the string theory is quite subtle in this context. It relies on consistency conditions, namely tadpole cancellation. For more details see Ref. [22].

for all T (cf. Eq. (4.3)). In general, we expect this simple form only for small loop-momenta $\ll \Lambda_{\text{UV}}$, corresponding to $T \gg 1$,

$$\lim_{T \gg 1} f = e^{-\frac{\tilde{k}^2 \Lambda_{\text{UV}}^2}{4T}}. \quad (4.6)$$

A crucial determinant of the behaviour is the interplay between f and Z . In order to describe this further, we will define two properties of the UV complete theory that we will consider to be necessary:

- *All couplings in the $k \rightarrow 0$ limit tend to the couplings of the $\theta = 0$ theory.*
- *All physics in the $\theta \rightarrow 0$ limit tends continuously to $\theta = 0$ physics.*

These we will take to be fundamental properties of a consistent UV completion and are certainly true for both string theory and the field theory with a cutoff. As stated in the introduction, the non-zero B field is a mild background field that we can dial continuously to zero; it would be very odd for there to be any sort of discontinuity at $B = 0$. At least the second of these assumptions is known to be false in noncommutative field theory, but then of course that theory does not provide a finite UV completion. Both properties are obviously true for the one-loop contribution above if the integral is finite and uniformly convergent, and if the function f is continuous.

Given these assumptions it is clear that the behaviour of the theory is essentially determined by whether it is the function f or Z which is doing the regulating of the integral. If the regularisation is controlled by Z , then the behaviour must by continuity be identical to the commutative string theory. However as we shall see more thoroughly at the end of this section, for momenta in the intermediate range $\Lambda_{\text{IR}} < k < \Lambda_{\text{UV}}$, the integral is regulated by f . This leads to the field theoretical behaviour where the integral is effectively regulated by the noncommutativity (cf. Sect. 2).

To examine the question of continuity further, it is instructive to consider taking the $\theta \rightarrow 0$ limit by scaling $\theta \rightarrow \lambda\theta$. The only place θ appears is in f ; we may redefine $\Lambda_{\text{UV}} \rightarrow \sqrt{\lambda}\Lambda_{\text{UV}}$ and $k \rightarrow \sqrt{\lambda}k$. The net result is

$$I(\lambda\theta, k, \Lambda_{\text{UV}}) = I(\theta, \sqrt{\lambda}k, \sqrt{\lambda}\Lambda_{\text{UV}}). \quad (4.7)$$

This equation looks a bit peculiar but on inspection it makes sense: it says that the effect of taking the commutative limit is the same as lowering the mass scales of all the modes of the UV completion to zero and leaving θ untouched. In other words, on the right hand side the threshold effects of an increasing number of the additional UV modes are included whilst $\tilde{k} \cdot p \ll 1$, and in the limit the one-loop correction to the gauge coupling includes all the same contributions as the commutative theory, thus proving the second property for gauge couplings, namely that as $\theta \rightarrow 0$ they tend to the commutative ones. Note that

this last statement is only true because of the assumed convergence of the integral in a UV finite theory.

However the second property we are demanding of our theory is actually a stronger requirement than this; noncommutativity introduces new operators where momenta are contracted with θ 's, and the second property says that they tend to zero in the IR. This is especially surprising given that in noncommutative field theory the very same operators are divergent in the IR. A typical operator is precisely the contribution to the vacuum polarisation tensor of the trace-U(1) photon,

$$\Pi_{\mu\nu} \supset \Pi_2(k^2, \tilde{k}^2) \frac{\tilde{k}_\mu \tilde{k}_\nu}{\tilde{k}^2}. \quad (4.8)$$

Π_2 has dimensions of $mass^2$, and in a generic non-supersymmetric field theory (with no matching between numbers of bosonic and fermionic degrees of freedom) $\Pi_2 \sim 1/\tilde{k}^2$ [16; 17]. In the UV complete theory, this contribution is of the general form

$$\Pi_{\mu\nu} \supset \tilde{k}_\mu \tilde{k}_\nu J(\theta, k, m_i) = \tilde{k}_\mu \tilde{k}_\nu \Lambda_{\text{UV}}^4 \int_0^\infty dT T g\left(\frac{\tilde{k}^2 \Lambda_{\text{UV}}^2}{4T}\right) Z\left(T, \frac{k}{\Lambda_{\text{UV}}}\right), \quad (4.9)$$

where g is a function with the same properties as f . Importantly, continuity removes the possibility that it could have any divergences in \tilde{k}^2 , and insists that in the limit $k \rightarrow 0$ the integral J converges to the value in the ‘‘commutative theory’’ which is of order unity. In the deep IR, therefore, we must have

$$\Pi_2 \sim \tilde{k}^2 \Lambda_{\text{UV}}^4, \quad (4.10)$$

rather than any sort of divergence. In presence of softly broken SUSY, the Λ_{UV}^4 is softened to $\Delta M_{\text{SUSY}}^2 \Lambda_{\text{UV}}^2$ (cf. Eq. (3.2)).

At what momentum scale does this behaviour take over from the usual noncommutative field theory behaviour? The extra UV modes can only contribute in the integral when $T < 1$. Outside this region, contributions from the UV modes in the Schwinger integral are exponentially suppressed, and the one loop contributions are approximately those of the UV divergent field theory. Here, in the diagrams sensitive to the noncommutativity, the UV divergence is tamed by the functions f, g , which act as a cutoff for modes with $T < 4\tilde{k}^2 \Lambda_{\text{UV}}^2$. When the second inequality saturates the first, that is when

$$\tilde{k}^2 > \frac{1}{4\Lambda_{\text{UV}}^2}, \quad \text{i.e.} \quad k^2 > \frac{M_{\text{NC}}^4}{4\Lambda_{\text{UV}}^2} \sim \Lambda_{\text{IR}}^2, \quad (4.11)$$

we never get contributions from UV modes in the integral and the behaviour is entirely field theoretical. On the other hand when \tilde{k}^2 is less than this value, there is a region $4\tilde{k}^2 \Lambda_{\text{UV}}^2 < T < 1$ where the UV modes are contributing significantly. In this regime, the integration tends to the values that we deduced from the convergence and continuity

properties of the UV completion and approaches a finite value as $k \rightarrow 0$. Thus we can define a “deep-IR” region,

$$|k| < \Lambda_{\text{IR}} = \frac{M_{\text{NC}}^2}{\Lambda_{\text{UV}}}, \quad (4.12)$$

in which one-loop integrals give approximately constant contributions, and Wilsonian behaviour is restored.

All of these properties are true for string theory, and by inspection, they are mimicked by the introduction of a cutoff in the Schwinger integral, thus justifying the approach that we have taken in the previous sections.

5 Conclusions

Noncommutative gauge theories are not universal. Therefore, any discussion of low energy effects requires the specification of the ultraviolet sector. In this work we considered a noncommutative field theory model where the fluctuations with momenta larger than an ultraviolet cutoff Λ_{UV} give an overall vanishing contribution. We argued that this is a good approximation to a large class of more fundamental UV finite theories, which includes string theory.

The presence of an ultraviolet cutoff Λ_{UV} induces an effective infrared scale $\Lambda_{\text{IR}} \sim M_{\text{NC}}^2/\Lambda_{\text{UV}}$ below which the running coupling behaves up to threshold corrections like that of a commutative gauge theory⁸. Only in the range $\Lambda_{\text{IR}} < k < \Lambda_{\text{UV}}$ do we observe full noncommutative behavior. However, for large noncommutativity scales $M_{\text{NC}} \gtrsim 10^{11}$ GeV and a cutoff $\Lambda_{\text{UV}} \sim M_{\text{P}}$ one easily finds that all known experiments are performed in the nearly commutative region $k < \Lambda_{\text{IR}}$.

If supersymmetry is broken, an additional Lorentz symmetry violating structure is present in the polarisation tensor. For scales $k > \Lambda_{\text{IR}}$ it leads to a mass term for the gauge boson in accord with Refs. [5; 6]. However, below Λ_{IR} the mass term turns into a modification of the phase velocity of plane wave solutions, leading to birefringence. If the trace-U(1) gauge boson is to be interpreted as (part of) the photon, a mass is not acceptable and birefringence must be smaller than the experimental limits. Using the most stringent limits from cosmological observations one obtains a rather strong limit of $M_{\text{NC}} \gtrsim 0.1M_{\text{P}}$. If we use the more conservative astrophysical or laboratory limits the same argument yields only $M_{\text{NC}} \gtrsim (10^{-7} - 10^{-5}) M_{\text{P}}$. In this setting high precision measurements of the properties of light are a wonderful tool to test (nearly) Planck scale physics.

⁸This is in stark contrast to a situation where the noncommutative gauge theory is assumed to be valid at all scales and no ultraviolet cutoff exists. There $\Lambda_{\text{IR}} = 0$ and the theory shows strong effects of noncommutativity at all scales. In such a situation a noncommutative U(1) can never be the photon as demonstrated in [5; 6].

Acknowledgements

We are indebted to Chong-Sun Chu and Mark Goodsell for discussions. SAA would like to thank CERN for hospitality. VVK thanks the Aspen Centre for Physics for hospitality and acknowledges support of PPARC through a Senior Fellowship.

A Low energy Π_2 for general regulators

The effects discussed in this paper originate from the introduction of a UV cutoff Λ_{UV} . Let us now check that the qualitative behavior is independent of the specific implementation of the cutoff, i.e. the choice of the function that suppresses fluctuations with $k \gtrsim \Lambda_{\text{UV}}$. In particular, let us check that the form given in (3.2) is generic for $k \ll \Lambda_{\text{IR}}$.

Since the first term in (2.3) does not contain θ it is obvious that only the second term can contribute to Π_2 . Similarly the term in the second line of Eq. (2.4) can only contribute at order $\mathcal{O}(k^2 \tilde{k}^2)$. Collecting the remaining terms of Eq. (2.4) which are $\propto \tilde{k}_\mu \tilde{k}_\nu$ one finds,

$$\Pi_2 = -2\tilde{k}^2 \sum_j \alpha_j d(j) \int \frac{d^4 q}{(2\pi)^4} \left[\frac{q^2}{P_j(q)} - \frac{2}{d} \frac{q^4}{P_j^2(q)} \right] \quad (\text{A.1})$$

where $P_j(q)$ is the inverse propagator of the particle j . (In absence of a cutoff, $P_j(q) = q^2 + m_j^2$.) In presence of any reasonable UV cutoff that acts for all particles identically (which, in particular, respects SUSY) one obtains⁹,

$$\int \frac{d^4 q}{(2\pi)^4} \left[\frac{q^2}{P_j(q)} - \frac{2}{d} \frac{q^4}{P_j^2(q)} \right] = \Lambda_{\text{UV}}^4 f\left(\frac{m_j^2}{\Lambda_{\text{UV}}^2}\right). \quad (\text{A.2})$$

As long as

$$\sum_j \alpha_j d(j) f\left(\frac{m_j^2}{\Lambda_{\text{UV}}^2}\right) \neq 0, \quad (\text{A.3})$$

we will observe a birefringence effect as discussed in Sect. 3. For $m_j^2 \ll \Lambda_{\text{UV}}^2$, we can further expand,

$$f\left(\frac{m_j^2}{\Lambda_{\text{UV}}^2}\right) = A + B \frac{m_j^2}{\Lambda_{\text{UV}}^2} + \dots \quad (\text{A.4})$$

Remembering that $\sum_j \alpha_j d(j) = 0$ we find

$$\Pi_2 \sim \tilde{k}^2 B \Delta M_{\text{SUSY}}^2 \Lambda_{\text{UV}}^2, \quad (\text{A.5})$$

as in (3.2). We have checked for several regulators that $B \neq 0$. However, let us remark that B depends on the choice of the regulator.

⁹A large class of different cutoff functions can be implemented by using ERGE scheme regularisation (see, e.g., Appendix C of [29]).

References

- [1] N. Seiberg and E. Witten, “String theory and noncommutative geometry,” JHEP **9909** (1999) 032 [hep-th/9908142].
- [2] M. R. Douglas and N. A. Nekrasov, “Noncommutative field theory,” Rev. Mod. Phys. **73** (2001) 977 [hep-th/0106048].
- [3] R. J. Szabo, “Quantum field theory on noncommutative spaces,” Phys. Rept. **378** (2003) 207 [hep-th/0109162].
- [4] C. S. Chu, “Non-commutative geometry from strings,” hep-th/0502167.
- [5] L. Alvarez-Gaume and M. A. Vazquez-Mozo, “General properties of noncommutative field theories,” Nucl. Phys. B **668** (2003) 293 [hep-th/0305093].
- [6] J. Jaeckel, V. V. Khoze and A. Ringwald, “Telltale traces of U(1) fields in noncommutative standard model extensions,” JHEP **0602** (2006) 028 [hep-ph/0508075].
- [7] S. A. Abel, J. Jaeckel, V. V. Khoze and A. Ringwald, “Photon Defects in Noncommutative Standard Model Candidates,” hep-ph/0606106.
- [8] V. V. Khoze and J. Levell, “Noncommutative standard modelling,” JHEP **0409** (2004) 019 [hep-th/0406178].
- [9] M. Chaichian, P. Presnajder, M. M. Sheikh-Jabbari and A. Tureanu, “Noncommutative standard model: Model building,” Eur. Phys. J. C **29** (2003) 413 hep-th/0107055.
- [10] K. Matsubara, “Restrictions on gauge groups in noncommutative gauge theory,” Phys. Lett. B **482** (2000) 417 [hep-th/0003294].
- [11] A. Armoni, “Comments on perturbative dynamics of non-commutative Yang-Mills theory,” Nucl. Phys. B **593** (2001) 229 [hep-th/0005208].
- [12] J. M. Gracia-Bondia and C. P. Martin, “Chiral gauge anomalies on noncommutative R^{*4} ,” Phys. Lett. B **479** (2000) 321 [hep-th/0002171].
- [13] S. Terashima, “A note on superfields and noncommutative geometry,” Phys. Lett. B **482** (2000) 276 [hep-th/0002119].
- [14] M. Chaichian, P. Presnajder, M. M. Sheikh-Jabbari and A. Tureanu, “Noncommutative gauge field theories: A no-go theorem,” Phys. Lett. B **526** (2002) 132 [hep-th/0107037].
- [15] M. Hayakawa, “Perturbative analysis on infrared aspects of noncommutative QED on R^{*4} ,” Phys. Lett. B **478** (2000) 394 hep-th/9912094; “Perturbative analysis on infrared and ultraviolet aspects of noncommutative QED on R^{*4} ,” hep-th/9912167.

- [16] S. Minwalla, M. Van Raamsdonk and N. Seiberg, “Noncommutative perturbative dynamics,” *JHEP* **0002** (2000) 020 [hep-th/9912072].
- [17] A. Matusis, L. Susskind and N. Toumbas, “The IR/UV connection in the non-commutative gauge theories,” *JHEP* **0012** (2000) 002 [hep-th/0002075].
- [18] V. V. Khoze and G. Travaglini, “Wilsonian effective actions and the IR/UV mixing in noncommutative gauge theories,” *JHEP* **0101** (2001) 026 [hep-th/0011218].
- [19] T. J. Hollowood, V. V. Khoze and G. Travaglini, “Exact results in noncommutative $N = 2$ supersymmetric gauge theories,” *JHEP* **0105** (2001) 051 [hep-th/0102045].
- [20] S. A. Abel, J. Jaeckel, V. V. Khoze and A. Ringwald, “Noncommutativity, extra dimensions, and power law running in the infrared,” *JHEP* **0601** (2006) 105 [hep-ph/0511197].
- [21] T. Filk, “Divergencies in a field theory on quantum space,” *Phys. Lett. B* **376** (1996) 53.
- [22] S. Abel, C. S. Chu and M. Goodsell, “Noncommutativity from the string perspective: Modification of gravity at a mm without mm sized extra dimensions,” hep-th/0606248.
- [23] M. Reuter and C. Wetterich, “Effective average action for gauge theories and exact evolution equations,” *Nucl. Phys. B* **417** (1994) 181.
- [24] U. Ellwanger, “Flow Equations And BRS Invariance For Yang-Mills Theories,” *Phys. Lett. B* **335** (1994) 364 [hep-th/9402077].
- [25] F. Freire, D. F. Litim and J. M. Pawłowski, “Gauge invariance and background field formalism in the exact renormalisation group,” *Phys. Lett. B* **495** (2000) 256 [hep-th/0009110].
- [26] V. A. Kostelecky and M. Mewes, “Signals for Lorentz violation in electrodynamics,” *Phys. Rev. D* **66** (2002) 056005 [hep-ph/0205211].
- [27] V. A. Kostelecky and M. Mewes, “Cosmological constraints on Lorentz violation in electrodynamics,” *Phys. Rev. Lett.* **87** (2001) 251304 [hep-ph/0111026].
- [28] A. Kostelecky and M. Mewes, “Sensitive polarimetric search for relativity violations in gamma-ray bursts,” hep-ph/0607084.
- [29] J. Jaeckel, “Effective actions for strongly interacting fermionic systems,” hep-ph/0309090.