

Categorification of algebraic quantum field theories

Marco Benini^{1,2,a}, Marco Perin^{3,b}, Alexander Schenkel^{3,c} and Lukas Woike^{4,d}

¹ Dipartimento di Matematica, Università di Genova,
Via Dodecaneso 35, 16146 Genova, Italy.

² INFN, Sezione di Genova,
Via Dodecaneso 33, 16146 Genova, Italy.

³ School of Mathematical Sciences, University of Nottingham,
University Park, Nottingham NG7 2RD, United Kingdom.

⁴ Fachbereich Mathematik, Universität Hamburg,
Bundesstr. 55, 20146 Hamburg, Germany.

Email: ^a benini@dima.unige.it
^b marco.perin@nottingham.ac.uk
^c alexander.schenkel@nottingham.ac.uk
^d lukas.jannik.woike@uni-hamburg.de

April 2020

Abstract

This paper develops a novel concept of 2-categorical algebraic quantum field theories (2AQFTs) that assign locally presentable linear categories to spacetimes. It is proven that ordinary AQFTs embed as a coreflective full 2-subcategory into the 2-category of 2AQFTs. Examples of 2AQFTs that do not come from ordinary AQFTs via this embedding are constructed by a local gauging construction for finite groups, which admits a physical interpretation in terms of orbifold theories. A categorification of Fredenhagen's universal algebra is developed and also computed for simple examples of 2AQFTs.

Report no.: ZMP-HH/20-8, Hamburger Beiträge zur Mathematik Nr. 830

Keywords: algebraic quantum field theory, 2-categories, 2-operads, quasi-coherent sheaves, locally presentable linear categories, categorified orbifold theories, local-to-global extensions

MSC 2020: 81Txx, 18M60, 18N10, 18N25

Contents

1	Introduction and summary	2
2	AQFTs and algebra-valued prefactorization algebras	4
3	Definition of 2AQFTs	8
4	Inclusion-truncation biadjunction	11
4.1	Truncation	11
4.2	Inclusion	12
4.3	Biadjunction	14
5	Gauging construction and orbifold 2AQFTs	17

6	Fredenhagen’s universal category	24
6.1	Preliminaries	24
6.2	Extension	26
6.3	Examples on $M = \mathbb{S}^1$	28
A	Basic theory of Cat-enriched colored operads	33

1 Introduction and summary

An algebraic quantum field theory (AQFT) is a functor $\mathfrak{A} : \mathbf{C} \rightarrow \mathbf{Alg}_{\mathbb{K}}$ from a category of spacetimes \mathbf{C} to the category of associative and unital algebras over a field \mathbb{K} , which satisfies certain physically motivated axioms such as Einstein causality and the time-slice axiom [HK64, BFV03, FV15]. The (non-commutative) algebra $\mathfrak{A}(c)$ that is assigned to an object $c \in \mathbf{C}$ is interpreted as the algebra of quantum observables of the theory on the spacetime c .

Describing quantum observables in terms of ordinary algebras in $\mathbf{Alg}_{\mathbb{K}}$ is however insufficient to capture the important, but rather subtle, higher categorical structures that feature in gauge theories. For instance, in the context of the BRST/BV formalism [FR12, FR13], the quantum observables of a gauge theory are described by differential graded algebras (dg-algebras) and the latter contain in general more information than their 0-th cohomology, which is the ordinary algebra of gauge invariant quantum observables. An axiomatic framework for homotopy-coherent AQFTs with values in dg-algebras was developed in [BSW19], see also [BS19] for a review and [BBS19] for concrete examples. In these works it was also shown that the higher structures encoded by dg-algebras are crucial for formalizing descent (i.e. local-to-global) properties of quantum gauge theories.

The main aim of this paper is to develop a novel kind of higher categorical AQFTs, which describe quantum observables in terms of locally presentable \mathbb{K} -linear categories. As we explain in detail below, such AQFTs are more sensitive to global aspects of quantum gauge theories, e.g. finite gauge transformations, than those assigning dg-algebras.

In order to motivate why it is reasonable to describe quantum observables of gauge theories by locally presentable \mathbb{K} -linear categories, let us first recall why ordinary AQFTs assign associative and unital \mathbb{K} -algebras. From the point of view of quantum theory, a (non-commutative) algebra $A \in \mathbf{Alg}_{\mathbb{K}}$ is interpreted as a quantized function algebra on the phase space X of a physical system, i.e. A arises as a (deformation) quantization of the commutative algebra $\mathcal{O}(X)$ of \mathbb{K} -valued functions on X . If X is a sufficiently “nice” space (in technical terms, X is an affine scheme over \mathbb{K}), there is no loss of information when passing from X to its function algebra $\mathcal{O}(X)$. This explains why it is justified to quantize the space X by quantizing its function algebra $\mathcal{O}(X)$.

However, many important examples of phase spaces that feature in physics are *not* of this “nice” kind. For instance, if the phase space X is a *stack*, as it happens to be in a gauge theory, it is in general not true that X is faithfully encoded by its function algebra $\mathcal{O}(X)$, which in this case is a dg-algebra, see e.g. [Toe06]. As an illustrative example, let G be a finite group and consider the quotient stack $\mathbf{BG} := \{*\}/G$, which is a non-trivial stack, namely the classifying stack of principal G -bundles. The corresponding function dg-algebra $\mathcal{O}(\mathbf{BG}) = C^\bullet(G, \mathbb{K})$ is then given by the group cochains with values in the trivial G -representation \mathbb{K} . Taking for example $G = \mathbb{Z}_2$, the cyclic group of order 2, all cohomology groups $H^n(\mathbb{Z}_2, \mathbb{K}) = 0$, for $n \neq 0$, are trivial if \mathbb{K} has characteristic zero. It then follows that $\mathcal{O}(\mathbf{BZ}_2) \simeq H^0(\mathbb{Z}_2, \mathbb{K}) \simeq \mathbb{K} = \mathcal{O}(\{*\})$ is quasi-isomorphic to the function algebra of the point $\{*\}$, i.e. all information about the group $G = \mathbb{Z}_2$ is lost when passing from the stack \mathbf{BG} to its function algebra. As a consequence, it is in general *not* reasonable to quantize a stack X by quantizing its function dg-algebra $\mathcal{O}(X)$. We would like to emphasize that these issues arise for *finite gauge transformations* and not for infinitesimal gauge transformations. In particular, every Lie algebroid $X = Y//\mathfrak{g}$ is completely determined by its function dg-algebra, which in this case is given by the Chevalley-Eilenberg

cochains $\mathcal{O}(X) = \mathbf{CE}^\bullet(\mathfrak{g}, \mathcal{O}(Y))$ on the Lie algebra \mathfrak{g} with values in the ordinary function algebra $\mathcal{O}(Y)$ of the affine scheme Y .

The feature that stacks are in general not completely determined by their function dg-algebras is well-known to algebraic geometers, see e.g. [Bra14] for an excellent overview, who have also proposed the following interesting solution: Instead of assigning a function dg-algebra $\mathcal{O}(X)$ to a space or stack X , it is better to assign the category $\mathbf{QCoh}(X)$ of quasi-coherent sheaves on X . The latter is a locally presentable symmetric monoidal \mathbb{K} -linear category that should be interpreted roughly as the category of vector bundles over X . This is indeed a better choice because, by a theorem of Lurie [Lur04], every geometric stack X can be reconstructed from its quasi-coherent sheaf category $\mathbf{QCoh}(X)$. This fact becomes evident in our illustrative example \mathbf{BG} from above: We observe that $\mathbf{QCoh}(\mathbf{BG}) = \mathbf{Rep}_{\mathbb{K}}(G)$ is the symmetric monoidal category of \mathbb{K} -linear representations of G . By Tannakian reconstruction, $\mathbf{Rep}_{\mathbb{K}}(G)$ encodes the full information about the group G , hence $\mathbf{QCoh}(\mathbf{BG})$ is indeed much richer than the function dg-algebra $\mathcal{O}(\mathbf{BG})$ considered in the previous paragraph (think of $G = \mathbb{Z}_2$ for example). Furthermore, for a “nice” space X , i.e. an affine scheme over \mathbb{K} , the usual function algebra $\mathcal{O}(X)$ can be recovered as follows: One finds that in this case $\mathbf{QCoh}(X) \simeq \mathbf{Mod}_{\mathcal{O}(X)}$ is the symmetric monoidal \mathbb{K} -linear category of (right) modules over the commutative algebra $\mathcal{O}(X)$, hence $\mathcal{O}(X)$ can be reconstructed from $\mathbf{QCoh}(X)$ as the endomorphism algebra $\mathrm{End}(\mathcal{O}(X)) \cong \mathcal{O}(X)$ of the rank 1 free module $\mathcal{O}(X) \in \mathbf{Mod}_{\mathcal{O}(X)}$, i.e. as the endomorphism algebra of the monoidal unit of $\mathbf{QCoh}(X)$. This means that for “nice” spaces the function algebra perspective and the quasi-coherent sheaf category perspective are compatible and carry the same information.

The previous paragraph explained the need to move from the function algebra perspective to the quasi-coherent sheaf category perspective. This, however, raises another question: What does it mean to quantize a quasi-coherent sheaf category? As an illustrative example, let us start with the case where the space X is “nice”, i.e. affine, and assume that we already have a non-commutative algebra $A \in \mathbf{Alg}_{\mathbb{K}}$ that quantizes the function algebra $\mathcal{O}(X)$. We may then form the locally presentable \mathbb{K} -linear category \mathbf{Mod}_A of right A -modules and interpret it as a quantization of the quasi-coherent sheaf category $\mathbf{QCoh}(X) \simeq \mathbf{Mod}_{\mathcal{O}(X)}$. It is important to observe the following structural difference between \mathbf{Mod}_A and $\mathbf{Mod}_{\mathcal{O}(X)}$: The tensor product $\otimes_{\mathcal{O}(X)}$ on $\mathbf{Mod}_{\mathcal{O}(X)}$ is only well-defined because $\mathcal{O}(X)$ is a *commutative* algebra and hence every right $\mathcal{O}(X)$ -module is automatically an $\mathcal{O}(X)$ -bimodule. Since the quantized algebra A is non-commutative, there is no counterpart on \mathbf{Mod}_A of the tensor product structure on $\mathbf{Mod}_{\mathcal{O}(X)}$. However, there is a counterpart on \mathbf{Mod}_A of the monoidal unit $\mathcal{O}(X) \in \mathbf{Mod}_{\mathcal{O}(X)}$, which is given by the pointing $A \in \mathbf{Mod}_A$ of the \mathbb{K} -linear category \mathbf{Mod}_A , see also [JF15]. This suggests that the quantization of the *symmetric monoidal* \mathbb{K} -linear category $\mathbf{QCoh}(X)$ should be a *pointed* \mathbb{K} -linear category. We would like to emphasize that this idea was made precise in the framework of derived algebraic geometry, see [PTVV13, CPTVV17] and also Toën’s ICM 2014 contribution [Toe14]. In this context, the quantization of a derived stack X endowed with an n -shifted symplectic structure is described by a quantization of the symmetric monoidal ∞ -category $\mathbf{QCoh}(X)$ of quasi-coherent sheaves as an E_n -monoidal ∞ -category. Since the phase space of a physical system is 0-shifted symplectic, we recover our intuition that one should quantize $\mathbf{QCoh}(X)$ as an E_0 -monoidal, i.e. pointed, \mathbb{K} -linear category.

Let us explain in more detail the content of the present paper and the results we obtain: In Section 2 we introduce an equivalent perspective on ordinary AQFTs as prefactorization algebras [CG17] with values in the symmetric monoidal category $\mathbf{Alg}_{\mathbb{K}}$ of associative and unital \mathbb{K} -algebras. This perspective will be used in Section 3 to introduce our novel concept of categorified AQFTs (called *2AQFTs*) that describe quantum observables by locally presentable \mathbb{K} -linear categories, in contrast to associative and unital algebras. In more detail, we define a 2AQFT as a (weak) prefactorization algebra on an orthogonal category $\overline{\mathbf{C}}$ (cf. Definition 2.1) with values in the symmetric monoidal 2-category $\mathbf{Pr}_{\mathbb{K}}$ of locally presentable \mathbb{K} -linear categories. In Section 4 we explore the relationship between ordinary AQFTs and our new concept of 2AQFTs. We

construct a biadjunction (cf. Theorem 4.3) that exhibits the 1-category of ordinary AQFTs as a coreflective full 2-subcategory of the 2-category of 2AQFTs, hence our framework for 2AQFTs includes ordinary AQFTs faithfully. In Section 5 we develop a local gauging construction for AQFTs with finite group actions, which allows us to construct concrete examples of 2AQFTs that admit an interpretation as categorified orbifold theories. In particular, we exhibit examples of 2AQFTs (cf. Example 5.7) that are not truncated, i.e. not equivalent to ordinary AQFTs, thus showing that our concept of 2AQFT is a genuine generalization of ordinary AQFTs. The main result of this section is Theorem 5.11: We prove that a categorified orbifold theory is truncated, i.e. equivalent to an ordinary AQFT, if and only if a suitable Hopf-Galois condition is fulfilled, which can be interpreted as a non-commutative analog of the condition that a G -action on a space is *free*. This matches with the intuition that an orbifold σ -model with a *quotient stack* $X//G$ as target boils down to an ordinary σ -model with the *quotient space* X/G as target whenever G acts freely on the space X (cf. Remark 5.12). In Section 6 we study a local-to-global extension for 2AQFTs (called *Fredenhagen's universal category*), which is a higher categorical analog of *Fredenhagen's universal algebra* [Fre90, Fre93, FRS92]. We develop concrete models for computing Fredenhagen's universal category and provide simple examples for extensions of 2AQFTs from intervals to the circle \mathbb{S}^1 . Appendix A introduces the relevant formalism for 2-categorical operad theory that we use throughout this paper.

2 AQFTs and algebra-valued prefactorization algebras

Let us fix once and for all a field \mathbb{K} of characteristic zero. We briefly review the definition of algebraic quantum field theories (AQFTs) on an orthogonal category $\overline{\mathbf{C}}$ in the sense of [BSW17]. We then prove that such theories admit an equivalent description in terms of prefactorization algebras on $\overline{\mathbf{C}}$ with values in the symmetric monoidal category $\mathbf{Alg}_{\mathbb{K}}$ of associative and unital \mathbb{K} -algebras. The latter perspective will be particularly useful for developing a categorification of AQFTs in Section 3.

Definition 2.1. An *orthogonal category* $\overline{\mathbf{C}} := (\mathbf{C}, \perp)$ is a small category \mathbf{C} together with a subset $\perp \subseteq \text{Mor}_{\mathbf{C}} \times_t \text{Mor}_{\mathbf{C}}$ of the set of pairs of morphisms with a common target, such that:

- (i) If $(f_1, f_2) \in \perp$, then $(f_2, f_1) \in \perp$.
- (ii) If $(f_1, f_2) \in \perp$, then $(g f_1 h_1, g f_2 h_2) \in \perp$, for all composable \mathbf{C} -morphisms g, h_1 and h_2 .

We shall denote orthogonal pairs $(f_1, f_2) \in \perp$ also by $f_1 \perp f_2$.

Example 2.2. Let $\mathbf{Open}(M)$ be the category of non-empty open subsets $U \subseteq M$ of a manifold M with morphisms $U \rightarrow V$ given by subset inclusions $U \subseteq V \subseteq M$. We introduce an orthogonality relation \perp_M by declaring two morphisms $U_1, U_2 \subseteq V \subseteq M$ to be orthogonal if and only if $U_1 \cap U_2 = \emptyset$. The orthogonal category $\overline{\mathbf{Open}(M)} := (\mathbf{Open}(M), \perp_M)$ features in factorization algebras [CG17] and, for $M = \mathbb{S}^1$ the circle, also in chiral conformal AQFT [Kaw15]. ∇

Example 2.3. The orthogonal category $\overline{\mathbf{Loc}}$ that features in locally covariant AQFT [BFV03, FV15] is given by the category \mathbf{Loc} of oriented and time-oriented globally hyperbolic Lorentzian manifolds, with orthogonality relation $\perp_{\mathbf{Loc}}$ determined by causal disjointness, see e.g. [BSW17] for the details. For a fixed $M \in \mathbf{Loc}$, consider the slice category \mathbf{Loc}/M and pull back the orthogonality relation $\perp_{\mathbf{Loc}}$ along the canonical functor $\mathbf{Loc}/M \rightarrow \mathbf{Loc}$. The resulting orthogonal category $\overline{\mathbf{Loc}/M}$ features in Haag-Kaster style AQFT [HK64] on a fixed $M \in \mathbf{Loc}$. ∇

Definition 2.4. An *algebraic quantum field theory* (AQFT) on an orthogonal category $\overline{\mathbf{C}}$ is a functor $\mathfrak{A} : \mathbf{C} \rightarrow \mathbf{Alg}_{\mathbb{K}}$ to the category of associative and unital \mathbb{K} -algebras that satisfies the \perp -commutativity property: For all orthogonal pairs $(f_1 : c_1 \rightarrow t) \perp (f_2 : c_2 \rightarrow t)$, the induced commutator

$$[\mathfrak{A}(f_1)(-), \mathfrak{A}(f_2)(-)]_{\mathfrak{A}(t)} : \mathfrak{A}(c_1) \otimes \mathfrak{A}(c_2) \longrightarrow \mathfrak{A}(t) \quad (2.1)$$

is zero. The category of AQFTs on $\overline{\mathbf{C}}$ is the full subcategory

$$\mathbf{AQFT}(\overline{\mathbf{C}}) \subseteq \mathbf{Fun}(\mathbf{C}, \mathbf{Alg}_{\mathbb{K}}) \quad (2.2)$$

of the functor category that consists of all \perp -commutative functors.

In preparation for our definition of categorified AQFTs in Section 3, we prove that the category $\mathbf{AQFT}(\overline{\mathbf{C}})$ is equivalent to the category of $\mathbf{Alg}_{\mathbb{K}}$ -valued prefactorization algebras on $\overline{\mathbf{C}}$. The following definition introduces a colored operad $\mathcal{P}_{\overline{\mathbf{C}}}$ that generalizes the prefactorization operad of Costello and Gwilliam [CG17] to an arbitrary orthogonal category $\overline{\mathbf{C}}$. The operad of Costello and Gwilliam is recovered by taking $\overline{\mathbf{C}} = \overline{\mathbf{Open}}(M)$ for a manifold M , see Example 2.2. For the relevant background and notations for colored operads we refer the reader to [Yau16, BSW17] and also to Appendix A.

Definition 2.5. The *prefactorization operad* $\mathcal{P}_{\overline{\mathbf{C}}}$ associated to an orthogonal category $\overline{\mathbf{C}}$ is the **Set**-valued colored operad defined by the following data:

- (1) The objects of $\mathcal{P}_{\overline{\mathbf{C}}}$ are the objects of the category \mathbf{C} .
- (2) The sets of operations are

$$\mathcal{P}_{\overline{\mathbf{C}}}(\underline{c}) := \left\{ \underline{f} := (f_1, \dots, f_n) \in \prod_{i=1}^n \mathbf{C}(c_i, t) : f_i \perp f_j \text{ for all } i \neq j \right\} \quad , \quad (2.3)$$

for each object $t \in \mathbf{C}$ and each tuple of objects $\underline{c} := (c_1, \dots, c_n) \in \mathbf{C}^n$. For the empty tuple $\underline{c} = \emptyset$, we set $\mathcal{P}_{\overline{\mathbf{C}}}(\emptyset) := \{*_t\}$ to be a singleton.

- (3) The composition maps $\gamma : \mathcal{P}_{\overline{\mathbf{C}}}(\underline{a}) \times \prod_{i=1}^n \mathcal{P}_{\overline{\mathbf{C}}}(\underline{b}_i) \rightarrow \mathcal{P}_{\overline{\mathbf{C}}}(\underline{b})$, where $\underline{b} := (b_1, \dots, b_n)$ denotes the concatenation of tuples, are given by composition in the category \mathbf{C} , i.e.

$$\gamma(\underline{f}, (\underline{g}_1, \dots, \underline{g}_n)) := \underline{f} \underline{g} := (f_1 g_{11}, \dots, f_1 g_{1k_1}, \dots, f_n g_{n1}, \dots, f_n g_{nk_n}) \quad . \quad (2.4)$$

- (4) The identity operations are $\mathbb{1} := \text{id}_t \in \mathcal{P}_{\overline{\mathbf{C}}}(\underline{t})$.
- (5) The permutation actions $\mathcal{P}_{\overline{\mathbf{C}}}(\sigma) : \mathcal{P}_{\overline{\mathbf{C}}}(\underline{c}) \rightarrow \mathcal{P}_{\overline{\mathbf{C}}}(\underline{c}\sigma)$ are given by

$$\mathcal{P}_{\overline{\mathbf{C}}}(\sigma)(\underline{f}) := \underline{f}\sigma := (f_{\sigma(1)}, \dots, f_{\sigma(n)}) \quad . \quad (2.5)$$

Let us endow the category $\mathbf{Alg}_{\mathbb{K}}$ of associative and unital \mathbb{K} -algebras with its standard symmetric monoidal structure. The tensor product of two algebras $A, B \in \mathbf{Alg}_{\mathbb{K}}$ is given by the tensor product algebra $A \otimes B$. Concretely, that is the tensor product of vector spaces with multiplication given by $(a \otimes b)(a' \otimes b') := (aa') \otimes (bb')$ and unit element $1_A \otimes 1_B \in A \otimes B$. The monoidal unit is $\mathbb{K} \in \mathbf{Alg}_{\mathbb{K}}$ and the symmetric braiding is given by the $\mathbf{Alg}_{\mathbb{K}}$ -morphisms $\tau : A \otimes B \rightarrow B \otimes A$, $a \otimes b \mapsto b \otimes a$. The symmetric monoidal category $\mathbf{Alg}_{\mathbb{K}}$ has an underlying **Set**-valued colored operad (see e.g. [EM09]) that we denote by the same symbol $\mathbf{Alg}_{\mathbb{K}}$. Concretely, the objects are the objects of $\mathbf{Alg}_{\mathbb{K}}$ and the sets of operations are given by

$$\mathbf{Alg}_{\mathbb{K}}(\underline{A}) := \mathbf{Alg}_{\mathbb{K}}\left(\bigotimes_{i=1}^n A_i, B\right) \quad . \quad (2.6)$$

The composition maps are determined by the monoidal structure, the identity operations are the identity morphisms, and the permutation actions are obtained from the symmetric braiding.

Definition 2.6. The category of $\mathbf{Alg}_{\mathbb{K}}$ -valued *prefactorization algebras* on $\overline{\mathbf{C}}$ is defined by

$$\mathbf{Alg}_{\mathcal{P}_{\overline{\mathbf{C}}}}(\mathbf{Alg}_{\mathbb{K}}) := [\mathcal{P}_{\overline{\mathbf{C}}}, \mathbf{Alg}_{\mathbb{K}}] \quad , \quad (2.7)$$

where $\mathbf{Alg}_{\mathbb{K}}$ is regarded as a colored operad (as explained above) and $[-, -]$ denotes the Hom-category from Remark A.6.

Remark 2.7. Let us unpack this definition by using the definitions from Appendix A. (These definitions simplify drastically in the present case because both $\mathcal{P}_{\overline{\mathbf{C}}}$ and $\mathbf{Alg}_{\mathbb{K}}$ are **Set**-valued colored operads. Hence, all coherence data are necessarily trivial. Non-trivial coherence data will be needed to describe categorified AQFTs in Remark 3.4.)

An $\mathbf{Alg}_{\mathbb{K}}$ -valued prefactorization algebra $\mathfrak{F} \in \mathbf{Alg}_{\mathcal{P}_{\overline{\mathbf{C}}}}(\mathbf{Alg}_{\mathbb{K}})$ is given by the following data:

- (1) For each $c \in \mathbf{C}$, an associative and unital \mathbb{K} -algebra $\mathfrak{F}(c) \in \mathbf{Alg}_{\mathbb{K}}$.
- (2) For each tuple $\underline{f} = (f_1, \dots, f_n) \in \mathcal{P}_{\overline{\mathbf{C}}}(\underline{t})$ of mutually orthogonal \mathbf{C} -morphisms, an $\mathbf{Alg}_{\mathbb{K}}$ -morphism (called *factorization product*)

$$\mathfrak{F}(\underline{f}) : \bigotimes_{i=1}^n \mathfrak{F}(c_i) \longrightarrow \mathfrak{F}(t) \quad (2.8)$$

from the tensor product algebra $\bigotimes_{i=1}^n \mathfrak{F}(c_i)$. For the empty tuple $\underline{c} = \emptyset$, the $\mathbf{Alg}_{\mathbb{K}}$ -morphism $\mathfrak{F}(*_t) : \mathbb{K} \rightarrow \mathfrak{F}(t)$ associated to the only element $*_t \in \mathcal{P}_{\overline{\mathbf{C}}}(\emptyset)$ is necessarily the unit of $\mathfrak{F}(t)$, because \mathbb{K} is the initial object in $\mathbf{Alg}_{\mathbb{K}}$.

These data are required to satisfy the following axioms:

$$\begin{array}{ccc} \bigotimes_{i=1}^n \bigotimes_{j=1}^{k_i} \mathfrak{F}(b_{ij}) & \xrightarrow{\mathfrak{F}(\underline{g}) := \bigotimes_i \mathfrak{F}(g_i)} & \bigotimes_{i=1}^n \mathfrak{F}(a_i) \\ & \searrow \mathfrak{F}(\underline{f}\underline{g}) & \downarrow \mathfrak{F}(\underline{f}) \\ & & \mathfrak{F}(t) \end{array} \quad (2.9a)$$

$$\mathfrak{F}(\text{id}_t) = \text{id}_{\mathfrak{F}(t)} : \mathfrak{F}(t) \longrightarrow \mathfrak{F}(t) \quad (2.9b)$$

$$\begin{array}{ccc} \bigotimes_{i=1}^n \mathfrak{F}(c_{\sigma(i)}) & \xrightarrow{\tau_\sigma} & \bigotimes_{i=1}^n \mathfrak{F}(c_i) \\ & \searrow \mathfrak{F}(\underline{f}\sigma) & \downarrow \mathfrak{F}(\underline{f}) \\ & & \mathfrak{F}(t) \end{array} \quad (2.9c)$$

In the last diagram, we have denoted by $\tau_\sigma : \bigotimes_{i=1}^n \mathfrak{F}(c_{\sigma(i)}) \rightarrow \bigotimes_{i=1}^n \mathfrak{F}(c_i)$, $a_{\sigma(1)} \otimes \dots \otimes a_{\sigma(n)} \mapsto a_1 \otimes \dots \otimes a_n$ the $\mathbf{Alg}_{\mathbb{K}}$ -morphism that permutes the factors of the tensor product algebra.

A morphism $\zeta : \mathfrak{F} \rightarrow \mathfrak{G}$ in $\mathbf{Alg}_{\mathcal{P}_{\overline{\mathbf{C}}}}(\mathbf{Alg}_{\mathbb{K}})$ is a family of $\mathbf{Alg}_{\mathbb{K}}$ -morphisms $\zeta_c : \mathfrak{F}(c) \rightarrow \mathfrak{G}(c)$, for all $c \in \mathbf{C}$, such that the diagrams

$$\begin{array}{ccc} \bigotimes_{i=1}^n \mathfrak{F}(c_i) & \xrightarrow{\bigotimes_i \zeta_{c_i}} & \bigotimes_{i=1}^n \mathfrak{G}(c_i) \\ \mathfrak{F}(\underline{f}) \downarrow & & \downarrow \mathfrak{G}(\underline{f}) \\ \mathfrak{F}(t) & \xrightarrow{\zeta_t} & \mathfrak{G}(t) \end{array} \quad (2.10)$$

commute, for all $\underline{f} \in \mathcal{P}_{\overline{\mathbf{C}}}(\underline{t})$. △

It is easy to see that every $\mathfrak{A} \in \mathbf{AQFT}(\overline{\mathbf{C}})$ defines an $\mathbf{Alg}_{\mathbb{K}}$ -valued prefactorization algebra on $\overline{\mathbf{C}}$ by introducing the factorization products

$$\mathfrak{A}(\underline{f}) : \bigotimes_{i=1}^n \mathfrak{A}(c_i) \xrightarrow{\bigotimes_i \mathfrak{A}(f_i)} \mathfrak{A}(t)^{\otimes n} \xrightarrow{\mu_{\mathfrak{A}(t)}^n} \mathfrak{A}(t) \quad , \quad (2.11)$$

where $\mu_{\mathfrak{A}(t)}^n : \mathfrak{A}(t)^{\otimes n} \rightarrow \mathfrak{A}(t)$, $a_1 \otimes \cdots \otimes a_n \mapsto a_1 \cdots a_n$ denotes the n -ary multiplication in the associative and unital algebra $\mathfrak{A}(t) \in \mathbf{Alg}_{\mathbb{K}}$. Using \perp -commutativity and $f_i \perp f_j$, for all $i \neq j$, one shows that $\mathfrak{A}(\underline{f})$ is indeed an $\mathbf{Alg}_{\mathbb{K}}$ -morphism on the tensor product algebra. Furthermore, every $\mathbf{AQFT}(\overline{\mathbf{C}})$ -morphism $\zeta : \mathfrak{A} \rightarrow \mathfrak{B}$ defines an $\mathbf{Alg}_{\mathcal{P}_{\overline{\mathbf{C}}}}(\mathbf{Alg}_{\mathbb{K}})$ -morphism between the corresponding prefactorization algebras, hence we obtain a functor $\mathbf{AQFT}(\overline{\mathbf{C}}) \rightarrow \mathbf{Alg}_{\mathcal{P}_{\overline{\mathbf{C}}}}(\mathbf{Alg}_{\mathbb{K}})$. Note that this functor is fully faithful.

Conversely, we have the following lemma showing that every $\mathbf{Alg}_{\mathbb{K}}$ -valued prefactorization algebra is completely determined by an underlying AQFT.

Lemma 2.8. *For every $\mathfrak{F} \in \mathbf{Alg}_{\mathcal{P}_{\overline{\mathbf{C}}}}(\mathbf{Alg}_{\mathbb{K}})$, the factorization product $\mathfrak{F}(\underline{f})$ for $\underline{f} = (f_1, \dots, f_n) \in \mathcal{P}_{\overline{\mathbf{C}}}(\underline{t})$ factorizes as*

$$\mathfrak{F}(\underline{f}) : \bigotimes_{i=1}^n \mathfrak{F}(c_i) \xrightarrow{\otimes_i \mathfrak{F}(f_i)} \mathfrak{F}(t)^{\otimes n} \xrightarrow{\mu_{\mathfrak{F}(t)}^n} \mathfrak{F}(t) \quad , \quad (2.12)$$

where $\mu_{\mathfrak{F}(t)}^n$ denotes the n -ary multiplication in the associative and unital algebra $\mathfrak{F}(t) \in \mathbf{Alg}_{\mathbb{K}}$. In particular, \mathfrak{F} is completely determined by its underlying functor $\mathfrak{F} : \mathbf{C} \rightarrow \mathbf{Alg}_{\mathbb{K}}$, which satisfies the \perp -commutativity property from Definition 2.4 and hence defines an AQFT.

Proof. Using the composition maps from Definition 2.5, we compute

$$\gamma(\underline{f}, (*_{c_1}, \dots, *_{c_{k-1}}, \text{id}_{c_k}, *_{c_{k+1}}, \dots, *_{c_n})) = f_k \quad , \quad (2.13)$$

for all $k = 1, \dots, n$, where $*_t \in \mathcal{P}_{\overline{\mathbf{C}}}(\underline{t})$ denotes the unique arity zero operation. The corresponding commutative diagram in (2.9a) then reads as

$$\begin{array}{ccc} \bigotimes_{i=1}^n \mathfrak{F}(c_i) & \xrightarrow{\mathfrak{F}(\underline{f})} & \mathfrak{F}(t) \\ \uparrow \otimes_i \mathfrak{F}(*_{c_i}) \otimes \mathfrak{F}(\text{id}_{c_k}) \otimes \otimes_i \mathfrak{F}(*_{c_i}) & & \uparrow \mathfrak{F}(f_k) \\ \bigotimes_{i=1}^{k-1} \mathbb{K} \otimes \mathfrak{F}(c_k) \otimes \bigotimes_{i=k+1}^n \mathbb{K} & \xrightarrow{\cong} & \mathfrak{F}(c_k) \end{array} \quad (2.14)$$

Using further that $\mathfrak{F}(\text{id}_{c_k}) = \text{id}_{\mathfrak{F}(c_k)}$ (cf. (2.9b)) and that $\mathfrak{F}(*_{c_i}) : \mathbb{K} \rightarrow \mathfrak{F}(c_i)$ is the unit of $\mathfrak{F}(c_i) \in \mathbf{Alg}_{\mathbb{K}}$ (cf. Remark 2.7), the commutative diagram (2.14) implies that

$$\mathfrak{F}(\underline{f})(1_{\mathfrak{F}(c_1)} \otimes \cdots \otimes 1_{\mathfrak{F}(c_{k-1})} \otimes a_k \otimes 1_{\mathfrak{F}(c_{k+1})} \otimes \cdots \otimes 1_{\mathfrak{F}(c_n)}) = \mathfrak{F}(f_k)(a_k) \quad , \quad (2.15)$$

for all $a_k \in \mathfrak{F}(c_k)$. By definition of the product of a tensor product algebra, it then follows that

$$\mathfrak{F}(\underline{f})(a_1 \otimes \cdots \otimes a_n) = \mathfrak{F}(f_1)(a_1) \cdots \mathfrak{F}(f_n)(a_n) \quad , \quad (2.16)$$

for all $a_1 \otimes \cdots \otimes a_n \in \bigotimes_{i=1}^n \mathfrak{F}(c_i)$, which proves (2.12). Using further that every two elements of the form $a \otimes 1_B$ and $1_A \otimes b$ commute in a tensor product algebra $A \otimes B$, it follows that the underlying functor $\mathfrak{F} : \mathbf{C} \rightarrow \mathbf{Alg}_{\mathbb{K}}$ is \perp -commutative. \square

Summing up, we have proven the following

Theorem 2.9. *For every orthogonal category $\overline{\mathbf{C}}$, there exists a canonical isomorphism*

$$\mathbf{AQFT}(\overline{\mathbf{C}}) \cong \mathbf{Alg}_{\mathcal{P}_{\overline{\mathbf{C}}}}(\mathbf{Alg}_{\mathbb{K}}) \quad (2.17)$$

between the category of AQFTs on $\overline{\mathbf{C}}$ and the category of $\mathbf{Alg}_{\mathbb{K}}$ -valued prefactorization algebras on $\overline{\mathbf{C}}$.

Remark 2.10. The equivalent description of AQFTs in terms of $\mathbf{Alg}_{\mathbb{K}}$ -valued prefactorization algebras provides an interesting conceptual interpretation of the \perp -commutativity property from Definition 2.4. From the prefactorization algebra point of view, every quantum field theory comes with two different kinds of “multiplications”, namely the object-wise products $\mu_{\mathfrak{F}(c)} : \mathfrak{F}(c) \otimes \mathfrak{F}(c) \rightarrow \mathfrak{F}(c)$, for every $c \in \mathbf{C}$, and the factorization products $\mathfrak{F}(\underline{f}) : \bigotimes_{i=1}^n \mathfrak{F}(c_i) \rightarrow \mathfrak{F}(t)$, for every tuple \underline{f} of mutually orthogonal \mathbf{C} -morphisms. These two kinds of “multiplications” are compatible with each other because the factorization products $\mathfrak{F}(\underline{f})$ are $\mathbf{Alg}_{\mathbb{K}}$ -morphisms. The \perp -commutativity property is thus a consequence of an Eckmann-Hilton argument. \triangle

3 Definition of 2AQFTs

The aim of this section is to introduce a categorification of the concept of AQFTs, which we shall call *2-categorical algebraic quantum field theories* (2AQFTs). While ordinary AQFTs assign associative and unital \mathbb{K} -algebras to the objects of an orthogonal category $\overline{\mathbf{C}}$, our novel concept of 2AQFTs will assign locally presentable \mathbb{K} -linear categories, cf. [AR94, BCJF15].

Recall that a \mathbb{K} -linear category is a category \mathbf{D} that is enriched over the symmetric monoidal category $\mathbf{Vec}_{\mathbb{K}}$ of vector spaces over \mathbb{K} . Concretely, this means that we have a vector space $\mathbf{D}(d, d') \in \mathbf{Vec}_{\mathbb{K}}$ of morphisms, for every pair of objects $d, d' \in \mathbf{D}$, and that the composition maps $\circ : \mathbf{D}(d', d'') \times \mathbf{D}(d, d') \rightarrow \mathbf{D}(d, d'')$ are \mathbb{K} -bilinear, for all $d, d', d'' \in \mathbf{D}$. Given two \mathbb{K} -linear categories \mathbf{D} and \mathbf{E} , a \mathbb{K} -linear functor $F : \mathbf{D} \rightarrow \mathbf{E}$ is a functor such that the maps $F : \mathbf{D}(d, d') \rightarrow \mathbf{E}(Fd, Fd')$ are \mathbb{K} -linear, for all $d, d' \in \mathbf{D}$.

A \mathbb{K} -linear category \mathbf{D} is called *locally presentable* if it is 1.) cocomplete, i.e. has all small colimits, and 2.) generated under small colimits by a set $\Gamma \subset \mathbf{D}_0$ of objects that are λ -presentable for some infinite cardinal λ , see e.g. [BCJF15] for a recollection of the relevant material on locally presentable categories. The natural concept of functors $F : \mathbf{D} \rightarrow \mathbf{E}$ between two locally presentable \mathbb{K} -linear categories \mathbf{D} and \mathbf{E} is given by *co-continuous \mathbb{K} -linear functors*, i.e. \mathbb{K} -linear functors that preserve all small colimits. Natural transformations in this context are just ordinary natural transformations.

Definition 3.1. The operad $\mathbf{Pr}_{\mathbb{K}}$ of *locally presentable \mathbb{K} -linear categories* is the \mathbf{Cat} -enriched colored operad (cf. Definition A.1) defined by the following data:

- (1) The objects are all locally presentable \mathbb{K} -linear categories.
- (2) For \mathbf{T} and $\underline{\mathbf{D}} = (\mathbf{D}_1, \dots, \mathbf{D}_n)$ locally presentable \mathbb{K} -linear categories, the category of operations is the full subcategory

$$\mathbf{Pr}_{\mathbb{K}}(\underline{\mathbf{D}}) := \mathbf{MultLin}_{\mathbb{K},c} \left(\prod_{i=1}^n \mathbf{D}_i, \mathbf{T} \right) \subseteq \mathbf{Fun} \left(\prod_{i=1}^n \mathbf{D}_i, \mathbf{T} \right) \quad (3.1)$$

of the functor category that consists of all functors $F : \prod_{i=1}^n \mathbf{D}_i \rightarrow \mathbf{T}$ that are \mathbb{K} -linear and co-continuous in each variable. For the empty tuple $\underline{\mathbf{D}} = \emptyset$, we set $\mathbf{Pr}_{\mathbb{K}}(\underline{\emptyset}) := \mathbf{Fun}(\mathbf{1}, \mathbf{T})$, where $\mathbf{1}$ is the category with only one object and its identity morphism.

- (3) The composition functors $\gamma : \mathbf{Pr}_{\mathbb{K}}(\underline{\mathbf{D}}) \times \prod_{i=1}^n \mathbf{Pr}_{\mathbb{K}}(\underline{\mathbf{E}}^i) \rightarrow \mathbf{Pr}_{\mathbb{K}}(\underline{\mathbf{E}})$ are given by composition of functors and (horizontal) composition of natural transformations, i.e.

$$\gamma(F, (G_1, \dots, G_n)) := F \underline{G} := F \prod_{i=1}^n G_i \quad , \quad (3.2a)$$

$$\gamma(\alpha, (\beta_1, \dots, \beta_n)) := \alpha * \underline{\beta} := \alpha * \prod_{i=1}^n \beta_i \quad . \quad (3.2b)$$

- (4) The identity 1-operations are the identity functors $\mathbf{1} := \text{id}_{\mathbf{T}} \in \mathbf{Pr}_{\mathbb{K}}(\underline{\mathbf{T}}) \subseteq \mathbf{Fun}(\mathbf{T}, \mathbf{T})$.

(5) The permutation action functors $\mathbf{Pr}_{\mathbb{K}}(\sigma) : \mathbf{Pr}_{\mathbb{K}}(\underline{\mathbf{D}}) \rightarrow \mathbf{Pr}_{\mathbb{K}}(\underline{\mathbf{D}}_{\sigma})$ are given by

$$\mathbf{Pr}_{\mathbb{K}}(\sigma)(F) := F \text{ flip}_{\sigma} \quad , \quad \mathbf{Pr}_{\mathbb{K}}(\sigma)(\alpha) := \alpha * \text{Id}_{\text{flip}_{\sigma}} \quad , \quad (3.3)$$

where $\text{flip}_{\sigma} : \prod_{i=1}^n \mathbf{D}_{\sigma(i)} \rightarrow \prod_{i=1}^n \mathbf{D}_i$ is the permutation functor and $\text{Id}_{\text{flip}_{\sigma}} : \text{flip}_{\sigma} \Rightarrow \text{flip}_{\sigma}$ its identity natural transformation.

Remark 3.2. With some abuse of notation, we will sometimes denote by the same symbol $\mathbf{Pr}_{\mathbb{K}}$ the underlying 2-category of 1-ary operations of the \mathbf{Cat} -enriched colored operad from Definition 3.1. It should be clear from the context whether we mean by the symbol $\mathbf{Pr}_{\mathbb{K}}$ a \mathbf{Cat} -enriched colored operad or a 2-category. The underlying 2-category $\mathbf{Pr}_{\mathbb{K}}$ is known to be (closed) symmetric monoidal with respect to the Kelly-Deligne tensor product $\mathbf{D} \boxtimes \mathbf{E}$ of locally presentable \mathbb{K} -linear categories, whose monoidal unit is given by the \mathbb{K} -linear category $\mathbf{Vec}_{\mathbb{K}}$ of vector spaces, see [Kel05] and also [BCJF15] for a review. This symmetric monoidal structure is linked as follows to our \mathbf{Cat} -enriched colored operad from Definition 3.1: By the universal property of the Kelly-Deligne tensor product, the categories of operations

$$\mathbf{Pr}_{\mathbb{K}}(\underline{\mathbf{D}}) \simeq \mathbf{Lin}_{\mathbb{K},c} \left(\bigboxtimes_{i=1}^n \mathbf{D}_i, \mathbf{T} \right) \quad (3.4)$$

are equivalent to the categories of co-continuous \mathbb{K} -linear functors out of the Kelly-Deligne tensor product. Hence, the \mathbf{Cat} -enriched colored operad $\mathbf{Pr}_{\mathbb{K}}$ can also be understood as the underlying operad of the symmetric monoidal 2-category $(\mathbf{Pr}_{\mathbb{K}}, \boxtimes, \mathbf{Vec}_{\mathbb{K}})$. This alternative perspective will become useful in some of our computations in Sections 4, 5 and 6. \triangle

Recalling Theorem 2.9, ordinary AQFTs on $\overline{\mathbf{C}}$ are equivalently $\mathbf{Alg}_{\mathbb{K}}$ -valued prefactorization algebras, i.e. $\mathbf{AQFT}(\overline{\mathbf{C}}) \cong \mathbf{Alg}_{\mathcal{P}_{\overline{\mathbf{C}}}}(\mathbf{Alg}_{\mathbb{K}})$. Replacing the target $\mathbf{Alg}_{\mathbb{K}}$ with $\mathbf{Pr}_{\mathbb{K}}$ suggests the following

Definition 3.3. The 2-category of 2-categorical algebraic quantum field theories (2AQFTs) on an orthogonal category $\overline{\mathbf{C}}$ is defined as the Hom-2-category (cf. Remark A.6)

$$\mathbf{2AQFT}(\overline{\mathbf{C}}) := \mathbf{Alg}_{\mathcal{P}_{\overline{\mathbf{C}}}}(\mathbf{Pr}_{\mathbb{K}}) := [\mathcal{P}_{\overline{\mathbf{C}}}, \mathbf{Pr}_{\mathbb{K}}] \quad , \quad (3.5)$$

where $\mathcal{P}_{\overline{\mathbf{C}}}$ is the prefactorization operad from Definition 2.5 and $\mathbf{Pr}_{\mathbb{K}}$ is the \mathbf{Cat} -enriched colored operad from Definition 3.1.

Remark 3.4. Let us unpack this definition by using the definitions from Appendix A.

An object $\mathfrak{A} \in \mathbf{2AQFT}(\overline{\mathbf{C}})$ is given by the following data:

- (1) For each $c \in \mathbf{C}$, a locally presentable \mathbb{K} -linear category $\mathfrak{A}(c) \in \mathbf{Pr}_{\mathbb{K}}$.
- (2) For each tuple $\underline{f} = (f_1, \dots, f_n) \in \mathcal{P}_{\overline{\mathbf{C}}}(\underline{c})$ of mutually orthogonal \mathbf{C} -morphisms, a functor (called *factorization product*)

$$\mathfrak{A}(\underline{f}) : \prod_{i=1}^n \mathfrak{A}(c_i) \longrightarrow \mathfrak{A}(t) \quad (3.6)$$

that is \mathbb{K} -linear and co-continuous in each variable. For the empty tuple $\underline{c} = \emptyset$, this is an object $\mathfrak{a}_t := \mathfrak{A}(*_t) \in \mathfrak{A}(t)$ (called *pointing*) that is associated to the only element $*_t \in \mathcal{P}_{\overline{\mathbf{C}}}(\emptyset)$.

- (3) For each $\underline{f} \in \mathcal{P}_{\overline{\mathbf{C}}}(\underline{a})$ and $\underline{g} = (g_1, \dots, g_n) \in \prod_{i=1}^n \mathcal{P}_{\overline{\mathbf{C}}}(\underline{b}_i)$, a natural isomorphism

$$\begin{array}{ccc} \prod_{i=1}^n \prod_{j=1}^{k_i} \mathfrak{A}(b_{ij}) & \xrightarrow{\mathfrak{A}(\underline{g}) := \prod_i \mathfrak{A}(g_i)} & \prod_{i=1}^n \mathfrak{A}(a_i) \\ & \searrow \mathfrak{A}(\underline{f}, \underline{g}) & \downarrow \mathfrak{A}(\underline{f}) \\ & \mathfrak{A}(\underline{f}, \underline{g}) & \mathfrak{A}(t) \end{array} \quad (3.7)$$

(4) For each $t \in \mathbf{C}$, a natural isomorphism

$$\begin{array}{ccc}
 & \text{id}_{\mathfrak{A}(t)} & \\
 & \curvearrowright & \\
 \mathfrak{A}(t) & \begin{array}{c} \Downarrow \\ \mathfrak{A}_t^0 \\ \Downarrow \end{array} & \mathfrak{A}(t) \\
 & \curvearrowleft & \\
 & \mathfrak{A}(\text{id}_t) &
 \end{array} \tag{3.8}$$

(5) For each $\underline{f} \in \mathcal{P}_{\overline{\mathbf{C}}}(t)$ and permutation $\sigma \in \Sigma_n$, a natural isomorphism

$$\begin{array}{ccc}
 \prod_{i=1}^n \mathfrak{A}(c_{\sigma(i)}) & \xrightarrow{\text{flip}_\sigma} & \prod_{i=1}^n \mathfrak{A}(c_i) \\
 & \searrow \mathfrak{A}(\underline{f}\sigma) & \downarrow \mathfrak{A}(\underline{f}) \\
 & & \mathfrak{A}(t)
 \end{array}
 \begin{array}{c}
 \swarrow \mathfrak{A}_f^\sigma \\
 \swarrow \mathfrak{A}(\underline{f})
 \end{array} \tag{3.9}$$

These data are required to satisfy the axioms from Definition A.2.

A 1-morphism $\zeta : \mathfrak{A} \rightarrow \mathfrak{B}$ in $\mathbf{2AQFT}(\overline{\mathbf{C}})$ is given by the following data:

- (1) For each $c \in \mathbf{C}$, a co-continuous \mathbb{K} -linear functor $\zeta_c : \mathfrak{A}(c) \rightarrow \mathfrak{B}(c)$.
- (2) For each $\underline{f} \in \mathcal{P}_{\overline{\mathbf{C}}}(t)$, a natural isomorphism

$$\begin{array}{ccc}
 \prod_{i=1}^n \mathfrak{A}(c_i) & \xrightarrow{\prod_i \zeta_{c_i}} & \prod_{i=1}^n \mathfrak{B}(c_i) \\
 \mathfrak{A}(\underline{f}) \downarrow & \swarrow \zeta_{\underline{f}} & \downarrow \mathfrak{B}(\underline{f}) \\
 \mathfrak{A}(t) & \xrightarrow{\zeta_t} & \mathfrak{B}(t)
 \end{array} \tag{3.10}$$

Note that, for $\underline{f} = *_t \in \mathcal{P}_{\overline{\mathbf{C}}}(t)$, this amounts to an isomorphism $\zeta_{*_t} : \mathbf{b}_t \xrightarrow{\cong} \zeta_t(\mathbf{a}_t)$ in $\mathfrak{B}(t)$ from the pointing $\mathbf{b}_t = \mathfrak{B}(*_t) \in \mathfrak{B}(t)$ to the image of the pointing $\mathbf{a}_t = \mathfrak{A}(*_t) \in \mathfrak{A}(t)$ under the functor $\zeta_t : \mathfrak{A}(t) \rightarrow \mathfrak{B}(t)$.

These data are required to satisfy the axioms from Definition A.4.

A 2-morphism $\Gamma : \zeta \Rightarrow \kappa$ between two 1-morphisms $\zeta, \kappa : \mathfrak{A} \rightarrow \mathfrak{B}$ in $\mathbf{2AQFT}(\overline{\mathbf{C}})$ is given by the following data:

- (1) For each $c \in \mathbf{C}$, a natural transformation

$$\begin{array}{ccc}
 & \zeta_c & \\
 & \curvearrowright & \\
 \mathfrak{A}(c) & \begin{array}{c} \Downarrow \\ \Gamma_c \\ \Downarrow \end{array} & \mathfrak{B}(c) \\
 & \curvearrowleft & \\
 & \kappa_c &
 \end{array} \tag{3.11}$$

These data are required to satisfy the axioms from Definition A.5. △

Remark 3.5. Category-valued prefactorization algebras were studied before in the context of factorization homology of 2-manifolds [BZBJ18a, BZBJ18b]. Our framework for 2AQFTs allows us to interpret the examples studied in these papers as 2-dimensional topological AQFTs. This is achieved by considering the orthogonal category $\overline{\mathbf{Man}}_2$ of 2-dimensional (oriented or framed) manifolds, with orthogonality relation given by disjointness, and restricting to topological theories by considering locally constant prefactorization algebras. △

4 Inclusion-truncation biadjunction

In this section we explore the relationship between ordinary AQFTs and our novel concept of 2AQFTs from Definition 3.3. We shall show that every $\mathfrak{A} \in \mathbf{2AQFT}(\overline{\mathbf{C}})$ has an underlying ordinary AQFT $\pi(\mathfrak{A}) \in \mathbf{AQFT}(\overline{\mathbf{C}})$, which we call the truncation of \mathfrak{A} . Our truncation construction is given by a 2-functor $\pi : \mathbf{2AQFT}(\overline{\mathbf{C}}) \rightarrow \mathbf{AQFT}(\overline{\mathbf{C}})$. We shall also define, for every $\mathfrak{A} \in \mathbf{AQFT}(\overline{\mathbf{C}})$, a 2AQFT $\iota(\mathfrak{A}) \in \mathbf{2AQFT}(\overline{\mathbf{C}})$ that assigns to each object $c \in \mathbf{C}$ the locally presentable \mathbb{K} -linear category $\iota(\mathfrak{A})(c) = \mathbf{Mod}_{\mathfrak{A}(c)}$ of right $\mathfrak{A}(c)$ -modules. This construction is given by an inclusion pseudo-functor $\iota : \mathbf{AQFT}(\overline{\mathbf{C}}) \rightarrow \mathbf{2AQFT}(\overline{\mathbf{C}})$. Inclusion and truncation are compatible with each other in the sense that they determine a biadjunction $\iota \dashv \pi$, see e.g. [Str80, Str87] and also [LN16] for the relevant bicategorical background. We prove that this biadjunction exhibits $\mathbf{AQFT}(\overline{\mathbf{C}})$ as a coreflective full 2-subcategory of $\mathbf{2AQFT}(\overline{\mathbf{C}})$. The conceptual meaning and relevance of this result is as follows: On the one hand, ordinary AQFTs can be studied equally well inside the 2-category of 2AQFTs by applying the fully faithful inclusion pseudo-functor $\iota : \mathbf{AQFT}(\overline{\mathbf{C}}) \rightarrow \mathbf{2AQFT}(\overline{\mathbf{C}})$. There is no loss of information in doing so, because the unit $\eta : \text{id} \Rightarrow \pi \iota$ of the biadjunction is a natural isomorphism and hence one can recover every $\mathfrak{A} \in \mathbf{AQFT}(\overline{\mathbf{C}})$ from its corresponding 2AQFT $\iota(\mathfrak{A})$ by applying the truncation 2-functor. On the other hand, the 2-category $\mathbf{2AQFT}(\overline{\mathbf{C}})$ has in general also objects that *do not* lie in the essential image of the inclusion pseudo-functor ι . These are the genuine 2AQFTs that are not fully determined by their truncation to an ordinary AQFT. We refer to Section 5 for concrete examples.

4.1 Truncation

Given any $\mathfrak{A} \in \mathbf{2AQFT}(\overline{\mathbf{C}})$, we define its truncation $\pi(\mathfrak{A}) \in \mathbf{AQFT}(\overline{\mathbf{C}})$ by providing the required data listed in Remark 2.7:

- (1) For each $c \in \mathbf{C}$, we set

$$\pi(\mathfrak{A})(c) := \text{End}(\mathbf{a}_c) := \mathfrak{A}(c)(\mathbf{a}_c, \mathbf{a}_c) \quad (4.1)$$

to be the endomorphism algebra of the pointing $\mathbf{a}_c \in \mathfrak{A}(c)$. (Note that this is an associative and unital \mathbb{K} -algebra, because $\mathfrak{A}(c)$ is a \mathbb{K} -linear category.)

- (2) For each non-empty tuple $\underline{f} = (f_1, \dots, f_n) \in \mathcal{P}_{\overline{\mathbf{C}}}(\underline{t})$ of mutually orthogonal \mathbf{C} -morphisms, the given functor $\mathfrak{A}(\underline{f}) : \prod_{i=1}^n \mathfrak{A}(c_i) \rightarrow \mathfrak{A}(t)$ restricts to endomorphism algebras as $\mathfrak{A}(\underline{f}) : \prod_{i=1}^n \text{End}(\mathbf{a}_{c_i}) \rightarrow \text{End}(\mathfrak{A}(\underline{f})(\mathbf{a}_{c_1}, \dots, \mathbf{a}_{c_n}))$. Because $\mathfrak{A}(\underline{f})$ is \mathbb{K} -linear in each variable, we obtain an $\mathbf{Alg}_{\mathbb{K}}$ -morphism $\mathfrak{A}(\underline{f}) : \otimes_{i=1}^n \text{End}(\mathbf{a}_{c_i}) \rightarrow \text{End}(\mathfrak{A}(\underline{f})(\mathbf{a}_{c_1}, \dots, \mathbf{a}_{c_n}))$ from the tensor product algebra. The coherence map in (3.7) that is associated to $(\underline{f}, *_{\underline{c}}) := (\underline{f}, (*_{c_1}, \dots, *_{c_n}))$ provides an isomorphism $\mathfrak{A}_{(\underline{f}, *_{\underline{c}})}^2 : \mathfrak{A}(\underline{f})(\mathbf{a}_{c_1}, \dots, \mathbf{a}_{c_n}) \rightarrow \mathbf{a}_t$ in the category $\mathfrak{A}(t)$, which we use to define the $\mathbf{Alg}_{\mathbb{K}}$ -morphism

$$\begin{aligned} \pi(\mathfrak{A})(\underline{f}) : \bigotimes_{i=1}^n \pi(\mathfrak{A})(c_i) &\longrightarrow \pi(\mathfrak{A})(t) \quad , \\ h_1 \otimes \dots \otimes h_n &\longmapsto \mathfrak{A}_{(\underline{f}, *_{\underline{c}})}^2 \circ \mathfrak{A}(\underline{f})(h_1, \dots, h_n) \circ (\mathfrak{A}_{(\underline{f}, *_{\underline{c}})}^2)^{-1} \quad , \end{aligned} \quad (4.2)$$

where \circ denotes composition in $\mathfrak{A}(t)$. As noted in Remark 2.7, the $\mathbf{Alg}_{\mathbb{K}}$ -morphism $\pi(\mathfrak{A})(*_t) : \mathbb{K} \rightarrow \pi(\mathfrak{A})(t)$ associated to the empty tuple $*_t \in \mathcal{P}_{\overline{\mathbf{C}}}(\underline{t})$ is the unit $\text{id}_{\mathbf{a}_t}$ of $\pi(\mathfrak{A})(t)$.

Using the axioms of 2AQFTs from Remark 3.4, it is easy to check that $\pi(\mathfrak{A})$ satisfies the axioms of $\mathbf{Alg}_{\mathbb{K}}$ -valued prefactorization algebras from Remark 2.7. Hence, $\pi(\mathfrak{A}) \in \mathbf{AQFT}(\overline{\mathbf{C}})$ is an AQFT by Theorem 2.9.

Let us consider now a 1-morphism $\zeta : \mathfrak{A} \rightarrow \mathfrak{B}$ in $\mathbf{2AQFT}(\overline{\mathbf{C}})$. For each $c \in \mathbf{C}$, the \mathbb{K} -linear functor $\zeta_c : \mathfrak{A}(c) \rightarrow \mathfrak{B}(c)$ restricts to endomorphism algebras as $\zeta_c : \text{End}(\mathbf{a}_c) \rightarrow \text{End}(\zeta_c(\mathbf{a}_c))$. The coherence map in (3.10) that is associated to $*_c$ provides an isomorphism $\zeta_{*c} : \mathbf{b}_c \rightarrow \zeta_c(\mathbf{a}_c)$ in the category $\mathfrak{B}(c)$, which we use to define the $\mathbf{Alg}_{\mathbb{K}}$ -morphism

$$\begin{aligned} \pi(\zeta)_c : \pi(\mathfrak{A})(c) &\longrightarrow \pi(\mathfrak{B})(c) \quad , \\ h &\longmapsto (\zeta_{*c})^{-1} \circ \zeta_c(h) \circ \zeta_{*c} \quad . \end{aligned} \quad (4.3)$$

Using the axioms of 1-morphisms of 2AQFTs from Remark 3.4, it is easy to check that $\pi(\zeta) : \pi(\mathfrak{A}) \rightarrow \pi(\mathfrak{B})$ is a morphism of $\mathbf{Alg}_{\mathbb{K}}$ -valued prefactorization algebras in the sense of Remark 2.7, and hence by Theorem 2.9 a morphism of AQFTs.

Let $\Gamma : \zeta \Rightarrow \kappa$ be a 2-morphism between two 1-morphisms $\zeta, \kappa : \mathfrak{A} \rightarrow \mathfrak{B}$ in $\mathbf{2AQFT}(\overline{\mathbf{C}})$. Using the axioms from Remark 3.4, we obtain a commutative diagram

$$\begin{array}{ccc} \mathbf{b}_c & \xrightarrow{=} & \mathbf{b}_c \\ \zeta_{*c} \downarrow \cong & & \cong \downarrow \kappa_{*c} \\ \zeta_c(\mathbf{a}_c) & \xrightarrow{\Gamma_c} & \kappa_c(\mathbf{a}_c) \end{array} \quad (4.4)$$

in the category $\mathfrak{B}(c)$, where isomorphisms are indicated by \cong . Hence Γ_c in this diagram is an isomorphism too. From (4.4) and (4.3), we compute

$$\begin{aligned} \pi(\kappa)_c(h) &= (\kappa_{*c})^{-1} \circ \kappa_c(h) \circ \kappa_{*c} = (\zeta_{*c})^{-1} \circ (\Gamma_c)^{-1} \circ \kappa_c(h) \circ \Gamma_c \circ \zeta_{*c} \\ &= (\zeta_{*c})^{-1} \circ \zeta_c(h) \circ \zeta_{*c} = \pi(\zeta)_c(h) \quad , \end{aligned} \quad (4.5)$$

where in the third step we used that (3.11) is a natural transformation. Hence, $\pi(\kappa) = \pi(\zeta) : \pi(\mathfrak{A}) \rightarrow \pi(\mathfrak{B})$ define the same morphism in $\mathbf{AQFT}(\overline{\mathbf{C}})$ and we can set $\pi(\Gamma) := \text{Id} : \pi(\zeta) \Rightarrow \pi(\kappa)$.

Proposition 4.1. *For every orthogonal category $\overline{\mathbf{C}}$, the construction above defines a 2-functor*

$$\pi : \mathbf{2AQFT}(\overline{\mathbf{C}}) \longrightarrow \mathbf{AQFT}(\overline{\mathbf{C}}) \quad , \quad (4.6)$$

which we call the truncation 2-functor.

4.2 Inclusion

Let $\mathfrak{A} \in \mathbf{AQFT}(\overline{\mathbf{C}})$ be an ordinary AQFT, regarded as an $\mathbf{Alg}_{\mathbb{K}}$ -valued prefactorization algebra via Theorem 2.9. We define its inclusion $\iota(\mathfrak{A}) \in \mathbf{2AQFT}(\overline{\mathbf{C}})$ by providing the data listed in Remark 3.4:

- (1) For each $c \in \mathbf{C}$, we set

$$\iota(\mathfrak{A})(c) := \mathbf{Mod}_{\mathfrak{A}(c)} \quad (4.7)$$

to be the \mathbb{K} -linear category of right $\mathfrak{A}(c)$ -modules. This is a locally presentable \mathbb{K} -linear category, see e.g. [BCJF15].

- (2) For each non-empty tuple $\underline{f} = (f_1, \dots, f_n) \in \mathcal{P}_{\overline{\mathbf{C}}}(\underline{c})$ of mutually orthogonal \mathbf{C} -morphisms, the given $\mathbf{Alg}_{\mathbb{K}}$ -morphism $\mathfrak{A}(\underline{f}) : \bigotimes_{i=1}^n \mathfrak{A}(c_i) \rightarrow \mathfrak{A}(t)$ induces a restriction functor $\mathfrak{A}(\underline{f})^* : \mathbf{Mod}_{\mathfrak{A}(t)} \rightarrow \mathbf{Mod}_{\bigotimes_{i=1}^n \mathfrak{A}(c_i)}$, which admits a left adjoint functor (called the *induced module functor*)

$$\mathfrak{A}(\underline{f})! = (-) \otimes_{\bigotimes_{i=1}^n \mathfrak{A}(c_i)} \mathfrak{A}(t) : \mathbf{Mod}_{\bigotimes_{i=1}^n \mathfrak{A}(c_i)} \longrightarrow \mathbf{Mod}_{\mathfrak{A}(t)} \quad . \quad (4.8)$$

The latter functor is clearly \mathbb{K} -linear and co-continuous. Observe further that the functor $\otimes^n : \prod_{i=1}^n \mathbf{Vec}_{\mathbb{K}} \rightarrow \mathbf{Vec}_{\mathbb{K}}$, $(V_1, \dots, V_n) \mapsto V_1 \otimes \dots \otimes V_n$ taking n -ary tensor products of vector spaces induces a functor

$$\otimes^n : \prod_{i=1}^n \mathbf{Mod}_{\mathfrak{A}(c_i)} \longrightarrow \mathbf{Mod}_{\otimes_{i=1}^n \mathfrak{A}(c_i)} \quad (4.9)$$

that is \mathbb{K} -linear and co-continuous in each variable. We define by composition

$$\iota(\mathfrak{A})(\underline{f}) : \prod_{i=1}^n \mathbf{Mod}_{\mathfrak{A}(c_i)} \xrightarrow{\otimes^n} \mathbf{Mod}_{\otimes_{i=1}^n \mathfrak{A}(c_i)} \xrightarrow{\mathfrak{A}(\underline{f})!} \mathbf{Mod}_{\mathfrak{A}(t)} \quad . \quad (4.10)$$

For the empty tuple $\underline{c} = \emptyset$, we set the pointing $\iota(\mathfrak{A})(*_t) := \mathfrak{A}(t) \in \mathbf{Mod}_{\mathfrak{A}(t)}$ to be the rank 1 free $\mathfrak{A}(t)$ -module.

(3) The coherence natural isomorphisms in (3.7) are given by pasting of

$$\begin{array}{ccc} \prod_{i=1}^n \prod_{j=1}^{k_i} \mathbf{Mod}_{\mathfrak{A}(b_{ij})} & \xrightarrow{\prod_i \otimes^{k_i}} & \prod_{i=1}^n \mathbf{Mod}_{\otimes_{j=1}^{k_i} \mathfrak{A}(b_{ij})} & \xrightarrow{\prod_i \mathfrak{A}(\underline{g}_i)!} & \prod_{i=1}^n \mathbf{Mod}_{\mathfrak{A}(a_i)} & (4.11) \\ & \searrow \scriptstyle (\star) & \downarrow \scriptstyle \otimes^n & \swarrow \scriptstyle (\star\star) & \downarrow \scriptstyle \otimes^n & \\ & \otimes^{\sum k_i} & \mathbf{Mod}_{\otimes_{i=1}^n \otimes_{j=1}^{k_i} \mathfrak{A}(b_{ij})} & \xrightarrow{(\otimes_i \mathfrak{A}(\underline{g}_i))!} & \mathbf{Mod}_{\otimes_{i=1}^n \mathfrak{A}(a_i)} & \\ & & & \swarrow \scriptstyle (\star\star\star) & \downarrow \scriptstyle \mathfrak{A}(\underline{f})! & \\ & & & \mathfrak{A}(\underline{f}\underline{g})! & \mathbf{Mod}_{\mathfrak{A}(t)} & \end{array}$$

The natural isomorphisms (\star) and $(\star\star)$ are canonically determined by the coherence isomorphisms for tensor products. (Recall that the induced module functor (4.8) is given by a relative tensor product.) The natural isomorphism $(\star\star\star)$ is canonically determined by uniqueness (up to a unique natural isomorphism) of left adjoint functors and the strict composition property $(\otimes_i \mathfrak{A}(\underline{g}_i))^* \mathfrak{A}(\underline{f})^* = (\mathfrak{A}(\underline{f}) \otimes_i \mathfrak{A}(\underline{g}_i))^* = \mathfrak{A}(\underline{f}\underline{g})^*$ of the right adjoints, see also (2.9a).

(4) The coherence natural isomorphisms in (3.8) are canonically determined by uniqueness of left adjoint functors and the strict identity property $\mathfrak{A}(\text{id}_t)^* = \text{id}_{\mathfrak{A}(t)}^* = \text{id}_{\mathbf{Mod}_{\mathfrak{A}(t)}}$ of the right adjoints, see also (2.9b).

(5) The coherence natural isomorphisms in (3.9) are given by pasting of

$$\begin{array}{ccc} \prod_{i=1}^n \mathbf{Mod}_{\mathfrak{A}(c_{\sigma(i)})} & \xrightarrow{\text{flip}_{\sigma}} & \prod_{i=1}^n \mathbf{Mod}_{\mathfrak{A}(c_i)} & (4.12) \\ & \searrow \scriptstyle (\star) & \downarrow \scriptstyle \otimes^n & \\ & \otimes^n & \mathbf{Mod}_{\otimes_{i=1}^n \mathfrak{A}(c_{\sigma(i)})} & \xrightarrow{(\tau_{\sigma})!} & \mathbf{Mod}_{\otimes_{i=1}^n \mathfrak{A}(c_i)} & \\ & & & \swarrow \scriptstyle (\star\star) & \downarrow \scriptstyle \mathfrak{A}(\underline{f})! & \\ & & & \mathfrak{A}(\underline{f}\sigma)! & \mathbf{Mod}_{\mathfrak{A}(t)} & \end{array}$$

The natural isomorphism (\star) is canonically determined by the coherence isomorphisms for tensor products and the natural isomorphism $(\star\star)$ is canonically determined by uniqueness of left adjoint functors and the strict permutation property $(\tau_{\sigma})^* \mathfrak{A}(\underline{f})^* = (\mathfrak{A}(\underline{f}) \tau_{\sigma})^* = \mathfrak{A}(\underline{f}\sigma)^*$ of the right adjoints, see also (2.9c).

Since the coherences in (3-5) are canonically given by coherence isomorphisms, one confirms that $\iota(\mathfrak{A}) \in \mathbf{2AQFT}(\overline{\mathbf{C}})$ satisfies the axioms of 2AQFTs from Remark 3.4.

Let us consider now a morphism $\zeta : \mathfrak{A} \rightarrow \mathfrak{B}$ in $\mathbf{AQFT}(\overline{\mathbf{C}})$. Then the following data defines a 1-morphism $\iota(\zeta) : \iota(\mathfrak{A}) \rightarrow \iota(\mathfrak{B})$ in $\mathbf{2AQFT}(\overline{\mathbf{C}})$, see also Remark 3.4:

(1) For each $c \in \mathbf{C}$, we set

$$\iota(\zeta)_c := (\zeta_c)! : \mathbf{Mod}_{\mathfrak{A}(c)} \longrightarrow \mathbf{Mod}_{\mathfrak{B}(c)} \quad (4.13)$$

to be the \mathbb{K} -linear and co-continuous induced module functor along the given $\mathbf{Alg}_{\mathbb{K}}$ -morphism $\zeta_c : \mathfrak{A}(c) \rightarrow \mathfrak{B}(c)$.

(2) The coherence natural isomorphisms in (3.10) are given by pasting of

$$\begin{array}{ccc} \prod_{i=1}^n \mathbf{Mod}_{\mathfrak{A}(c_i)} & \xrightarrow{\prod_i (\zeta_{c_i})!} & \prod_{i=1}^n \mathbf{Mod}_{\mathfrak{B}(c_i)} \\ \downarrow \otimes^n & \swarrow (\star) & \downarrow \otimes^n \\ \mathbf{Mod}_{\otimes_{i=1}^n \mathfrak{A}(c_i)} & \xrightarrow{(\otimes_i \zeta_{c_i})!} & \mathbf{Mod}_{\otimes_{i=1}^n \mathfrak{B}(c_i)} \\ \downarrow \mathfrak{A}(\underline{f})! & \swarrow (\star\star) & \downarrow \mathfrak{B}(\underline{f})! \\ \mathbf{Mod}_{\mathfrak{A}(t)} & \xrightarrow{(\zeta_t)!} & \mathbf{Mod}_{\mathfrak{B}(t)} \end{array} \quad (4.14)$$

where (\star) is canonically determined by the coherence isomorphisms for tensor products and $(\star\star)$ is canonically determined by uniqueness of left adjoint functors and the strict naturality property $(\otimes_i \zeta_{c_i})^* \mathfrak{B}(\underline{f})^* = (\mathfrak{B}(\underline{f}) \otimes_i \zeta_{c_i})^* = (\zeta_t \mathfrak{A}(\underline{f}))^* = \mathfrak{A}(\underline{f})^* (\zeta_t)^*$ of the right adjoints, see also (2.10).

Proposition 4.2. *For every orthogonal category $\overline{\mathbf{C}}$, the construction above defines a pseudo-functor*

$$\iota : \mathbf{AQFT}(\overline{\mathbf{C}}) \longrightarrow \mathbf{2AQFT}(\overline{\mathbf{C}}) \quad , \quad (4.15)$$

which we call the inclusion pseudo-functor.

4.3 Biadjunction

We now prove that the pseudo-functors in Propositions 4.1 and 4.2 determine a biadjunction, with the inclusion $\iota : \mathbf{AQFT}(\overline{\mathbf{C}}) \rightarrow \mathbf{2AQFT}(\overline{\mathbf{C}})$ as the left adjoint and the truncation $\pi : \mathbf{2AQFT}(\overline{\mathbf{C}}) \rightarrow \mathbf{AQFT}(\overline{\mathbf{C}})$ as the right adjoint.

We describe first the unit $\eta : \text{id} \Rightarrow \pi \iota$ of this biadjunction, which is easier than the counit $\epsilon : \iota \pi \Rightarrow \text{id}$ because $\mathbf{AQFT}(\overline{\mathbf{C}})$ is just a 1-category, hence η is a natural transformation between ordinary functors. Let $\mathfrak{A} \in \mathbf{AQFT}(\overline{\mathbf{C}})$ be an ordinary AQFT. From the explicit descriptions of π and ι in Sections 4.1 and 4.2, we observe that

$$(\pi \iota(\mathfrak{A}))(c) = \text{End}(\mathfrak{A}(c)) = \mathbf{Mod}_{\mathfrak{A}(c)}(\mathfrak{A}(c), \mathfrak{A}(c)) \quad (4.16)$$

is the endomorphism algebra of the rank 1 free module $\mathfrak{A}(c) \in \mathbf{Mod}_{\mathfrak{A}(c)}$, for every $c \in \mathbf{C}$. We define the \mathfrak{A} -component $\eta_{\mathfrak{A}} : \mathfrak{A} \rightarrow \pi \iota(\mathfrak{A})$ of the unit η as the $\mathbf{AQFT}(\overline{\mathbf{C}})$ -morphism determined by the component $\mathbf{Alg}_{\mathbb{K}}$ -morphisms

$$(\eta_{\mathfrak{A}})_c : \mathfrak{A}(c) \longrightarrow \text{End}(\mathfrak{A}(c)) \quad , \quad a \longmapsto \mu_{\mathfrak{A}(c)}(a \otimes -) \quad , \quad (4.17)$$

for all $c \in \mathbf{C}$, where explicitly $\mu_{\mathfrak{A}(c)}(a \otimes -) : \mathfrak{A}(c) \rightarrow \mathfrak{A}(c)$, $a' \mapsto a a'$ is the right $\mathfrak{A}(c)$ -module endomorphism given by left multiplication by $a \in \mathfrak{A}(c)$. Naturality of $\eta_{\mathfrak{A}}$ in $\mathfrak{A} \in \mathbf{AQFT}(\overline{\mathbf{C}})$

is obvious, hence we have constructed the desired natural transformation $\eta : \text{id} \Rightarrow \pi \iota$. We further observe that η is a natural isomorphism because each of its components (4.17) is an isomorphism, with inverse given by the $\mathbf{Alg}_{\mathbb{K}}$ -morphism $(\eta_{\mathfrak{A}})_c^{-1} : \text{End}(\mathfrak{A}(c)) \rightarrow \mathfrak{A}(c)$, $h \mapsto h(1_{\mathfrak{A}(c)})$ that evaluates an endomorphism h on the unit element $1_{\mathfrak{A}(c)} \in \mathfrak{A}(c)$.

Using the natural transformation $\eta : \text{id} \Rightarrow \pi \iota$, we can define, for every $\mathfrak{A} \in \mathbf{AQFT}(\overline{\mathbf{C}})$ and $\mathfrak{B} \in \mathbf{2AQFT}(\overline{\mathbf{C}})$, a functor between Hom-categories

$$\widetilde{(-)} : \mathbf{2AQFT}(\overline{\mathbf{C}})(\iota(\mathfrak{A}), \mathfrak{B}) \longrightarrow \mathbf{AQFT}(\overline{\mathbf{C}})(\mathfrak{A}, \pi(\mathfrak{B})) \quad , \quad (4.18)$$

where we note that the target is a set, which we regard as a category with only identity morphisms. To a 1-morphism $\zeta : \iota(\mathfrak{A}) \rightarrow \mathfrak{B}$ in $\mathbf{2AQFT}(\overline{\mathbf{C}})$, this functor assigns the $\mathbf{AQFT}(\overline{\mathbf{C}})$ -morphism

$$\widetilde{\zeta} : \mathfrak{A} \xrightarrow{\eta_{\mathfrak{A}}} \pi \iota(\mathfrak{A}) \xrightarrow{\pi(\zeta)} \pi(\mathfrak{B}) \quad . \quad (4.19)$$

Given any 2-morphism $\Gamma : \zeta \Rightarrow \kappa$ between 1-morphisms $\zeta, \kappa : \iota(\mathfrak{A}) \rightarrow \mathfrak{B}$ in $\mathbf{2AQFT}(\overline{\mathbf{C}})$, we have already seen in Section 4.1 that $\pi(\zeta) = \pi(\kappa)$, hence setting $\widetilde{\Gamma} = \text{Id} : \widetilde{\zeta} \Rightarrow \widetilde{\kappa} = \widetilde{\kappa}$ consistently defines the functor (4.18).

Theorem 4.3. *Let $\overline{\mathbf{C}}$ be any orthogonal category. Then the functor (4.18) is an equivalence of categories, for every $\mathfrak{A} \in \mathbf{AQFT}(\overline{\mathbf{C}})$ and $\mathfrak{B} \in \mathbf{2AQFT}(\overline{\mathbf{C}})$. As a consequence, we obtain a biadjunction*

$$\iota : \mathbf{AQFT}(\overline{\mathbf{C}}) \rightleftarrows \mathbf{2AQFT}(\overline{\mathbf{C}}) : \pi \quad , \quad (4.20)$$

whose left adjoint is the inclusion pseudo-functor from Proposition 4.2 and whose right adjoint is the truncation 2-functor from Proposition 4.1. Because the unit $\eta : \text{id} \Rightarrow \pi \iota$ is a natural isomorphism, this biadjunction exhibits the category $\mathbf{AQFT}(\overline{\mathbf{C}})$ of ordinary AQFTs as a coreflective full 2-subcategory of the 2-category $\mathbf{2AQFT}(\overline{\mathbf{C}})$.

Proof. Let us recall from [BCJF15] the following fact: For any associative and unital \mathbb{K} -algebra $A \in \mathbf{Alg}_{\mathbb{K}}$, denote by $\mathbf{BEnd}(A)$ the full \mathbb{K} -linear subcategory of $\mathbf{Mod}_A \in \mathbf{Pr}_{\mathbb{K}}$ on the object $A \in \mathbf{Mod}_A$. (Note that $\mathbf{BEnd}(A)$ is just the endomorphism algebra $\text{End}(A)$ regarded as a \mathbb{K} -linear category with only one object.) Then, for any locally presentable \mathbb{K} -linear category $\mathbf{D} \in \mathbf{Pr}_{\mathbb{K}}$, the restriction along the inclusion $\mathbf{BEnd}(A) \subseteq \mathbf{Mod}_A$ of \mathbb{K} -linear categories induces an equivalence (i.e. a fully faithful and essentially surjective functor)

$$\mathbf{Lin}_{\mathbb{K},c}(\mathbf{Mod}_A, \mathbf{D}) \xrightarrow{\cong} \mathbf{Lin}_{\mathbb{K}}(\mathbf{BEnd}(A), \mathbf{D}) \quad (4.21)$$

from the full subcategory of $\mathbf{Fun}(\mathbf{Mod}_A, \mathbf{D})$ that consists of \mathbb{K} -linear and co-continuous functors to the full subcategory of $\mathbf{Fun}(\mathbf{BEnd}(A), \mathbf{D})$ that consists of \mathbb{K} -linear functors. Using this result, we can check that the functor (4.18) is fully faithful and essentially surjective as claimed.

Faithful: Let $\Gamma, \Delta : \zeta \Rightarrow \kappa$ be 2-morphisms between the 1-morphisms $\zeta, \kappa : \iota(\mathfrak{A}) \rightarrow \mathfrak{B}$ in $\mathbf{2AQFT}(\overline{\mathbf{C}})$. (Note that $\widetilde{\Gamma} = \widetilde{\Delta}$ is automatic.) From (4.4) we deduce that, for every $c \in \mathbf{C}$, the morphisms $\Gamma_c = \Delta_c : \zeta_c(\mathfrak{A}(c)) \rightarrow \kappa_c(\mathfrak{A}(c))$ in $\mathfrak{B}(c)$ coincide. This means that the two natural transformations $\Gamma_c, \Delta_c : \zeta_c \Rightarrow \kappa_c$ between the co-continuous \mathbb{K} -linear functors $\zeta_c, \kappa_c : \mathbf{Mod}_{\mathfrak{A}(c)} \rightarrow \mathfrak{B}(c)$ have the same restriction along the inclusion $\mathbf{BEnd}(\mathfrak{A}(c)) \subseteq \mathbf{Mod}_{\mathfrak{A}(c)}$. Recalling that the restriction functor (4.21) is faithful, we conclude that $\Gamma_c = \Delta_c : \zeta_c \Rightarrow \kappa_c$ coincide as natural transformations, for all $c \in \mathbf{C}$, and hence that $\Gamma = \Delta : \zeta \Rightarrow \kappa$ coincide as 2-morphisms in $\mathbf{2AQFT}(\overline{\mathbf{C}})$. This shows that the functor (4.18) is faithful.

Full: Let $\zeta, \kappa : \iota(\mathfrak{A}) \rightarrow \mathfrak{B}$ be 1-morphisms in $\mathbf{2AQFT}(\overline{\mathbf{C}})$ such that $\widetilde{\zeta} = \widetilde{\kappa} : \mathfrak{A} \rightarrow \pi(\mathfrak{B})$ in $\mathbf{AQFT}(\overline{\mathbf{C}})$. (Recall that $\mathbf{AQFT}(\overline{\mathbf{C}})$ only has identity 2-morphisms.) For each $c \in \mathbf{C}$, consider the morphism $\kappa_{*c} \circ (\zeta_{*c})^{-1} : \zeta_c(\mathfrak{A}(c)) \rightarrow \kappa_c(\mathfrak{A}(c))$ in $\mathfrak{B}(c)$. Using $\widetilde{\zeta} = \widetilde{\kappa}$, one shows that this defines a natural transformation between the restrictions along the inclusion functor $\mathbf{BEnd}(\mathfrak{A}(c)) \subseteq$

$\mathbf{Mod}_{\mathfrak{A}(c)}$ of the co-continuous \mathbb{K} -linear functors $\zeta_c, \kappa_c : \mathbf{Mod}_{\mathfrak{A}(c)} \rightarrow \mathfrak{B}(c)$. Recalling that the restriction functor (4.21) is full, there exists a natural transformation $\Gamma_c : \zeta_c \Rightarrow \kappa_c$ whose $\mathfrak{A}(c)$ -component is $\kappa_{*c} \circ (\zeta_{*c})^{-1}$. We still have to prove that the collection Γ_c , for all $c \in \mathbf{C}$, defines a 2-morphism $\Gamma : \zeta \Rightarrow \kappa$ between the 1-morphisms $\zeta, \kappa : \iota(\mathfrak{A}) \rightarrow \mathfrak{B}$ in $\mathbf{2AQFT}(\overline{\mathbf{C}})$. This amounts to showing that the diagram

$$\begin{array}{ccc} \mathfrak{B}(\underline{f}) \prod_i \zeta_{c_i} & \xrightarrow{\text{Id} * \prod_i \Gamma_{c_i}} & \mathfrak{B}(\underline{f}) \prod_i \kappa_{c_i} \\ \zeta_{\underline{f}} \downarrow & & \downarrow \kappa_{\underline{f}} \\ \zeta_t \iota(\mathfrak{A})(\underline{f}) & \xrightarrow{\Gamma_t * \text{Id}} & \kappa_t \iota(\mathfrak{A})(\underline{f}) \end{array} \quad (4.22)$$

of natural transformations commutes, for all $\underline{f} \in \mathcal{P}_{\overline{\mathbf{C}}}(t)$. Since this diagram lives in the category $\mathbf{Pr}_{\mathbb{K}}(\mathfrak{B}_{\iota(\mathfrak{A})(\underline{c})}^{(t)})$, i.e. all functors are \mathbb{K} -linear and co-continuous in each variable, we deduce from the equivalences in (3.4) and (4.21) that the diagram (4.22) of natural transformations commutes if and only if the corresponding component on the object $\prod_{i=1}^n \mathfrak{A}(c_i) \in \prod_{i=1}^n \mathbf{Mod}_{\mathfrak{A}(c_i)}$ commutes. This can be checked directly by using that $\zeta, \kappa : \iota(\mathfrak{A}) \rightarrow \mathfrak{B}$ are 1-morphisms in $\mathbf{2AQFT}(\overline{\mathbf{C}})$. (Here the axioms of Definition A.4 enter explicitly.) This shows that the functor (4.18) is full.

Essentially surjective: Let $\zeta : \mathfrak{A} \rightarrow \pi(\mathfrak{B})$ be any $\mathbf{AQFT}(\overline{\mathbf{C}})$ -morphism. We denote its component $\mathbf{Alg}_{\mathbb{K}}$ -morphisms by $\zeta_c : \mathfrak{A}(c) \rightarrow \text{End}(\mathbf{b}_c)$, for all $c \in \mathbf{C}$. Recalling that $\mathfrak{A}(c) \in \mathbf{Alg}_{\mathbb{K}}$ is naturally isomorphic via η (cf. (4.17)) to the endomorphism algebra $\text{End}(\mathfrak{A}(c))$ of the object $\mathfrak{A}(c) \in \mathbf{Mod}_{\mathfrak{A}(c)}$, we define a functor $\widehat{\zeta}_c : \mathbf{BEnd}(\mathfrak{A}(c)) \rightarrow \mathfrak{B}(c)$ that sends the only object $\mathfrak{A}(c) \in \mathbf{BEnd}(\mathfrak{A}(c))$ to $\mathbf{b}_c \in \mathfrak{B}(c)$ and each $\mathbf{BEnd}(\mathfrak{A}(c))$ -morphism $h \in \text{End}(\mathfrak{A}(c))$ to the $\mathfrak{B}(c)$ -morphism $\widehat{\zeta}_c(h) := \zeta_c((\eta_{\mathfrak{A}})_c^{-1}(h))$. This functor is by construction \mathbb{K} -linear, i.e. $\widehat{\zeta}_c \in \mathbf{Lin}_{\mathbb{K}}(\mathbf{BEnd}(\mathfrak{A}(c)), \mathfrak{B}(c))$. Since the functor (4.21) is essentially surjective, there exists a \mathbb{K} -linear and co-continuous functor $\kappa_c \in \mathbf{Lin}_{\mathbb{K},c}(\mathbf{Mod}_{\mathfrak{A}(c)}, \mathfrak{B}(c))$ and a natural isomorphism κ_{*c} from the functor $\widehat{\zeta}_c$ to the restriction along the inclusion $\mathbf{BEnd}(\mathfrak{A}(c)) \subseteq \mathbf{Mod}_{\mathfrak{A}(c)}$ of the functor κ_c . Because $\mathfrak{A}(c) \in \mathbf{BEnd}(\mathfrak{A}(c))$ is the only object, the natural isomorphism κ_{*c} consists of a single $\mathfrak{B}(c)$ -isomorphism $\kappa_{*c} : \mathbf{b}_c \rightarrow \kappa_c(\mathfrak{A}(c))$, with naturality being encoded in the condition $\kappa_c(h) \circ \kappa_{*c} = \kappa_{*c} \circ \widehat{\zeta}_c(h)$, for all $h \in \text{End}(\mathfrak{A}(c))$. Note that we have just constructed part of the data defining a 1-morphism $\kappa : \iota(\mathfrak{A}) \rightarrow \mathfrak{B}$ in $\mathbf{2AQFT}(\overline{\mathbf{C}})$ (cf. Remark 3.4). To complete the data, we have to construct, for each $\underline{f} \in \mathcal{P}_{\overline{\mathbf{C}}}(t)$, a natural isomorphism $\kappa_{\underline{f}} : \mathfrak{B}(\underline{f}) \prod_i \kappa_{c_i} \Rightarrow \kappa_t \iota(\mathfrak{A})(\underline{f})$ between functors from $\prod_{i=1}^n \mathbf{Mod}_{\mathfrak{A}(c_i)}$ to $\mathfrak{B}(t)$ that are \mathbb{K} -linear and co-continuous in each variable. Using again the equivalences in (3.4) and (4.21), this problem is equivalent to constructing a $\mathfrak{B}(t)$ -isomorphism, denoted with a slight abuse of notation also by $\kappa_{\underline{f}} : \mathfrak{B}(\underline{f}) (\prod_i \kappa_{c_i} (\prod_i \mathfrak{A}(c_i))) \rightarrow \kappa_t (\iota(\mathfrak{A})(\underline{f}) (\prod_i \mathfrak{A}(c_i)))$, fulfilling the naturality condition $\kappa_t (\iota(\mathfrak{A})(\underline{f})(\underline{h})) \circ \kappa_{\underline{f}} = \kappa_{\underline{f}} \circ \mathfrak{B}(\underline{f}) (\prod_i \kappa_{c_i} (\underline{h}))$, for all $\underline{h} \in \prod_{i=1}^n \text{End}(\mathfrak{A}(c_i))$. We define the $\mathfrak{B}(t)$ -isomorphism $\kappa_{\underline{f}}$ according to

$$\begin{array}{ccc} \mathfrak{B}(\underline{f}) (\prod_i \kappa_{c_i} (\prod_i \mathfrak{A}(c_i))) & \overset{\kappa_{\underline{f}}}{\dashrightarrow} & \kappa_t (\iota(\mathfrak{A})(\underline{f}) (\prod_i \mathfrak{A}(c_i))) \\ \mathfrak{B}(\underline{f}) (\prod_i \kappa_{*c_i}) \uparrow \cong & & \cong \downarrow \kappa_t (\iota(\mathfrak{A})_{(\underline{f}, *_{\underline{c}})}^2) \\ \mathfrak{B}(\underline{f}) (\prod_i \mathbf{b}_{c_i}) & \xrightarrow{\mathfrak{B}_{(\underline{f}, *_{\underline{c}})}^2} \mathbf{b}_t \xrightarrow{\kappa_{*t}} & \kappa_t (\mathfrak{A}(t)) \end{array} \quad (4.23)$$

and observe that the required naturality condition for $\kappa_{\underline{f}}$ follows from naturality of κ_{*c} and of ζ . This provides us with the desired natural isomorphism $\kappa_{\underline{f}} : \mathfrak{B}(\underline{f}) \prod_i \kappa_{c_i} \Rightarrow \kappa_t \iota(\mathfrak{A})(\underline{f})$ and hence completes the data needed to define a 1-morphism $\kappa : \iota(\mathfrak{A}) \rightarrow \mathfrak{B}$ in $\mathbf{2AQFT}(\overline{\mathbf{C}})$. It remains to check that the relevant axioms hold (cf. Remark 3.4 and Definition A.4). Using once again the equivalences in (3.4) and (4.21), confirming these axioms can be reduced to checking that certain diagrams in $\mathfrak{B}(t)$ commute. This can be done directly by using that $\iota(\mathfrak{A})$ and \mathfrak{B} are objects

in $\mathbf{2AQFT}(\overline{\mathbf{C}})$. (Here the axioms of Definition A.2 enter explicitly.) Since by construction the $\mathbf{AQFT}(\overline{\mathbf{C}})$ -morphisms $\tilde{\kappa} = \zeta : \mathfrak{A} \rightarrow \pi(\mathfrak{B})$ coincide, this shows that the functor (4.18) is essentially surjective. \square

Remark 4.4. The counit $\epsilon : \iota\pi \Rightarrow \text{id}$ of the inclusion-truncation biadjunction is determined implicitly by Theorem 4.3, see e.g. [LN16, Definition 2.5 and Remark 2.6] for further details on biadjunctions. Its \mathfrak{B} -component $\epsilon_{\mathfrak{B}} : \iota\pi(\mathfrak{B}) \rightarrow \mathfrak{B}$ is a 1-morphism in $\mathbf{2AQFT}(\overline{\mathbf{C}})$ which maps under the equivalence of categories in (4.18) to the identity $\tilde{\epsilon}_{\mathfrak{B}} = \text{id}_{\pi(\mathfrak{B})} : \pi(\mathfrak{B}) \rightarrow \pi(\mathfrak{B})$ in $\mathbf{AQFT}(\overline{\mathbf{C}})$. (Note that the latter property determines the 1-morphism $\epsilon_{\mathfrak{B}}$ up to invertible 2-morphisms in $\mathbf{2AQFT}(\overline{\mathbf{C}})$.)

The counit allows us to detect whether an object $\mathfrak{B} \in \mathbf{2AQFT}(\overline{\mathbf{C}})$ lies in the essential image of the inclusion pseudo-functor $\iota : \mathbf{AQFT}(\overline{\mathbf{C}}) \rightarrow \mathbf{2AQFT}(\overline{\mathbf{C}})$. We say that $\mathfrak{B} \in \mathbf{2AQFT}(\overline{\mathbf{C}})$ is *truncated* if the corresponding component $\epsilon_{\mathfrak{B}} : \iota\pi(\mathfrak{B}) \rightarrow \mathfrak{B}$ of the counit is an equivalence in $\mathbf{2AQFT}(\overline{\mathbf{C}})$. This means that a truncated $\mathbf{2AQFT}$ \mathfrak{B} is fully determined by its truncation $\pi(\mathfrak{B}) \in \mathbf{AQFT}(\overline{\mathbf{C}})$ as it can be reconstructed (up to equivalence) by applying the inclusion pseudo-functor ι . Our goal in Section 5 is to construct examples of $\mathbf{2AQFT}$ s that are *not* truncated in the above sense. \triangle

5 Gauging construction and orbifold $\mathbf{2AQFT}$ s

We present a construction of $\mathbf{2AQFT}$ s from the data of an ordinary \mathbf{AQFT} $\mathfrak{A} \in \mathbf{AQFT}(\overline{\mathbf{C}})$ that is endowed with an action of a finite group G . This construction can be interpreted physically as a local gauging of \mathfrak{A} with respect to G and the resulting $\mathbf{2AQFT}$ as the corresponding categorified orbifold theory, see Proposition 5.4 and Remark 5.6 below. Let us start with introducing some relevant terminology.

Definition 5.1. Let $\overline{\mathbf{C}}$ be an orthogonal category and G a finite group. A G -equivariant \mathbf{AQFT} on $\overline{\mathbf{C}}$ is a pair (\mathfrak{A}, ρ) consisting of an object $\mathfrak{A} \in \mathbf{AQFT}(\overline{\mathbf{C}})$ and a representation $\rho : G \rightarrow \text{Aut}(\mathfrak{A})$ of G as automorphisms of \mathfrak{A} . A morphism $\zeta : (\mathfrak{A}, \rho) \rightarrow (\mathfrak{B}, \sigma)$ of G -equivariant \mathbf{AQFT} s is an $\mathbf{AQFT}(\overline{\mathbf{C}})$ -morphism $\zeta : \mathfrak{A} \rightarrow \mathfrak{B}$ that commutes with the G -actions, i.e. $\zeta \rho(g) = \sigma(g) \zeta$, for all $g \in G$. We denote by $G\text{-}\mathbf{AQFT}(\overline{\mathbf{C}})$ the category of G -equivariant \mathbf{AQFT} s on $\overline{\mathbf{C}}$.

Remark 5.2. Our choice of terminology in Definition 5.1 is motivated by the following equivalent perspective on G -equivariant \mathbf{AQFT} s. Let us denote by $\mathbf{Rep}_{\mathbb{K}}(G)$ the category of \mathbb{K} -linear representations of G . Recall that this is a (closed) symmetric monoidal category with monoidal product given by the tensor product $V \otimes W$ of representations, monoidal unit given by the trivial representation \mathbb{K} and symmetric braiding given by the flip map. Hence, we can introduce the category $G\text{-}\mathbf{Alg}_{\mathbb{K}} := \mathbf{Alg}_{\text{As}}(\mathbf{Rep}_{\mathbb{K}}(G))$ of associative and unital algebras in $\mathbf{Rep}_{\mathbb{K}}(G)$, which are also called G -equivariant associative and unital \mathbb{K} -algebras. (Note that for the trivial group $G = \{e\}$, this is just the category $\mathbf{Alg}_{\mathbb{K}}$ that we considered in Section 2.) It is then easy to check that a G -equivariant \mathbf{AQFT} (\mathfrak{A}, ρ) , as introduced in Definition 5.1, is the same data as a functor $\mathbf{C} \rightarrow G\text{-}\mathbf{Alg}_{\mathbb{K}}$ to the category of G -equivariant associative and unital \mathbb{K} -algebras that satisfies the \perp -commutativity property from Definition 2.4. From this perspective, morphisms of G -equivariant \mathbf{AQFT} s are just natural transformations of functors from \mathbf{C} to $G\text{-}\mathbf{Alg}_{\mathbb{K}}$. \triangle

Given any G -equivariant \mathbf{AQFT} $(\mathfrak{A}, \rho) \in G\text{-}\mathbf{AQFT}(\overline{\mathbf{C}})$, one can construct its associated *orbifold theory* $\mathfrak{A}_0^G \in \mathbf{AQFT}(\overline{\mathbf{C}})$ by taking the invariants of the action $\rho : G \rightarrow \text{Aut}(\mathfrak{A})$, see e.g. [Xu00, Mug05]. We have added a subscript 0 to emphasize that, as we shall show in Proposition 5.4, the traditional orbifold theory $\mathfrak{A}_0^G \cong \pi(\mathfrak{A}^G)$ is only the truncation of a richer (in general not truncated) *categorified orbifold theory* $\mathfrak{A}^G \in \mathbf{2AQFT}(\overline{\mathbf{C}})$. The latter will be described by a gauging construction that we develop in this section. We also refer to Remark 5.6 for a physical interpretation of our gauging construction and the resulting categorified orbifold theory.

As a preparation, let us briefly recall some standard facts and constructions from the theory of equivariant algebras and modules. As already mentioned in Remark 5.2, the representation category $\mathbf{Rep}_{\mathbb{K}}(G)$ of a finite group G is a (closed) symmetric monoidal category, hence we can introduce the category $G\text{-}\mathbf{Alg}_{\mathbb{K}}$ of G -equivariant associative and unital \mathbb{K} -algebras. Analogously to the non-equivariant case $\mathbf{Alg}_{\mathbb{K}}$ from Section 2, this category is symmetric monoidal with monoidal product the tensor product algebra $A \otimes B$ (endowed with the tensor product G -action), monoidal unit the algebra \mathbb{K} (endowed with the trivial G -action) and symmetric braiding given by the flip map. For every object $A \in G\text{-}\mathbf{Alg}_{\mathbb{K}}$, one can introduce the locally presentable \mathbb{K} -linear category $G\text{-}\mathbf{Mod}_A := \mathbf{Mod}_A(\mathbf{Rep}_{\mathbb{K}}(G))$ of G -equivariant right A -modules. An object in this category is an object $V \in \mathbf{Rep}_{\mathbb{K}}(G)$ together with a $\mathbf{Rep}_{\mathbb{K}}(G)$ -morphism $V \otimes A \rightarrow V$ that satisfies the usual axioms of a right A -action. Morphisms are $\mathbf{Rep}_{\mathbb{K}}(G)$ -morphism that preserve the right A -actions. Similarly to the non-equivariant case, given any morphism $\kappa : A \rightarrow B$ in $G\text{-}\mathbf{Alg}_{\mathbb{K}}$, one can define a \mathbb{K} -linear induced module functor $\kappa_! = (-) \otimes_A B : G\text{-}\mathbf{Mod}_A \rightarrow G\text{-}\mathbf{Mod}_B$. This functor has a right adjoint given by the restriction functor $\kappa^* : G\text{-}\mathbf{Mod}_B \rightarrow G\text{-}\mathbf{Mod}_A$. As a consequence, $\kappa_!$ is a co-continuous \mathbb{K} -linear functor between locally presentable \mathbb{K} -linear categories, i.e. a 1-morphism in the 2-category $\mathbf{Pr}_{\mathbb{K}}$.

Let now $(\mathfrak{A}, \rho) \in G\text{-}\mathbf{AQFT}(\overline{\mathbf{C}})$ be any G -equivariant AQFT on $\overline{\mathbf{C}}$. We define its gauging $\mathfrak{A}^G \in \mathbf{2AQFT}(\overline{\mathbf{C}})$ by a G -equivariant generalization of the inclusion pseudo-functor ι from Section 4.2. Concretely, \mathfrak{A}^G is described by the following data as listed in Remark 3.4:

- (1) For each $c \in \mathbf{C}$, we set

$$\mathfrak{A}^G(c) := G\text{-}\mathbf{Mod}_{\mathfrak{A}(c)} \quad (5.1)$$

to be the locally presentable \mathbb{K} -linear category of G -equivariant right modules over the G -equivariant associative and unital \mathbb{K} -algebra $\mathfrak{A}(c) \in G\text{-}\mathbf{Alg}_{\mathbb{K}}$.

- (2) For each non-empty tuple $\underline{f} = (f_1, \dots, f_n) \in \mathcal{P}_{\overline{\mathbf{C}}}(\underline{c})$ of mutually orthogonal \mathbf{C} -morphisms, we define the functor

$$\mathfrak{A}^G(\underline{f}) : \prod_{i=1}^n G\text{-}\mathbf{Mod}_{\mathfrak{A}(c_i)} \xrightarrow{\otimes^n} G\text{-}\mathbf{Mod}_{\otimes_{i=1}^n \mathfrak{A}(c_i)} \xrightarrow{\mathfrak{A}(\underline{f})_!} G\text{-}\mathbf{Mod}_{\mathfrak{A}(t)} \quad , \quad (5.2)$$

where $\otimes^n : (V_1, \dots, V_n) \mapsto V_1 \otimes \dots \otimes V_n$ is the functor assigning the n -ary tensor product of representations (equipped with the induced structure of a G -equivariant module over the tensor product of algebras) and $\mathfrak{A}(\underline{f})_!$ is the induced module functor for G -equivariant modules along the $G\text{-}\mathbf{Alg}_{\mathbb{K}}$ -morphism $\mathfrak{A}(\underline{f}) : \otimes_{i=1}^n \mathfrak{A}(c_i) \rightarrow \mathfrak{A}(t)$. Note that the functor (5.2) is \mathbb{K} -linear and co-continuous in each variable, i.e. it defines a 1-operation in $\mathbf{Pr}_{\mathbb{K}}$. For the empty tuple $\underline{c} = \emptyset$, we set the pointing to be $\mathfrak{A}^G(*_t) := \mathfrak{A}(t) \in G\text{-}\mathbf{Mod}_{\mathfrak{A}(t)}$.

- (3-5) The coherence isomorphisms for \mathfrak{A}^G are completely analogous to the ones for the inclusion $\iota(\mathfrak{A}) \in \mathbf{2AQFT}(\overline{\mathbf{C}})$ from Section 4.2.

Let us consider now a morphism $\zeta : (\mathfrak{A}, \rho) \rightarrow (\mathfrak{B}, \sigma)$ in $G\text{-}\mathbf{AQFT}(\overline{\mathbf{C}})$. Then the following data defines a 1-morphism $\zeta^G : \mathfrak{A}^G \rightarrow \mathfrak{B}^G$ in $\mathbf{2AQFT}(\overline{\mathbf{C}})$, see also Remark 3.4:

- (1) For each $c \in \mathbf{C}$, we set

$$(\zeta^G)_c := (\zeta_c)_! : G\text{-}\mathbf{Mod}_{\mathfrak{A}(c)} \longrightarrow G\text{-}\mathbf{Mod}_{\mathfrak{B}(c)} \quad (5.3)$$

to be the \mathbb{K} -linear and co-continuous induced module functor for G -equivariant modules along the $G\text{-}\mathbf{Alg}_{\mathbb{K}}$ -morphism $\zeta_c : \mathfrak{A}(c) \rightarrow \mathfrak{B}(c)$.

- (2) The coherence isomorphisms for $\zeta^G : \mathfrak{A}^G \rightarrow \mathfrak{B}^G$ are completely analogous to the ones for $\iota(\zeta) : \iota(\mathfrak{A}) \rightarrow \iota(\mathfrak{B})$ from Section 4.2.

Proposition 5.3. *For every orthogonal category $\overline{\mathbf{C}}$ and finite group G , the construction above defines a pseudo-functor*

$$(-)^G : G\text{-AQFT}(\overline{\mathbf{C}}) \longrightarrow \mathbf{2AQFT}(\overline{\mathbf{C}}) \quad , \quad (5.4)$$

which we call the gauging construction.

The following result relates our novel gauging construction to the traditional concept of orbifold theories from [Xu00, Mug05].

Proposition 5.4. *For every G -equivariant AQFT $(\mathfrak{A}, \rho) \in G\text{-AQFT}(\overline{\mathbf{C}})$, there exists a natural isomorphism $\mathfrak{A}_0^G \cong \pi(\mathfrak{A}^G)$ in $\mathbf{AQFT}(\overline{\mathbf{C}})$ between the traditional orbifold theory \mathfrak{A}_0^G (that assigns subalgebras of G -invariants) and the truncation (cf. Section 4.1) of the gauging construction $\mathfrak{A}^G \in \mathbf{2AQFT}(\overline{\mathbf{C}})$ from Proposition 5.3.*

Proof. From the description of the truncation 2-functor in Section 4.1, we obtain that $\pi(\mathfrak{A}^G)(c) = \text{End}(\mathfrak{A}(c))$ is the endomorphism algebra of the G -equivariant module $\mathfrak{A}(c) \in G\text{-Mod}_{\mathfrak{A}(c)}$, for each $c \in \mathbf{C}$. Since morphisms in $G\text{-Mod}_{\mathfrak{A}(c)}$ preserve by definition the G -action, it follows that $\text{End}(\mathfrak{A}(c))$ is isomorphic to the subalgebra of G -invariants in $\mathfrak{A}(c)$, hence $\pi(\mathfrak{A}^G)(c) \cong \mathfrak{A}_0^G(c)$ is isomorphic to the algebra that is assigned by the traditional orbifold theory \mathfrak{A}_0^G . Using further the explicit description of the factorization products of $\pi(\mathfrak{A}^G)$ from Section 4.1, one checks that this family of $\mathbf{Alg}_{\mathbb{K}}$ -isomorphisms defines an $\mathbf{AQFT}(\overline{\mathbf{C}})$ -isomorphism $\pi(\mathfrak{A}^G) \cong \mathfrak{A}_0^G$. Naturality of this isomorphism in $(\mathfrak{A}, \rho) \in G\text{-AQFT}(\overline{\mathbf{C}})$ is obvious. \square

The previous proposition provides a justification for the following terminology.

Definition 5.5. We call the gauging construction $\mathfrak{A}^G \in \mathbf{2AQFT}(\overline{\mathbf{C}})$ the *categorified orbifold theory* associated to the G -equivariant AQFT $(\mathfrak{A}, \rho) \in G\text{-AQFT}(\overline{\mathbf{C}})$.

Remark 5.6. In addition to our result in Proposition 5.4, there is further justification for calling the 2AQFT \mathfrak{A}^G a categorified orbifold theory. The presentation in this remark is intentionally rather informal, which is convenient to convey our main message.

Let us recall that the field configurations of a classical σ -model are given by maps $\phi : \Sigma \rightarrow X$ from a world-sheet Σ to a target space X . When a finite group G acts on the target space X , one can form the orbifold (i.e. quotient stack) $X//G$ and consider the corresponding *orbifold σ -model* whose field configurations are now maps $\phi : \Sigma \rightarrow X//G$ with values in a stack. As a consequence, the “space” of field configurations is a stack too, namely the mapping stack

$$\text{Fields}(\Sigma) := \text{Map}(\Sigma, X//G) \quad . \quad (5.5)$$

In order to study local aspects of this field theory, let us introduce the category $\mathbf{Disk}(\Sigma) \subseteq \mathbf{Open}(\Sigma)$ of open subsets $U \subseteq \Sigma$ that are diffeomorphic to a Cartesian space $U \cong \mathbb{R}^m$. (Global aspects of such field theories will be discussed later in Section 6.) Restricting to such subsets, the stack of fields is equivalent to the stacky quotient

$$\text{Fields}(U) \simeq \text{Map}(U, X)//G \quad (5.6)$$

of the *ordinary* mapping space $\text{Map}(U, X)$ by the finite group G . If we ignore for the moment the stacky quotient by G , we are in the familiar scenario where the space of fields $\text{Map}(U, X)$ is just an *ordinary* space and not a stack. Formal deformation quantization of such a field theory leads to an *ordinary* AQFT $\mathfrak{A} \in \mathbf{AQFT}(\overline{\mathbf{Disk}(\Sigma)})$, which in the case there are no anomalies will carry a G -action, i.e. $(\mathfrak{A}, \rho) \in G\text{-AQFT}(\overline{\mathbf{Disk}(\Sigma)})$. By construction, $\mathfrak{A}(U)$ is a deformation quantization of a suitable G -equivariant function algebra $\mathcal{O}(\text{Map}(U, X))$.

Things get more interesting when we consider the stacky quotient by G in (5.6). From the perspective of [Bra14, Lur04], which we recalled in Section 1, it is better to assign to the stack $\mathbf{Fields}(U)$ in (5.6) its category of quasi-coherent sheaves

$$\mathbf{QCoh}(\mathbf{Fields}(U)) \simeq \mathbf{QCoh}(\mathrm{Map}(U, X)//G) \simeq G\text{-}\mathbf{Mod}_{\mathcal{O}(\mathrm{Map}(U, X))} \quad , \quad (5.7)$$

which is the category of G -equivariant modules over the classical G -equivariant function algebra $\mathcal{O}(\mathrm{Map}(U, X))$. The G -equivariant quantization $\mathfrak{A}(U)$ of $\mathcal{O}(\mathrm{Map}(U, X))$ from the previous paragraph then determines a quantization of this category $\mathfrak{A}^G(U) = G\text{-}\mathbf{Mod}_{\mathfrak{A}(U)}$, which we recognize as our gauging construction. Hence, our gauging construction encodes the local aspects of orbifold σ -models. \triangle

We still have to address the important question whether or not it is possible to obtain genuine non-truncated $\mathfrak{A}^G \in \mathbf{2AQFT}(\overline{\mathbf{C}})$ from our gauging construction. This will of course depend on details of the group G and its action $\rho : G \rightarrow \mathrm{Aut}(\mathfrak{A})$ on $\mathfrak{A} \in \mathbf{AQFT}(\overline{\mathbf{C}})$. For example, if $G = \{e\}$ is the trivial group, then the gauging construction from Proposition 5.3 coincides with the inclusion pseudo-functor ι from Proposition 4.2, hence gauging the trivial group $G = \{e\}$ always leads to truncated 2AQFTs in the sense of Remark 4.4. On the other hand, it is very easy to give simple examples of gaugings that define non-truncated 2AQFTs.

Example 5.7. Let G be a finite group and consider the trivial AQFT $\mathfrak{A} = \mathbb{K} \in \mathbf{AQFT}(\overline{\mathbf{C}})$ that assigns $\mathfrak{A}(c) = \mathbb{K} \in \mathbf{Alg}_{\mathbb{K}}$ to every $c \in \mathbf{C}$. When endowed with the trivial G -action $\rho : G \rightarrow \mathrm{Aut}(\mathbb{K})$, $g \mapsto \mathrm{id}_{\mathbb{K}}$, this defines a G -equivariant AQFT $(\mathbb{K}, \rho) \in G\text{-}\mathbf{AQFT}(\overline{\mathbf{C}})$. Note that the corresponding traditional orbifold theory $\mathbb{K}_0^G = \mathbb{K}$ is of course the trivial AQFT too. In contrast, the categorified orbifold theory $\mathbb{K}^G \in \mathbf{2AQFT}(\overline{\mathbf{C}})$ that is obtained from our gauging construction is much more interesting. It assigns to every $c \in \mathbf{C}$ the representation category $\mathbb{K}^G(c) = G\text{-}\mathbf{Mod}_{\mathbb{K}} = \mathbf{Rep}_{\mathbb{K}}(G)$ of the group G and its factorization products $\mathbb{K}^G(f) = \otimes^n : \prod_{i=1}^n \mathbf{Rep}_{\mathbb{K}}(G) \rightarrow \mathbf{Rep}_{\mathbb{K}}(G)$ are given by the n -ary tensor products of representations. The pointing $\mathbb{K}^G(*_t) = \mathbb{K} \in \mathbf{Rep}_{\mathbb{K}}(G)$ is given by the trivial representation. By Proposition 5.4, the truncation $\pi(\mathbb{K}^G) \cong \mathbb{K}_0^G = \mathbb{K}$ of this 2AQFT is the trivial theory

Our claim is that the categorified orbifold theory $\mathbb{K}^G \in \mathbf{2AQFT}(\overline{\mathbf{C}})$ is not truncated, whenever $G \neq \{e\}$ is non-trivial. To prove this claim, we consider as explained in Remark 4.4 the corresponding component $\epsilon_{\mathbb{K}^G} : \iota \pi(\mathbb{K}^G) \rightarrow \mathbb{K}^G$ of the counit of the inclusion-truncation biadjunction. This is a 1-morphism in $\mathbf{2AQFT}(\overline{\mathbf{C}})$ whose components $(\epsilon_{\mathbb{K}^G})_c : \iota \pi(\mathbb{K}^G)(c) \simeq \mathbf{Vec}_{\mathbb{K}} \rightarrow \mathbb{K}^G(c) = \mathbf{Rep}_{\mathbb{K}}(G)$ are given by co-continuous \mathbb{K} -linear functors from the category of vector spaces to the representation category of G . Because 1-morphisms in $\mathbf{2AQFT}(\overline{\mathbf{C}})$ preserve the pointings (up to coherence isomorphisms), we know that the 1-dimensional vector space $\mathbb{K} \in \mathbf{Vec}_{\mathbb{K}}$ is mapped to a *trivial* representation $(\epsilon_{\mathbb{K}^G})_c(\mathbb{K}) \cong \mathbb{K} \in \mathbf{Rep}_{\mathbb{K}}(G)$. Using further that every vector space $V \cong \bigoplus_{b \in B} \mathbb{K} \in \mathbf{Vec}_{\mathbb{K}}$ is isomorphic to a coproduct of the 1-dimensional vector space \mathbb{K} (by choosing a basis B) and co-continuity of the functor $\epsilon_{\mathbb{K}^G}$, we observe that the essential image of $(\epsilon_{\mathbb{K}^G})_c : \mathbf{Vec}_{\mathbb{K}} \rightarrow \mathbf{Rep}_{\mathbb{K}}(G)$ lies in the full subcategory of trivial G -representations, hence it cannot be an equivalence of categories as every finite group $G \neq \{e\}$ admits non-trivial \mathbb{K} -linear representations. As a consequence, the component $\epsilon_{\mathbb{K}^G} : \iota \pi(\mathbb{K}^G) \rightarrow \mathbb{K}^G$ of the counit is not an equivalence in $\mathbf{2AQFT}(\overline{\mathbf{C}})$ and hence the categorified orbifold theory $\mathbb{K}^G \in \mathbf{2AQFT}(\overline{\mathbf{C}})$ is not truncated. ∇

Quite remarkably, it is possible to characterize precisely those G -equivariant AQFTs $(\mathfrak{A}, \rho) \in G\text{-}\mathbf{AQFT}(\overline{\mathbf{C}})$ whose associated gauging construction $\mathfrak{A}^G \in \mathbf{2AQFT}(\overline{\mathbf{C}})$ is truncated. Our arguments below make use of some standard terminology and results from Hopf-Galois theory, see e.g. [DT89] and also the review article [Mon09]. Let H be a Hopf algebra over \mathbb{K} . (In our applications below, $H = \mathcal{O}(G) = \mathrm{Map}(G, \mathbb{K})$ is the function Hopf algebra of a finite group G .) A *right H -comodule algebra* is an algebra $A \in \mathbf{Alg}_{\mathbb{K}}$ endowed with a right H -coaction $\delta : A \rightarrow A \otimes H$ that is an $\mathbf{Alg}_{\mathbb{K}}$ -morphism. We denote by $B := A^{\mathrm{co}H} := \{a \in A : \delta(a) = a \otimes 1_H\} \subseteq A$ the subalgebra of H -coaction invariants.

Definition 5.8. The algebra extension $B = A^{\text{co}H} \subseteq A$ is called *H-Hopf-Galois* if the canonical map

$$\beta : A \otimes_B A \longrightarrow A \otimes H, \quad a \otimes_B a' \longmapsto (a \otimes 1_H) \delta(a') \quad (5.8)$$

is bijective.

Associated to any right H -comodule algebra A are two \mathbb{K} -linear categories of interest: First, we have the category \mathbf{Mod}_A^H of *right (H, A)-Hopf modules*. An object in this category is a right A -module $V \in \mathbf{Mod}_A$ that is endowed with a compatible right H -comodule structure $\delta^V : V \rightarrow V \otimes H$, i.e. $\delta^V(va) = \delta^V(v) \delta(a)$, for all $v \in V$ and $a \in A$. The morphisms in \mathbf{Mod}_A^H are \mathbb{K} -linear maps that preserve both the A -actions and the H -coactions. Second, we have the \mathbb{K} -linear category \mathbf{Mod}_B of right modules over the subalgebra $B = A^{\text{co}H} \subseteq A$ of H -coaction invariants. These two categories are related by a \mathbb{K} -linear adjunction

$$\Phi : \mathbf{Mod}_B \rightleftarrows \mathbf{Mod}_A^H : \Psi, \quad (5.9)$$

whose left adjoint $\Phi = (-) \otimes_B A$ is the induced module functor, where $W \otimes_B A$ is endowed with the right H -coaction $\text{id} \otimes_B \delta$, for all $W \in \mathbf{Mod}_B$, and whose right adjoint $\Psi = (-)^{\text{co}H}$ is the functor taking H -coaction invariants $V^{\text{co}H} := \{v \in V : \delta^V(v) = v \otimes 1_H\}$, for all $V \in \mathbf{Mod}_A^H$. We shall need the following result (cf. [Mon09, Theorem 5.6]), which is originally due to Doi and Takeuchi [DT89].

Theorem 5.9. *Let H be finite dimensional. Then $B = A^{\text{co}H} \subseteq A$ is H -Hopf-Galois if and only if the counit $\epsilon : \Phi \Psi \Rightarrow \text{id}$ of the adjunction (5.9) is a natural isomorphism.*

In the case of interest to us, the Hopf algebra $H = \mathcal{O}(G) = \text{Map}(G, \mathbb{K})$ is given by the function Hopf algebra of a finite group G . In particular, H is finite dimensional. One easily observes that a right $\mathcal{O}(G)$ -coaction $\delta^V : V \rightarrow V \otimes \mathcal{O}(G)$ is the same datum as a group action $\rho : G \rightarrow \text{Aut}(V)$ by the relationship $\rho(g)(v) = v_{(0)} \langle v_{(1)}, g \rangle$, where we used Sweedler notation $\delta^V(v) = v_{(0)} \otimes v_{(1)}$ and the duality pairing $\langle \cdot, \cdot \rangle : \mathcal{O}(G) \otimes \mathbb{K}[G] \rightarrow \mathbb{K}$. In particular, right $\mathcal{O}(G)$ -comodule algebras are precisely G -equivariant algebras $A \in G\text{-Alg}_{\mathbb{K}}$ and the \mathbb{K} -linear category $\mathbf{Mod}_A^{\mathcal{O}(G)}$ of right $(\mathcal{O}(G), A)$ -Hopf modules is the locally presentable \mathbb{K} -linear category $G\text{-Mod}_A$ of G -equivariant right A -modules. Moreover, the subalgebra $B = A^{\text{co}H} = A_0^G \subseteq A$ of $\mathcal{O}(G)$ -coaction invariants is precisely the subalgebra of G -invariants.

Corollary 5.10. *Let G be a finite group and $H = \mathcal{O}(G)$ the function Hopf algebra of G . In this case, the adjunction in (5.9) reads as*

$$\Phi : \mathbf{Mod}_B \rightleftarrows G\text{-Mod}_A : \Psi. \quad (5.10)$$

This is an (adjoint) equivalence in the 2-category $\mathbf{Pr}_{\mathbb{K}}$ of locally presentable \mathbb{K} -linear categories if and only if $B = A^{\text{co}H} = A_0^G \subseteq A$ is $\mathcal{O}(G)$ -Hopf-Galois.

Proof. The left adjoint functor $\Phi = (-) \otimes_B A$ is clearly \mathbb{K} -linear and co-continuous, i.e. a 1-morphism in $\mathbf{Pr}_{\mathbb{K}}$. The right adjoint functor $\Psi = (-)^{\text{co}H} = (-)_0^G$ assigns the G -invariants (given by a categorical limit), which for actions of finite groups G and $\text{char}(\mathbb{K}) = 0$ coincides with the G -coinvariants (i.e. a categorical colimit). Hence, the right adjoint Ψ is a \mathbb{K} -linear and co-continuous functor too and the adjunction (5.10) is in the 2-category $\mathbf{Pr}_{\mathbb{K}}$.

The unit $\eta : \text{id} \Rightarrow \Psi \Phi$ of the adjunction (5.10) is given by the components $\eta_W : W \rightarrow (W \otimes_B A)_0^G$, $w \mapsto w \otimes_B 1_A$, for all $W \in \mathbf{Mod}_B$. Using again that forming G -invariants coincides with forming G -coinvariants, we find that $\eta : \text{id} \Rightarrow \Psi \Phi$ is a natural isomorphism. Our claim then follows from Theorem 5.9. \square

Theorem 5.11. *Let G be a finite group and $(\mathfrak{A}, \rho) \in G\text{-AQFT}(\overline{\mathbf{C}})$ a G -equivariant AQFT. Then the categorified orbifold theory $\mathfrak{A}^G \in \mathbf{2AQFT}(\overline{\mathbf{C}})$ is truncated if and only if the algebra extension $\mathfrak{A}_0^G(c) \subseteq \mathfrak{A}(c)$ is $\mathcal{O}(G)$ -Hopf-Galois, for all $c \in \mathbf{C}$.*

Proof. Recalling Remark 4.4, the $\mathbf{2AQFT} \mathfrak{A}^G \in \mathbf{2AQFT}(\overline{\mathbf{C}})$ is by definition truncated if the corresponding component $\epsilon_{\mathfrak{A}^G} : \iota \pi(\mathfrak{A}^G) \rightarrow \mathfrak{A}^G$ of the counit of the inclusion-truncation biadjunction from Theorem 4.3 is an equivalence in $\mathbf{2AQFT}(\overline{\mathbf{C}})$. The component $\epsilon_{\mathfrak{A}^G}$ of the counit is determined uniquely (up to invertible 2-morphisms in $\mathbf{2AQFT}(\overline{\mathbf{C}})$) by the condition $\widetilde{\epsilon_{\mathfrak{A}^G}} = \text{id}_{\pi(\mathfrak{A}^G)} : \pi(\mathfrak{A}^G) \rightarrow \pi(\mathfrak{A}^G)$ on its adjunct under (4.18). Using the explicit description of the inclusion and truncation pseudo-functors from Section 4 and the one of the gauging construction from the present section, one observes that the induced module functors $\Phi_c = (-) \otimes_{\mathfrak{A}_0^G(c)} \mathfrak{A}(c) : \iota \pi(\mathfrak{A}^G)(c) \cong \mathbf{Mod}_{\mathfrak{A}_0^G(c)} \rightarrow \mathfrak{A}^G(c) = G\text{-Mod}_{\mathfrak{A}(c)}$ (together with the obvious coherence isomorphisms) define a 1-morphism $\Phi : \iota \pi(\mathfrak{A}^G) \rightarrow \mathfrak{A}^G$ in $\mathbf{2AQFT}(\overline{\mathbf{C}})$ that satisfies $\widetilde{\Phi} = \text{id}_{\pi(\mathfrak{A}^G)} : \pi(\mathfrak{A}^G) \rightarrow \pi(\mathfrak{A}^G)$. Hence, $\Phi \cong \epsilon_{\mathfrak{A}^G}$ and we can equivalently investigate if Φ is an equivalence in $\mathbf{2AQFT}(\overline{\mathbf{C}})$.

By a straightforward but slightly lengthy calculation, one proves that a 1-morphism in $\mathbf{2AQFT}(\overline{\mathbf{C}})$ is an equivalence if and only if all its components are equivalences in the 2-category $\mathbf{Pr}_{\mathbb{K}}$. (In this proof one uses that every equivalence in any 2-category (here $\mathbf{Pr}_{\mathbb{K}}$) can be upgraded to an adjoint equivalence in order to define the quasi-inverse 1-morphism in $\mathbf{2AQFT}(\overline{\mathbf{C}})$.) Thus, to prove that $\mathfrak{A}^G \in \mathbf{2AQFT}(\overline{\mathbf{C}})$ is truncated we can equivalently study the components $\Phi_c = (-) \otimes_{\mathfrak{A}_0^G(c)} \mathfrak{A}(c) : \mathbf{Mod}_{\mathfrak{A}_0^G(c)} \rightarrow G\text{-Mod}_{\mathfrak{A}(c)}$, for all $c \in \mathbf{C}$. By Corollary 5.10, these components are equivalences in $\mathbf{Pr}_{\mathbb{K}}$ if and only if the algebra extension $\mathfrak{A}_0^G(c) \subseteq \mathfrak{A}(c)$ is $\mathcal{O}(G)$ -Hopf-Galois, for all $c \in \mathbf{C}$. This completes the proof. \square

Remark 5.12. We would like to emphasize that our result in Theorem 5.11 matches perfectly our physical interpretation of the gauging construction $\mathfrak{A}^G \in \mathbf{2AQFT}(\overline{\mathbf{C}})$ in terms of orbifold σ -models from Remark 5.6. The $H = \mathcal{O}(G)$ -Hopf-Galois condition from Definition 5.8 should be interpreted as a non-commutative algebraic generalization of a free G -action on a space, see e.g. [Mon09, Examples 2.11 and 2.12]. Because the quotient stack $X//G \simeq X/G$ corresponding to a free G -action is equivalent to the ordinary quotient space, the resulting ‘‘orbifold’’ σ -model in this case is just an ordinary σ -model with target space X/G . In particular, for free G -actions one does not expect higher categorical features in the corresponding ‘‘orbifold’’ σ -model. This is precisely what we have proven in Theorem 5.11 for orbifold quantum field theories. \triangle

We conclude this section by presenting more examples of non-truncated and also truncated categorified orbifold theories $\mathfrak{A}^G \in \mathbf{2AQFT}(\overline{\mathbf{C}})$.

Example 5.13. Let us denote by $\mathbf{Disk}(\mathbb{S}^1) \subset \mathbf{Open}(\mathbb{S}^1)$ the full subcategory of all non-empty open intervals $I \subset \mathbb{S}^1$ in the circle \mathbb{S}^1 . Restricting the orthogonality relation $\perp_{\mathbb{S}^1}$ from Example 2.2, we obtain a full orthogonal subcategory $\overline{\mathbf{Disk}(\mathbb{S}^1)} \subset \overline{\mathbf{Open}(\mathbb{S}^1)}$. Objects $\mathfrak{A} \in \mathbf{AQFT}(\overline{\mathbf{Disk}(\mathbb{S}^1)})$ are interpreted as chiral conformal AQFTs [Kaw15]. In this example we set $\mathbb{K} = \mathbb{C}$ to be the field of complex numbers. Let us consider the following specific theory, which is called the *chiral free boson*. To each interval $I \subset \mathbb{S}^1$, we assign the canonical commutation relations (CCR) algebra

$$\mathfrak{A}(I) := T_{\mathbb{C}}^{\otimes} C_c^{\infty}(I) \left/ \left\langle \varphi_1 \otimes \varphi_2 - \varphi_2 \otimes \varphi_1 - i\hbar \int_I \varphi_1 d\varphi_2 \right. \right\rangle \in \mathbf{Alg}_{\mathbb{C}} \quad , \quad (5.11)$$

where $C_c^{\infty}(I)$ denotes the vector space of compactly supported real-valued functions on $I \subset \mathbb{S}^1$ and $T_{\mathbb{C}}^{\otimes} C_c^{\infty}(I) := \bigoplus_{n=0}^{\infty} (C_c^{\infty}(I) \otimes_{\mathbb{R}} \mathbb{C})^{\otimes n} \in \mathbf{Alg}_{\mathbb{C}}$ is the complexified free algebra. (For later use, we have made the deformation parameter $\hbar \in \mathbb{R}$ explicit in the commutation relations for $\varphi_1, \varphi_2 \in C_c^{\infty}(I)$.) To each interval inclusion $\iota_I^J : I \rightarrow J$, we assign the $\mathbf{Alg}_{\mathbb{K}}$ -morphism $\mathfrak{A}(\iota_I^J) : \mathfrak{A}(I) \rightarrow \mathfrak{A}(J)$ that is defined on the generators by pushforward (i.e. extension by zero) of compactly supported functions. This defines an AQFT $\mathfrak{A} \in \mathbf{AQFT}(\overline{\mathbf{Disk}(\mathbb{S}^1)})$ in the sense of Definition 2.4.

Let us consider the representation $\rho : G = \mathbb{Z}_2 \rightarrow \text{Aut}(\mathfrak{A})$ of the cyclic group of order 2 that is defined on the generators of $\mathfrak{A}(I)$ by multiplication with ± 1 , i.e. $\rho(\pm 1)(\varphi) = \pm \varphi$, for all $\varphi \in \mathfrak{A}(I)$. This defines a \mathbb{Z}_2 -equivariant AQFT (\mathfrak{A}, ρ) and we can form the corresponding categorified orbifold theory $\mathfrak{A}^{\mathbb{Z}_2} \in \mathbf{2AQFT}(\overline{\text{Disk}}(\mathbb{S}^1))$ from Definition 5.5. To find out whether this theory is truncated or not, we use our results from Theorem 5.11. Let us consider an arbitrary interval $I \subset \mathbb{S}^1$ and set $A := \mathfrak{A}(I)$. Observe that the subalgebra $B := A_0^{\mathbb{Z}_2} \subset A$ of \mathbb{Z}_2 -invariants is the even part of the algebra (5.11). Regarding $A = \mathfrak{A}(I)$ as a B -bimodule, we obtain a direct sum decomposition $A = B \oplus V$, where V is the odd part of (5.11). Hence, the source of the canonical map (5.8) is isomorphic to $A \otimes_B A \cong B \oplus (V \otimes_B V) \oplus V \oplus V$. Using further that $A \otimes \mathcal{O}(G) \cong \prod_{g \in G} A$, the canonical map (5.8) explicitly reads as

$$\begin{aligned} \beta : B \oplus (V \otimes_B V) \oplus V \oplus V &\longrightarrow \prod_{g \in \mathbb{Z}_2} A \quad , \\ b + v \otimes_B v' + v_1 + v_2 &\longmapsto \begin{pmatrix} b + v v' + v_1 + v_2 \\ b - v v' + v_1 - v_2 \end{pmatrix} \quad . \end{aligned} \quad (5.12)$$

Note that the canonical map β is bijective if and only if the map $\mu : V \otimes_B V \rightarrow B$, $v \otimes_B v' \mapsto v v'$ that is induced by the multiplication $\mu : A \otimes A \rightarrow A$ on (5.11) is bijective.

Let us consider first the case where the deformation parameter $\hbar = 0$ is zero, which describes a classical (i.e. not quantized) field theory. In this case (5.11) is a complexified symmetric algebra over $C_c^\infty(I)$ and the map $\mu : V \otimes_B V \rightarrow B$ is *not* surjective because its image is at least quadratic in the generators. This implies that the canonical map β in (5.12) is not bijective, hence by Theorem 5.11 the categorified orbifold theory $\mathfrak{A}^{\mathbb{Z}_2} \in \mathbf{2AQFT}(\overline{\text{Disk}}(\mathbb{S}^1))$ for $\hbar = 0$ is non-truncated.

The situation changes drastically in the quantum case $\hbar \neq 0$. From the canonical commutation relations in (5.11), we deduce that one can always find two generators $\varphi_1, \varphi_2 \in C_c^\infty(I) \subseteq V \subseteq A$ that satisfy $[\varphi_1, \varphi_2] = i\hbar 1$. Dividing by \hbar , which is possible because we assumed that $\hbar \neq 0$, we can now prove that the map $\mu : V \otimes_B V \rightarrow B$ is bijective. For surjectivity, consider an arbitrary $b \in B$ and observe that

$$\mu\left(b \frac{1}{i\hbar}(\varphi_1 \otimes_B \varphi_2 - \varphi_2 \otimes_B \varphi_1)\right) = b \frac{1}{i\hbar}[\varphi_1, \varphi_2] = b \quad . \quad (5.13)$$

For injectivity, consider $\sum_j v_j \otimes_B v'_j \in V \otimes_B V$ such that $\sum_j v_j v'_j = 0$ and observe that

$$\begin{aligned} \sum_j v_j \otimes_B v'_j &= \frac{1}{i\hbar} \sum_j [\varphi_1, \varphi_2] v_j \otimes_B v'_j \\ &= \frac{1}{i\hbar} \varphi_1 \otimes_B \varphi_2 \sum_j v_j v'_j - \frac{1}{i\hbar} \varphi_2 \otimes_B \varphi_1 \sum_j v_j v'_j = 0 \quad , \end{aligned} \quad (5.14)$$

where we also used that $\varphi_1 v_j \in B$ and $\varphi_2 v_j \in B$. Theorem 5.11 then implies that the categorified orbifold theory $\mathfrak{A}^{\mathbb{Z}_2} \in \mathbf{2AQFT}(\overline{\text{Disk}}(\mathbb{S}^1))$ for $\hbar \neq 0$ is truncated.

Summing up, we have seen an example of a *non-truncated* classical orbifold field theory that is quantized to a *truncated* orbifold quantum field theory. We would like to emphasize that this result crucially relies on inverting the deformation parameter $0 \neq \hbar \in \mathbb{R}$ and hence it does not arise in formal deformation quantization. (In fact, treating \hbar in (5.11) as a formal parameter, the categorified orbifold theory $\mathfrak{A}^{\mathbb{Z}_2} \in \mathbf{2AQFT}(\overline{\text{Disk}}(\mathbb{S}^1))$ is non-truncated as in the classical case $\hbar = 0$.) A similar interplay between quantization and orbifold singularities was observed before within a different framework [Brz14, BS17]. \square

Remark 5.14. We would like to emphasize that the results of Example 5.13 hold true in much greater generality. Let $\overline{\mathbf{C}}$ be any orthogonal category and $\mathfrak{A} \in \mathbf{AQFT}(\overline{\mathbf{C}})$ any AQFT that assigns, to every $c \in \mathbf{C}$, a CCR-algebra $\mathfrak{A}(c) = \mathcal{CCR}(\mathcal{L}(c), \sigma_c)$ of a symplectic vector space $(\mathcal{L}(c), \sigma_c)$, i.e. σ_c is non-degenerate. Using similar arguments as in Example 5.13, one shows that the categorified

orbifold theory corresponding to the \mathbb{Z}_2 -action $\rho(\pm 1) : (\mathcal{L}(c), \sigma_c) \rightarrow (\mathcal{L}(c), \sigma_c)$, $\varphi \mapsto \pm\varphi$ is truncated, provided that $\mathbb{R} \ni \hbar \neq 0$. The same holds true for AQFTs assigning canonical anticommutation relation (CAR) algebras of non-degenerate inner product spaces. \triangle

6 Fredenhagen's universal category

The goal of this section is to present a categorified version of Fredenhagen's universal algebra. Let us briefly recall the original 1-categorical construction for ordinary AQFTs from [Fre90, Fre93, FRS92], see also [Lan14, BSW17] for more details. Given a full orthogonal subcategory embedding $J : \overline{\mathbf{C}} \rightarrow \overline{\mathbf{D}}$ and any ordinary AQFT $\mathfrak{A} \in \mathbf{AQFT}(\overline{\mathbf{C}})$ on $\overline{\mathbf{C}}$, operadic left Kan extension along the induced operad morphism $J : \mathcal{P}_{\overline{\mathbf{C}}} \rightarrow \mathcal{P}_{\overline{\mathbf{D}}}$ determines a canonical extension $J_! \mathfrak{A} \in \mathbf{AQFT}(\overline{\mathbf{D}})$ of \mathfrak{A} to the larger orthogonal category $\overline{\mathbf{D}}$. The algebra $J_! \mathfrak{A}(d) \in \mathbf{Alg}_{\mathbb{K}}$ that is assigned by the extended AQFT $J_! \mathfrak{A}$ to an object $d \in \mathbf{D}$ is usually referred to as Fredenhagen's universal algebra. This extension prescription is canonical in the sense that it is part of an adjunction $J_! : \mathbf{AQFT}(\overline{\mathbf{C}}) \rightleftarrows \mathbf{AQFT}(\overline{\mathbf{D}}) : J^*$, where the right adjoint functor J^* is given by restriction of AQFTs along J . The following two examples of full orthogonal subcategory embeddings $J : \overline{\mathbf{C}} \rightarrow \overline{\mathbf{D}}$ are typically considered in applications to physics.

Example 6.1. Recall from Example 2.2 the orthogonal category $\overline{\mathbf{Open}(M)} = (\mathbf{Open}(M), \perp_M)$ of non-empty open subsets of a manifold M . Consider the full subcategory $\mathbf{Disk}(M) \subseteq \mathbf{Open}(M)$ of all disks in M , i.e. all open subsets $U \subseteq M$ such that $U \cong \mathbb{R}^m$ is diffeomorphic to a Cartesian space, and endow it with the restricted orthogonality relation. This defines an orthogonal category $\overline{\mathbf{Disk}(M)}$ together with a full orthogonal subcategory embedding $J : \overline{\mathbf{Disk}(M)} \rightarrow \overline{\mathbf{Open}(M)}$. For the case of the circle $M = \mathbb{S}^1$, the corresponding extension functor $J_! : \mathbf{AQFT}(\overline{\mathbf{Disk}(\mathbb{S}^1)}) \rightarrow \mathbf{AQFT}(\overline{\mathbf{Open}(\mathbb{S}^1)})$ is studied in the context of chiral conformal AQFT [Fre90, Fre93, FRS92]. ∇

Example 6.2. Recall from Example 2.3 the orthogonal category $\overline{\mathbf{Loc}} = (\mathbf{Loc}, \perp_{\mathbf{Loc}})$ of oriented and time-oriented globally hyperbolic Lorentzian manifolds. Consider the full subcategory $\mathbf{Loc}_{\diamond} \subseteq \mathbf{Loc}$ of all objects $M \in \mathbf{Loc}$ whose underlying manifold is diffeomorphic to a Cartesian space and endow it with the restricted orthogonality relation. This defines an orthogonal category $\overline{\mathbf{Loc}_{\diamond}}$ together with a full orthogonal subcategory embedding $J : \overline{\mathbf{Loc}_{\diamond}} \rightarrow \overline{\mathbf{Loc}}$. The corresponding extension functor $J_! : \mathbf{AQFT}(\overline{\mathbf{Loc}_{\diamond}}) \rightarrow \mathbf{AQFT}(\overline{\mathbf{Loc}})$ is studied in the context of locally covariant AQFT [Lan14, BSW17]. ∇

We will study a generalization of this extension construction to 2AQFTs, which is based on the biadjunction

$$J_! : \mathbf{2AQFT}(\overline{\mathbf{C}}) \rightleftarrows \mathbf{2AQFT}(\overline{\mathbf{D}}) : J^* \quad , \quad (6.1)$$

where the right adjoint 2-functor J^* is given by restriction of 2AQFTs along J . Hence, the left adjoint pseudo-functor $J_!$ is a 2-categorical generalization of operadic left Kan extension. Given any $\mathfrak{A} \in \mathbf{2AQFT}(\overline{\mathbf{C}})$ on $\overline{\mathbf{C}}$, this determines a canonical extension $J_! \mathfrak{A} \in \mathbf{2AQFT}(\overline{\mathbf{D}})$ to the larger orthogonal category $\overline{\mathbf{D}}$. Following AQFT terminology, we shall refer to the locally presentable \mathbb{K} -linear category $J_! \mathfrak{A}(d) \in \mathbf{Pr}_{\mathbb{K}}$ that is assigned by the extended 2AQFT $J_! \mathfrak{A}$ to an object $d \in \mathbf{D}$ as *Fredenhagen's universal category*. In the context of Example 6.1, we will provide examples of such categories for 2AQFTs on the circle $M = \mathbb{S}^1$.

6.1 Preliminaries

Our construction of the extension pseudo-functor $J_! : \mathbf{2AQFT}(\overline{\mathbf{C}}) \rightarrow \mathbf{2AQFT}(\overline{\mathbf{D}})$ associated to a full orthogonal subcategory embedding $J : \overline{\mathbf{C}} \rightarrow \overline{\mathbf{D}}$ uses the *monoidal envelope* $\mathcal{P}_{\overline{\mathbf{C}}}^{\otimes}$ of the prefactorization operad $\mathcal{P}_{\overline{\mathbf{C}}}$ from Definition 2.5. We refer the reader to [EM09, Theorem 4.2] for details on monoidal envelopes for colored operads. In our case of interest, $\mathcal{P}_{\overline{\mathbf{C}}}^{\otimes}$ is given by the following symmetric monoidal category:

Objects: (Possibly empty) tuples $\underline{c} = (c_1, \dots, c_n) \in \mathbf{C}^n$ of objects in $\mathcal{P}_{\overline{\mathbf{C}}}$.

Morphisms: Pairs $(\alpha, \underline{f}) : \underline{c} = (c_1, \dots, c_n) \rightarrow \underline{t} = (t_1, \dots, t_m)$ with $\alpha : \{1, \dots, n\} \rightarrow \{1, \dots, m\}$ a map of sets and $\underline{f} = (\underline{f}_1, \dots, \underline{f}_m)$ a tuple of operations $\underline{f}_j = (f_{j1}, \dots, f_{jk_j}) \in \mathcal{P}_{\overline{\mathbf{C}}}(\underline{c}_{\alpha,j})$, for $j = 1, \dots, m$, where $\underline{c}_{\alpha,j}$ denotes the (possibly empty) sub-tuple of \underline{c} containing only the c_i 's satisfying $\alpha(i) = j$ and k_j denotes the length of $\underline{c}_{\alpha,j}$.

Identities and composition: The identity morphism for $\underline{c} = (c_1, \dots, c_n) \in \mathcal{P}_{\overline{\mathbf{C}}}^{\otimes}$ is given by $\text{id}_{\underline{c}} := (\text{id}, (\text{id}_{c_1}, \dots, \text{id}_{c_n})) : \underline{c} \rightarrow \underline{c}$. The composition of two morphisms $(\alpha, \underline{f}) : \underline{b} \rightarrow \underline{a}$ and $(\beta, \underline{g}) : \underline{a} \rightarrow \underline{t}$ in $\mathcal{P}_{\overline{\mathbf{C}}}^{\otimes}$ is given by $(\beta\alpha, \underline{h}) := (\beta\alpha, \underline{h}) : \underline{b} \rightarrow \underline{t}$, where $\beta\alpha$ is the usual composition of maps of sets and $\underline{h} := (\underline{h}_1, \dots, \underline{h}_\ell)$ is the tuple of operations $\underline{h}_k := \underline{g}_k \underline{f}_{\beta,k} \in \mathcal{P}_{\overline{\mathbf{C}}}(\underline{b}_{\beta\alpha,k})$ determined by operadic composition, for $k = 1, \dots, \ell$, where $\underline{f}_{\beta,k}$ is the sub-tuple of $\underline{f} = (\underline{f}_1, \dots, \underline{f}_m)$ containing only the \underline{f}_j 's satisfying $\beta(j) = k$.

Symmetric monoidal structure: The tensor product $\underline{c} \otimes \underline{c}' := (\underline{c}, \underline{c}')$ is defined by concatenation of tuples, the monoidal unit is the empty tuple \emptyset and the symmetric braiding is given by the $\mathcal{P}_{\overline{\mathbf{C}}}^{\otimes}$ -morphisms $(\alpha_{n,n'}, (\text{id}_{c_1}, \dots, \text{id}_{c_n}, \text{id}_{c'_1}, \dots, \text{id}_{c'_{n'}})) : \underline{c} \otimes \underline{c}' \rightarrow \underline{c}' \otimes \underline{c}$, where $\alpha_{n,n'} : \{1, \dots, n+n'\} \rightarrow \{1, \dots, n+n'\}$ is defined by $\alpha_{n,n'}(i) = n' + i$, for $i = 1, \dots, n$, and $\alpha_{n,n'}(n+i) = i$, for $i = 1, \dots, n'$.

Applying the same construction to $\mathcal{P}_{\overline{\mathbf{D}}}$ defines a symmetric monoidal category $\mathcal{P}_{\overline{\mathbf{D}}}^{\otimes}$. Furthermore, the orthogonal functor $J : \overline{\mathbf{C}} \rightarrow \overline{\mathbf{D}}$ induces an operad morphism $J : \mathcal{P}_{\overline{\mathbf{C}}} \rightarrow \mathcal{P}_{\overline{\mathbf{D}}}$ and hence a symmetric monoidal functor $J^{\otimes} : \mathcal{P}_{\overline{\mathbf{C}}}^{\otimes} \rightarrow \mathcal{P}_{\overline{\mathbf{D}}}^{\otimes}$ between the monoidal envelopes. The latter reads explicitly as follows:

On objects: For $\underline{c} = (c_1, \dots, c_n) \in \mathcal{P}_{\overline{\mathbf{C}}}^{\otimes}$, we set $J^{\otimes}(\underline{c}) := (J(c_1), \dots, J(c_n)) \in \mathcal{P}_{\overline{\mathbf{D}}}^{\otimes}$.

On morphisms: For $(\alpha, \underline{f}) : \underline{c} \rightarrow \underline{t}$ in $\mathcal{P}_{\overline{\mathbf{C}}}^{\otimes}$, we set $J^{\otimes}(\alpha, \underline{f}) := (\alpha, J(\underline{f})) : J^{\otimes}(\underline{c}) \rightarrow J^{\otimes}(\underline{t})$ in $\mathcal{P}_{\overline{\mathbf{D}}}^{\otimes}$, where $J(\underline{f}) := ((J(f_{11}), \dots, J(f_{1k_1})), \dots, (J(f_{m1}), \dots, J(f_{mk_m})))$.

Symmetric monoidal structure: Since $J^{\otimes}(\underline{c}) \otimes J^{\otimes}(\underline{c}') = J^{\otimes}(\underline{c} \otimes \underline{c}')$ and $\emptyset = J^{\otimes}(\emptyset)$, it is straightforward to equip J^{\otimes} with a symmetric monoidal structure.

Recall from Definition 3.3 that 2AQFTs on $\overline{\mathbf{C}}$ are by definition $\mathcal{P}_{\overline{\mathbf{C}}}$ -algebras. Hence, by the universal property of monoidal envelopes, we can associate to every $\mathfrak{A} \in \mathbf{2AQFT}(\overline{\mathbf{C}})$ a symmetric monoidal pseudo-functor

$$\underline{\mathfrak{A}} : \mathcal{P}_{\overline{\mathbf{C}}}^{\otimes} \longrightarrow \mathbf{Pr}_{\mathbb{K}} \quad (6.2a)$$

from the monoidal envelope of $\mathcal{P}_{\overline{\mathbf{C}}}$. This pseudo-functor acts on objects $\underline{c} = (c_1, \dots, c_n) \in \mathcal{P}_{\overline{\mathbf{C}}}^{\otimes}$ as the n -ary Kelly-Deligne tensor product

$$\underline{\mathfrak{A}}(\underline{c}) := \bigotimes_{i=1}^n \underline{\mathfrak{A}}(c_i) \quad (6.2b)$$

of the locally presentable \mathbb{K} -linear categories $\underline{\mathfrak{A}}(c_i) \in \mathbf{Pr}_{\mathbb{K}}$, cf. Remark 3.2. (By convention, we set $\underline{\mathfrak{A}}(\emptyset) := \mathbf{Vec}_{\mathbb{K}}$ to be the monoidal unit of $\mathbf{Pr}_{\mathbb{K}}$.) On morphisms $(\alpha, \underline{f}) : \underline{c} \rightarrow \underline{t}$ in $\mathcal{P}_{\overline{\mathbf{C}}}^{\otimes}$, this pseudo-functor acts as

$$\underline{\mathfrak{A}}(\alpha, \underline{f}) : \underline{\mathfrak{A}}(\underline{c}) = \bigotimes_{i=1}^n \underline{\mathfrak{A}}(c_i) \xrightarrow{\simeq_{\alpha}} \bigotimes_{j=1}^m \underline{\mathfrak{A}}(\underline{c}_{\alpha,j}) \xrightarrow{\bigotimes_j \underline{\mathfrak{A}}(\underline{f}_j)} \bigotimes_{j=1}^m \underline{\mathfrak{A}}(t_j) = \underline{\mathfrak{A}}(\underline{t}) \quad , \quad (6.2c)$$

where \simeq_{α} is the equivalence in the symmetric monoidal 2-category $\mathbf{Pr}_{\mathbb{K}}$ that is associated to the displayed permutation determined by α . The coherence data for the symmetric monoidal pseudo-functor $\underline{\mathfrak{A}} : \mathcal{P}_{\overline{\mathbf{C}}}^{\otimes} \rightarrow \mathbf{Pr}_{\mathbb{K}}$ are canonically given by the coherence data for $\mathfrak{A} \in \mathbf{2AQFT}(\overline{\mathbf{C}})$ and the symmetric monoidal structure on $\mathbf{Pr}_{\mathbb{K}}$.

6.2 Extension

The extension pseudo-functor $J_! : \mathbf{2AQFT}(\overline{\mathbf{C}}) \rightarrow \mathbf{2AQFT}(\overline{\mathbf{D}})$ in the biadjunction (6.1) is obtained canonically via operadic left pseudo-Kan extension along $J : \mathcal{P}_{\overline{\mathbf{C}}} \rightarrow \mathcal{P}_{\overline{\mathbf{D}}}$. Passing from colored operads to their monoidal envelopes, $J_!$ can be obtained via (categorical) left pseudo-Kan extension along $J^\otimes : \mathcal{P}_{\overline{\mathbf{C}}}^\otimes \rightarrow \mathcal{P}_{\overline{\mathbf{D}}}^\otimes$, cf. [Hor17]. Furthermore, the latter left pseudo-Kan extension can be computed in terms of suitable bicolimits [Lac10, LN16]. Using this approach, we can now describe the extension $J_!(\mathfrak{A}) \in \mathbf{2AQFT}(\overline{\mathbf{D}})$ of a $\mathbf{2AQFT} \mathfrak{A} \in \mathbf{2AQFT}(\overline{\mathbf{C}})$. For each $d \in \mathbf{D}$, Fredenhagen's universal category is the locally presentable \mathbb{K} -linear category

$$J_!(\mathfrak{A})(d) := \text{bicolim} \left(J^\otimes / (d) \xrightarrow{\text{forget}} \mathcal{P}_{\overline{\mathbf{C}}}^\otimes \xrightarrow{\mathfrak{A}} \mathbf{Pr}_{\mathbb{K}} \right) \quad (6.3)$$

obtained as a bicolimit in $\mathbf{Pr}_{\mathbb{K}}$, where $J^\otimes / (d)$ denotes the slice category for the functor $J^\otimes : \mathcal{P}_{\overline{\mathbf{C}}}^\otimes \rightarrow \mathcal{P}_{\overline{\mathbf{D}}}^\otimes$ over the object $(d) \in \mathcal{P}_{\overline{\mathbf{D}}}^\otimes$. Recall also (6.2) for the construction of the pseudo-functor $\mathfrak{A} : \mathcal{P}_{\overline{\mathbf{C}}}^\otimes \rightarrow \mathbf{Pr}_{\mathbb{K}}$. (To avoid confusion, let us stress that the symbol (d) stands for the tuple of length one that is defined by the object $d \in \mathbf{D}$, i.e. $(d) \in \mathcal{P}_{\overline{\mathbf{D}}}^\otimes$ is an object in the monoidal envelope.) This bicolimit always exists because $\mathbf{Pr}_{\mathbb{K}}$ is bicategorically cocomplete, see e.g. [BCJF15, Lemma 2.5]. For each tuple $\underline{g} = (g_1, \dots, g_n) \in \mathcal{P}_{\overline{\mathbf{D}}}^\otimes(\underline{s})$ of mutually orthogonal \mathbf{D} -morphisms, we set the factorization product

$$J_!(\mathfrak{A})(\underline{g}) : \prod_{i=1}^n J_!(\mathfrak{A})(d_i) \rightarrow J_!(\mathfrak{A})(s) \quad (6.4a)$$

to be the functor that is defined below, which is co-continuous and \mathbb{K} -linear in each entry: Consider the diagram

$$\begin{array}{ccccc} \prod_{i=1}^n J^\otimes / (d_i) & \xrightarrow{\prod_i \text{forget}} & \prod_{i=1}^n \mathcal{P}_{\overline{\mathbf{C}}}^\otimes & \xrightarrow{\prod_i \mathfrak{A}} & \prod_{i=1}^n \mathbf{Pr}_{\mathbb{K}} \\ \underline{g}_* \downarrow & & \otimes^n \downarrow & \swarrow (\star) & \downarrow \boxtimes^n \\ J^\otimes / (s) & \xrightarrow{\text{forget}} & \mathcal{P}_{\overline{\mathbf{C}}}^\otimes & \xrightarrow{\mathfrak{A}} & \mathbf{Pr}_{\mathbb{K}} \end{array} \quad (6.4b)$$

where $\underline{g}_* : \prod_{i=1}^n J^\otimes / (d_i) \rightarrow J^\otimes / (s)$ is the functor induced by post-composition with \underline{g} in the colored operad $\mathcal{P}_{\overline{\mathbf{D}}}$. By direct inspection, the left square commutes. In the right square, instead, the clockwise and counter-clockwise paths give functors that are related by the natural isomorphism (\star) determined by the symmetric monoidal structure on the pseudo-functor \mathfrak{A} . Passing to bicolimits and recalling that the Kelly-Deligne tensor product \boxtimes commutes with bicolim (in each entry) provides a co-continuous \mathbb{K} -linear functor $\boxtimes_{i=1}^n J_!(\mathfrak{A})(d_i) \rightarrow J_!(\mathfrak{A})(s)$. Pre-composition with the canonical functor $\prod_{i=1}^n J_!(\mathfrak{A})(d_i) \rightarrow \boxtimes_{i=1}^n J_!(\mathfrak{A})(d_i)$, which is co-continuous and \mathbb{K} -linear in each entry, completes the construction of (6.4a). For the empty tuple $\underline{d} = \emptyset$, the pointing $J_!(\mathfrak{A})(*_s) \in J_!(\mathfrak{A})(s)$ of Fredenhagen's universal category $J_!(\mathfrak{A})(s)$ is obtained in the same fashion from (6.4b). (Notice that empty products are initial categories, while \otimes^0 and \boxtimes^0 assign the respective monoidal units.) The coherence data, cf. Remark 3.4, for the extended $\mathbf{2AQFT} J_!(\mathfrak{A}) \in \mathbf{2AQFT}(\overline{\mathbf{D}})$ are obtained canonically from the construction above and the symmetric monoidal pseudo-functor $\mathfrak{A} : \mathcal{P}_{\overline{\mathbf{C}}}^\otimes \rightarrow \mathbf{Pr}_{\mathbb{K}}$.

For an arbitrary $d \in \mathbf{D}$, we shall now describe Fredenhagen's universal category $J_!(\mathfrak{A})(d)$ in fully explicit terms, using the prescription in [BCJF15, Lemma 2.5] to compute the relevant bicolimit (6.3). This is a two-step procedure:

- 1.) Every co-continuous \mathbb{K} -linear functor between two locally presentable \mathbb{K} -linear categories admits a right adjoint by the special adjoint functor theorem, cf. [AR94, BCJF15]. Hence, from the pseudo-functor $\mathfrak{A} : \mathcal{P}_{\overline{\mathbf{C}}}^\otimes \rightarrow \mathbf{Pr}_{\mathbb{K}}$, we obtain a new pseudo-functor $\mathfrak{A}^R : (\mathcal{P}_{\overline{\mathbf{C}}}^\otimes)^{\text{op}} \rightarrow$

\mathbf{Cat} that acts on objects as $\underline{\mathfrak{A}}$, i.e. $\underline{\mathfrak{A}}^{\mathbf{R}}(\underline{c}) := \underline{\mathfrak{A}}(\underline{c})$ for all $\underline{c} \in \mathcal{P}_{\mathbb{C}}^{\otimes}$, and that assigns to a morphism $(\alpha, \underline{f}) : \underline{c} \rightarrow \underline{t}$ in $\mathcal{P}_{\mathbb{C}}^{\otimes}$ the right adjoint of the co-continuous \mathbb{K} -linear functor assigned by $\underline{\mathfrak{A}}$, i.e. $\underline{\mathfrak{A}}(\alpha, \underline{f}) \dashv \underline{\mathfrak{A}}^{\mathbf{R}}(\alpha, \underline{f}) : \underline{\mathfrak{A}}^{\mathbf{R}}(\underline{t}) \rightarrow \underline{\mathfrak{A}}^{\mathbf{R}}(\underline{c})$. (Note that $\underline{\mathfrak{A}}^{\mathbf{R}}$ is just a pseudo-functor to \mathbf{Cat} and not necessarily to $\mathbf{Pr}_{\mathbb{K}}$ because the right adjoint functors $\underline{\mathfrak{A}}^{\mathbf{R}}(\alpha, \underline{f})$ may fail to be co-continuous.)

- 2.) The category underlying the bicolimit (6.3) of $\underline{\mathfrak{A}} \circ \text{forget} : J^{\otimes}/(d) \rightarrow \mathbf{Pr}_{\mathbb{K}}$ can be computed as a bilimit of the pseudo-functor $\underline{\mathfrak{A}}^{\mathbf{R}} \circ \text{forget} : (J^{\otimes}/(d))^{\text{op}} \rightarrow \mathbf{Cat}$. The outcome is a locally presentable \mathbb{K} -linear category in a canonical way, cf. [BCJF15].

Using the explicit model [Str80, LN16] for computing bilimits of pseudo-functors to \mathbf{Cat} , we obtain the following description of Fredenhagen's universal category $J_!(\underline{\mathfrak{A}})(d)$ in terms of explicit data and conditions:

Objects: An object

$$(V, \xi^V) := (\{V_{\underline{h}}\}, \{\xi_{(\alpha, \underline{f})}^V\}) \in J_!(\underline{\mathfrak{A}})(d) \quad (6.5)$$

consists of the following data:

- (1) For each object $(\underline{h} := (*, \underline{h}) : \underline{c} \rightarrow (d)) \in J^{\otimes}/(d)$, where $* : \{1, \dots, n\} \rightarrow \{1\}$ denotes the unique map of sets to the singleton $\{1\}$, an object

$$V_{\underline{h}} \in \underline{\mathfrak{A}}(\underline{c}) = \bigotimes_{i=1}^n \underline{\mathfrak{A}}(c_i) \quad (6.6a)$$

- (2) For each morphism $(\alpha, \underline{f}) : \underline{h} \rightarrow \underline{h}'$ in $J^{\otimes}/(d)$, an isomorphism

$$\xi_{(\alpha, \underline{f})}^V : \underline{\mathfrak{A}}^{\mathbf{R}}(\alpha, \underline{f})(V_{\underline{h}'}) \longrightarrow V_{\underline{h}} \quad (6.6b)$$

in the category $\underline{\mathfrak{A}}(\underline{c})$.

These data have to satisfy the following cocycle conditions:

- (i) For all objects $(\underline{h} : \underline{c} \rightarrow (d)) \in J^{\otimes}/(d)$, the diagram

$$\begin{array}{ccc} \underline{\mathfrak{A}}^{\mathbf{R}}(\text{id}_{\underline{h}})(V_{\underline{h}}) & \xrightarrow{\xi_{\text{id}_{\underline{h}}}^V} & V_{\underline{h}} \\ \underline{\mathfrak{A}}_{\underline{c}}^{\mathbf{R}0} \downarrow \cong & \searrow \text{id}_{V_{\underline{h}}} & \\ V_{\underline{h}} & & \end{array} \quad (6.7a)$$

in $\underline{\mathfrak{A}}(\underline{c})$ commutes, where $\underline{\mathfrak{A}}_{\underline{c}}^{\mathbf{R}0}$ denotes the coherence isomorphisms for identities that are associated with the pseudo-functor $\underline{\mathfrak{A}}^{\mathbf{R}}$.

- (ii) For all composable pairs of morphisms $(\alpha, \underline{f}) : \underline{h} \rightarrow \underline{h}'$ and $(\beta, \underline{g}) : \underline{h}' \rightarrow \underline{h}''$ in $J^{\otimes}/(d)$, the diagram

$$\begin{array}{ccc} \underline{\mathfrak{A}}^{\mathbf{R}}(\alpha, \underline{f}) \underline{\mathfrak{A}}^{\mathbf{R}}(\beta, \underline{g})(V_{\underline{h}''}) & \xrightarrow{\underline{\mathfrak{A}}^{\mathbf{R}}(\alpha, \underline{f})(\xi_{(\beta, \underline{g})}^V)} & \underline{\mathfrak{A}}^{\mathbf{R}}(\alpha, \underline{f})(V_{\underline{h}'}) \\ \underline{\mathfrak{A}}_{((\beta, \underline{g}), (\alpha, \underline{f}))}^{\mathbf{R}2} \downarrow \cong & & \downarrow \xi_{(\alpha, \underline{f})}^V \\ \underline{\mathfrak{A}}^{\mathbf{R}}((\beta, \underline{g}) \circ (\alpha, \underline{f}))(V_{\underline{h}''}) & \xrightarrow{\xi_{(\beta, \underline{g}) \circ (\alpha, \underline{f})}^V} & V_{\underline{h}} \end{array} \quad (6.7b)$$

in $\underline{\mathfrak{A}}(\underline{c})$ commutes, where $\underline{\mathfrak{A}}_{((\beta, \underline{g}), (\alpha, \underline{f}))}^{\mathbf{R}2}$ denotes the coherence isomorphisms for compositions that are associated with the pseudo-functor $\underline{\mathfrak{A}}^{\mathbf{R}}$.

Morphisms: A morphism

$$\Gamma := \{\Gamma_{\underline{h}}\} : (V, \xi^V) \longrightarrow (W, \xi^W) \quad (6.8)$$

in $J_!(\mathfrak{A})(d)$ consists of a family of $\mathfrak{A}(\underline{c})$ -morphisms

$$\Gamma_{\underline{h}} : V_{\underline{h}} \longrightarrow W_{\underline{h}} \quad , \quad (6.9a)$$

for all $(\underline{h} : \underline{c} \rightarrow (d)) \in J^\otimes/(d)$, such that the diagrams

$$\begin{array}{ccc} \mathfrak{A}^R(\alpha, \underline{f})(V_{\underline{h}'}) & \xrightarrow{\mathfrak{A}^R(\alpha, \underline{f})(\Gamma_{\underline{h}'})} & \mathfrak{A}^R(\alpha, \underline{f})(W_{\underline{h}'}) \\ \xi_{(\alpha, \underline{f})}^V \downarrow & & \downarrow \xi_{(\alpha, \underline{f})}^W \\ V_{\underline{h}} & \xrightarrow{\Gamma_{\underline{h}}} & W_{\underline{h}} \end{array} \quad (6.9b)$$

in $\mathfrak{A}(\underline{c})$ commute, for all morphisms $(\alpha, \underline{f}) : \underline{h} \rightarrow \underline{h}'$ in $J^\otimes/(d)$.

Identities and composition: Identities and composition are defined component-wise.

6.3 Examples on $M = \mathbb{S}^1$

The aim of this subsection is to study examples of Fredenhagen's universal category for the full orthogonal subcategory embedding $J : \overline{\mathbf{Disk}}(\mathbb{S}^1) \rightarrow \overline{\mathbf{Open}}(\mathbb{S}^1)$ introduced in Example 6.1. Given any $\mathfrak{A} \in \mathbf{2AQFT}(\overline{\mathbf{Disk}}(\mathbb{S}^1))$, which is by definition only defined on open intervals in \mathbb{S}^1 , we are particularly interested in the locally presentable \mathbb{K} -linear category

$$J_!(\mathfrak{A})(\mathbb{S}^1) = \text{bicolim} \left(J^\otimes/(\mathbb{S}^1) \xrightarrow{\text{forget}} \mathcal{P}_{\overline{\mathbf{Disk}}(\mathbb{S}^1)}^\otimes \xrightarrow{\mathfrak{A}} \mathbf{Pr}_{\mathbb{K}} \right) \quad (6.10)$$

that is assigned to the whole circle \mathbb{S}^1 . The slice category $J^\otimes/(\mathbb{S}^1)$ in the present case admits the following simple description: An object is a tuple $\underline{I} = (I_1, \dots, I_n)$ of mutually disjoint open intervals $I_i \subset \mathbb{S}^1$, i.e. $I_i \cap I_j = \emptyset$ for all $i \neq j$. A morphism $\alpha : \underline{I} = (I_1, \dots, I_n) \rightarrow \underline{J} = (J_1, \dots, J_m)$ is a map of sets $\alpha : \{1, \dots, n\} \rightarrow \{1, \dots, m\}$ such that $I_i \subseteq J_{\alpha(i)}$, for all $i = 1, \dots, n$.

Example 6.3. Let us consider first the simplest case where \mathfrak{A} is truncated, i.e. $\mathfrak{A} = \iota(\mathfrak{A})$ with $\mathfrak{A} \in \mathbf{AQFT}(\overline{\mathbf{Disk}}(\mathbb{S}^1))$ an ordinary AQFT. We have the following square of biadjunctions

$$\begin{array}{ccc} \mathbf{AQFT}(\overline{\mathbf{Disk}}(\mathbb{S}^1)) & \begin{array}{c} \xrightarrow{\iota} \\ \xleftarrow{\pi} \end{array} & \mathbf{2AQFT}(\overline{\mathbf{Disk}}(\mathbb{S}^1)) \\ J_! \updownarrow J^* & & J_! \updownarrow J^* \\ \mathbf{AQFT}(\overline{\mathbf{Open}}(\mathbb{S}^1)) & \begin{array}{c} \xrightarrow{\iota} \\ \xleftarrow{\pi} \end{array} & \mathbf{2AQFT}(\overline{\mathbf{Open}}(\mathbb{S}^1)) \end{array} \quad (6.11)$$

The horizontal biadjunctions are the inclusion-truncation biadjunctions from Theorem 4.3, the left vertical (bi)adjunction is an ordinary operadic left Kan extension and the right vertical biadjunction is the operadic left pseudo-Kan extension (6.1). By direct inspection, one confirms that the square formed by the right adjoint 2-functors commutes, i.e. $\pi J^* = J^* \pi$, hence the square formed by the left adjoint pseudo-functors commutes up to an equivalence, i.e. $\iota J_! \simeq J_! \iota$. As a consequence, Fredenhagen's universal category for a truncated 2AQFT

$$J_!(\iota(\mathfrak{A}))(\mathbb{S}^1) \simeq \mathbf{Mod}_{J_!(\mathfrak{A})(\mathbb{S}^1)} \quad (6.12)$$

is equivalent to the category of right modules over Fredenhagen's universal algebra $J_!(\mathfrak{A})(\mathbb{S}^1) \in \mathbf{Alg}_{\mathbb{K}}$. The latter is given by the ordinary colimit

$$J_!(\mathfrak{A})(\mathbb{S}^1) = \operatorname{colim} \left(J^{\otimes}/(\mathbb{S}^1) \xrightarrow{\text{forget}} \mathcal{P}_{\overline{\mathbf{Disk}(\mathbb{S}^1)}}^{\otimes} \xrightarrow{\mathfrak{A}} \mathbf{Alg}_{\mathbb{K}} \right) , \quad (6.13)$$

where $\mathfrak{A} : \mathcal{P}_{\overline{\mathbf{Disk}(\mathbb{S}^1)}}^{\otimes} \rightarrow \mathbf{Alg}_{\mathbb{K}}$ is the symmetric monoidal functor from the monoidal envelope that is determined by $\mathfrak{A} \in \mathbf{AQFT}(\overline{\mathbf{Disk}(\mathbb{S}^1)})$.

To obtain a better understanding of the objects and morphisms in our general presentation of Fredenhagen's universal category $J_!(\iota(\mathfrak{A}))(\mathbb{S}^1)$, we construct explicitly a functor $\mathbf{Mod}_{J_!(\mathfrak{A})(\mathbb{S}^1)} \rightarrow J_!(\iota(\mathfrak{A}))(\mathbb{S}^1)$ that implements the equivalence (6.12). Let us first describe this functor on objects. Given any right module $V \in \mathbf{Mod}_{J_!(\mathfrak{A})(\mathbb{S}^1)}$ over Fredenhagen's universal algebra, we use the canonical $\mathbf{Alg}_{\mathbb{K}}$ -morphisms $\chi_{\underline{I}} : \mathfrak{A}(\underline{I}) = \bigotimes_{i=1}^n \mathfrak{A}(I_i) \rightarrow J_!(\mathfrak{A})(\mathbb{S}^1)$ to the colimit (6.13) to define

$$V_{\underline{I}} := \chi_{\underline{I}}^*(V) \in \mathbf{Mod}_{\mathfrak{A}(\underline{I})} \quad (6.14)$$

by restriction of modules, for each $\underline{I} \in J^{\otimes}/(\mathbb{S}^1)$. Given any morphism $\alpha : \underline{I} \rightarrow \underline{J}$ in $J^{\otimes}/(\mathbb{S}^1)$, the functor $\iota(\mathfrak{A})^{\mathbf{R}}(\alpha) = \mathfrak{A}(\alpha)^* : \mathbf{Mod}_{\mathfrak{A}(\underline{J})} \rightarrow \mathbf{Mod}_{\mathfrak{A}(\underline{I})}$ is given by restriction of modules along the $\mathbf{Alg}_{\mathbb{K}}$ -morphism $\mathfrak{A}(\alpha) : \mathfrak{A}(\underline{I}) \rightarrow \mathfrak{A}(\underline{J})$. Because $\{\chi_{\underline{I}}\}_{\underline{I}}$ is a co-cone, we obtain

$$\iota(\mathfrak{A})^{\mathbf{R}}(\alpha)(V_{\underline{J}}) = \mathfrak{A}(\alpha)^* \chi_{\underline{J}}^*(V) = (\chi_{\underline{J}} \mathfrak{A}(\alpha))^*(V) = \chi_{\underline{I}}^*(V) = V_{\underline{I}} \quad (6.15a)$$

and therefore we can set

$$\xi_{\alpha}^V := \operatorname{id}_{V_{\underline{I}}} : \iota(\mathfrak{A})^{\mathbf{R}}(\alpha)(V_{\underline{J}}) \longrightarrow V_{\underline{I}} \quad (6.15b)$$

to be the identity morphism. One easily checks that the coherence conditions (6.7) are satisfied, hence we have defined an object $(V, \xi^V) \in J_!(\iota(\mathfrak{A}))(\mathbb{S}^1)$. Let us now define the functor $\mathbf{Mod}_{J_!(\mathfrak{A})(\mathbb{S}^1)} \rightarrow J_!(\iota(\mathfrak{A}))(\mathbb{S}^1)$ on morphisms. Given any morphism $L : V \rightarrow W$ in $\mathbf{Mod}_{J_!(\mathfrak{A})(\mathbb{S}^1)}$, consider the restrictions

$$L_{\underline{I}} := \chi_{\underline{I}}^*(L) : V_{\underline{I}} = \chi_{\underline{I}}^*(V) \longrightarrow \chi_{\underline{I}}^*(W) = W_{\underline{I}} , \quad (6.16)$$

for all $\underline{I} \in J^{\otimes}/(\mathbb{S}^1)$. One easily checks that the coherence conditions (6.9) are satisfied, hence we have defined a morphism $L : (V, \xi^V) \rightarrow (W, \xi^W)$ in $J_!(\iota(\mathfrak{A}))(\mathbb{S}^1)$. Using the universal property of the colimit (6.13), one checks that the resulting functor $\mathbf{Mod}_{J_!(\mathfrak{A})(\mathbb{S}^1)} \rightarrow J_!(\iota(\mathfrak{A}))(\mathbb{S}^1)$ implements the equivalence (6.12).

Summing up, we have found that, in the case of a truncated 2AQFT $\mathfrak{A} = \iota(\mathfrak{A})$, the objects of Fredenhagen's universal category can be described as families of right modules (6.14) over the local algebras $\mathfrak{A}(\underline{I}) = \bigotimes_{i=1}^n \mathfrak{A}(I_i)$ on disjoint unions of intervals, whose restrictions along inclusions $\alpha : \underline{I} \rightarrow \underline{J}$ coincide (6.15). Morphisms in Fredenhagen's universal category can be described by locally defined module morphisms (6.16), whose restrictions along inclusions $\alpha : \underline{I} \rightarrow \underline{J}$ coincide. ∇

Example 6.4. Let us consider the gauging $\mathbb{K}^G \in \mathbf{2AQFT}(\overline{\mathbf{Disk}(\mathbb{S}^1)})$ of the trivial AQFT $\mathbb{K} \in \mathbf{AQFT}(\overline{\mathbf{Disk}(\mathbb{S}^1)})$ with respect to the trivial action of a finite group G , which is a non-truncated 2AQFT for every non-trivial group $G \neq \{e\}$, see Example 5.7. Because 2AQFTs are by definition prefactorization algebras with values in $\mathbf{Pr}_{\mathbb{K}}$ (cf. Definition 3.3) and $\mathbb{K}^G \in \mathbf{2AQFT}(\overline{\mathbf{Disk}(\mathbb{S}^1)})$ is also *locally constant*, we can compute Fredenhagen's universal category $J_!(\mathbb{K}^G)(\mathbb{S}^1)$ for this particular example by factorization homology [AF15]. Using in particular [AF15, Theorem 3.19], we obtain that

$$J_!(\mathbb{K}^G)(\mathbb{S}^1) \simeq \mathbf{HH}_{\bullet}(\mathbf{Rep}_{\mathbb{K}}(G)) \quad (6.17)$$

is equivalent to the Hochschild homology of the associative and unital algebra $(\mathbf{Rep}_{\mathbb{K}}(G), \otimes, \mathbb{K}) \in \mathbf{Alg}_{\text{As}}(\mathbf{Pr}_{\mathbb{K}})$ in $\mathbf{Pr}_{\mathbb{K}}$. (The latter is just the usual monoidal category of \mathbb{K} -linear representations of G , regarded internally in the symmetric monoidal 2-category $\mathbf{Pr}_{\mathbb{K}}$.) Hochschild homology can be computed as a bicolimit (in $\mathbf{Pr}_{\mathbb{K}}$)

$$\mathrm{HH}_{\bullet}(\mathbf{Rep}_{\mathbb{K}}(G)) = \mathrm{bicolim} \left(\mathbf{Rep}_{\mathbb{K}}(G) \xleftarrow{\quad} \mathbf{Rep}_{\mathbb{K}}(G)^{\boxtimes 2} \xleftarrow{\quad} \mathbf{Rep}_{\mathbb{K}}(G)^{\boxtimes 3} \xleftarrow{\quad} \dots \right) \quad (6.18)$$

of the simplicial diagram associated with $(\mathbf{Rep}_{\mathbb{K}}(G), \otimes, \mathbb{K}) \in \mathbf{Alg}_{\text{As}}(\mathbf{Pr}_{\mathbb{K}})$, see e.g. [BZFN10, Section 5.1]. (As usual, we suppress the degeneracy maps in (6.18).) Since we are working in a 2-categorical context, this simplicial diagram may be truncated after $\mathbf{Rep}_{\mathbb{K}}(G)^{\boxtimes 3}$.

We will now compute the bicolimit (6.18) explicitly by using the techniques of [BCJF15], see also the end of Section 6.2 for a short summary. A more conceptual explanation of the obtained result is given in Remark 6.5 below. By [BCJF15, Lemma 2.5], we can compute this bicolimit in terms of the bilimit (in the 2-category \mathbf{Cat} of categories)

$$\mathrm{HH}_{\bullet}(\mathbf{Rep}_{\mathbb{K}}(G)) = \mathrm{bilim} \left(\mathbf{Rep}_{\mathbb{K}}(G) \rightrightarrows \mathbf{Rep}_{\mathbb{K}}(G^2) \rightrightarrows \mathbf{Rep}_{\mathbb{K}}(G^3) \right) \quad (6.19)$$

of the truncated cosimplicial diagram obtained by taking right adjoints of the face and degeneracy maps in (6.18). In this expression we have also used that $\mathbf{Rep}_{\mathbb{K}}(G)^{\boxtimes n} \simeq \mathbf{Rep}_{\mathbb{K}}(G^n)$ is equivalent to the representation category of the product group G^n . The coface and codegeneracy maps in (6.19) are given by coinduced representation functors $\phi_* : \mathbf{Rep}_{\mathbb{K}}(G') \rightarrow \mathbf{Rep}_{\mathbb{K}}(G'')$ for suitable group maps $\phi : G' \rightarrow G''$. Concretely, we have that

$$\delta^0 = \delta^1 = \Delta_* : \mathbf{Rep}_{\mathbb{K}}(G) \longrightarrow \mathbf{Rep}_{\mathbb{K}}(G^2) \quad (6.20)$$

for the diagonal map $\Delta : G \rightarrow G^2$, $g \mapsto (g, g)$, and that

$$\delta^i = \phi_*^i : \mathbf{Rep}_{\mathbb{K}}(G^2) \longrightarrow \mathbf{Rep}_{\mathbb{K}}(G^3) \quad (6.21a)$$

for

$$\phi^i : G^2 \longrightarrow G^3, \quad (g_1, g_2) \mapsto \begin{cases} (g_1, g_1, g_2) & , \text{ for } i = 0, \\ (g_1, g_2, g_2) & , \text{ for } i = 1, \\ (g_1, g_2, g_1) & , \text{ for } i = 2. \end{cases} \quad (6.21b)$$

The codegeneracy map $\epsilon^0 : \mathbf{Rep}_{\mathbb{K}}(G^2) \rightarrow \mathbf{Rep}_{\mathbb{K}}(G)$ is given by the coinduced representation functor for $G^2 \rightarrow G$, $(g_1, g_2) \mapsto g_1$.

We are now ready to describe the bilimit (6.19) and hence the category $\mathrm{HH}_{\bullet}(\mathbf{Rep}_{\mathbb{K}}(G))$ in more explicit terms:

- An object is a tuple (V, θ^V) , where $V \in \mathbf{Rep}_{\mathbb{K}}(G)$ and $\theta^V : \delta^1(V) \rightarrow \delta^0(V)$ is an isomorphism in $\mathbf{Rep}_{\mathbb{K}}(G^2)$, such that $\epsilon^0(\theta^V) = \mathrm{id}_V$ and $\delta^0(\theta^V) \circ \delta^2(\theta^V) = \delta^1(\theta^V)$ in $\mathbf{Rep}_{\mathbb{K}}(G^3)$.
- A morphism $L : (V, \theta^V) \rightarrow (W, \theta^W)$ is a morphism $L : V \rightarrow W$ in $\mathbf{Rep}_{\mathbb{K}}(G)$, such that the diagram

$$\begin{array}{ccc} \delta^1(V) & \xrightarrow{\delta^1(L)} & \delta^1(W) \\ \theta^V \downarrow & & \downarrow \theta^W \\ \delta^0(V) & \xrightarrow{\delta^0(L)} & \delta^0(W) \end{array} \quad (6.22)$$

in $\mathbf{Rep}_{\mathbb{K}}(G^2)$ commutes.

We can simplify this description further by using explicit models for the coinduced representation functors $\phi_* : \mathbf{Rep}_{\mathbb{K}}(G') \rightarrow \mathbf{Rep}_{\mathbb{K}}(G'')$ for group maps $\phi : G' \rightarrow G''$. Since we consider only *finite* groups and a base field \mathbb{K} of characteristic 0, there exists a natural isomorphism between the coinduced and the induced representation functors $\phi_* \cong \phi_! : \mathbf{Rep}_{\mathbb{K}}(G') \rightarrow \mathbf{Rep}_{\mathbb{K}}(G'')$. The latter is easy to describe: For $V \in \mathbf{Rep}_{\mathbb{K}}(G')$, we set $\phi_!(V) := \mathbb{K}[G''] \otimes_{\mathbb{K}[G']} V \in \mathbf{Rep}_{\mathbb{K}}(G'')$ to be the relative tensor product, where $\mathbb{K}[G']$ and $\mathbb{K}[G'']$ denote the group Hopf algebras associated with the finite groups G' and G'' . (Recall that $\mathbf{Rep}_{\mathbb{K}}(G) = \mathbb{K}[G]\mathbf{Mod}$ is the category of left $\mathbb{K}[G]$ -modules, and similar for G'' .) Given any object $(V, \theta^V) \in \mathbf{HH}_{\bullet}(\mathbf{Rep}_{\mathbb{K}}(G))$, we use this explicit description to deduce that $\theta^V : \mathbb{K}[G^2] \otimes_{\mathbb{K}[G]} V \rightarrow \mathbb{K}[G^2] \otimes_{\mathbb{K}[G]} V$ is completely determined by a \mathbb{K} -linear map $\vartheta^V : V \rightarrow \mathbb{K}[G] \otimes V$ via $\theta^V(1 \otimes 1 \otimes v) = 1 \otimes \vartheta^V(v)$, which is G -equivariant with respect to the adjoint action on $\mathbb{K}[G]$ and satisfies the axioms of a left $\mathbb{K}[G]$ -coaction. Moreover, we deduce that a morphism in $\mathbf{HH}_{\bullet}(\mathbf{Rep}_{\mathbb{K}}(G))$ is a G -equivariant map that preserves these $\mathbb{K}[G]$ -coactions. In summary, we have obtained the following chain of equivalences

$$J_!(\mathbb{K}^G)(\mathbb{S}^1) \simeq \mathbf{HH}_{\bullet}(\mathbf{Rep}_{\mathbb{K}}(G)) \simeq G\text{-}\mathbb{K}[G]\mathbf{Mod} \simeq G\text{-}\mathbf{Mod}_{\mathcal{O}(G)} \quad , \quad (6.23)$$

where in the last step we have used that $\mathbb{K}[G]$ -comodules are equivalent to modules over the dual Hopf algebra $\mathcal{O}(G)$ of functions on G . (The G -action on $\mathcal{O}(G)$ is again the adjoint action.)

Let us briefly explain the physical interpretation of this result. By Remark 5.6, we can interpret $\mathbb{K}^G \in \mathbf{2AQFT}(\overline{\mathbf{Disk}(\mathbb{S}^1)})$ as an orbifold σ -model that is defined on intervals and whose target is the classifying stack $\mathbf{BG} = \{*\}/G$ of G . Indeed, the stack of fields on an interval $I \subset \mathbb{S}^1$ is $\mathbf{Fields}(I) = \mathbf{Map}(I, \mathbf{BG}) \simeq \{*\}/G$ and its category of quasi-coherent sheaves is $\mathbf{QCoh}(\mathbf{Fields}(I)) \simeq \mathbf{Rep}_{\mathbb{K}}(G)$, which coincides with the category that the 2AQFT \mathbb{K}^G assigns to intervals. On the whole circle \mathbb{S}^1 , the stack of fields of this orbifold σ -model is given by the loop stack $\mathbf{Fields}(\mathbb{S}^1) = \mathbf{Map}(\mathbb{S}^1, \mathbf{BG}) \simeq \mathbf{Bun}_G(\mathbb{S}^1)$, which is equivalent to the stack of principal G -bundles on \mathbb{S}^1 . The non-trivial bundles can be interpreted physically as “twisted sectors” of this orbifold σ -model, see e.g. [DVVV89]. The category of quasi-coherent sheaves on this stack is given by $\mathbf{QCoh}(\mathbf{Fields}(\mathbb{S}^1)) \simeq G\text{-}\mathbf{Mod}_{\mathcal{O}(G)}$, which coincides with our result for Fredenhagen’s universal category (6.23). Hence, Fredenhagen’s universal category successfully detects all “twisted sectors” for this simple example of an orbifold σ -model. ∇

Remark 6.5. The category (6.23) that we obtain for the circle is the representation category of the groupoid of principal G -bundles over the circle, i.e. the representation category of the loop groupoid $G//G$ of G (the action groupoid of the action of G on itself by conjugation). This category is also the Drinfeld center of $\mathbf{Rep}_{\mathbb{K}}(G)$, i.e. the Hochschild cohomology. As a consequence, the Hochschild homology and Hochschild cohomology for $\mathbf{Rep}_{\mathbb{K}}(G)$ are equivalent. More general conditions under which one finds such an equivalence are given in [DSPS13, Corollary 3.1.5] within the framework of finite tensor categories and in [BZFN10, Theorem 1.7] within the framework of derived algebraic geometry. \triangle

Example 6.6. As a last example, we discuss briefly the gauging $\mathfrak{A}^G \in \mathbf{2AQFT}(\overline{\mathbf{Disk}(\mathbb{S}^1)})$ of an arbitrary G -equivariant AQFT $(\mathfrak{A}, \rho) \in G\text{-}\mathbf{AQFT}(\overline{\mathbf{Disk}(\mathbb{S}^1)})$, which includes Examples 6.3 and 6.4 as very special cases. Unfortunately, it seems to be very hard to simplify our explicit description of Fredenhagen’s universal category $J_!(\mathfrak{A}^G)(\mathbb{S}^1)$ in this general case. (Note that computing this category as in Example 6.4 by importing techniques from factorization homology is in general not possible, because we are also interested in 2AQFTs that are *not* locally constant.) In order to develop a better understanding of the category $J_!(\mathfrak{A}^G)(\mathbb{S}^1)$, we shall specialize our general description of Fredenhagen’s universal category from the end of Section 6.2 to our example at hand. Concretely, an object $(V, \xi^V) \in J_!(\mathfrak{A}^G)(\mathbb{S}^1)$ consists of the following data:

- (1) For each tuple $\underline{I} = (I_1, \dots, I_n) \in \mathcal{J}^{\otimes}/(\mathbb{S}^1)$ of mutually disjoint intervals, a G^n -equivariant module

$$V_{\underline{I}} \in G^n\text{-}\mathbf{Mod}_{\underline{\mathfrak{A}}(\underline{I})} \quad (6.24)$$

over the tensor product algebra $\underline{\mathfrak{A}}(\underline{I}) = \bigotimes_{i=1}^n \mathfrak{A}(I_i)$. (The G^n -action on the tensor product algebra is given by the component-wise G -actions.)

- (2) For each morphism $\alpha : \underline{I} = (I_1, \dots, I_n) \rightarrow \underline{J} = (J_1, \dots, J_m)$ in $J^\otimes/(\mathbb{S}^1)$, a G^n - $\mathbf{Mod}_{\underline{\mathfrak{A}}(\underline{I})}$ -isomorphism

$$\xi_\alpha^V : (\mathfrak{A}^G)^{\mathbf{R}}(\alpha)(V_{\underline{J}}) \longrightarrow V_{\underline{I}} \quad . \quad (6.25)$$

Here $(\mathfrak{A}^G)^{\mathbf{R}}(\alpha) : G^m\text{-}\mathbf{Mod}_{\underline{\mathfrak{A}}(\underline{J})} \rightarrow G^n\text{-}\mathbf{Mod}_{\underline{\mathfrak{A}}(\underline{I})}$ is the right adjoint of the functor

$$\begin{array}{ccc} G^m\text{-}\mathbf{Mod}_{\underline{\mathfrak{A}}(\underline{I})} & \xrightarrow{\mathfrak{A}^G(\alpha)} & G^m\text{-}\mathbf{Mod}_{\underline{\mathfrak{A}}(\underline{J})} \\ & \searrow \Delta_\alpha^* & \nearrow \underline{\mathfrak{A}}(\alpha)_! \\ & G^m\text{-}\mathbf{Mod}_{\Delta_\alpha^*(\underline{\mathfrak{A}}(\underline{I}))} & \end{array} \quad (6.26)$$

where $\Delta_\alpha : G^m \rightarrow G^n$, $(g_1, \dots, g_m) \mapsto (g_{\alpha(1)}, \dots, g_{\alpha(n)})$ is the group map determined by $\alpha : \{1, \dots, n\} \rightarrow \{1, \dots, m\}$, $\Delta_\alpha^* : \mathbf{Rep}_{\mathbb{K}}(G^n) \rightarrow \mathbf{Rep}_{\mathbb{K}}(G^m)$ denotes the corresponding restricted representation functor and $\underline{\mathfrak{A}}(\alpha)_!$ is the induced module functor for the G^m -equivariant algebra morphism $\underline{\mathfrak{A}}(\alpha) : \Delta_\alpha^*(\underline{\mathfrak{A}}(\underline{I})) \rightarrow \underline{\mathfrak{A}}(\underline{J})$. Explicitly, one finds that $(\mathfrak{A}^G)^{\mathbf{R}}(\alpha)$ is given by the composition

$$\begin{array}{ccc} G^m\text{-}\mathbf{Mod}_{\underline{\mathfrak{A}}(\underline{J})} & \xrightarrow{(\mathfrak{A}^G)^{\mathbf{R}}(\alpha)} & G^n\text{-}\mathbf{Mod}_{\underline{\mathfrak{A}}(\underline{I})} \\ \underline{\mathfrak{A}}(\alpha)^* \downarrow & & \uparrow \eta_{\underline{\mathfrak{A}}(\underline{I})}^* \\ G^m\text{-}\mathbf{Mod}_{\Delta_\alpha^*(\underline{\mathfrak{A}}(\underline{I}))} & \xrightarrow{\Delta_{\alpha*}} & G^n\text{-}\mathbf{Mod}_{\Delta_{\alpha*}\Delta_\alpha^*(\underline{\mathfrak{A}}(\underline{I}))} \end{array} \quad (6.27)$$

where η denotes the unit of the adjunction $\Delta_\alpha^* : \mathbf{Rep}_{\mathbb{K}}(G^n) \rightleftarrows \mathbf{Rep}_{\mathbb{K}}(G^m) : \Delta_{\alpha*}$.

These data have to satisfy the coherence conditions (6.7).

Observe from (6.24) that $V_{\underline{I}}$ is a module over the tensor product algebra $\underline{\mathfrak{A}}(\underline{I}) = \bigotimes_{i=1}^n \mathfrak{A}(I_i)$ associated to a tuple of mutually disjoint intervals together with a *separate* G -action for each *connected component*. In other words, the group G is allowed to act differently on different intervals, which is a characteristic feature of a *local* gauge symmetry. To understand better the coherence maps (6.25), let us consider the case where we include two intervals into a single bigger interval, i.e. $\alpha : \underline{I} = (I_1, I_2) \rightarrow J$. In this case $\Delta_\alpha = \Delta : G \rightarrow G^2$ is the diagonal map and (6.25) is given by a G^2 - $\mathbf{Mod}_{\underline{\mathfrak{A}}(\underline{I})}$ -isomorphism

$$\xi_\alpha^V : \eta_{\underline{\mathfrak{A}}(\underline{I})}^* \Delta_* \underline{\mathfrak{A}}(\alpha)^*(V_J) \longrightarrow V_{\underline{I}} \quad . \quad (6.28)$$

Using as in Example 6.4 that $\Delta_* \cong \Delta_! : \mathbf{Rep}_{\mathbb{K}}(G) \rightarrow \mathbf{Rep}_{\mathbb{K}}(G^2)$ is naturally isomorphic to the induced representation functor, we obtain that ξ_α^V is completely determined by a \mathbb{K} -linear map $\kappa_\alpha^V : V_J \rightarrow V_{\underline{I}}$ via $\xi_\alpha^V(1 \otimes 1 \otimes v) = \kappa_\alpha^V(v)$, for all $v \in V_J$. This \mathbb{K} -linear map has to satisfy the following conditions: 1.) G -equivariance: $\kappa_\alpha^V(gv) = (g, g) \kappa_\alpha^V(v)$, for all $v \in V_J$ and $g \in G$. 2.) Preservation of the $\underline{\mathfrak{A}}(\underline{I})$ -actions:

$$\kappa_\alpha^V(v) \cdot (a_1 \otimes a_2) = \sum_{(g_1, g_2) \in G^2} (g_1^{-1}, g_2^{-1}) \kappa_\alpha^V \left(v \cdot \left(\underline{\mathfrak{A}}(\iota_{I_1}^J)(g_1 a_1) \underline{\mathfrak{A}}(\iota_{I_2}^J)(g_2 a_2) \right) \right) \quad , \quad (6.29)$$

for all $a_1 \otimes a_2 \in \mathfrak{A}(I_1) \otimes \mathfrak{A}(I_2)$ and $v \in V_J$, where $\iota_{I_i}^J : I_i \rightarrow J$ denote the interval inclusions. (The sum over G^2 comes from the unit η of the adjunction $\Delta^* \dashv \Delta_*$ when we use $\Delta_!$ as a model for Δ_* .) Comparing (6.29) with the truncated case from Example 6.3, we observe that there is a component-wise G^2 -action on the algebra element $a_1 \otimes a_2 \in \mathfrak{A}(I_1) \otimes \mathfrak{A}(I_2)$ on a pair of intervals before it acts on the module element $v \in V_J$ on the single bigger interval. From a superficial point of view, this behavior resembles the twisted representations of G -equivariant AQFTs by M\"uger [Mug05]. Unfortunately, we do not understand at the moment if there exists a precise relationship between Fredenhagen's universal category $J_!(\mathfrak{A}^G)(\mathbb{S}^1)$ for categorified orbifold theories and the results in [Mug05]. ∇

Acknowledgments

We would like to thank Simen Bruinsma, Chris Fewster, Ignacio Lopez Franco, Klaus Fredenhagen, Owen Gwilliam, Theo Johnson-Freyd and Robert Laugwitz for useful discussions related to this work. M.B. gratefully acknowledges the financial support of the National Group of Mathematical Physics GNFM-INdAM (Italy). M.P. is supported by a PhD scholarship of the Royal Society (UK). A.S. gratefully acknowledges the financial support of the Royal Society (UK) through a Royal Society University Research Fellowship (UF150099), a Research Grant (RG160517) and two Enhancement Awards (RGF\EA\180270 and RGF\EA\201051). L.W. is supported by the RTG 1670 “Mathematics inspired by String Theory and Quantum Field Theory”.

A Basic theory of Cat-enriched colored operads

The aim of this appendix is to set up a suitable framework for **Cat**-enriched colored operads (which one could also call 2-operads or 2-multicategories) that will be used in this work. Our definitions of pseudo-morphisms, pseudo-transformations and modifications are a relatively straightforward generalization of the analogous concepts from 2-category theory (see e.g. [Lei04, Lac10, SP09]) to the theory of colored operads (see e.g. [Yau16, BSW17]). We would like to note that our approach is slightly more flexible than the earlier one by Corner and Gurski [CG13], because we allow our pseudo-morphisms to preserve permutation actions only up to coherent isomorphisms. This generalization is necessary to capture the quantum field theoretical examples that we study in this work. See also Remark A.3 for precise comment on the relationship to [CG13].

Definition A.1. A *Cat-enriched colored operad* \mathcal{O} consists of the following data:

- (1) A collection \mathcal{O}_0 . Elements are called objects and are denoted by symbols like $a, b, c \in \mathcal{O}$.
- (2) Categories $\mathcal{O}(\underline{c})^t$, for each $t \in \mathcal{O}$ and each tuple $\underline{c} := (c_1, \dots, c_n) \in \mathcal{O}^n$. Objects of $\mathcal{O}(\underline{c})^t$ are called 1-operations and are denoted by symbols like ϕ, ψ . Morphisms of $\mathcal{O}(\underline{c})^t$ are called 2-operations and are denoted by symbols like α, β . We write Id for the identity 2-operations and $\alpha \beta$ for the (vertical) composition of 2-operations.
- (3) Composition functors $\gamma : \mathcal{O}(\underline{a})^t \times \prod_{i=1}^n \mathcal{O}(\underline{b}_i)^{a_i} \rightarrow \mathcal{O}(\underline{b})^t$, for each $t \in \mathcal{O}$, $\underline{a} \in \mathcal{O}^n$ and $\underline{b}_i \in \mathcal{O}^{k_i}$, for $i = 1, \dots, n$, where $\underline{b} := (\underline{b}_1, \dots, \underline{b}_n)$. We write $\phi \underline{\psi} := \gamma(\phi, (\psi_1, \dots, \psi_n))$ for the composition of 1-operations and $\alpha * \underline{\beta} := \gamma(\alpha, (\beta_1, \dots, \beta_n))$ for the (horizontal) composition of 2-operations.
- (4) Functors $\mathbf{1} : \mathbf{1} \rightarrow \mathcal{O}(\underline{c})^t$, for each $t \in \mathcal{O}$, where $\mathbf{1}$ is the category with only one object and its identity morphism. We also write $\mathbf{1} \in \mathcal{O}(\underline{c})^t$ for the corresponding identity 1-operation.
- (5) Permutation functors $\mathcal{O}(\sigma) : \mathcal{O}(\underline{c})^t \rightarrow \mathcal{O}(\underline{c\sigma})^t$, for each $t \in \mathcal{O}$, $\underline{c} \in \mathcal{O}^n$ and permutation $\sigma \in \Sigma_n$, where $\underline{c\sigma} := (c_{\sigma(1)}, \dots, c_{\sigma(n)})$. We write $\phi \cdot \sigma := \mathcal{O}(\sigma)(\phi)$ and $\alpha \cdot \sigma := \mathcal{O}(\sigma)(\alpha)$ for the permutation action on 1- and 2-operations.

These data are required to satisfy the usual permutation action, associativity, unitality and equivariance axioms, see e.g. [Yau16, Definition 11.2.1].

Definition A.2. Let \mathcal{O} and \mathcal{P} be **Cat**-enriched colored operads. A *pseudo-morphism* $F : \mathcal{O} \rightarrow \mathcal{P}$ consists of the following data:

- (1) A function $F : \mathcal{O}_0 \rightarrow \mathcal{P}_0$.
- (2) Functors $F : \mathcal{O}(\underline{c})^t \rightarrow \mathcal{P}(\underline{F\underline{c}})^{Ft}$, for each $t \in \mathcal{O}$ and $\underline{c} \in \mathcal{O}^n$, where $F\underline{c} := (F c_1, \dots, F c_n)$.

(3) Natural isomorphisms

$$\begin{array}{ccc}
\mathcal{O}(\underline{t}) \times \prod_{i=1}^n \mathcal{O}(\underline{b}_i) & \xrightarrow{F \times \prod_i F} & \mathcal{P}(\underline{Ft}) \times \prod_{i=1}^n \mathcal{P}(\underline{Fb}_i) \\
\gamma^\sigma \downarrow & \swarrow F^2 & \downarrow \gamma^{\mathcal{P}} \\
\mathcal{O}(\underline{t}) & \xrightarrow{F} & \mathcal{P}(\underline{Ft})
\end{array} \tag{A.1}$$

for each $t \in \mathcal{O}$, $\underline{a} \in \mathcal{O}^n$ and $\underline{b}_i \in \mathcal{O}^{k_i}$, for $i = 1, \dots, n$.

(4) Natural isomorphisms

$$\begin{array}{ccc}
\mathbf{1} & & \\
\mathbf{1}^\sigma \downarrow & \swarrow \mathbf{1}^{\mathcal{P}} & \\
\mathcal{O}(\underline{t}) & \xrightarrow{F} & \mathcal{P}(\underline{Ft})
\end{array} \tag{A.2}$$

for each $t \in \mathcal{O}$.

(5) Natural isomorphisms

$$\begin{array}{ccc}
\mathcal{O}(\underline{t}) & \xrightarrow{F} & \mathcal{P}(\underline{Ft}) \\
\mathcal{O}(\sigma) \downarrow & \swarrow F^\sigma & \downarrow \mathcal{P}(\sigma) \\
\mathcal{O}(\underline{t}) & \xrightarrow{F} & \mathcal{P}(\underline{Ft})
\end{array} \tag{A.3}$$

for each $t \in \mathcal{O}$, $\underline{c} \in \mathcal{O}^n$ and $\sigma \in \Sigma_n$.

These data are required to satisfy the following axioms:

$$\begin{array}{ccc}
(F\phi) (F\underline{\psi}) (F\underline{\rho}) & \xrightarrow{F^2 * \prod \text{Id}} & F(\phi \underline{\psi}) (F\underline{\rho}) \\
\text{Id} * \prod F^2 \downarrow & & \downarrow F^2 \\
(F\phi) F(\underline{\psi} \underline{\rho}) & \xrightarrow{F^2} & F(\phi \underline{\psi} \underline{\rho})
\end{array} \tag{A.4a}$$

$$\begin{array}{ccc}
\mathbf{1}^{\mathcal{P}} (F\phi) & & (F\phi) \prod \mathbf{1}^{\mathcal{P}} \\
F^0 * \text{Id} \downarrow & \searrow \text{Id} & \downarrow \text{Id} \\
(F\mathbf{1}^\sigma) (F\phi) & \xrightarrow{F^2} & F(\mathbf{1}^\sigma \phi) \\
\text{Id} * \prod F^0 \downarrow & & \downarrow \text{Id} \\
(F\phi) \prod F\mathbf{1}^\sigma & \xrightarrow{F^2} & F(\phi \prod \mathbf{1}^\sigma)
\end{array} \tag{A.4b}$$

$$\begin{array}{ccc}
((F\phi) \cdot \sigma) \cdot \sigma' & \xrightarrow{F^\sigma \cdot \sigma'} & (F(\phi \cdot \sigma)) \cdot \sigma' \\
\text{Id} \downarrow & & \downarrow F^{\sigma'} \\
(F\phi) \cdot (\sigma\sigma') & \xrightarrow{F^{\sigma\sigma'}} & F(\phi \cdot (\sigma\sigma'))
\end{array} \tag{A.4c}$$

$$\begin{array}{ccc}
F\phi & & \\
\text{Id} \downarrow & \searrow \text{Id} & \\
(F\phi) \cdot e & \xrightarrow{F^e} & F(\phi \cdot e)
\end{array}$$

$$\begin{array}{ccc}
((F\phi)(F\underline{\psi})) \cdot \sigma \langle k_1, \dots, k_n \rangle & \xrightarrow{F^2 \cdot \sigma \langle k_1, \dots, k_n \rangle} & (F(\phi \underline{\psi})) \cdot \sigma \langle k_1, \dots, k_n \rangle \\
\text{Id} \downarrow & & \downarrow_{F^\sigma \langle k_1, \dots, k_n \rangle} \\
((F\phi) \cdot \sigma)(F\underline{\psi}\sigma) & & F((\phi \underline{\psi}) \cdot \sigma \langle k_1, \dots, k_n \rangle) \\
F^\sigma * \prod \text{Id} \downarrow & & \downarrow \text{Id} \\
F(\phi \cdot \sigma)(F\underline{\psi}\sigma) & \xrightarrow{F^2} & F((\phi \cdot \sigma)(\underline{\psi}\sigma))
\end{array} \tag{A.4d}$$

$$\begin{array}{ccc}
((F\phi)(F\underline{\psi})) \cdot (\sigma_1 \oplus \dots \oplus \sigma_n) & \xrightarrow{F^2 \cdot (\sigma_1 \oplus \dots \oplus \sigma_n)} & (F(\phi \underline{\psi})) \cdot (\sigma_1 \oplus \dots \oplus \sigma_n) \\
\text{Id} \downarrow & & \downarrow_{F^{\sigma_1 \oplus \dots \oplus \sigma_n}} \\
(F\phi)((F\underline{\psi}) \cdot (\sigma_1 \oplus \dots \oplus \sigma_n)) & & F((\phi \underline{\psi}) \cdot (\sigma_1 \oplus \dots \oplus \sigma_n)) \\
\text{Id} * \prod F^{\sigma_i} \downarrow & & \downarrow \text{Id} \\
(F\phi)(F(\underline{\psi} \cdot (\sigma_1 \oplus \dots \oplus \sigma_n))) & \xrightarrow{F^2} & F(\phi(\underline{\psi} \cdot (\sigma_1 \oplus \dots \oplus \sigma_n)))
\end{array} \tag{A.4e}$$

Remark A.3. In the case all coherences F^σ for permutation actions in (A.3) are identities, our concept of pseudo-morphisms specializes to [CG13, Definition 2.2]. We however require the more flexible Definition A.2 in the present paper, because our quantum field theoretical examples of interest generically come with non-trivial coherences F^σ . \triangle

Definition A.4. Let \mathcal{O} and \mathcal{P} be **Cat**-enriched colored operads and $F, G : \mathcal{O} \rightarrow \mathcal{P}$ pseudo-morphisms. A *pseudo-transformation* $\zeta : F \Rightarrow G$ consists of the following data:

- (1) Functors $\zeta_c : \mathbf{1} \rightarrow \mathcal{P}(\frac{G_c}{F_c})$, for each $c \in \mathcal{O}$. We also write $\zeta_c \in \mathcal{P}(\frac{G_c}{F_c})$ for the corresponding 1-operation.
- (2) Natural isomorphisms

$$\begin{array}{ccc}
\mathcal{O}(\underline{c}) \times \prod_{i=1}^n \mathbf{1} & \xrightarrow{G \times \prod_i \zeta_{c_i}} & \mathcal{P}(\frac{G_t}{G_c}) \times \prod_{i=1}^n \mathcal{P}(\frac{G_{c_i}}{F_{c_i}}) \\
\cong \downarrow & \nearrow \zeta^\bullet & \downarrow \gamma^{\mathcal{P}} \\
\mathbf{1} \times \mathcal{O}(\underline{c}) & & \\
\zeta_t \times F \downarrow & & \downarrow \gamma^{\mathcal{P}} \\
\mathcal{P}(\frac{G_t}{F_t}) \times \mathcal{P}(\frac{F_t}{F_c}) & \xrightarrow{\gamma^{\mathcal{P}}} & \mathcal{P}(\frac{G_t}{F_c})
\end{array} \tag{A.5}$$

for each $t \in \mathcal{O}$ and $\underline{c} \in \mathcal{O}^n$.

These data are required to satisfy the following axioms:

$$\begin{array}{ccc}
(G\phi)(G\underline{\psi}) \prod \zeta_{b_{ij}} & \xrightarrow{\text{Id} * \prod \zeta^\bullet} & (G\phi) \prod \zeta_{a_i}(F\underline{\psi}) \xrightarrow{\zeta^\bullet * \prod \text{Id}} \zeta_t(F\phi)(F\underline{\psi}) \\
G^2 * \prod \text{Id} \downarrow & & \downarrow \text{Id} * F^2 \\
G(\phi \underline{\psi}) \prod \zeta_{b_{ij}} & \xrightarrow{\zeta^\bullet} & \zeta_t F(\phi \underline{\psi})
\end{array} \tag{A.6a}$$

$$\begin{array}{ccc}
\mathbb{1}^{\mathcal{P}} \zeta_t & \xrightarrow{G^0 * \text{Id}} & (G\mathbb{1}^{\mathcal{O}}) \zeta_t \\
\text{Id} \downarrow & & \downarrow \zeta^\bullet \\
\zeta_t \mathbb{1}^{\mathcal{P}} & \xrightarrow{\text{Id} * F^0} & \zeta_t (F\mathbb{1}^{\mathcal{O}})
\end{array} \tag{A.6b}$$

$$\begin{array}{ccc}
((G\phi) \prod \zeta_{c_i}) \cdot \sigma & \xrightarrow{\zeta^\bullet \cdot \sigma} & (\zeta_t (F\phi)) \cdot \sigma \\
\text{Id} \downarrow & & \downarrow \text{Id} \\
((G\phi) \cdot \sigma) \prod \zeta_{c_{\sigma(i)}} & & \zeta_t ((F\phi) \cdot \sigma) \\
G^\sigma * \prod \text{Id} \downarrow & & \downarrow \text{Id} * F^\sigma \\
G(\phi \cdot \sigma) \prod \zeta_{c_{\sigma(i)}} & \xrightarrow{\zeta^\bullet} & \zeta_t F(\phi \cdot \sigma)
\end{array} \tag{A.6c}$$

Definition A.5. Let \mathcal{O} and \mathcal{P} be **Cat**-enriched colored operads, $F, G : \mathcal{O} \rightarrow \mathcal{P}$ pseudo-morphisms and $\zeta, \kappa : F \Rightarrow G$ pseudo-transformations. A *modification* $\Gamma : \zeta \Rrightarrow \kappa$ consists of the following data:

(1) Natural transformations

$$\begin{array}{ccc}
& \zeta_c & \\
& \curvearrowright & \\
\mathbf{1} & \Downarrow \Gamma_c & \mathcal{P} \begin{pmatrix} G_c \\ F_c \end{pmatrix} \\
& \curvearrowleft & \\
& \kappa_c &
\end{array} \tag{A.7}$$

for each $c \in \mathcal{O}$.

These data are required to satisfy the following axioms:

$$\begin{array}{ccc}
(G\phi) \prod \zeta_{c_i} & \xrightarrow{\text{Id} * \prod \Gamma_{c_i}} & (G\phi) \prod \kappa_{c_i} \\
\zeta^\bullet \downarrow & & \downarrow \kappa^\bullet \\
\zeta_t (F\phi) & \xrightarrow{\Gamma_t * \text{Id}} & \kappa_t (F\phi)
\end{array} \tag{A.8}$$

Remark A.6. **Cat**-enriched colored operads, pseudo-morphisms, pseudo-transformations and modifications assemble into a tricategory. The various compositions are similar to the case of the tricategory of bicategories and hence will not be displayed in full detail here. We refer the reader to [SP09, Appendix A.1] for a brief review of the tricategory of bicategories and to [GPS95] for the details.

Let us nevertheless fix the relevant notations that will appear in the bulk of this paper. Given two **Cat**-enriched colored operads \mathcal{O} and \mathcal{P} , the tricategory structure implies that there exists a Hom-2-category

$$\mathbf{Alg}_{\mathcal{O}}(\mathcal{P}) := [\mathcal{O}, \mathcal{P}] \in \mathbf{2Cat} \quad , \tag{A.9}$$

whose objects are pseudo-morphisms from \mathcal{O} to \mathcal{P} , 1-morphisms are pseudo-transformations and 2-morphisms are modifications. Following the usual terminology of operad theory, we shall call $\mathbf{Alg}_{\mathcal{O}}(\mathcal{P})$ the *2-category of \mathcal{O} -algebras with values in \mathcal{P}* . Given pseudo-morphisms $F : \mathcal{O} \rightarrow \mathcal{O}'$ and $G : \mathcal{P} \rightarrow \mathcal{P}'$, there exist pseudo-functors

$$F^* : [\mathcal{O}', \mathcal{P}] \longrightarrow [\mathcal{O}, \mathcal{P}] \quad , \quad G_* : [\mathcal{O}, \mathcal{P}] \longrightarrow [\mathcal{O}, \mathcal{P}'] \quad , \tag{A.10}$$

which we call *pullback* and *pushforward*.

Let us note that in the case \mathcal{O} and \mathcal{P} are **Set**-valued colored operads, i.e. all categories of operations in Definition A.1 are sets, then $\mathbf{Alg}_{\mathcal{O}}(\mathcal{P}) = [\mathcal{O}, \mathcal{P}] \in \mathbf{Cat}$ is an ordinary category that coincides with the usual category of \mathcal{O} -algebras with values in \mathcal{P} , see e.g. [Yau16, BSW17]. \triangle

References

- [AR94] J. Adámek and J. Rosický, *Locally presentable and accessible categories*, London Math. Soc. Lect. Note Series **189**, Cambridge University Press, Cambridge (1994).
- [AF15] D. Ayala and J. Francis, “Factorization homology of topological manifolds,” *J. Topol.* **8**, no. 4, 1045–1084 (2015) [arXiv:1206.5522 [math.AT]].
- [BBS19] M. Benini, S. Bruinsma and A. Schenkel, “Linear Yang-Mills theory as a homotopy AQFT,” *to appear in Communications in Mathematical Physics* [arXiv:1906.00999 [math-ph]].
- [BS19] M. Benini and A. Schenkel, “Higher structures in algebraic quantum field theory,” *Fortsch. Phys.* **67**, no. 8-9, 1910015 (2019) [arXiv:1903.02878 [hep-th]].
- [BSW17] M. Benini, A. Schenkel and L. Woike, “Operads for algebraic quantum field theory,” *to appear in Communications in Contemporary Mathematics* [arXiv:1709.08657 [math-ph]].
- [BSW19] M. Benini, A. Schenkel and L. Woike, “Homotopy theory of algebraic quantum field theories,” *Lett. Math. Phys.* **109**, no. 7, 1487 (2019) [arXiv:1805.08795 [math-ph]].
- [BZBJ18a] D. Ben-Zvi, A. Brochier and D. Jordan, “Integrating quantum groups over surfaces,” *J. Topol.* **11**, no. 4, 874–917 (2018) [arXiv:1501.04652 [math.QA]].
- [BZBJ18b] D. Ben-Zvi, A. Brochier and D. Jordan, “Quantum character varieties and braided module categories,” *Sel. Math. New Ser.* **24**, 4711–4748 (2018) [arXiv:1606.04769 [math.QA]].
- [BZFN10] D. Ben-Zvi, J. Francis and D. Nadler, “Integral transforms and Drinfeld centers in derived algebraic geometry,” *J. Amer. Math. Soc.* **23**, no. 4, 909–966 (2010) [arXiv:0805.0157 [math.AG]].
- [Bra14] M. Brandenburg, *Tensor categorical foundations of algebraic geometry*, PhD thesis, University of Münster (2014) [arXiv:1410.1716v1 [math.AG]].
- [BCJF15] M. Brandenburg, A. Chirvasitu and T. Johnson-Freyd, “Reflexivity and dualizability in categorified linear algebra,” *Theory Appl. Categ.* **30**, 808–835 (2015) [arXiv:1409.5934 [math.CT]].
- [BFV03] R. Brunetti, K. Fredenhagen and R. Verch, “The generally covariant locality principle: A new paradigm for local quantum field theory,” *Commun. Math. Phys.* **237**, 31 (2003) [math-ph/0112041].
- [Brz14] T. Brzeziński, “On the Smoothness of the Noncommutative Pillow and Quantum Teardrops,” *SIGMA* **10**, 015 (2014) [arXiv:1311.4758 [math.QA]].
- [BS17] T. Brzeziński and A. Sitarz, “Smooth geometry of the noncommutative pillow, cones and lens spaces,” *J. Noncommut. Geom.* **11**, 413–449 (2017) [arXiv:1410.6587 [math.QA]].
- [CPTVV17] D. Calaque, T. Pantev, B. Toën, M. Vaquié and G. Vezzosi, “Shifted Poisson structures and deformation quantization,” *J. Topol.* **10**, no. 2, 483 (2017) [arXiv:1506.03699 [math.AG]].
- [CG13] A. S. Corner and N. Gurski, “Operads with general groups of equivariance, and some 2-categorical aspects of operads in *Cat*,” arXiv:1312.5910 [math.CT].
- [CG17] K. Costello and O. Gwilliam, *Factorization algebras in quantum field theory*, New Mathematical Monographs **31**, Cambridge University Press, Cambridge (2017).

- [DVVV89] R. Dijkgraaf, C. Vafa, E. P. Verlinde and H. L. Verlinde, “The operator algebra of orbifold models,” *Commun. Math. Phys.* **123**, 485 (1989).
- [DT89] Y. Doi and M. Takeuchi, “Hopf-Galois extensions of algebras, the Miyashita-Ulbrich action, and Azumaya algebras,” *J. Algebra* **121**, 488–516 (1989).
- [DSPS13] C. L. Douglas, C. Schommer-Pries and N. Snyder, “Dualizable tensor categories,” *to appear in Memoirs of the AMS* [arXiv:1312.7188 [math.QA]].
- [EM09] A.D. Elmendorf and M. A. Mandell, “Permutative categories, multicategories and algebraic K -theory,” *Algebr. Geom. Topol.* **9**, no. 4, 2391–2441 (2009) [arXiv:0710.0082 [math.KT]].
- [FV15] C. J. Fewster and R. Verch, “Algebraic quantum field theory in curved spacetimes,” in: R. Brunetti, C. Dappiaggi, K. Fredenhagen and J. Yngvason (eds.), *Advances in algebraic quantum field theory*, 125–189, Springer Verlag, Heidelberg (2015) [arXiv:1504.00586 [math-ph]].
- [Fre90] K. Fredenhagen, “Generalizations of the theory of superselection sectors,” in: D. Kastler (ed.), *The algebraic theory of superselection sectors: Introduction and recent results*, World Scientific Publishing, 379 (1990).
- [Fre93] K. Fredenhagen, “Global observables in local quantum physics,” in: H. Araki, K. R. Ito, A. Kishimoto and I. Ojima (eds.), *Quantum and non-commutative analysis: Past, present and future perspectives*, 41–51, Kluwer Academic Publishers (1993).
- [FRS92] K. Fredenhagen, K.-H. Rehren and B. Schroer, “Superselection sectors with braid group statistics and exchange algebras II: Geometric aspects and conformal covariance,” *Rev. Math. Phys.* **4**, 113 (1992).
- [FR12] K. Fredenhagen and K. Rejzner, “Batalin-Vilkovisky formalism in the functional approach to classical field theory,” *Commun. Math. Phys.* **314**, 93 (2012) [arXiv:1101.5112 [math-ph]].
- [FR13] K. Fredenhagen and K. Rejzner, “Batalin-Vilkovisky formalism in perturbative algebraic quantum field theory,” *Commun. Math. Phys.* **317**, 697 (2013) [arXiv:1110.5232 [math-ph]].
- [GPS95] R. Gordon, A. J. Power and R. Street, “Coherence for tricategories,” *Mem. Amer. Math. Soc.* **117**, no. 558 (1995).
- [HK64] R. Haag and D. Kastler, “An algebraic approach to quantum field theory,” *J. Math. Phys.* **5**, 848 (1964).
- [Hor17] G. Horel, “Factorization homology and calculus à la Kontsevich-Soibelman,” *J. Noncommut. Geom.* **11**, 703 (2017) [arXiv:1307.0322 [math.AT]].
- [JF15] T. Johnson-Freyd, “Heisenberg–picture quantum field theory,” *to appear in Progress in Mathematics* [arXiv:1508.05908 [math-ph]].
- [Kaw15] Y. Kawahigashi, “Conformal field theory, tensor categories and operator algebras,” *J. Phys. A* **48**, no. 30, 303001 (2015) [arXiv:1503.05675 [math-ph]].
- [Kel05] G. M. Kelly, “Basic concepts of enriched category theory,” *Repr. Theory Appl. Categ.* No. **10** (2005).
- [Lac10] S. Lack, “A 2-categories companion,” in: J. C. Baez and J. P. May (eds.), *Towards higher categories*, IMA Vol. Math. Appl. **152**, 105–191, Springer, New York (2010) [arXiv:math/0702535 [math.CT]].

- [Lan14] B. Lang, *Universal constructions in algebraic and locally covariant quantum field theory*, PhD thesis, University of York (2014).
- [Lei04] T. Leinster, *Higher operads, higher categories*, London Math. Soc. Lect. Note Series **298**, Cambridge University Press, Cambridge (2004).
- [LN16] F. Lucatelli Nunes, “On biadjoint triangles,” *Theory Appl. Categ.* **31**, no. 9, 217–256 (2016) [arXiv:1606.05009 [math.CT]].
- [Lur04] J. Lurie, “Tannaka duality for geometric stacks,” arXiv:math/0412266 [math.AG].
- [Mon09] S. Montgomery, “Hopf Galois theory: A survey,” *Geometry & Topology Monographs* **16**, 367–400 (2009).
- [Mug05] M. Müger, “Conformal orbifold theories and braided crossed G -categories,” *Commun. Math. Phys.* **260**, 727 (2005) Erratum: [*Commun. Math. Phys.* **260**, 763 (2005)] [math/0403322 [math.QA]].
- [PTVV13] T. Pantev, B. Toën, M. Vaquié and G. Vezzosi, “Shifted symplectic structures,” *Publ. Math. Inst. Hautes Études Sci.* **117**, 271 (2013) [arXiv:1111.3209 [math.AG]].
- [SP09] C. Schommer-Pries, *The classification of two-dimensional extended topological field theories*, PhD thesis, UC Berkeley (2009) [arXiv:1112.1000 [math.AT]].
- [Str80] R. Street, “Fibrations in bicategories,” *Cahiers Topologie Géom. Différentielle* **21**, no. 2, 111–160 (1980).
- [Str87] R. Street, “Correction to: “Fibrations in bicategories” [*Cahiers Topologie Géom. Différentielle* **21**, no. 2, 111–160 (1980)],” *Cahiers Géom. Différentielle Catég.* **28**, no. 1, 53–56 (1987).
- [Toe06] B. Toën, “Affine stacks (Champs affines),” *Selecta Math. (N.S.)* **12**, no. 1, 39–135 (2006) [arXiv:math/0012219 [math.AG]].
- [Toe14] B. Toën, “Derived algebraic geometry and deformation quantization,” *Proceedings of the International Congress of Mathematicians, Seoul* (2014) [arXiv:1403.6995 [math.AG]].
- [Xu00] F. Xu, “Algebraic orbifold conformal field theories,” *Proc. Natl. Acad. Sci. USA* **97**, no. 26, 14069–14073 (2000) [arXiv:math/0004150 [math.QA]].
- [Yau16] D. Yau, *Colored operads*, Graduate Studies in Mathematics **170**, American Mathematical Society, Providence, RI (2016).