

Dynamical Emergence of Scalaron in Higgs Inflation

Yohei Ema

DESY, Notkestraße 85, D-22607 Hamburg, Germany

Abstract

We point out that a light scalaron dynamically emerges if scalar fields have a sizable non-minimal coupling to the Ricci scalar as in the Higgs inflation model. We support this claim in two ways. One is based on the renormalization group equation; the non-minimal coupling inevitably induces a Ricci scalar quadratic term due to the renormalization group running. The other is based on scattering amplitudes; a scalar four-point amplitude develops a pole after summing over a certain class of diagrams, which we identify as the scalaron. Our result implies that the Higgs inflation is actually a two-field inflationary model. Another implication is that the Higgs inflation does not suffer from the unitarity issue since the scalaron pushes up the cut-off scale to the Planck scale.

Contents

1	Introduction	2
2	Renormalization group equation and scalaron	3
3	Self-healing mechanism and scalaron	5
3.1	Unitarity of Higgs inflation	5
3.2	Self-healing mechanism	6
3.3	Self-healed amplitude as scalaron emergence	8
4	Equivalence of Jordan and Einstein frames	10
4.1	Jordan frame	11
4.2	Einstein frame	12
5	Single field case	13
6	Summary and discussions	15
A	Computational details	17
A.1	Preliminary	18
A.2	Feynman rules	19
A.3	Tree-level amplitude	20
A.4	Scalar one-loop amplitude	21
A.5	Renormalization group equation	22
B	Higgs scalaron inflationary model	23
B.1	Einstein frame action and unitarity	23
B.2	Inflationary predictions	24
B.3	Feynman rules around the vacuum	25

1 Introduction

Inflation plays an essential role in the modern cosmology. It not only solves the initial condition problems such as the flatness and horizon problems, but also provides seeds for the large scale structure and the anisotropy of the cosmic microwave background (CMB) in the present universe.

Among many candidates, the Higgs inflation model [1–3] is probably one of the most popular inflationary models because of its minimality and consistency with the CMB observation [4]. It identifies the (radial component of the) standard model Higgs doublet as the inflaton. The original Higgs quartic potential needs to be modified at the large field value region to be consistent with the CMB, and hence the following interaction

$$\mathcal{L}_\xi = \xi |H|^2 R, \quad (1.1)$$

is introduced, where ξ is a non-minimal coupling between the Higgs doublet H and the Ricci scalar R . The CMB normalization requires ξ and the Higgs quartic coupling λ to satisfy $\xi^2 \simeq 2 \times 10^9 \lambda$, indicating that $\xi \sim \mathcal{O}(10^4)$ unless λ is tiny. Such a large ξ triggers a lot of discussions, most importantly concerning the unitarity issue of the Higgs inflation [5–9]. In the Higgs inflation, the tree-level unitarity is violated at the energy scale of $\mathcal{O}(M_P/\xi)$ around the vacuum (where the Higgs field value vanishes). It does not necessarily spoil the inflationary prediction since the scale of the violation depends on the Higgs field value and is larger during inflation [10]. Nevertheless, several issues, such as possible UV completion of the Higgs inflation [11–14] or violent phenomena caused by the large non-minimal coupling after inflation [15–17], have been discussed so far.

In this paper, we study consequences of the non-minimal coupling one step further, assuming that $\xi \gg 1$. In particular, we see that a scalaron degree of freedom (i.e. the spin-0 part of the metric) becomes dynamical once we properly take into account quantum effects. Our argument is based on two distinct (yet closely related) observations. First, we study the renormalization group (RG) equation of this model. It is well-known that, once we treat the gravity as an effective field theory, we have to introduce the Ricci scalar quadratic term,

$$\mathcal{L}_{\alpha_1} = \alpha_1 R^2, \quad (1.2)$$

as a counter term to renormalize divergences at the one-loop level [18, 19]. It results in the RG running of α_1 . The scalar loop contribution to the beta function is given by

$$\beta_{\alpha_1} \equiv \frac{d\alpha_1}{d \ln \mu} = -\frac{N_s}{1152\pi^2} (1 + 6\xi)^2, \quad (1.3)$$

where N_s is the number of real scalar fields ($N_s = 4$ in the Higgs inflation). It means that we cannot keep $|\alpha_1| \lesssim \mathcal{O}(1)$ for all energy scales for $\xi \gg 1$. Since α_1 is inversely proportional to the scalaron mass squared, a light scalaron inevitably shows up in the theory. This line of argument is also emphasized in Refs. [20–23]. Second, and more interestingly, we see that the scalaron dynamically emerges as an intermediate state in 2-to-2 scattering amplitudes.^{†1} If the above argument based on the RG equation is true, the unitarity issue of the Higgs inflation has to be gone after including quantum effects. This is because, once the scalaron exists, the cut-off scale of the theory is pushed up to the Planck scale [22, 24]. Such a mechanism of healing the unitarity is indeed discussed in Refs. [25, 26], called the self-healing mechanism. We show that the

^{†1} In this paper, we use the word “dynamical” to indicate that the scalaron emerges after resumming over a certain class of Feynman diagrams (see Eqs. (3.5) and (4.16)).

self-healed scattering amplitude is equivalent to an amplitude with the scalaron propagating as an intermediate state. It strongly supports that the scalaron dynamically emerges once we take into account quantum effects properly.

This paper is organized as follows. In Sec. 2, we argue that the scalaron inevitably shows up in the theory based on the RG equation. It also provides some notations used in this paper. In Sec. 3, we study a 2-to-2 scattering amplitude between the scalar fields. We first review the unitarity issue of the Higgs inflation based on the scattering amplitude, and the self-healing mechanism. We then move to our main point, identifying the self-healing mechanism as the dynamical emergence of the scalaron. Sec. 4 shows the equivalence of our results between the Jordan and Einstein frames. We consider only the case $N_s \geq 2$ in Secs. 2–4. The case $N_s = 1$ is tricky and is separately discussed in Sec. 5. Note that the subtlety discussed there is irrelevant for the Higgs inflation that has $N_s = 4$. Finally Sec. 6 is devoted to summary and discussions. The appendix is composed of two parts. In App. A, we provide some computational details for the sake of clarity and completeness. We review the Higgs scalaron inflationary model, i.e. an inflationary model with the Higgs and the scalaron, in App. B.

2 Renormalization group equation and scalaron

In this paper, we consider the following action:

$$S = \int d^4x \sqrt{-g} \left[\frac{M_P^2}{2} R + \frac{1}{2} g^{\mu\nu} \partial_\mu \phi_i \partial_\nu \phi_i + \frac{\xi}{2} R \phi_i^2 - \frac{\lambda \phi_i^4}{4} \right] + S_{\text{c.t.}}, \quad (2.1)$$

where M_P is the (reduced) Planck scale, ϕ_i are the real scalar fields, $i = 1, 2, \dots, N_s$, $g_{\mu\nu}$ is the metric with g its determinant, R is the Ricci scalar, and ξ and λ are the coupling constants. We ignore the electroweak scale throughout this paper. It is justified as the energy scale of our interest is significantly higher than the electroweak scale. We focus on the case $N_s \geq 2$ in this section, Secs. 3 and 4. The Higgs inflation corresponds to $N_s = 4$.^{‡2} The case $N_s = 1$ is tricky and is discussed separately in Sec. 5. The last term $S_{\text{c.t.}}$ is the counter term required to renormalize this theory. It plays an essential role in the following.

The inflationary prediction of this model (without the counter term) is studied in detail in literature, especially in the case that ϕ_i are (the components of) the standard model Higgs doublet. It reproduces the normalization of the CMB anisotropy when

$$\xi^2 \simeq 2 \times 10^9 \lambda. \quad (2.2)$$

It requires that $\xi \gg 1$ unless λ is tiny, and hence we assume that $\xi \gg 1$ throughout this paper.^{‡3} The spectral index is consistent with the current CMB observation [4], and the tensor-to-scalar ratio is within the reach of the future observations [29]. Because of these features and its minimality, the Higgs inflation is one among the most popular inflationary models so far.

The inflationary prediction is usually studied at the classical level.^{‡4} Once we consider quantum effects, however, we have to take into account counter terms. Since this theory contains the gravity,

^{‡2} As far as the beta functions or scattering amplitudes around the vacuum at high energy scale are concerned, it is appropriate to treat the Higgs doublet as four real scalar fields.

^{‡3} The case with tiny λ is called the critical Higgs inflation model [27, 28]. Our discussion in the following does not apply to this case as long as $\xi \lesssim \mathcal{O}(1)$.

^{‡4} See Refs. [30–39] and references therein for quantum effects on the Higgs potential and the non-minimal coupling. They are also studied based on the exact RG group approach in Ref. [40].

the following counter terms (among others) are required to cancel divergences at the one-loop level [18]:

$$S_{\text{c.t.}} = \int d^4x \sqrt{-g} \left[\alpha_1 R^2 + \alpha_2 \left(R_{\mu\nu} R^{\mu\nu} - \frac{1}{3} R^2 \right) \right]. \quad (2.3)$$

As a result, the coefficients α_1 and α_2 run according to the energy scale of the system. The (scalar loop contributions to the) beta functions are given by^{h5}

$$\beta_{\alpha_1} \equiv \frac{d\alpha_1}{d \ln \mu} = -\frac{N_s}{1152\pi^2} (1 + 6\xi)^2, \quad (2.4)$$

$$\beta_{\alpha_2} \equiv \frac{d\alpha_2}{d \ln \mu} = -\frac{N_s}{960\pi^2}. \quad (2.5)$$

See App. A for the derivation. Two important properties are read off. First, the runnings of α_1 and α_2 are additive, not multiplicative. That is, the operators associated with α_1 and α_2 inevitably emerge at other energy scales even if they are absent at one specific energy scale. Second, the size of α_1 is generically of $\mathcal{O}(\xi^2)$ (or larger) that is significantly larger than unity since we assume $\xi \gg 1$. It is well-known that the R^2 operator introduces an additional scalar degree of freedom [45–48], which we call “scalaron” in this paper. Its mass is given by

$$m_s^2 = \frac{M_P^2}{12\alpha_1}, \quad (2.6)$$

which is lighter as α_1 is larger. Thus, the beta function indicates that the Higgs inflation model with the large non-minimal coupling ξ necessarily contains the scalaron as a light degree of freedom.^{h6} This line of argument is also emphasized in Refs. [20–23].

We emphasize that we cannot simply ignore the scalaron. In order to make the theory well-defined at the quantum level, we have to introduce the α_1 -operator and hence the scalaron from the beginning. We can make the scalaron heavy at some specific energy scale by taking α_1 small there, but it soon becomes as light as $\mathcal{O}(M_P/\xi)$ once one goes below that scale.^{h7} In other words, one cannot keep the scalaron heavy for all energy scales. It is in contrast to the case of the α_2 -operator. It is known that the α_2 -operator introduces a spin-2 ghost degree of freedom whose mass squared is [49, 50]

$$m_2^2 = -\frac{M_P^2}{2\alpha_2}. \quad (2.7)$$

It is kept to be of $\mathcal{O}(M_P^2)$ for all scales once we take $|\alpha_2| \lesssim \mathcal{O}(1)$ at some scale, since β_{α_2} does not depend on ξ . Then we might forget about it by, e.g., believing that some UV completion of the gravity cures the pathology associated with the ghost. The same argument does not apply to the scalaron. Since the scalaron mass is in general significantly smaller than the Planck scale for $\xi \gg 1$, it has to be distinguishable from the effects of the UV completion of the gravity.

Although this argument might be already convincing, we will support it from a different point of view in the following, since the statement that the scalaron inevitably exists has a huge impact

^{h5} Eq. (2.4) agrees with Refs. [20, 23, 41–43], while Refs. [21, 44] have an opposite sign. Note that the sign convention of the Ricci tensor does not affect that of α_1 since the latter depends quadratically on the former.

^{h6} Here the word “light” means that it is significantly lighter than the Planck scale.

^{h7} See Sec. 6 for more details on the RG running of the scalaron mass.

on both the theoretical and phenomenological sides of the Higgs inflation. To be specific, in Sec. 3, we consider a 2-to-2 scattering amplitude $\phi_i\phi_i \rightarrow \phi_j\phi_j$ with $i \neq j$. We will see that the scalaron dynamically emerges as an intermediate state in the amplitude even if we do not include it at the beginning. It also reveals that the dynamical emergence of the scalaron is closely related to the unitarity structure of the Higgs inflation.

3 Self-healing mechanism and scalaron

In this section, we consider the 2-to-2 scattering amplitude $\phi_i\phi_i \rightarrow \phi_j\phi_j$ with $i \neq j$ around the vacuum $\phi_i = 0$, and show that a scalaron degree of freedom dynamically emerges as an intermediate state in this process. This phenomena is closely related to the self-healing mechanism discussed in Refs. [25, 26]. Since the self-healing mechanism is studied in the context of the unitarity issue in the Higgs inflation, we first review the latter in Sec. 3.1. In Sec. 3.2, we review the self-healing mechanism. The self-healing mechanism in practice sums over a certain class of Feynman diagrams (see Eqs. (3.5) and (4.16)). As a result, the scattering amplitude develops a pole structure. We see in Sec. 3.3 that it can be identified with the scalaron propagating as an intermediate state.

3.1 Unitarity of Higgs inflation


As we have discussed in Sec. 2, the Higgs inflation model requires $\xi \gg 1$. It triggers huge discussions, especially concerning the unitarity of the Higgs inflation model. Let us consider the tree-level 2-to-2 scattering amplitude $\phi_i\phi_i \rightarrow \phi_j\phi_j$ with $i \neq j$ in the Higgs inflation. It is given as (see App. A for the derivation)

$$A_{\text{tree}}^{(ii \rightarrow jj)} = \frac{1}{M_P^2 s} \left[\frac{(1 + 6\xi)^2}{6} s^2 - \left(\frac{s^2}{6} - tu \right) \right], \quad (3.1)$$

around $\phi_i = 0$, where s, t and u are the standard Mandelstam variables. Here and henceforth we ignore the contribution from the Higgs potential since it is irrelevant for our discussion. It indicates that the cut-off of the theory is $\mathcal{O}(M_P/\xi)$, significantly lower than the Planck scale [5–9]. It does not necessarily spoil the inflationary prediction of this model since the cut-off scale depends on the Higgs field value, and is higher than the typical energy scale during inflation [10]. It still seems to require that the Higgs inflation should be UV completed below the Planck scale. More interestingly, the unitarity could be violated during preheating [15–17], resulting in the necessity of UV completion to describe the reheating dynamics. These are the (very) rough sketch of the unitarity issue on the Higgs inflation in literature so far.

In this paper, our viewpoint on the unitarity of the Higgs inflation is totally different. In Sec. 2, we argue that the scalaron inevitably shows up in the Higgs inflation model due to the quantum effects. If it is true, the above discussion on the unitarity has to be just an illusion, and the unitarity of the Higgs inflation has to be remedied below the Planck scale once quantum effects are properly taken into account. This is because the cut-off scale of the Higgs inflation is pushed up to the Planck scale once the scalaron is included, as first pointed out in Ref. [22] and further studied in Ref. [24] (see also App. B.1). Indeed, such a healing mechanism of the unitarity, or the self-healing mechanism, is discussed in Refs. [25, 26]. In particular, Ref. [26] discusses the self-healing mechanism of the Higgs inflation, concluding that it has no unitarity issue once quantum effects are included. In the following, we shed a new light on the self-healing mechanism; we

μ_2 , respectively. The one-loop contribution is larger than the tree-level one for $\sqrt{s} \gtrsim M_P/\sqrt{N_s}\xi$, and hence we may define a dressed amplitude by summing over the scalar loop diagrams as



$$A_{\text{dressed}}^{(ii \rightarrow jj)} \equiv \text{[Tree-level diagram]} + \text{[One-loop diagram]} + \text{[Two-loop diagram]} + \dots \quad (3.5)$$

Note that it is the leading order contribution of the large N_s expansion, since the loops involving the graviton and ghost (from the gauge fixing) are suppressed by N_s compared to the scalar loops.

Now we study the unitarity structure of this dressed amplitude. We consider an elastic scattering amplitude between flavor-singlet states,

$$|\mathbf{1}\rangle \equiv \frac{1}{\sqrt{N_s}} \sum_{i=1}^{N_s} |\phi_i \phi_i\rangle, \quad (3.6)$$

where $|\phi_i \phi_i\rangle$ denotes the state with two ϕ_i particles. It is given at the leading order in $1/N_s$ by

$$A^{(\mathbf{1} \rightarrow \mathbf{1})} = N_s A_{\text{dressed}}^{(ii \rightarrow jj)}, \quad (3.7)$$

as the diagonal parts are suppressed by $1/N_s$. The factor N_s originates from the combinatorial factor N_s^2 divided by the normalization of the state (the prefactor in Eq. (3.6)). We now perform the partial wave expansion,⁹

$$A^{(\mathbf{1} \rightarrow \mathbf{1})} = 32\pi \sum_l (2l+1) a^{(l)}(s) P_l(\cos\theta), \quad (3.8)$$

where l is the angular momentum, P_l is the Legendre polynomial of degree l with $P_l(1) = 1$, and θ is the scattering angle. The Mandelstam variables t and u are given in terms of s and θ as

$$t = -\frac{s}{2}(1 - \cos\theta), \quad u = -\frac{s}{2}(1 + \cos\theta). \quad (3.9)$$

Since the different spin parts do not mix with each other in the summation (3.5), we obtain the s -wave part of the partial wave amplitude as

$$a^{(0)} = \frac{a_{\text{tree}}^{(0)}}{1 - a_{\text{1-loop}}^{(0)}/a_{\text{tree}}^{(0)}}, \quad (3.10)$$

where the s -wave parts of the tree-level and one-loop amplitudes are respectively given as

$$a_{\text{tree}}^{(0)} = \frac{N_s(1+6\xi)^2}{192\pi M_P^2} s, \quad (3.11)$$

$$a_{\text{1-loop}}^{(0)} = -\frac{N_s^2(1+6\xi)^4}{36864\pi^3 M_P^4} s^2 \left[\ln\left(\frac{s}{\mu_1^2}\right) - i\pi \right]. \quad (3.12)$$

⁹ We extract the factor 32π instead of 16π since the final state particles are identical bosons. The unitarity requires $\text{Im}[a^{(l)}] = |a^{(l)}|^2$ with this convention in our case, since the cross section is divided by the combinatorial factor of the identical final state particles.

We can check that it exactly satisfies the elastic unitarity,

$$\text{Im} [a^{(0)}] = |a^{(0)}|^2. \quad (3.13)$$

In this sense, the unitarity is maintained at the leading order in $1/N_s$. This is the self-healing mechanism discussed in Refs. [25, 26]. In the following, we call the dressed amplitude as the self-healed amplitude.

3.3 Self-healed amplitude as scalaron emergence

In the previous subsection, we see that the (s -wave part of the) self-healed amplitude satisfies the elastic unitarity. Now we study its structure in more detail. The s -wave part is given by

$$a^{(0)} = -\frac{N_s (1 + 6\xi)^2 s}{2304\pi\alpha_1} \left[s \left(1 - \frac{i\pi}{\ln(s/\mu_1^2)} \right) - \frac{M_P^2}{12\alpha_1} \right]^{-1}, \quad (3.14)$$

where we have defined

$$\alpha_1 \equiv -\frac{N_s (1 + 6\xi)^2}{2304\pi^2} \ln \left(\frac{s}{\mu_1^2} \right). \quad (3.15)$$

It is indeed understood as the coupling α_1 at the energy scale \sqrt{s} , since it is consistent with the beta function β_{α_1} with μ_1 being the energy scale at which α_1 vanishes. We ignore the imaginary part of the amplitude for now, whose meaning will be clarified soon. The amplitude is then written as

$$a^{(0)} = -\frac{N_s (1 + 6\xi)^2}{2304\pi\alpha_1} \left[\frac{m_s^2}{s - m_s^2} + 1 \right], \quad (3.16)$$

where we have defined

$$m_s^2 \equiv \frac{M_P^2}{12\alpha_1}. \quad (3.17)$$

We now see that it is equivalent to the tree-level scattering amplitude with the scalaron as a dynamical degree of freedom. We consider the following action:

$$S = \int d^4x \sqrt{-g} \left[\frac{M_P^2}{2} R + \bar{\alpha}_1 R^2 + \frac{\xi}{2} R \phi_i^2 + \frac{1}{2} g^{\mu\nu} \partial_\mu \phi_i \partial_\nu \phi_i - \frac{\lambda}{4} \phi_i^4 \right]. \quad (3.18)$$

By introducing an auxiliary field and performing the Weyl transformation, it is equivalent to (see App. B.1 for details)

$$S = \int d^4x \sqrt{-g} \left[\frac{M_P^2}{2} R + \frac{1}{2} (\partial s)^2 + \frac{1}{2} \left(\partial \phi_i + \frac{1}{\sqrt{6}} \frac{\phi_i}{M_P} \partial s \right)^2 - \frac{M_P^4}{16\bar{\alpha}_1} \left(1 - \exp \left(-\sqrt{\frac{2}{3}} \frac{s}{M_P} \right) - \frac{\xi \phi_i^2}{M_P^2} \right)^2 - \frac{\lambda}{4} \phi_i^4 \right], \quad (3.19)$$

where s is the scalaron degree of freedom that shows up due to the $\bar{\alpha}_1$ -term as advertised.^{†10} We compute the 2-to-2 scattering amplitude $\phi_i\phi_i \rightarrow \phi_j\phi_j$ with $i \neq j$ around the vacuum $\phi_i = s = 0$ in this model (see App. B.3 for the Feynman rules). The tree-level amplitude is diagrammatically given by

$$i\bar{A}_{\text{tree}}^{(ii \rightarrow jj)} = \text{diagram 1} + \text{diagram 2} + \text{diagram 3}, \quad (3.20)$$

where the solid line denotes the scalaron. We have added the “bar” to clarify that it is derived from the action (3.19). It is easily computed, and the s -wave part of the flavor-singlet elastic scattering amplitude is given by

$$\bar{a}_{\text{tree}}^{(0)} = -\frac{N_s (1 + 6\xi)^2}{2304\pi\bar{\alpha}_1} \left[\frac{\bar{m}_s^2}{s - \bar{m}_s^2} + 1 \right], \quad (3.21)$$

to the leading order in N_s , where the scalaron mass squared \bar{m}_s^2 is given by

$$\bar{m}_s^2 = \frac{M_P^2}{12\bar{\alpha}_1}, \quad (3.22)$$

and we have again omitted the contribution from the potential. The factor N_s again comes from the combinatorial factor and the normalization of the state. Thus, it coincides with the s -wave part of the self-healed amplitude with $\bar{\alpha}_1 = \alpha_1$. Remember that this identification is consistent with the beta function β_{α_1} . Based on this observation, we conclude that the self-healed amplitude can be understood as the amplitude with the scalaron as the intermediate state. It strongly supports that the Higgs inflation inevitably contains the scalaron as a dynamical degree of freedom.

The physical meaning of ignoring the imaginary part in Eq. (3.14) is now clear. When we compute the barred amplitude (3.21), we have included only the tree-level contribution. Once we include loop corrections, however, the scalaron propagator develops an imaginary part that corresponds to the decay rate. In this sense, it is the tree-level approximation to ignore the imaginary part. We can indeed verify that the imaginary part of Eq. (3.14) is consistent with the decay rate of the scalaron. Assuming that $|\ln(s/\mu_1^2)| \gg \pi$ and treating the logarithm as a constant, the imaginary part shifts the position of the pole as

$$s = m_s^2 - im_s\Gamma_s, \quad (3.23)$$

where we have defined

$$\Gamma_s \equiv -\frac{\pi m_s}{\ln(s/\mu_1^2)} = \frac{N_s (1 + 6\xi)^2}{192\pi} \frac{m_s^3}{M_P^2}. \quad (3.24)$$

On the other hand, the decay rate of the scalaron is computed from the action (3.19) as^{†11}

$$\bar{\Gamma}_s = \frac{N_s (1 + 6\xi)^2}{192\pi} \frac{\bar{m}_s^3}{M_P^2}. \quad (3.25)$$

^{†10} It should not be confused with the Mandelstam variable.

^{†11} The scalaron does not decay into a graviton pair although it linearly couples to R in the Jordan frame [52, 53].

They coincide with each other under the identification $\alpha_1 = \bar{\alpha}_1$. The condition $|\ln(s/\mu_1^2)| \gg \pi$ guarantees that the scalaron has a narrow width. Otherwise the perturbativity of the action (3.19) is lost. It is also naturally expected that loop corrections induce a logarithmic dependence of $\bar{\alpha}_1$ on s as in Eq. (3.15). We leave an explicit confirmation of it as a future work.

In hindsight, the result may be naturally understood as follows. It is the scalar loop contribution to the graviton vacuum polarization that induces the α_1 -term. The resummation of this contribution is equivalent to dealing with the α_1 -term as a zero-th order term in perturbation, thus equivalent to treating the scalaron as a fundamental degree of freedom. The above computation confirms this intuition.

Here we emphasize that the limit $\xi \gg 1$ and $N_s \gg 1$ is important since it justifies to include the α_1 -term to all orders while drop other terms including the α_2 -term. The α_2 -term is suppressed by ξ compared to the α_1 -term, while other (higher dimensional) terms are suppressed at least by N_s . The large ξ limit is probably good for $\xi = \mathcal{O}(10^4)$, while the validity of the large N_s limit could be a subject of discussion for the Higgs inflation with $N_s = 4$. We leave a detailed study on this point as a future work (see also Sec. 6).

Before closing this section, let us clarify differences between our analysis and Ref. [26]. The latter also discusses the self-healing mechanism of the Higgs inflation, concluding that the unitarity is self-healed. However, it does not discuss any physical interpretation of the self-healed amplitude.^{†12} Our main point in this section is that the self-healed amplitude can be understood as the amplitude with the scalaron as the intermediate state, thus to provide a physical interpretation to the self-healing mechanism of the Higgs inflation.

4 Equivalence of Jordan and Einstein frames

So far we have discussed the self-healing mechanism and the emergence of the scalaron in the Jordan frame, i.e. in the frame where the non-minimal coupling to the Ricci scalar is present. It can be cast into the Einstein frame where the non-minimal coupling is absent by the Weyl transformation. Since people often discuss physics in the Einstein frame, it is valuable to explicitly check that the same results are obtained in the Einstein frame. For this purpose, we show the equivalence of the Jordan and Einstein frames of our results in this section.

We consider our action:

$$S = \int d^4x \sqrt{-g_J} \left[\frac{M_P^2}{2} R_J + \frac{1}{2} g_J^{\mu\nu} \partial_\mu \phi_i \partial_\nu \phi_i + \frac{\xi}{2} R_J \phi_i^2 \right] + S_{\text{g.f.}}, \quad (4.1)$$

where $S_{\text{g.f.}}$ is the gauge fixing term that is important in the following discussion, and we have ignored the potential as usual. We assign the subscripts J and E to the quantities in the Jordan frame and the Einstein frame, respectively. We perform the Weyl transformation as

$$g_{J\mu\nu} = \Omega^{-2} g_{E\mu\nu}, \quad \Omega^2 = 1 + \frac{\xi \phi_i^2}{M_P^2}. \quad (4.2)$$

^{†12} Refs. [21, 26] rather claim that there is no physical pole in the s -wave sector, saying that the denominator is of the form $1 - sF_1(s)/2$ with $F_1(s) < 0$. We disagree for two reasons. First, their F_1 can be positive depending on μ_1 (or the parameter choice of the theory) since it is proportional to $\ln(s/\mu_1^2)$. Second, the amplitude has a pole even if F_1 is negative, although the corresponding particle (i.e. the scalaron) is tachyonic in such a case.

The Ricci scalar is transformed as

$$R_J = \Omega^2 \left[R_E + \frac{3}{2} g_E^{\mu\nu} \partial_\mu \ln \Omega^2 \partial_\nu \ln \Omega^2 - 3 \square_E \ln \Omega^2 \right]. \quad (4.3)$$

As a result, the action in the Einstein frame is given by


$$S = \int d^4x \sqrt{-g_E} \left[\frac{M_P^2}{2} R_E + \frac{1}{2\Omega^4} \left(\Omega^2 \delta_{ij} + \frac{6\xi^2 \phi_i \phi_j}{M_P^2} \right) g_E^{\mu\nu} \partial_\mu \phi_i \partial_\nu \phi_j \right] + S_{\text{g.f.}} \quad (4.4)$$

The metric of the scalar kinetic term is curved for $N_s \geq 2$, and hence ξ is physical. The situation is different for $N_s = 1$, which is discussed separately in Sec. 5.

The Weyl transformation is merely a field redefinition, and physics is expected to be the same in both frames [54–56]. In this section, we explicitly check that this equivalence holds in our case. In particular, we will see that the Feynman rules around $\phi_i = 0$ for 2-to-2 scattering are exactly the same even at *off-shell* level, once we properly take into account contributions from the gauge fixing term. It enables us to see the equivalence parts by parts in Feynman diagrams.

4.1 Jordan frame

First, we summarize the Feynman rules in the Jordan frame. Note that all the diagrams of our interest are constructed from the *off-shell* 2-to-2 scattering diagram,




$$\quad (4.5)$$

We may take the de Donder gauge in the Jordan frame, namely,

$$S_{\text{g.f.}} = \int d^4x \left[\left(\partial_\nu h_J^{\mu\nu} - \frac{1}{2} \partial^\mu h_J \right) \left(\partial^\rho h_{J\mu\rho} - \frac{1}{2} \partial_\mu h_J \right) \right], \quad (4.6)$$

at the quadratic order, where $g_{J\mu\nu} = \eta_{\mu\nu} + (2/M_P)h_{J\mu\nu}$, and $h_J \equiv \eta_{\mu\nu} h_J^{\mu\nu}$. Then the Feynman rule for the above diagram is



$$= -\frac{i}{2M_P^2} \frac{1}{q^2} V_{\mu\nu}^{(0)}(p_1, p_2) P^{\mu\nu\rho\sigma} V_{\rho\sigma}^{(0)}(p_3, p_4) + \frac{6i\xi^2}{M_P^2} q^2$$

$$+ \frac{4i\xi}{M_P^2} \left[(p_1 \cdot p_2) + (p_3 \cdot p_4) - \frac{(p_1 \cdot q)(p_2 \cdot q) + (p_3 \cdot q)(p_4 \cdot q)}{q^2} \right], \quad (4.7)$$

where $q \equiv p_1 + p_2 = p_3 + p_4$, we take the external momenta p_1, p_2, p_3 and p_4 to be off-shell, and

$$V_{\mu\nu}^{(0)}(p, q) \equiv (p \cdot q) \eta_{\mu\nu} - (p_\mu q_\nu + p_\nu q_\mu), \quad (4.8)$$

$$P_{\mu\nu\rho\sigma} \equiv \eta_{\mu\rho} \eta_{\nu\sigma} + \eta_{\mu\sigma} \eta_{\nu\rho} - \eta_{\mu\nu} \eta_{\rho\sigma}. \quad (4.9)$$

The function $V_{\mu\nu}^{(0)}$ corresponds to the scalar-scalar-graviton vertex for $\xi = 0$. See App. A for more details. Below we see that the same off-shell Feynman rule is obtained in the Einstein frame once we properly take into account contributions from the gauge fixing term.

4.2 Einstein frame

Now we derive the Feynman rules for off-shell 2-to-2 scattering diagrams in the Einstein frame. The action (4.4) contains scalar four-point vertices,

$$S \ni \int d^4x \left[-\frac{\xi\phi_j^2}{2M_P^2} \partial^\mu \phi_i \partial_\mu \phi_i + \frac{3\xi^2}{4M_P^2} (\partial^\mu \phi_i^2) (\partial_\mu \phi_j^2) \right], \quad (4.10)$$

as well as the standard scalar-scalar-graviton vertex. There are additional contributions from the gauge fixing term. The gravitons in the Jordan and Einstein frames are related as

$$\begin{aligned} h_{J\mu\nu} &= \frac{1}{\Omega^2} \left(\frac{M_P}{2} \eta_{\mu\nu} + h_{E\mu\nu} \right) - \frac{M_P}{2} \eta_{\mu\nu} \\ &= h_{E\mu\nu} - \frac{\xi\phi_i^2}{2M_P} \eta_{\mu\nu} + \dots, \end{aligned} \quad (4.11)$$

and hence the de Donder gauge in the Jordan frame results in additional vertices in the Einstein frame:

$$S_{\text{g.f.}} \ni \int d^4x \left[-\frac{\xi}{M_P} \left(\partial^\mu \partial^\nu \phi_i^2 - \frac{1}{2} \eta^{\mu\nu} \partial^2 \phi_i^2 \right) h_{E\mu\nu} + \frac{\xi^2}{4M_P^2} (\partial^\mu \phi_i^2) (\partial_\mu \phi_j^2) \right]. \quad (4.12)$$

The Feynman rules are thus given by

$$\begin{array}{c} p_{1,i} \\ \diagdown \\ \text{---} \\ \diagup \\ p_{2,i} \end{array} = -\frac{i}{M_P} V_{\mu\nu}^{(0)}(p_1, p_2) + \frac{2i\xi}{M_P} \left(q_\mu q_\nu - \frac{\eta_{\mu\nu}}{2} q^2 \right), \quad (4.13)$$

for the scalar-scalar-graviton vertex, and

$$\begin{array}{ccc} p_{1,i} & & p_{3,j} \\ & \diagdown & / \\ & \text{---} & \\ & / & \diagdown \\ p_{2,i} & & p_{4,j} \end{array} = \frac{2i\xi}{M_P^2} ((p_1 \cdot p_2) + (p_3 \cdot p_4)) + \frac{8i\xi^2}{M_P^2} q^2, \quad (4.14)$$

for the scalar four-point vertex with $i \neq j$, where again $q \equiv p_1 + p_2$. We now obtain the Feynman rule for the off-shell 2-to-2 scattering diagrams as

$$\begin{array}{ccc} p_{1,i} & & p_{3,j} \\ \diagdown & & / \\ \text{---} & & \\ \diagup & & \diagdown \\ p_{2,i} & & p_{4,j} \end{array} + \begin{array}{ccc} p_{1,i} & & p_{3,j} \\ & \diagdown & / \\ & \text{---} & \\ & / & \diagdown \\ p_{2,i} & & p_{4,j} \end{array} = -\frac{i}{2M_P^2} \frac{1}{q^2} V_{\mu\nu}^{(0)}(p_1, p_2) P^{\mu\nu\rho\sigma} V_{\rho\sigma}^{(0)}(p_3, p_4) + \frac{6i\xi^2}{M_P^2} q^2 \\ + \frac{4i\xi}{M_P^2} \left[(p_1 \cdot p_2) + (p_3 \cdot p_4) - \frac{(p_1 \cdot q)(p_2 \cdot q) + (p_3 \cdot q)(p_4 \cdot q)}{q^2} \right], \quad (4.15)$$

for $i \neq j$, which is exactly the same as the Feynman rule (4.7) in the Jordan frame. We can show the equivalence for $i = j$ in a similar manner once we include not only the s -channel but also the t - and u -channels, as they all contribute for $i = j$. The equivalence of the Feynman rules associated with the counter terms can be confirmed in the same way.^{†13} Thus, we have shown the equivalence of the Jordan and Einstein frames at the diagrammatic level. The equivalence of our argument, especially the self-healing mechanism, follows straightforwardly.

In the above argument we have carefully taken into account the vertices arising from the gauge fixing term. The choice of the gauge fixing term is unphysical, and hence we will obtain the same result even if we do not include those vertices as far as physical quantities are concerned.^{†14} Nevertheless, we feel it meaningful to treat the gauge fixing term carefully, since it enables us to see the equivalence not only at the on-shell level, but also at the off-shell (i.e. Feynman diagrammatic) level. For instance, we now easily see that the self-healed amplitude (3.5) in the Jordan frame roughly corresponds to

$$A_{\text{dressed}}^{(ii \rightarrow jj)} = \text{diagram 1} + \text{diagram 2} + \text{diagram 3} + \dots, \quad (4.16)$$

in the Einstein frame. In other words, the self-healing mechanism in the Einstein frame comes dominantly from the non-renormalizable interaction in the scalar kinetic terms. This observation has an interesting implication, which we will briefly discuss in Sec. 6.

5 Single field case

So far we have focused on the case $N_s \geq 2$. The single field case $N_s = 1$ is tricky and is discussed separately in this section. Remember that $N_s = 4$ in the Higgs inflation, and hence the subtlety discussed in the following does not apply to that case. We ignore the potential first, and briefly comment on effects of the potential in the end.

It is known that the non-minimal coupling ξ is unphysical for $N_s = 1$ once we ignore the potential, as emphasized in Ref. [8]. It is best described by the Weyl transformation. We consider the single field action with the non-minimal coupling:

$$S = \int d^4x \sqrt{-g_J} \left[\frac{M_P^2}{2} R_J + \frac{1}{2} g_J^{\mu\nu} \partial_\mu \phi \partial_\nu \phi + \frac{\xi}{2} R_J \phi^2 \right]. \quad (5.1)$$

We can move to the Einstein frame in the same way to obtain Eq. (4.4) from Eq. (4.1). The action in the Einstein frame is

$$S = \int d^4x \sqrt{-g_E} \left[\frac{M_P^2}{2} R_E + \frac{1}{2\Omega^4} \left(\Omega^2 + \frac{6\xi^2 \phi^2}{M_P^2} \right) g_E^{\mu\nu} \partial_\mu \phi \partial_\nu \phi \right]. \quad (5.2)$$

Now we redefine the scalar field as

$$d\chi = \frac{d\phi}{\Omega^2} \sqrt{\Omega^2 + \frac{6\xi^2 \phi^2}{M_P^2}}, \quad (5.3)$$

^{†13} To the order of our interest, the α_1 -term in the Jordan frame results in a scalar-scalar-graviton vertex and a scalar four-point vertex in addition to the graviton quadratic term (A.21) in the Einstein frame. The α_2 -term in the Jordan frame does not induce vertices involving the scalar fields in the Einstein frame.

^{†14} For instance, we can start with the de Donder gauge in the Einstein frame, and compute quantities such as the scalar loop to the graviton vacuum polarization. We have checked that it reproduces the same result as expected.

so that the action is recast into the Einstein-Hilbert term and a canonically normalized scalar field. The non-minimal coupling ξ can be erased by the field redefinition, and hence is unphysical. The crucial difference between Eq. (4.4) and (5.2) is that the target space of the scalar fields is curved for $N_s \geq 2$, while it is not curved for $N_s = 1$. The curvature depends on ξ for $N_s \geq 2$, indicating that ξ is physical as the curvature is invariant under field redefinition (that corresponds to coordinate transformation of the target space).


It is possible to check explicitly that physical quantities such as scattering amplitudes do not depend on ξ . At the tree-level, the s -channel scattering amplitude of $\phi\phi \rightarrow \phi\phi$ is given by Eq. (3.1), i.e.,

$$A_{\text{tree}}^{(s)} = \frac{1}{M_P^2 s} \left[\frac{(1 + 6\xi)^2}{6} s^2 - \left(\frac{s^2}{6} - tu \right) \right]. \quad (5.4)$$


In the present case, the t - and u -channels also contribute to the scattering amplitude, and hence the tree-level amplitude is given by

$$A_{\text{tree}} = A_{\text{tree}}^{(s)} + A_{\text{tree}}^{(t)} + A_{\text{tree}}^{(u)} = \frac{1}{M_P^2} \left[\frac{tu}{s} + \frac{us}{t} + \frac{st}{u} \right]. \quad (5.5)$$

In particular, the ξ -dependent part drops since the Mandelstam variables satisfy $s + t + u = 0$. At the one-loop level, it is probably easiest to work with the de Donder gauge in the Einstein frame (that results in additional vertices in the Jordan frame, in a similar way as discussed in Sec. 4). For instance, we can easily see in the Einstein frame (before the field redefinition) that the following one-loop diagram vanishes:

$$iA_{1\text{-loop}}^{(s)} = \text{diagram} = 0. \quad (5.6)$$


In a similar way as in Sec. 4, the scalar four-point vertex in the Einstein frame is equivalent at the off-shell level to the diagrams in the Jordan frame as

$$\text{diagram} \sim \text{diagram} + (s \leftrightarrow t) + (s \leftrightarrow u) + \text{diagram}, \quad (5.7)$$


where we have used the similarity since the right hand side includes the contributions from the spin-2 part of the graviton as well.^{‡15} It is thus clear that the one-loop diagram (5.6) in the Einstein frame corresponds to many diagrams including the vacuum polarization, the vertex correction and the box diagrams in the Jordan frame. It means that the ξ -dependence cancels out among these diagrams in the Jordan frame. The diagrams with the scalar loop depend on N_s differently from the other diagrams, and hence this cancellation holds only for $N_s = 1$.^{‡16}

It follows that the cut-off scale of the theory with $N_s = 1$ does not depend on ξ , and hence the self-healing mechanism is not at work (at least below the Planck scale). Thus, the dynamical

^{‡15} The de Donder gauge in one frame generates a scalar four-point vertex in the other frame, as in Eq. (4.12).

^{‡16} The beta functions in Sec. 2 take into account only the scalar loop. While there may be additional contributions from the vertex correction and the box diagrams, the argument in Sec. 2 is intact since the cancellation does not occur for $N_s \geq 2$. Also we can ignore these terms in Sec. 3 as they are sub-leading in the large N_s expansion.

emergence of the scalaron does not happen in the single field case without the potential. Once we include the potential, the story could be totally different. The field redefinition (5.3) results in higher dimensional operators in the potential, whose cut-off scale does depend on ξ [8]. We emphasize that this case requires a separate discussion from the case $N_s \geq 2$. At least, there is no scalar four-point derivative interaction in the Einstein frame (after field redefinition), and hence the self-healing mechanism (if at work) will not be in the form of Eq. (4.16). Moreover, we may expect that a particle that emerges due to the self-healing mechanism (if any) is not the scalaron, since the scalaron may not cure the low cut-off scale associated with the higher dimensional operators in the potential sector. A detailed study on this case, albeit definitely interesting, is beyond the scope of this paper.

6 Summary and discussions

In this paper, we have argued that a light scalaron dynamically emerges if there is a sizable non-minimal coupling between scalar fields and the Ricci scalar $\xi\phi_i^2 R$ as in the Higgs inflation model, as long as the number of the scalar fields N_s satisfies $N_s \geq 2$. This claim is based on two (closely related) observations. First, the R^2 -term necessarily emerges due to the RG running whose coefficient is generically of $\mathcal{O}(\xi^2)$ for $\xi \gg 1$. Since the coefficient of the R^2 -term is inversely proportional to the scalaron mass squared, it implies that a light scalaron inevitably exists in the theory (see also Refs. [20–23]). Second, we have seen that the 2-to-2 scattering amplitude $\phi_i\phi_i \rightarrow \phi_j\phi_j$ with $i \neq j$ develops a pole structure after resumming over the diagrams that are leading order in the large N_s expansion. We have explicitly checked that the resultant scattering amplitude is equivalent to the amplitude with the scalaron in the intermediate state. The resummation heals the unitarity of the Higgs inflation, called the self-healing mechanism in Refs. [25,26], and hence our result identifies the self-healing mechanism as the dynamical emergence of the scalaron. We have confirmed that our results do not depend on whether we work in the Jordan frame or in the Einstein frame.

Two implications immediately follow. First, the Higgs inflation is actually a two-field inflationary model, the (radial component of the) Higgs and the scalaron. In particular, the inflaton is in general an admixture of the Higgs and the scalaron. The inflationary dynamics of this theory is discussed in App. B and references therein. Second, contrary to the common wisdom, the Higgs inflation does not suffer from the unitarity issue since the self-healing mechanism is at work. It is nothing but the fact that the scalaron pushes up the cut-off scale to the Planck scale, as pointed out in Refs. [22,24]. Thus, we can follow all the dynamics from the inflation till the end of the reheating within the validity of the theory.

There are several other points that are not discussed in detail in this paper. Below we list some of them before ending this paper.

Self-healing in other inflationary models

In Sec. 4, we see that the self-healing mechanism of the Higgs inflation is caused by the higher dimensional operators in the scalar kinetic sector in the Einstein frame (see Eq. (4.16)). There are other inflationary models that have non-trivial scalar kinetic terms (or a curved target space), such as the running kinetic model [57,58], the α -attractor model [59], and the Higgs-dilaton model [60]. The Palatini Higgs inflation model [61–65] could also be within this class of models in the Einstein frame, among probably many others. Since these non-trivial kinetic terms are interpreted as higher




Figure 1: Schematic picture of the RG running of the scalaron mass squared m_s^2 .

dimensional operators once expanded around the vacuum, a similar self-healing mechanism may be at work and a new degree of freedom may show up dynamically in these models. This new degree of freedom, if any, is not necessarily be the scalaron. In particular, it is possible in some models that the new degree of freedom is pathological, either ghost-like or tachyonic, that may spoil the inflationary prediction. Also it is possible that there appear more than one new degrees of freedom, which might drastically modify the inflationary prediction. These points will be clarified once we study pole structures of scattering amplitudes resulting from resummation similar to Eq. (4.16). For instance, the position of the pole tells whether a particle is tachyonic or not, while the sign of the residue does whether it is ghost-like or not.^{‡17} We believe that it is of great importance to re-analyze the other inflationary models from this point of view, which will be done elsewhere.

Running of scalaron mass

The running of the scalaron mass squared shows a peculiar behavior. The (scalar loop contribution to the) RG equation of α_1 is given by

$$\beta_{\alpha_1} \equiv \frac{d\alpha_1}{d \ln \mu} = -\frac{N_s}{1152\pi^2} (1 + 6\xi)^2. \quad (6.1)$$

Since the scalaron mass squared m_s^2 in terms of α_1 is

$$m_s^2 = \frac{M_P^2}{12\alpha_1}, \quad (6.2)$$

its RG running is schematically given in Fig. 1.^{‡18} An interesting feature is that its mass squared diverges at the energy scale μ_1 at which α_1 vanishes, and becomes tachyonic above μ_1 . It might

^{‡17} The spin-2 (*d*-wave) part of the self-healed amplitude in our theory also develops a pole. Its residue has an opposite sign from that of the spin-0 (*s*-wave) part, as long as the poles are not tachyonic. Since the unitarity requires that the residues are expanded by the Legendre polynomials whose coefficients have a definite sign, it indicates that a ghost exists in the spin-2 sector. The position of the pole is consistent with Eq. (2.7).

^{‡18} It crucially depends on the sign of the beta function. See the footnote ‡5.

be hard to imagine that the low energy theory is healthy above μ_1 . An easy way to avoid such a situation is to assume that UV completion of the gravity provides $\alpha_1 > 0$ as the boundary condition, i.e. $\mu_1 > \Lambda_{\text{UV}}$ where Λ_{UV} is the scale at which the UV completion comes into play. We may further require $\ln(\mu_1^2/\Lambda_{\text{UV}}^2) \gg \pi$ to guarantee the perturbativity of the action (3.19).

The RG equation indicates that it has some unease to assume that the scalaron is so heavy at the energy scale of the Higgs inflation (which we call μ_{HI}) that it does not affect the inflationary dynamics. If we assume $\alpha_1 \ll \xi^2$ at μ_{HI} , the scalaron becomes tachyonic just above μ_{HI} . It probably requires UV completion just above μ_{HI} that is significantly lower than the Planck scale. Also the scalaron has to be taken into account after inflation such as reheating even if it is heavy during inflation.

Large N_s expansion

In Sec. 3, we have discussed the self-healing mechanism relying on the large N_s limit. Based on the discussion in Sec. 2, however, it is natural to expect that the dynamical emergence of the scalaron is at work for finite N_s as well. A first step to check this expectation may be to include next-to-leading order (NLO) terms in the large N_s expansion. At the NLO level, one is probably required to include diagrams with one graviton or ghost inside loops, and diagonal parts of the scattering amplitude between the flavor-singlet states. Computation will be of course more tedious, but it is nevertheless valuable to see whether the self-healing mechanism works or not at the NLO level. We leave it as a future work.

Amplitudes around $\phi_i \neq 0$

In this paper we have focused on the scattering amplitudes around the vacuum $\phi_i = 0$. However, the self-healing mechanism of the Higgs inflation is discussed also around the finite Higgs field value $\phi_i \neq 0$ in Ref. [26], which concludes that it works in the same way for $\phi_i \neq 0$. Thus we expect that our discussion in this paper can be extended to the amplitudes around $\phi_i \neq 0$ as well. It probably requires some cares including the prescription issue (see e.g. Refs. [32, 33, 54, 55, 66] and references therein) to see the equivalence between the Jordan and Einstein frames in this case.

Acknowledgement

YE thanks Kyohei Mukaida for valuable discussions and motivating him to write this paper. The Feynman diagrams in this paper are generated by TikZ-Feynman [67].

A Computational details

In this appendix, for the sake of clarity and completeness, we give our derivation of the scattering amplitudes and the RG equations in detail.

A.1 Preliminary

Gravity sector

Here we summarize our convention in this paper. We use the almost-minus convention for the metric (denoted as $g_{\mu\nu}$). In particular, the flat spacetime metric is given by

$$\eta_{\mu\nu} = \text{diag}(+1, -1, -1, -1). \quad (\text{A.1})$$

The Christoffel symbol is given by

$$\Gamma^\rho{}_{\mu\nu} = \frac{g^{\rho\sigma}}{2} (g_{\mu\sigma,\nu} + g_{\nu\sigma,\mu} - g_{\mu\nu,\sigma}), \quad (\text{A.2})$$

where $g_{\mu\nu,\rho} \equiv \partial_\rho g_{\mu\nu}$. The sign convention for the Ricci tensor is chosen such that

$$R_{\mu\rho} = \Gamma^\nu{}_{\nu\rho,\mu} - \Gamma^\nu{}_{\mu\rho,\nu} + \Gamma^\alpha{}_{\nu\rho}\Gamma^\nu{}_{\alpha\mu} - \Gamma^\alpha{}_{\mu\rho}\Gamma^\nu{}_{\alpha\nu}, \quad (\text{A.3})$$

and the Ricci scalar is given by

$$R = g^{\mu\nu} R_{\mu\nu}. \quad (\text{A.4})$$

It follows from these conventions that the variations are given by

$$\delta\sqrt{-g} = -\frac{\sqrt{-g}}{2} g_{\mu\nu} \delta g^{\mu\nu}, \quad (\text{A.5})$$

$$\delta R = R_{\mu\nu} \delta g^{\mu\nu} - g_{\mu\nu} \square \delta g^{\mu\nu} + \nabla_\mu \nabla_\nu \delta g^{\mu\nu}, \quad (\text{A.6})$$

where $g \equiv \det(g_{\mu\nu})$, ∇_μ is the covariant derivative and $\square = \nabla^\alpha \nabla_\alpha$.

Energy momentum tensor

In this paper we focus on the following action:^{‡19}

$$S = \int d^4x \sqrt{-g} \left[\frac{M_P^2}{2} R + \frac{1}{2} g^{\mu\nu} \partial_\mu \phi_i \partial_\nu \phi_i + \frac{\xi}{2} R \phi_i^2 - V(\phi_i) \right], \quad (\text{A.7})$$

where $i = 1, 2, \dots, N_s$ is the flavor index ($N_s = 4$ in the Higgs inflation). With this definition, $\xi = -1/6$ corresponds to the conformal coupling. The potential $V(\phi_i)$ is not important in our discussion, and is ignored henceforth. The energy stress tensor is defined as

$$T_{\mu\nu} = -\frac{2}{\sqrt{-g}} \frac{\delta S_m}{\delta g^{\mu\nu}}, \quad (\text{A.8})$$

where S_m is the matter part of the action. It is given in our case as

$$\begin{aligned} T_{\mu\nu} = & \frac{g_{\mu\nu}}{2} (1 + 4\xi) g^{\alpha\beta} \partial_\alpha \phi_i \partial_\beta \phi_i - (1 + 2\xi) \partial_\mu \phi_i \partial_\nu \phi_i \\ & + 2\xi (g_{\mu\nu} \phi_i \square \phi_i - \phi_i \nabla_\mu \nabla_\nu \phi_i) - \xi \left(R_{\mu\nu} - \frac{g_{\mu\nu}}{2} R \right) \phi_i \phi_i. \end{aligned} \quad (\text{A.9})$$

In the flat spacetime it reduces to

$$T_{\mu\nu}^{(\text{flat})} = \frac{\eta_{\mu\nu}}{2} (1 + 4\xi) \partial^\alpha \phi_i \partial_\alpha \phi_i - (1 + 2\xi) \partial_\mu \phi_i \partial_\nu \phi_i + 2\xi (\eta_{\mu\nu} \phi_i \partial^2 \phi_i - \phi_i \partial_\mu \partial_\nu \phi_i). \quad (\text{A.10})$$

^{‡19} The sign of the Einstein Hilbert action is fixed once the sign convention of the Ricci scalar is fixed.

A.2 Feynman rules

Here we summarize the Feynman rules in the flat spacetime derived from the action (2.1). The ghost degree of freedom associated with the gauge fixing is irrelevant for our analysis, and hence we ignore it in the following.

Propagator

The scalar propagator is given by

$$\text{-----} = \frac{i}{p^2 + i0}. \quad (\text{A.11})$$

We expand the metric around the flat spacetime as

$$g_{\mu\nu} = \eta_{\mu\nu} + \frac{2}{M_P} h_{\mu\nu}. \quad (\text{A.12})$$

We take the de Donder gauge in the Jordan frame in this appendix, given by

$$S_{\text{g.f.}} = \int d^4x \left[\left(\partial_\nu h^{\mu\nu} - \frac{1}{2} \partial^\mu h \right) \left(\partial^\rho h_{\mu\rho} - \frac{1}{2} \partial_\mu h \right) \right], \quad (\text{A.13})$$

at the quadratic order, and hence the graviton propagator is given by

$$\begin{array}{c} \mu\nu \\ \text{~~~~~} \\ \rho\sigma \end{array} = \frac{i}{2p^2 + i0} P_{\mu\nu\rho\sigma}, \quad (\text{A.14})$$

where

$$P_{\mu\nu\rho\sigma} \equiv \eta_{\mu\rho}\eta_{\nu\sigma} + \eta_{\mu\sigma}\eta_{\nu\rho} - \eta_{\mu\nu}\eta_{\rho\sigma}. \quad (\text{A.15})$$

Scalar-scalar-graviton interaction

The interaction between the scalar fields and the graviton is given by

$$S_{\text{int}} = \frac{1}{M_P} \int d^4x \sqrt{-g} h^{\mu\nu} T_{\mu\nu}. \quad (\text{A.16})$$

From this expression, the scalar-scalar-graviton interaction reads

$$\begin{array}{c} p_1, i \\ \text{---} \\ \text{---} \\ p_2, j \end{array} \begin{array}{c} \text{~~~~~} \\ \mu\nu \end{array} = -\frac{i\delta_{ij}}{M_P} V_{\mu\nu}(p_1, p_2), \quad (\text{A.17})$$

where the vertex function is given by

$$\begin{aligned} V_{\mu\nu}(p_1, p_2) &= (1 + 4\xi) (p_1 \cdot p_2) \eta_{\mu\nu} - (1 + 2\xi) (p_{1\mu} p_{2\nu} + p_{1\nu} p_{2\mu}) \\ &\quad + 2\xi [(p_1^2 + p_2^2) \eta_{\mu\nu} - (p_{1\mu} p_{1\nu} + p_{2\mu} p_{2\nu})], \end{aligned} \quad (\text{A.18})$$

and the momenta are both outgoing or both incoming.^{‡20} Note that the first term in the second line does not vanish for off-shell momenta.

^{‡20} The factor two from the exchange of the scalars is included in this rule.

Counter terms

As we will see in the following, we need to include counter terms to renormalize divergences from the scalar one-loop diagram. Thus we include the following terms in our action:

$$S_{\text{c.t.}} = \int d^4x \sqrt{-g} \left[\alpha_1 R^2 + \alpha_2 \left(R_{\mu\nu} R^{\mu\nu} - \frac{1}{3} R^2 \right) \right]. \quad (\text{A.19})$$

The term $R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma}$ is absorbed into the above two terms up to the total derivative, and hence we ignore it in the following. To the leading order in $h_{\mu\nu}$, the Ricci tensor is given as

$$R_{\mu\rho} \simeq \frac{1}{M_P} \left[\partial_\mu \partial_\rho h^\alpha{}_\alpha + \partial^2 h_{\mu\rho} - \partial^\nu \partial_\rho h_{\mu\nu} - \partial^\nu \partial_\mu h_{\rho\nu} \right]. \quad (\text{A.20})$$

The Feynman rules are read off as

$$\begin{array}{c} \alpha_1 \\ \text{~~~~~} \\ \mu\nu \text{~~~~~} \times \text{~~~~~} \rho\sigma \end{array} = \frac{8i\alpha_1}{M_P^2} q^4 P_{\mu\nu} P_{\rho\sigma}, \quad (\text{A.21})$$

for the α_1 -term, and

$$\begin{array}{c} \alpha_2 \\ \text{~~~~~} \\ \mu\nu \text{~~~~~} \times \text{~~~~~} \rho\sigma \end{array} = \frac{i\alpha_2}{M_P^2} q^4 \left[P_{\mu\rho} P_{\nu\sigma} + P_{\mu\sigma} P_{\nu\rho} - \frac{2}{3} P_{\mu\nu} P_{\rho\sigma} \right], \quad (\text{A.22})$$

for the α_2 -term, where we define the projection operator as

$$P_{\mu\nu} = \eta_{\mu\nu} - \frac{q_\mu q_\nu}{q^2}. \quad (\text{A.23})$$

A.3 Tree-level amplitude

We consider the 2-to-2 scattering amplitude $\phi_i \phi_i \rightarrow \phi_j \phi_j$ with $i \neq j$. Only the s -channel process contributes in this case, which is given by

$$\begin{aligned} iA_{\text{tree}}^{(ii \rightarrow jj)} &= \begin{array}{c} p_{1,i} \qquad p_{3,j} \\ \text{---} \quad \text{---} \\ \text{---} \times \text{---} \\ \text{---} \quad \text{---} \\ p_{2,i} \qquad p_{4,j} \end{array} \\ &= \left(-\frac{i}{M_P} \right) V_{\mu\nu}(p_1, p_2) \frac{i}{2(p_1 + p_2)^2} P^{\mu\nu\rho\sigma} \left(-\frac{i}{M_P} \right) V_{\rho\sigma}(p_3, p_4). \end{aligned} \quad (\text{A.24})$$

It is simplified as

$$A_{\text{tree}}^{(ii \rightarrow jj)} = \frac{1}{M_P^2 s} \left[\frac{(1 + 6\xi)^2}{6} s^2 - \left(\frac{s^2}{6} - tu \right) \right], \quad (\text{A.25})$$

where the Mandelstam variables are defined as

$$s \equiv (p_1 + p_2)^2 = (p_3 + p_4)^2, \quad (\text{A.26})$$

$$t \equiv (p_1 - p_3)^2 = (p_2 - p_4)^2, \quad (\text{A.27})$$

$$u \equiv (p_1 - p_4)^2 = (p_2 - p_3)^2. \quad (\text{A.28})$$

It correctly reproduces the result in Ref. [25].

A.4 Scalar one-loop amplitude

Now we compute the one-loop correction to the 2-to-2 scattering amplitude. We focus on the scalar loop here. It is given by

$$\begin{aligned}
 iA_{1\text{-loop}}^{(ii \rightarrow jj)} &= \text{Diagram: A scalar loop diagram with external momenta } p_1, p_2, p_3, p_4 \text{ and loop momenta } l, l-q. \\
 &= \frac{N_s}{2} \int \frac{d^4 l}{(2\pi)^4} \left(-\frac{i}{M_P} \right)^4 \left(\frac{i}{2s} \right)^2 \frac{i}{l^2 + i0} \frac{i}{(l-q)^2 + i0} \\
 &\quad \times [V_{\mu\nu}(p_1, p_2) P^{\mu\nu\rho\sigma} V_{\rho\sigma}(l, -l+q)] [V_{\alpha\beta}(l, -l+q) P^{\alpha\beta\gamma\delta} V_{\gamma\delta}(p_3, p_4)], \quad (\text{A.29})
 \end{aligned}$$

where $q = p_1 + p_2 = p_3 + p_4$, $s = q^2$ is the Mandelstam variable, and the factor $1/2$ in front of the second line is the symmetry factor. We first focus on the real (divergent) part. By using the Feynman's trick,

$$\frac{1}{AB} = \int_0^1 dx \frac{1}{(A + (B-A)x)^2}, \quad (\text{A.30})$$

and moving to the Euclidean momentum space, we obtain

$$\begin{aligned}
 \text{Re} [A_{1\text{-loop}}^{(ii \rightarrow jj)}] &= \frac{N_s \mu^{4-d}}{2M_P^4 s^2} \int_0^1 dx \int \frac{d^d l_E}{(2\pi)^d} \frac{1}{(l_E^2 + x(1-x)s)^2} \\
 &\quad \times [(1+4\xi) s l_E^2 - 4(\xi(q \cdot l_E)^2 + (p_1 \cdot l_E)(p_2 \cdot l_E)) - s^2 \xi(1+6\xi)] \\
 &\quad \times [(1+4\xi) s l_E^2 - 4(\xi(q \cdot l_E)^2 + (p_3 \cdot l_E)(p_4 \cdot l_E)) - s^2 \xi(1+6\xi)], \quad (\text{A.31})
 \end{aligned}$$

where we have performed the dimensional regularization. We can simplify the integrals as

$$\int \frac{d^d l_E}{(2\pi)^d} l_{E\mu} l_{E\nu} f(l_E^2) = \frac{\eta_{\mu\nu}}{d} \int \frac{d^d l_E}{(2\pi)^d} l_E^2 f(l_E^2), \quad (\text{A.32})$$

$$\int \frac{d^d l_E}{(2\pi)^d} l_{E\mu} l_{E\nu} l_{E\rho} l_{E\sigma} f(l_E^2) = \frac{\eta_{\mu\nu} \eta_{\rho\sigma} + \eta_{\mu\rho} \eta_{\nu\sigma} + \eta_{\mu\sigma} \eta_{\nu\rho}}{d(d+2)} \int \frac{d^d l_E}{(2\pi)^d} l_E^4 f(l_E^2), \quad (\text{A.33})$$

where f is an arbitrary function of l_E^2 . After applying these formulas and some computation, we obtain the divergent part of the amplitude as

$$A_{1\text{-loop}}^{(ii \rightarrow jj)} \Big|_{\text{div}} = \frac{N_s}{960\pi^2 M_P^4} \frac{1}{\epsilon} \left[\frac{5}{6} (1+6\xi)^4 s^2 + \left(\frac{s^2}{6} - tu \right) \right], \quad (\text{A.34})$$

where ϵ is defined as $d = 4 - 2\epsilon$. It is renormalized by the α_1 - and α_2 -terms. The amplitude from the counter terms is

$$\begin{aligned}
 iA_{\text{c.t.}}^{(ii \rightarrow jj)} &= \text{Diagram: Two counterterm diagrams with external momenta } p_1, p_2, p_3, p_4 \text{ and vertices } \alpha_1, \alpha_2. \\
 &= \frac{2i}{M_P^4} \left[(1+6\xi)^2 s^2 \alpha_1 + \left(\frac{s^2}{6} - tu \right) \alpha_2 \right], \quad (\text{A.35})
 \end{aligned}$$

and hence the divergent parts of α_1 and α_2 are given by

$$\alpha_1|_{\text{div}} = -\frac{N_s (1 + 6\xi)^2}{2304\pi^2} \frac{1}{\epsilon}, \quad (\text{A.36})$$

$$\alpha_2|_{\text{div}} = -\frac{N_s}{1920\pi^2} \frac{1}{\epsilon}. \quad (\text{A.37})$$

They coincide with Ref. [18] for $\xi = 0$. Noting that the amplitude is associated with the factor $\mu^{4-d} \simeq 1 - \epsilon \ln \mu^{-2}$, we obtain the real part of the one-loop amplitude after the renormalization as

$$\text{Re} \left[A_{1\text{-loop}}^{(ii \rightarrow jj)} \right] = -\frac{N_s}{960\pi^2 M_P^4} \left[\frac{5}{6} (1 + 6\xi)^4 s^2 \ln \left(\frac{s}{\mu_1^2} \right) + \left(\frac{s^2}{6} - tu \right) \ln \left(\frac{s}{\mu_2^2} \right) \right]. \quad (\text{A.38})$$

Here μ_1 and μ_2 are the energy scales at which α_1 and α_2 are taken to vanish, respectively. They correspond to the parameter choices of our theory. Next we consider the imaginary part. By the cutting rule, it is given by

$$\text{Im} \left[A_{1\text{-loop}}^{(ii \rightarrow jj)} \right] = \frac{N_s}{16M_P^4 s^2} \int \frac{d^3 l_1}{(2\pi)^3 2l_1^0} \frac{d^3 l_2}{(2\pi)^3 2l_2^0} (2\pi)^4 \delta^{(4)}(l_1 + l_2 - q) \\ \times [V_{\mu\nu}(p_1, p_2) P^{\mu\nu\rho\sigma} V_{\rho\sigma}(l_1, l_2)] [V_{\alpha\beta}(l_1, l_2) P^{\alpha\beta\gamma\delta} V_{\gamma\delta}(p_3, p_4)], \quad (\text{A.39})$$

where the momenta l_1 and l_2 are now on-shell, i.e. $l_1^2 = l_2^2 = 0$. The two-body phase space integral is reduced in the standard way as

$$\int \frac{d^3 l_1}{(2\pi)^3 2l_1^0} \frac{d^3 l_2}{(2\pi)^3 2l_2^0} (2\pi)^4 \delta^{(4)}(l_1 + l_2 - q) = \frac{1}{32\pi^2} \int d\Omega_l, \quad (\text{A.40})$$

where Ω_l is the solid angle between \vec{l}_1 and \vec{p}_1 . After performing the integral, we obtain

$$\text{Im} \left[A_{1\text{-loop}}^{(ii \rightarrow jj)} \right] = \frac{N_s}{960\pi M_P^4} \left[\frac{5}{6} (1 + 6\xi)^4 s^2 + \left(\frac{s^2}{6} - tu \right) \right]. \quad (\text{A.41})$$

By combining the real and imaginary parts, the one-loop amplitude is given by

$$A_{1\text{-loop}}^{(ii \rightarrow jj)} = -\frac{N_s}{960\pi^2 M_P^4} \left[\frac{5}{6} (1 + 6\xi)^4 s^2 \left(\ln \left(\frac{s}{\mu_1^2} \right) - i\pi \right) + \left(\frac{s^2}{6} - tu \right) \left(\ln \left(\frac{s}{\mu_2^2} \right) - i\pi \right) \right]. \quad (\text{A.42})$$

It again agrees with Ref. [25].

A.5 Renormalization group equation

The scalar contributions to the RG equations for α_1 and α_2 are readily derived from Eqs. (A.36) and (A.37). The operators R^2 and $R_{\mu\nu} R^{\mu\nu}$ have the mass dimension of 4 in the d -dimensional spacetime, and hence the bare couplings associated with these operators have the mass dimension of $d - 4 = -2\epsilon$. Since the bare couplings do not depend on μ , we obtain

$$\beta_{\alpha_1} = -\frac{N_s}{1152\pi^2} (1 + 6\xi)^2, \quad (\text{A.43})$$

$$\beta_{\alpha_2} = -\frac{N_s}{960\pi^2}, \quad (\text{A.44})$$

at the one-loop level. They coincide with, e.g., (the scalar parts of) Ref. [42], noting that their couplings f_0^2 and f_2^2 are related to α_1 and α_2 as $f_0^{-2} = 6\alpha_1$ and $f_2^{-2} = -\alpha_2$, respectively. The RG equations of α_1 and α_2 are also discussed, e.g. in Refs. [20, 21, 23, 41–44, 68–70] (although the sign of β_{α_1} is opposite in some references).

B Higgs scalaron inflationary model

In this appendix, we summarize basic properties of the Higgs scalaron inflationary model.

B.1 Einstein frame action and unitarity

We consider the following action:

$$S = \int d^4x \sqrt{-g_J} \left[\frac{M_P^2}{2} R_J + \alpha_1 R_J^2 + \frac{\xi}{2} R_J \tilde{\phi}_i^2 + \frac{1}{2} g_J^{\mu\nu} \partial_\mu \tilde{\phi}_i \partial_\nu \tilde{\phi}_i - V(\tilde{\phi}_i) \right], \quad (\text{B.1})$$

where $i = 1, 2, \dots, N_s$ and we add the subscript J for the quantities in the Jordan frame. With an auxiliary field $\tilde{\chi}$, the above action is equivalent to

$$S = \int d^4x \sqrt{-g_J} \left[\frac{M_P^2}{2} R_J \left(1 + \frac{\xi \tilde{\phi}_i^2 + 4\alpha_1 \tilde{\chi}}{M_P^2} \right) - \alpha_1 \tilde{\chi}^2 + \frac{1}{2} g_J^{\mu\nu} \partial_\mu \tilde{\phi}_i \partial_\nu \tilde{\phi}_i - V(\tilde{\phi}_i) \right]. \quad (\text{B.2})$$

One can indeed see that it returns to Eq. (B.1) after integrating out $\tilde{\chi}$. Now we perform the Weyl transformation,

$$g_{J\mu\nu} = \Omega^{-2} g_{E\mu\nu}, \quad \Omega^2 = 1 + \frac{\xi \tilde{\phi}_i^2 + 4\alpha_1 \tilde{\chi}}{M_P^2}, \quad (\text{B.3})$$

where the subscript E indicates the Einstein frame. The Ricci scalar is transformed as

$$R_J = \Omega^2 \left[R_E + \frac{3}{2} g_E^{\mu\nu} \partial_\mu \ln \Omega^2 \partial_\nu \ln \Omega^2 - 3 \square_E \ln \Omega^2 \right]. \quad (\text{B.4})$$

We redefine the fields as

$$s \equiv \sqrt{\frac{3}{2}} M_P \ln \Omega^2, \quad \phi_i \equiv \frac{\tilde{\phi}_i}{\Omega}. \quad (\text{B.5})$$

The action is given by

$$S = \int d^4x \sqrt{-g_E} \left[\frac{M_P^2}{2} R_E + \frac{1}{2} (\partial s)^2 + \frac{1}{2} \left(\partial \phi_i + \frac{1}{\sqrt{6}} \frac{\phi_i}{M_P} \partial s \right)^2 - \frac{M_P^4}{16\alpha_1} \left(1 - \exp \left(-\sqrt{\frac{2}{3}} \frac{s}{M_P} \right) - \frac{\xi \phi_i^2}{M_P^2} \right)^2 - \frac{V(\Omega \phi_i)}{\Omega^4} \right]. \quad (\text{B.6})$$

In particular, for the quartic potential $V(\tilde{\phi}_i) = \lambda \tilde{\phi}_i^4/4$, it is given by

$$S = \int d^4x \sqrt{-g_E} \left[\frac{M_P^2}{2} R_E + \frac{1}{2} (\partial s)^2 + \frac{1}{2} \left(\partial \phi_i + \frac{1}{\sqrt{6}} \frac{\phi_i}{M_P} \partial s \right)^2 - \frac{M_P^4}{16\alpha_1} \left(1 - \exp \left(-\sqrt{\frac{2}{3}} \frac{s}{M_P} \right) - \frac{\xi \phi_i^2}{M_P^2} \right)^2 - \frac{\lambda}{4} \phi_i^4 \right]. \quad (\text{B.7})$$

This theory is unitary up to the Planck scale since all the higher dimensional operators are suppressed by M_P . In particular, neither ξ nor α_1 shows up in the cut-off scale of the higher dimensional operators. Still the perturbativity of the couplings requires

$$\frac{\xi^2}{4\alpha_1} \lesssim 4\pi. \quad (\text{B.8})$$

These features are first pointed out in Ref. [22] and further studied in Ref. [24]. They can be seen in the Jordan frame (B.2) as follows. In order to discuss the cut-off scale and the size of the couplings, we first have to make the fields canonically normalized. The scalar fields $\tilde{\phi}_i$ are already canonical around $\tilde{\phi}_i = 0$,^{‡21} but $\tilde{\chi}$ is not canonical as it does not have the standard kinetic term in the Jordan frame. We have to redefine it as

$$\chi \equiv \frac{4\alpha_1 \tilde{\chi} + \xi \tilde{\phi}_i^2}{M_P}, \quad (\text{B.9})$$

because the kinetic term of $\tilde{\chi}$ is supplied from the non-minimal coupling to the Ricci scalar (that induces the kinetic mixing between $\tilde{\chi}$ and the scalar part of the metric).^{‡22} The couplings ξ and α_1 now enter the action only in the form

$$\mathcal{L}_\chi = -\frac{M_P^2}{16\alpha_1} \left(\chi - \frac{\xi \tilde{\phi}_i^2}{M_P} \right)^2, \quad (\text{B.10})$$

and hence they do not affect the cut-off scale. It is also seen that the perturbativity requires $\xi^2/4\alpha_1 \lesssim 4\pi$. The field redefinitions (B.3) and (B.5), and the resultant action (B.6), are a refined version of this argument, since the Weyl transformation is essentially solving the kinetic mixing between $\tilde{\chi}$ and the scalar part of the metric.

B.2 Inflationary predictions

Here we briefly summarize the inflationary prediction of the Higgs scalaron model. We may take the unitary gauge for the Higgs, assuming that the Higgs has a large field value during inflation. Then the action is given by Eq. (B.7) with $N_s = 1$. It contains two scalar fields, but effectively reduces to a single field model for $\xi, \alpha_1 \gg 1$, as the other mode is heavy in this limit.^{‡23} After integrating out the heavy mode, the inflaton potential is given by

$$U(\phi) = \frac{M_P^4}{4} \frac{1}{\xi^2/\lambda + 4\alpha_1} \left[1 - \exp \left(-\sqrt{\frac{2}{3}} \frac{\phi}{M_P} \right) \right]^2, \quad (\text{B.11})$$

where ϕ is the canonically normalized inflaton which is an admixture of the scalaron and the Higgs. The CMB normalization requires [4]

$$\frac{\xi^2}{\lambda} + 4\alpha_1 \simeq 2 \times 10^9. \quad (\text{B.12})$$

^{‡21} If we instead expand $\tilde{\phi}_i$ around $\tilde{\phi}_i \neq 0$, the fields $\tilde{\phi}_i$ mix with the scalar part of the metric. In this case, we have to make them canonically normalized as well, as in Ref. [10].

^{‡22} The field χ is canonical only up to an $\mathcal{O}(1)$ factor, but is enough to estimate the cut-off scale and the perturbativity condition.

^{‡23} We assume that λ , ξ and α_1 are all positive.

The spectral index n_s and tensor-to-scalar ratio r are given as

$$n_s \simeq 1 - \frac{2}{N_e}, \quad r \simeq \frac{12}{N_e^2}, \quad (\text{B.13})$$

where N_e is the number of e-folds after inflation. An interesting feature of this model is that in principle we can determine N_e precisely, as the reheating dynamics is determined from the couplings to the standard model particles that are known, and the cut-off scale is the Planck scale. Indeed, Refs. [71, 72] have studied the reheating dynamics of this model in detail. The former studies particle production at the first oscillation for a generic parameter space, while the latter finds that the reheating temperature is as high as 10^{15} GeV (or $N_e \simeq 59$) at least for a tuned parameter space. A complete study on the reheating dynamics of this model is yet to be done. Assuming that N_e is between 50 and 60, the spectral index is consistent with the current CMB observation [4], and the tensor-to-scalar ratio is within the reach of the future observations [29]. For more comprehensive analysis on the inflationary perturbation, see Refs. [22, 73–77].

B.3 Feynman rules around the vacuum

Here we derive the Feynman rules of the Higgs scalaron model around the vacuum $\phi_i = 0$ and $s = 0$ that are used in Sec. 3.3. We expand the action (B.6) around $\phi_i = 0$ and $s = 0$ as

$$S \simeq \int d^4x \sqrt{-g} \left[\frac{M_P^2}{2} R + \frac{1}{2} \left((\partial s)^2 + (\partial \phi_i)^2 + \frac{\partial \phi_i^2 \partial s}{\sqrt{6} M_P} \right) - \frac{m_s^2}{2} s^2 + \frac{\xi m_s}{\sqrt{8\alpha_1}} s \phi_i^2 - \frac{\xi^2}{16\alpha_1} \phi_i^4 \right], \quad (\text{B.14})$$

where we have defined the scalaron mass as

$$m_s^2 = \frac{M_P^2}{12\alpha_1}, \quad (\text{B.15})$$

and we have retained only the terms relevant for our computation. As advertised, the scalaron mass squared is inversely proportional to α_1 . The Feynman rules for the scalaron and ϕ_i are easily read off from this action. We take the de Donder gauge in the Einstein frame in this subsection. The propagators for ϕ_i and the graviton are then given by Eqs. (A.11) and (A.14), respectively. The propagator for the scalaron s is

$$\text{—————} = \frac{i}{p^2 - m_s^2 + i\epsilon}. \quad (\text{B.16})$$

We denote the scalaron propagator by a solid line to distinguish it from ϕ_i . The scalar-scalar-scalar vertex is given by

$$\begin{array}{c} p_{1,i} \\ \text{---} \\ \text{---} \\ p_{2,j} \end{array} \text{---} = i \left[\frac{\xi m_s}{\sqrt{2\alpha_1}} + \frac{1}{\sqrt{6} M_P} (p_1 + p_2)^2 \right] \delta_{ij}, \quad (\text{B.17})$$

where we have included the factor 2 from the exchange of the scalar fields, and the momenta are both outgoing or both incoming. The scalar four-point vertex is given by

$$\begin{array}{c} p_1 & & p_3 \\ & \diagdown & / \\ & \times & \\ & / & \diagdown \\ p_2 & & p_4 \end{array} = \begin{cases} -i \frac{3\xi^2}{2\alpha_1} & \text{for } \phi_i\phi_i \rightarrow \phi_i\phi_i, \\ -i \frac{\xi^2}{2\alpha_1} & \text{for } \phi_i\phi_i \rightarrow \phi_j\phi_j \text{ or } \phi_i\phi_j \rightarrow \phi_i\phi_j \text{ with } i \neq j, \end{cases} \quad (\text{B.18})$$

where we have included the combinatorial factors, and ignored the contribution from the Higgs potential that is irrelevant for our discussion. Finally the scalar-scalar-graviton vertex is given by Eq. (A.17) with $\xi = 0$.

References

- [1] T. Futamase and K.-i. Maeda, “Chaotic Inflationary Scenario in Models Having Nonminimal Coupling With Curvature,” *Phys. Rev.* **D39** (1989) 399–404.
- [2] J. L. Cervantes-Cota and H. Dehnen, “Induced gravity inflation in the standard model of particle physics,” *Nucl. Phys.* **B442** (1995) 391–412, [arXiv:astro-ph/9505069 \[astro-ph\]](#).
- [3] F. L. Bezrukov and M. Shaposhnikov, “The Standard Model Higgs boson as the inflaton,” *Phys. Lett.* **B659** (2008) 703–706, [arXiv:0710.3755 \[hep-th\]](#).
- [4] **Planck** Collaboration, Y. Akrami *et al.*, “Planck 2018 results. X. Constraints on inflation,” [arXiv:1807.06211 \[astro-ph.CO\]](#).
- [5] C. P. Burgess, H. M. Lee, and M. Trott, “Power-counting and the Validity of the Classical Approximation During Inflation,” *JHEP* **09** (2009) 103, [arXiv:0902.4465 \[hep-ph\]](#).
- [6] J. L. F. Barbon and J. R. Espinosa, “On the Naturalness of Higgs Inflation,” *Phys. Rev.* **D79** (2009) 081302, [arXiv:0903.0355 \[hep-ph\]](#).
- [7] C. P. Burgess, H. M. Lee, and M. Trott, “Comment on Higgs Inflation and Naturalness,” *JHEP* **07** (2010) 007, [arXiv:1002.2730 \[hep-ph\]](#).
- [8] M. P. Hertzberg, “On Inflation with Non-minimal Coupling,” *JHEP* **11** (2010) 023, [arXiv:1002.2995 \[hep-ph\]](#).
- [9] A. Kehagias, A. Moradinezhad Dizgah, and A. Riotto, “Remarks on the Starobinsky model of inflation and its descendants,” *Phys. Rev.* **D89** no. 4, (2014) 043527, [arXiv:1312.1155 \[hep-th\]](#).
- [10] F. Bezrukov, A. Magnin, M. Shaposhnikov, and S. Sibiryakov, “Higgs inflation: consistency and generalisations,” *JHEP* **01** (2011) 016, [arXiv:1008.5157 \[hep-ph\]](#).
- [11] R. N. Lerner and J. McDonald, “A Unitarity-Conserving Higgs Inflation Model,” *Phys. Rev.* **D82** (2010) 103525, [arXiv:1005.2978 \[hep-ph\]](#).

- [12] G. F. Giudice and H. M. Lee, “Unitarizing Higgs Inflation,” *Phys. Lett.* **B694** (2011) 294–300, [arXiv:1010.1417 \[hep-ph\]](#).
- [13] M. P. Hertzberg, “Can Inflation be Connected to Low Energy Particle Physics?,” *JCAP* **1208** (2012) 008, [arXiv:1110.5650 \[hep-ph\]](#).
- [14] J. L. F. Barbon, J. A. Casas, J. Elias-Miro, and J. R. Espinosa, “Higgs Inflation as a Mirage,” *JHEP* **09** (2015) 027, [arXiv:1501.02231 \[hep-ph\]](#).
- [15] Y. Ema, R. Jinno, K. Mukaida, and K. Nakayama, “Violent Preheating in Inflation with Nonminimal Coupling,” *JCAP* **1702** no. 02, (2017) 045, [arXiv:1609.05209 \[hep-ph\]](#).
- [16] M. P. DeCross, D. I. Kaiser, A. Prabhu, C. Prescod-Weinstein, and E. I. Sfakianakis, “Preheating after Multifield Inflation with Nonminimal Couplings, I: Covariant Formalism and Attractor Behavior,” *Phys. Rev.* **D97** no. 2, (2018) 023526, [arXiv:1510.08553 \[astro-ph.CO\]](#).
- [17] E. I. Sfakianakis and J. van de Vis, “Preheating after Higgs Inflation: Self-Resonance and Gauge boson production,” *Phys. Rev.* **D99** no. 8, (2019) 083519, [arXiv:1810.01304 \[hep-ph\]](#).
- [18] G. ’t Hooft and M. J. G. Veltman, “One loop divergencies in the theory of gravitation,” *Ann. Inst. H. Poincare Phys. Theor.* **A20** (1974) 69–94.
- [19] J. F. Donoghue, “Introduction to the effective field theory description of gravity,” in *Advanced School on Effective Theories Almunecar, Spain, June 25-July 1, 1995*. 1995. [arXiv:gr-qc/9512024 \[gr-qc\]](#).
- [20] A. Salvio and A. Mazumdar, “Classical and Quantum Initial Conditions for Higgs Inflation,” *Phys. Lett.* **B750** (2015) 194–200, [arXiv:1506.07520 \[hep-ph\]](#).
- [21] X. Calmet and I. Kuntz, “Higgs Starobinsky Inflation,” *Eur. Phys. J.* **C76** no. 5, (2016) 289, [arXiv:1605.02236 \[hep-th\]](#).
- [22] Y. Ema, “Higgs Scalon Mixed Inflation,” *Phys. Lett.* **B770** (2017) 403–411, [arXiv:1701.07665 \[hep-ph\]](#).
- [23] D. M. Ghilencea, “Two-loop corrections to Starobinsky-Higgs inflation,” *Phys. Rev.* **D98** no. 10, (2018) 103524, [arXiv:1807.06900 \[hep-ph\]](#).
- [24] D. Gorbunov and A. Tokareva, “Scaloron the healer: removing the strong-coupling in the Higgs- and Higgs-dilaton inflations,” *Phys. Lett.* **B788** (2019) 37–41, [arXiv:1807.02392 \[hep-ph\]](#).
- [25] U. Aydemir, M. M. Anber, and J. F. Donoghue, “Self-healing of unitarity in effective field theories and the onset of new physics,” *Phys. Rev.* **D86** (2012) 014025, [arXiv:1203.5153 \[hep-ph\]](#).
- [26] X. Calmet and R. Casadio, “Self-healing of unitarity in Higgs inflation,” *Phys. Lett.* **B734** (2014) 17–20, [arXiv:1310.7410 \[hep-ph\]](#).

- [27] Y. Hamada, H. Kawai, K.-y. Oda, and S. C. Park, “Higgs Inflation is Still Alive after the Results from BICEP2,” *Phys. Rev. Lett.* **112** no. 24, (2014) 241301, [arXiv:1403.5043 \[hep-ph\]](#).
- [28] F. Bezrukov and M. Shaposhnikov, “Higgs inflation at the critical point,” *Phys. Lett.* **B734** (2014) 249–254, [arXiv:1403.6078 \[hep-ph\]](#).
- [29] **CMB-S4** Collaboration, K. N. Abazajian *et al.*, “CMB-S4 Science Book, First Edition,” [arXiv:1610.02743 \[astro-ph.CO\]](#).
- [30] A. O. Barvinsky, A. Yu. Kamenshchik, and A. A. Starobinsky, “Inflation scenario via the Standard Model Higgs boson and LHC,” *JCAP* **0811** (2008) 021, [arXiv:0809.2104 \[hep-ph\]](#).
- [31] A. De Simone, M. P. Hertzberg, and F. Wilczek, “Running Inflation in the Standard Model,” *Phys. Lett.* **B678** (2009) 1–8, [arXiv:0812.4946 \[hep-ph\]](#).
- [32] F. L. Bezrukov, A. Magnin, and M. Shaposhnikov, “Standard Model Higgs boson mass from inflation,” *Phys. Lett.* **B675** (2009) 88–92, [arXiv:0812.4950 \[hep-ph\]](#).
- [33] F. Bezrukov and M. Shaposhnikov, “Standard Model Higgs boson mass from inflation: Two loop analysis,” *JHEP* **07** (2009) 089, [arXiv:0904.1537 \[hep-ph\]](#).
- [34] A. O. Barvinsky, A. Yu. Kamenshchik, C. Kiefer, A. A. Starobinsky, and C. Steinwachs, “Asymptotic freedom in inflationary cosmology with a non-minimally coupled Higgs field,” *JCAP* **0912** (2009) 003, [arXiv:0904.1698 \[hep-ph\]](#).
- [35] A. O. Barvinsky, A. Yu. Kamenshchik, C. Kiefer, A. A. Starobinsky, and C. F. Steinwachs, “Higgs boson, renormalization group, and naturalness in cosmology,” *Eur. Phys. J.* **C72** (2012) 2219, [arXiv:0910.1041 \[hep-ph\]](#).
- [36] K. Allison, “Higgs xi-inflation for the 125-126 GeV Higgs: a two-loop analysis,” *JHEP* **02** (2014) 040, [arXiv:1306.6931 \[hep-ph\]](#).
- [37] D. P. George, S. Mooij, and M. Postma, “Quantum corrections in Higgs inflation: the Standard Model case,” *JCAP* **1604** no. 04, (2016) 006, [arXiv:1508.04660 \[hep-th\]](#).
- [38] J. Fumagalli and M. Postma, “UV (in)sensitivity of Higgs inflation,” *JHEP* **05** (2016) 049, [arXiv:1602.07234 \[hep-ph\]](#).
- [39] T. Markkanen, T. Tenkanen, V. Vaskonen, and H. Veermäe, “Quantum corrections to quartic inflation with a non-minimal coupling: metric vs. Palatini,” *JCAP* **1803** no. 03, (2018) 029, [arXiv:1712.04874 \[gr-qc\]](#).
- [40] I. D. Saltas, “Higgs inflation and quantum gravity: An exact renormalisation group approach,” *JCAP* **1602** (2016) 048, [arXiv:1512.06134 \[hep-th\]](#).
- [41] E. S. Fradkin and A. A. Tseytlin, “Renormalizable asymptotically free quantum theory of gravity,” *Nucl. Phys.* **B201** (1982) 469–491.
- [42] A. Salvio and A. Strumia, “Agravity,” *JHEP* **06** (2014) 080, [arXiv:1403.4226 \[hep-ph\]](#).

- [43] T. Markkanen, S. Nurmi, A. Rajantie, and S. Stopyra, “The 1-loop effective potential for the Standard Model in curved spacetime,” *JHEP* **06** (2018) 040, [arXiv:1804.02020 \[hep-ph\]](#).
- [44] A. Codello and R. K. Jain, “On the covariant formalism of the effective field theory of gravity and leading order corrections,” *Class. Quant. Grav.* **33** no. 22, (2016) 225006, [arXiv:1507.06308 \[gr-qc\]](#).
- [45] A. A. Starobinsky, “A New Type of Isotropic Cosmological Models Without Singularity,” *Phys. Lett.* **B91** (1980) 99–102. [771(1980)].
- [46] J. D. Barrow and A. C. Ottewill, “The Stability of General Relativistic Cosmological Theory,” *J. Phys.* **A16** (1983) 2757.
- [47] B. Whitt, “Fourth Order Gravity as General Relativity Plus Matter,” *Phys. Lett.* **145B** (1984) 176–178.
- [48] J. D. Barrow and S. Cotsakis, “Inflation and the Conformal Structure of Higher Order Gravity Theories,” *Phys. Lett.* **B214** (1988) 515–518.
- [49] J. Julve and M. Tonin, “Quantum Gravity with Higher Derivative Terms,” *Nuovo Cim.* **B46** (1978) 137–152.
- [50] A. Salvio, “Quadratic Gravity,” *Front.in Phys.* **6** (2018) 77, [arXiv:1804.09944 \[hep-th\]](#).
- [51] T. Han and S. Willenbrock, “Scale of quantum gravity,” *Phys. Lett.* **B616** (2005) 215–220, [arXiv:hep-ph/0404182 \[hep-ph\]](#).
- [52] Y. Ema, R. Jinno, K. Mukaida, and K. Nakayama, “Gravitational Effects on Inflaton Decay,” *JCAP* **1505** no. 05, (2015) 038, [arXiv:1502.02475 \[hep-ph\]](#).
- [53] Y. Ema, R. Jinno, K. Mukaida, and K. Nakayama, “Gravitational particle production in oscillating backgrounds and its cosmological implications,” *Phys. Rev.* **D94** no. 6, (2016) 063517, [arXiv:1604.08898 \[hep-ph\]](#).
- [54] D. P. George, S. Mooij, and M. Postma, “Quantum corrections in Higgs inflation: the real scalar case,” *JCAP* **1402** (2014) 024, [arXiv:1310.2157 \[hep-th\]](#).
- [55] M. Postma and M. Volponi, “Equivalence of the Einstein and Jordan frames,” *Phys. Rev.* **D90** no. 10, (2014) 103516, [arXiv:1407.6874 \[astro-ph.CO\]](#).
- [56] A. Karam, T. Pappas, and K. Tamvakis, “Frame-dependence of higher-order inflationary observables in scalar-tensor theories,” *Phys. Rev.* **D96** no. 6, (2017) 064036, [arXiv:1707.00984 \[gr-qc\]](#).
- [57] F. Takahashi, “Linear Inflation from Running Kinetic Term in Supergravity,” *Phys. Lett.* **B693** (2010) 140–143, [arXiv:1006.2801 \[hep-ph\]](#).
- [58] K. Nakayama and F. Takahashi, “Running Kinetic Inflation,” *JCAP* **1011** (2010) 009, [arXiv:1008.2956 \[hep-ph\]](#).
- [59] R. Kallosh, A. Linde, and D. Roest, “Superconformal Inflationary α -Attractors,” *JHEP* **11** (2013) 198, [arXiv:1311.0472 \[hep-th\]](#).

- [60] J. Garcia-Bellido, J. Rubio, M. Shaposhnikov, and D. Zenhausern, “Higgs-Dilaton Cosmology: From the Early to the Late Universe,” *Phys. Rev.* **D84** (2011) 123504, [arXiv:1107.2163 \[hep-ph\]](#).
- [61] F. Bauer and D. A. Demir, “Higgs-Palatini Inflation and Unitarity,” *Phys. Lett.* **B698** (2011) 425–429, [arXiv:1012.2900 \[hep-ph\]](#).
- [62] S. Rasanen and P. Wahlman, “Higgs inflation with loop corrections in the Palatini formulation,” *JCAP* **1711** no. 11, (2017) 047, [arXiv:1709.07853 \[astro-ph.CO\]](#).
- [63] I. Antoniadis, A. Karam, A. Lykkas, and K. Tamvakis, “Palatini inflation in models with an R^2 term,” *JCAP* **1811** no. 11, (2018) 028, [arXiv:1810.10418 \[gr-qc\]](#).
- [64] I. Antoniadis, A. Karam, A. Lykkas, T. Pappas, and K. Tamvakis, “Rescuing Quartic and Natural Inflation in the Palatini Formalism,” *JCAP* **1903** no. 03, (2019) 005, [arXiv:1812.00847 \[gr-qc\]](#).
- [65] T. Tenkanen, “Minimal Higgs inflation with an R^2 term in Palatini gravity,” *Phys. Rev.* **D99** no. 6, (2019) 063528, [arXiv:1901.01794 \[astro-ph.CO\]](#).
- [66] K. Falls and M. Herrero-Valea, “Frame (In)equivalence in Quantum Field Theory and Cosmology,” *Eur. Phys. J.* **C79** no. 7, (2019) 595, [arXiv:1812.08187 \[hep-th\]](#).
- [67] J. Ellis, “TikZ-Feynman: Feynman diagrams with TikZ,” *Comput. Phys. Commun.* **210** (2017) 103–123, [arXiv:1601.05437 \[hep-ph\]](#).
- [68] E. Elizalde and S. D. Odintsov, “Renormalization group improved effective potential for gauge theories in curved space-time,” *Phys. Lett.* **B303** (1993) 240–248, [arXiv:hep-th/9302074 \[hep-th\]](#). [Russ. Phys. J.37,25(1994)].
- [69] E. Elizalde and S. D. Odintsov, “Renormalization group improved effective Lagrangian for interacting theories in curved space-time,” *Phys. Lett.* **B321** (1994) 199–204, [arXiv:hep-th/9311087 \[hep-th\]](#).
- [70] R. Myrzakulov, S. Odintsov, and L. Sebastiani, “Inflationary universe from higher derivative quantum gravity coupled with scalar electrodynamics,” *Nucl. Phys.* **B907** (2016) 646–663, [arXiv:1604.06088 \[hep-th\]](#).
- [71] M. He, R. Jinno, K. Kamada, S. C. Park, A. A. Starobinsky, and J. Yokoyama, “On the violent preheating in the mixed Higgs- R^2 inflationary model,” *Phys. Lett.* **B791** (2019) 36–42, [arXiv:1812.10099 \[hep-ph\]](#).
- [72] F. Bezrukov, D. Gorbunov, C. Shepherd, and A. Tokareva, “Some like it hot: R^2 heals Higgs inflation, but does not cool it,” [arXiv:1904.04737 \[hep-ph\]](#).
- [73] Y.-C. Wang and T. Wang, “Primordial perturbations generated by Higgs field and R^2 operator,” *Phys. Rev.* **D96** no. 12, (2017) 123506, [arXiv:1701.06636 \[gr-qc\]](#).
- [74] S. Pi, Y.-l. Zhang, Q.-G. Huang, and M. Sasaki, “Scalaron from R^2 -gravity as a heavy field,” *JCAP* **1805** no. 05, (2018) 042, [arXiv:1712.09896 \[astro-ph.CO\]](#).

- [75] M. He, A. A. Starobinsky, and J. Yokoyama, “Inflation in the mixed Higgs- R^2 model,” *JCAP* **1805** no. 05, (2018) 064, [arXiv:1804.00409 \[astro-ph.CO\]](#).
- [76] A. Gundhi and C. F. Steinwachs, “Scalaron-Higgs inflation,” [arXiv:1810.10546 \[hep-th\]](#).
- [77] V.-M. Enckell, K. Enqvist, S. Rasanen, and L.-P. Wahlman, “Higgs- R^2 inflation - full slow-roll study at tree-level,” [arXiv:1812.08754 \[astro-ph.CO\]](#).