

DEUTSCHES ELEKTRONEN-SYNCHROTRON
Ein Forschungszentrum der Helmholtz-Gemeinschaft



DESY 19-180
CERN-TH-2019-204
IPPP/19/86
UUITP-48/19
SLAC-PUB-17491
arXiv:1912.00188
November 2019

All-Order Amplitudes at Any Multiplicity in the Multi-Regge Limit

V. Del Duca

*Institut für Theoretische Physik, ETH Zürich, Switzerland
and*

INFN, Laboratori Nazionali di Frascati, Italy

S. Drube, J. M. Drummond

School of Physics and Astronomy, University of Southampton, UK

C. Duhr

CERN, Geneva, Switzerland

F. Dulat

SLAC, Stanford University, USA

R. Marzucca

IPPP, Department of Physics, University of Durham, UK

G. Papathanasiou

Deutsches Elektronen-Synchrotron DESY, Hamburg

B. Verbeek

Department of Physics and Astronomy, Uppsala University, Sweden

ISSN 0418-9833

NOTKESTRASSE 85 – 22607 HAMBURG

DESY behält sich alle Rechte für den Fall der Schutzrechtserteilung und für die wirtschaftliche Verwertung der in diesem Bericht enthaltenen Informationen vor.

DESY reserves all rights for commercial use of information included in this report, especially in case of filing application for or grant of patents.

To be sure that your reports and preprints are promptly included in the
HEP literature database
send them to (if possible by air mail):

DESY Zentralbibliothek Notkestraße 85 22607 Hamburg Germany	DESY Bibliothek Platanenallee 6 15738 Zeuthen Germany
---	---

All-order amplitudes at any multiplicity in the multi-Regge limit

V. Del Duca^{1,2}, S. Druc³, J. M. Drummond³, C. Duhr⁴, F. Dulat⁵, R. Marzucca⁶, G. Papathanasiou⁷, B. Verbeek⁸

¹*Institute for Theoretical Physics, ETH Zürich, 8093 Zürich, Switzerland*

²*INFN, Laboratori Nazionali di Frascati, 00044 Frascati (RM), Italy*

³*School of Physics and Astronomy, University of Southampton, Highfield, SO17 1BJ, United Kingdom*

⁴*Theoretical Physics Department, CERN, CH-1211 Geneva 23, Switzerland*

⁵*SLAC National Accelerator Laboratory, Stanford University, Stanford, CA 94309, USA*

⁶*IPPP, Department of Physics, Durham University, Durham DH1 3LE, United Kingdom*

⁷*DESY Theory Group, DESY Hamburg, Notkestraße 85, D-22607 Hamburg, Germany*

⁸*Department of Physics and Astronomy, Uppsala University, 75108 Uppsala, Sweden*

We propose an all-loop expression for scattering amplitudes in planar $\mathcal{N} = 4$ super Yang-Mills theory in multi-Regge kinematics valid for all multiplicities, all helicity configurations and arbitrary logarithmic accuracy. Our expression is arrived at from comparing explicit perturbative results with general expectations from the integrable structure of a closely related collinear limit. A crucial ingredient of the analysis is an all-order extension for the central emission vertex that we recently computed at next-to-leading logarithmic accuracy. As an application, we use our all-order formula to prove that all amplitudes in this theory in multi-Regge kinematics are single-valued multiple polylogarithms of uniform transcendental weight.

Recent years have seen tremendous progress in our understanding of multi-loop multi-leg scattering amplitudes in planar $\mathcal{N} = 4$ Super Yang-Mills (SYM) theory. Its S -matrix exhibits a hidden dual conformal (DC) symmetry [1], which closes with the ordinary conformal symmetry into a Yangian algebra [2].

The DC symmetry is broken by infrared (IR) divergences. Such divergences are universal and independent of the hard scattering process and it is possible to construct DC-invariant functions by considering ratios where all IR-divergences cancel. We denote by \mathcal{R}_N the IR-finite ratio of the N -point color-ordered amplitude and the Bern-Dixon-Smirnov (BDS) amplitude [3], defined (loosely) as the exponential of the one-loop amplitude multiplied by the cusp anomalous dimension Γ_{cusp} [4]. DC-invariance dictates that \mathcal{R}_N only depends on $3N - 15$ independent cross-ratios. In particular, \mathcal{R}_N is trivial for $N \leq 5$ [5], and is known analytically in general kinematics for $N = 6$ through seven loops [6–17] and for $N = 7$ through four loops [18–22].

Explicit data for small N reveals that the perturbative expansion of \mathcal{R}_N can often be expressed in terms of a class of iterated integrals known as *multiple polylogarithms* (MPLs) [23]. Moreover only MPLs of (transcendental) weight $2L$ contribute to an L -loop amplitude, where weight is the number of iterated integrations.

The mathematical beauty and simplicity of the available perturbative results hint at some deeper structure governing amplitudes in planar $\mathcal{N} = 4$ SYM. This is corroborated by the fact that infinite-dimensional symmetries, like the Yangian symmetry of $\mathcal{N} = 4$ SYM, are a hallmark of integrability. One should then be able to




Figure 1. Fourier-Mellin factorisation of $2 \rightarrow N - 2$ gluon amplitude in multi-Regge kinematics.

compute \mathcal{R}_N at any value of the coupling. A major step in this direction was taken in [24–28], where it was argued that amplitudes (or their dual Wilson loops [29–33]) can be computed through an integrable flux-tube picture. The dream of computing amplitudes analytically at any value of the coupling constant g^2 , or at least at any order in perturbation theory, has not yet been achieved.

Here we present for the first time a way to compute scattering amplitudes in planar $\mathcal{N} = 4$ SYM to any order in the coupling, for any helicity configuration and any number of external legs, albeit in the simplified kinematic setup of multi-Regge kinematics (MRK) where the produced particles are strongly ordered in rapidity and have comparable transverse momenta. While in Euclidean kinematics the ratios \mathcal{R}_N become trivial in the limit [34–39], they develop a non-trivial kinematic de-

pendence when some of the energies of the produced gluons are analytically continued to negative values [35, 37]. Here we focus on the situation where all the centrally-produced gluons have a negative energy, and we propose a formula for any amplitude in MRK in this theory.

THE N -PARTICLE DISPERSION INTEGRAL

In MRK a subset of $N - 5$ cross-ratios, denoted by τ_i , approach zero. \mathcal{R}_N can then be expressed at each order as a polynomial in large logarithms $\log \tau_i$, multiplied by functions of the $2N - 10$ remaining real degrees of freedom. The latter are conveniently described by $N - 5$ complex variables z_i , see [40] and references therein for these standard conventions. We conjecture that, to all orders, \mathcal{R}_N can be written as a Fourier-Mellin (FM) integral with a factorised form, as also depicted in fig. 1,

$$\frac{\mathcal{R}_N e^{i\Gamma\delta}}{2\pi i} = \prod_{r=1}^{N-5} \left[\sum_{n_r} \left(\frac{z_r}{\bar{z}_r} \right)^{\frac{n_r}{2}} \int_{\mathcal{C}} \frac{d\nu_r}{2\pi} \frac{|z_r|^{2i\nu_r} \tilde{\Phi}_r}{(-\tau_r + i0)^{\omega_r}} \right] \times I_1^{h_1} \tilde{C}_{12}^{h_2} \dots \tilde{C}_{N-6, N-5}^{h_{N-5}} \bar{I}_{N-5}^{h_{N-4}}. \quad (1)$$

Equation (1) extends similar formulas in the literature for restricted subsets of amplitudes at leading logarithmic accuracy (LLA) and beyond [37, 40–46], see also [47] for an application. The ratio \mathcal{R}_N depends on the helicities h_r of all centrally-produced particles. The building blocks in the integrand are functions of the FM variables (ν_r, n_r) , and we use a shorthand notation $\omega_r = \omega(\nu_r, n_r)$ and $\tilde{C}_{r, r+1}^{h_{r+1}} = \tilde{C}^{h_{r+1}}(\nu_r, n_r, \nu_{r+1}, n_{r+1})$ etc. The phase $e^{i\Gamma\delta}$, where $\Gamma \equiv \Gamma_{\text{cusp}}/4$, captures terms in the BDS amplitude that do not vanish after analytic continuation in MRK [48].

In the limit where one of the centrally-produced gluons becomes soft, \mathcal{R}_N should reduce to \mathcal{R}_{N-1} . Provided that the building blocks have at most simple poles on the integration axis, this then dictates that the contour \mathcal{C} must take the form shown in fig. 2, and implies the following *exact bootstrap conditions* [46, 49],

$$\omega(\pm\pi\Gamma, 0) = 0, \quad \text{Res}_{\nu=\pm\pi\Gamma} \left(\tilde{\Phi}(\nu, 0) \right) = \pm \frac{1}{2\pi}, \quad (2)$$

$$\tilde{C}^h(\pi\Gamma, 0, \nu_2, n_2) = 2\pi i I^h(\nu_2, n_2), \quad (3)$$

$$\tilde{C}^h(\nu_1, n_1, -\pi\Gamma, 0) = -2\pi i \bar{I}^h(\nu_1, n_1), \quad (4)$$

$$\text{Res}_{\nu_1=\nu_2} \tilde{C}^h(\nu_1, n_2, \nu_2, n_2) = \frac{-i(-1)^{n_2} e^{i\pi\omega(\nu_2, n_2)}}{\tilde{\Phi}(\nu_2, n_2)}, \quad (5)$$

$$\tilde{C}^h(-\pi\Gamma, 0, \nu_2, n_2) = \tilde{C}^h(\nu_1, n_1, \pi\Gamma, 0) = 0. \quad (6)$$

Let us now proceed to fully specify the integral (1), by providing explicit expressions for its building blocks. The quantities known as BFKL eigenvalue ω_r , impact factor product $\tilde{\Phi}_r$ and helicity flip kernel I_r have already been determined to all loops [28], by means of an analytic




Figure 2. Contour of integration \mathcal{C} for the integral (1), with $\nu_{N-4} = -\pi\Gamma$, $\nu_0 = \pi\Gamma$ corresponding to the boundary cases.

continuation from the collinear limit. The latter limit is also described by a dispersion integral very similar to (1), whose building blocks are governed by an integrable flux tube, and may thus be computed at finite coupling within the Pentagon operator product expansion (OPE) [24–27] approach. Then, the authors of [28] were able to connect the multi-Regge and collinear integrands by analytically continuing in the integration variable, and in particular obtain ω_r , $\tilde{\Phi}_r$ and I_r from their OPE counterparts, the gluonic excitation energy, measure and NMHV impact factor respectively. A feature of this analysis is that at finite coupling it is more natural to use rapidities u_r rather than ν_r as integration variables, giving rise to the following implicit all-loop dispersion relation,

$$\nu_r = u_r - 2g(\mathbb{Q} \cdot \mathbb{M} \cdot \tilde{\kappa})_1, \quad \omega_r = -4g(\mathbb{Q} \cdot \mathbb{M} \cdot \kappa)_1. \quad (7)$$

The sources κ and $\tilde{\kappa}$ are infinite-dimensional vectors and are described explicitly in the Appendix along with the matrices \mathbb{Q} and \mathbb{M} which essentially encode the Beisert-Eden-Staudacher kernel [4, 50]. The subscript 1 in (7) means the first component of the vector.

CENTRAL EMISSION VERTEX

The only quantity in (1) only known at leading order (LO) [43] and next-to-LO [46] is the central emission vertex $C_{r, r+1}^\pm$. A main result of this paper is a conjecture for $C_{r, r+1}^\pm$ to all orders in the coupling, as we now move on to describe. We focus on the vertex for the emission of a positive helicity gluon. The case of negative helicity is then recovered from the helicity flip kernel [51],

$$\tilde{C}_{r, r+1}^- = \tilde{C}_{r, r+1}^+ \bar{I}_r^- I_{r+1}^-. \quad (8)$$

Our analysis parallels that of [28] for $N = 6$. We assume that also for $N = 7$, the dispersion integral (1) can be obtained by analytically continuing the contribution of gluon excitations to the pentagon OPE through the branch cut at $u_r = -in_r/2 \pm 2g$ in the rapidity plane. It follows that the central emission vertex is the analytic continuation of the new OPE building block appearing

at this multiplicity, known as the *gluon pentagon transition* [27]. Performing the analytic continuation in full generality is quite complicated, but we are able to present a conjectural all-orders form for the central emission vertex by continuing certain factors of the pentagon transition, and fixing the remaining proportionality coefficient by consistency with known perturbative data in MRK. More precisely, our conjecture reads

$$\tilde{C}_{12}^+ = \frac{\tilde{C}_{12}^{(0)}}{g^2} h_{12} Z_{12} \exp(f_{12} - f_{\bar{1}\bar{2}} - i f_{\bar{1}2} + i f_{1\bar{2}} - A). \quad (9)$$

Here $\tilde{C}_{12}^{(0)}$ denotes the LO central emission vertex of ref. [43], with the ν_r replaced with the rapidities u_r ,

$$\tilde{C}_{12}^{(0)} = \frac{\Gamma(1-iu_1-\frac{n_1}{2})\Gamma(1+iu_2+\frac{n_2}{2})\Gamma(iu_1-iu_2-\frac{n_1-n_2}{2})}{\Gamma(iu_1-\frac{n_1}{2})\Gamma(-iu_2+\frac{n_2}{2})\Gamma(1-iu_1+iu_2-\frac{n_1-n_2}{2})}. \quad (10)$$

The exponential factor and Z_{12} in (9) are obtained by analytically continuing the corresponding functions appearing in the pentagon transition [52]. The functions f_{rs} are given by

$$f_{rs} = 4\kappa(u_r, n_r) \cdot \mathbb{Q} \cdot \mathbb{M} \cdot \kappa(u_s, n_s), \quad (11)$$

similarly $f_{\bar{r}\bar{s}}$ ($f_{\bar{r}s}$) for $\kappa_r \rightarrow \bar{\kappa}_r$ (and $\kappa_s \rightarrow \bar{\kappa}_s$), in terms of the same sources appearing in (32). The constant A is given by

$$A = 2 \int_0^\infty \frac{dt}{t} \frac{1 - J_0(2gt)^2}{e^t - 1} - \pi^2 \Gamma. \quad (12)$$

For Z_{12} we have

$$Z_{12} = \sqrt{\frac{(x_1^- x_2^- - g^2)(x_1^+ x_2^+ - g^2)}{(x_1^+ x_2^- - g^2)(x_1^- x_2^+ - g^2)}}, \quad (13)$$

where we introduce the Zhukowski variables

$$x_r^\pm = x(u_r \pm i\frac{n_r}{2}), \quad x(u_r) = \frac{1}{2}(u_r + \sqrt{u_r^2 - 4g^2}). \quad (14)$$

The quantity h_{12} in (9) collects all the factors we have not addressed so far, and is a priori unknown. Nevertheless, it is constrained by the exact bootstrap condition (5) to be free of poles at $u_r = u_s$, and this condition also fixes the value of h_{12} at $(u_2, n_2) = (u_1, n_1)$ to be

$$\begin{aligned} h_{12|(u_2, n_2)=(u_1, n_1)} &= \frac{x^+ x^-}{u_1^2 + \frac{n_1^2}{4}} e^{i\pi\omega_1} \\ &= e^2 \int_0^\infty \frac{dt}{t} [1 - J_0(2gt)] \cos(u_1 t) e^{-\frac{n_1}{2}t + i\pi\omega_1}. \end{aligned} \quad (15)$$

There could be many functions h_{12} that satisfy (15), but there is a particularly simple solution where h_{12} takes a factorised form,

$$h_{12} = h_1 \check{h}_2, \quad \check{h}(u, n) = h(-u, -n). \quad (16)$$

This form is motivated by the fact that it reproduces the perturbative expansion of the same quantity to three

loops, extracted from the corresponding seven-particle MHV amplitude [20] with the method described in [46]. We conjecture that this minimal form persists to all orders in perturbation theory. Inserting the factorised form into (15), we find

$$h_1 = \sqrt{\frac{x^+ x^-}{u_1^2 + \frac{n_1^2}{4}}} e^{\frac{i}{2}\pi\omega_1} h_{o1}, \quad \check{h}_{o1} = h_{o1}^{-1}. \quad (17)$$

The remaining freedom h_{o1} can be determined by solving the exact bootstrap condition (4) order-by-order in perturbation theory. We observe empirically that the perturbative expansion of h_{o1} is consistent with an exponential form for h_{o1} very reminiscent of (15),

$$h_{o1} = e^{i \int_0^\infty \frac{dt}{t} \frac{(J_0(2gt)-1)(e^t+1)}{(e^t-1)} \sin(u_1 t) e^{-\frac{n_1}{2}t + \pi(u_1 - \nu_1)}}. \quad (18)$$

This concludes our conjecture for the all-order structure of \mathcal{R}_N in MRK. In fact, the dispersion integral (1) is valid also at finite coupling, and so is the central emission block (9), for all integer angular momenta n_r different from zero. As noted in [28], a subtlety that appears when $n_r = 0$ is that one needs two sheets in the rapidity u_r in order to cover the entire real ν_r line, with the expressions (10)-(18) only covering the interval $|\nu_r| \geq \tilde{\nu}_r = \nu(u_r = 2g)$ (this is not an issue at weak coupling, where we can express all building blocks as functions of ν_r directly). Covering also the $|\nu_r| < \tilde{\nu}_r$ interval would additionally serve as a starting point for analyzing the strong-coupling limit, and making contact with the string-theoretic description of the same regime [53].

The perturbative expansion of all quantities entering (1) is simple to obtain [25, 27, 54], since at fixed order only a finite number of components of the vectors (32) contribute. The coefficients of the perturbative expansion take a very special form; the ratio to their leading-order contribution is always a polynomial in the following *FM building blocks*, first introduced in [46, 55],

$$\begin{aligned} V_i &= \frac{i\nu_i}{\nu_i^2 + \frac{n_i^2}{4}}, \quad N_i = \frac{n_i}{\nu_i^2 + \frac{n_i^2}{4}}, \quad D_i = -i \frac{\partial}{\partial \nu_i}, \\ E_i &= \psi\left(1 + i\nu_i + \frac{|n_i|}{2}\right) + \psi\left(1 - i\nu_i + \frac{|n_i|}{2}\right) \\ &\quad - 2\psi(1) - \frac{1}{2} \frac{|n_i|}{\nu_i^2 + \frac{n_i^2}{4}}, \\ M_{ij} &= \psi\left(i\nu_{ij} - \frac{n_{ij}}{2}\right) + \psi\left(1 - i\nu_{ij} - \frac{n_{ij}}{2}\right) - 2\psi(1), \end{aligned} \quad (19)$$

where $\nu_{ij} = \nu_i - \nu_j$ and $n_{ij} = n_i - n_j$.

We implement the general expansion of \tilde{C}_{12}^+ , and provide explicit results through five loops, as ancillary material with the arXiv version of the paper. As independent checks, we have verified that inserting it to the dispersion integral (1) and evaluating, we find perfect agreement for the imaginary part of the four-loop seven-particle MHV symbol [21], as well as for the two-loop MHV amplitude at any multiplicity [46, 56]. More details on the integral evaluation step are provided in the next section.

ANALYTIC LOOP AMPLITUDES IN MRK

In this section we provide the last ingredient needed to compute amplitudes from the dispersive representation in eq. (1), and we discuss how the integrals can be efficiently performed in terms of the relevant class of functions in the limit, known as *single-valued MPLs* (SVMPLs) [40, 57, 58]. As an application, we will give for the first time a proof of the principle of uniform transcendentality in MRK:

An L -loop gluon amplitude in MRK in planar $\mathcal{N} = 4$ SYM is a linear combination of products of $\log \tau_i$, SVMPLs, zeta values and powers of $2\pi i$ of uniform weight $2L$, for any helicity configuration and any number of legs.

The proof is constructive, thereby providing an important algorithm to compute any scattering amplitude in MRK order by order in the coupling, as we now sketch. For $N = 6$ gluons, similar proofs for the relevant classes of functions in the collinear and LLA multi-Regge limit have appeared in [55, 59, 60] and [61, 62] respectively, see also [40] for an extension of the latter to any N .

We start by noting that at order $\mathcal{O}(g^2)$, the MHV amplitude will be the $(N - 5)$ -fold FM transform of the *vacuum ladder*,

$$\varpi = \prod_{r=1}^{N-5} \frac{1}{\nu_r^2 + \frac{n_r^2}{4}} \prod_{r=1}^{N-6} \tilde{C}_{r,r+1}^{(0)}. \quad (20)$$

Letting $\mathcal{F}[X_r]$ denote the FM transform of X_r , we have in particular that $\mathcal{F}[\varpi] = \delta/(4\pi)$, with δ as in (1) being of uniform weight one.

At higher loops, the integrand will be a product of (20) with sums of polynomials of the FM building blocks (19). If we assign weight 1 to them, and given that the polynomial coefficients are \mathbb{Q} -linear combinations of Riemann zeta values $\zeta_n = \zeta(n)$, whose weight is n , then we observe that these polynomials have uniform transcendentality weight. In other words, we see that the all-order formulæ obtained from integrability imply the principle of uniform transcendentality in FM space.

To go to momentum space, we then make use of the FM transform's property to map products to convolutions,

$$\mathcal{F}[f \cdot g] = \mathcal{F}[f] * \mathcal{F}[g], \quad (21)$$

where

$$(F * G)(z) = \int \frac{d^2 w}{|w|^2} F(w) G\left(\frac{z}{w}\right). \quad (22)$$

Every higher-loop amplitude in MRK can thus be built iteratively by convolving the vacuum ladder (20) with a finite number of FM building blocks (19). While the evaluation of the convolution integral seems a daunting task, it was shown in [63] (see also [40]) that, in the case

where the integrand only involves rational functions and SVMPLs, the integral can easily be evaluated in terms of residues.

The proof now proceeds by induction: Assume we have a pure linear combination of SVMPLs of uniform weight. We will show that convolution with any FM building block raises the weight by 1 and preserves purity. This implies that all MHV amplitudes in MRK satisfy the principle of uniform transcendentality.

More concretely, assume that $f(z)$ is a pure linear combination of SVMPLs of uniform weight n and let

$$\mathcal{K}(z) = |z|^2 \sum_{i,j} \frac{a_{ij}}{(z - \alpha_i)(\bar{z} - \beta_j)}, \quad (23)$$

with $a_{ij}, \alpha_i, \beta_j \in \mathbb{Q}$. One can show using Stoke's theorem [63] that $(f * \mathcal{K})(z)$ is again pure and has uniform weight $n + 1$. The FM transform of the building blocks E_r, N_r, V_r match the form in (23) [40, 46, 47]

$$\mathcal{F}[E_r] = -\frac{z_r + \bar{z}_r}{2|1 - z_r|^2}, \quad (24)$$

$$\mathcal{F}[V_r] = \frac{2 - z_r - \bar{z}_r}{2|1 - z_r|^2}, \quad (25)$$

$$\mathcal{F}[N_r] = \frac{z_r - \bar{z}_r}{|1 - z_r|^2}. \quad (26)$$

Hence they raise the weight of the function they are convolved with by 1. We may similarly show that the same holds true for the derivative D_r , by using integration by parts to let it act on the factor $|z_r|^{2\nu_r}$ in the definition of the FM transform,

$$\mathcal{F}[D_r X_r] = -\log |z_r|^2 \mathcal{F}[X_r]. \quad (27)$$

Finally, let us note that the FM building block $M_{r,s}$ obeys

$$M_{r,s} = D_r \log(\tilde{C}_{rs}^{(0)}) + E_r + V_r \quad (28)$$

$$= -D_s \log(\tilde{C}_{rs}^{(0)}) + E_s - V_s. \quad (29)$$

This allows us to shift occurrences of $M_{r,s}$ in its FM transform with the vacuum ladder to either end,

$$\mathcal{F}[\varpi M_{r,r+1}] = \mathcal{F}[\varpi M_{r-1,r}] + \mathcal{F}[D_r \varpi] \quad (30)$$

$$\mathcal{F}[\varpi M_{12}] = \mathcal{F}[\varpi E_1] - \mathcal{F}[\varpi V_1] + \mathcal{F}[D_1 \varpi], \quad (31)$$

and in this manner replace it by a combination of E, V, D . Hence, $M_{r,s}$ raises the weight of the integral by 1 as well. Finally, our proof may be immediately extended to non-MHV amplitudes as well. The latter can be obtained by convoluting MHV amplitudes with the helicity-flip kernel I^- , and the only difference is that at LO the latter does not raise the weight, and it does not preserve the purity of the function [40]. We therefore conclude that non-MHV amplitudes have the same weight as their MHV counterparts, but are no longer pure functions.

CONCLUSIONS

We have presented a dispersion integral for all gluon amplitudes of arbitrary multiplicity, helicity configuration in MRK. By combining our results with [40] we obtain an efficient algorithm to evaluate any scattering amplitude in MRK, for any number of loops or legs, and for arbitrary helicity configurations.

We believe that our results, while complete for the sector of planar $\mathcal{N} = 4$ super Yang-Mills theory that we have studied, should serve as the basis for many future generalisations in various directions. Firstly it should be straightforward to include the fermions and scalars into our expression, or to consider more general Mandelstam regions [56, 64–66]. We believe that a similar structure will survive for general gauge theories, at least in the planar limit, though the details will differ because in general dual conformal symmetry is broken. It would be very interesting to understand how the form of the amplitude generalises beyond the planar limit.

ACKNOWLEDGEMENTS

We would like to thank Jochen Bartels for comments on the manuscript. This work was supported in part by the ERC Consolidator grant 648630 IQFT and the ERC Starting grants 637019 MathAm and 804286 UNIS-CAMP, as well as the U.S. Department of Energy (DOE) under contract DE-AC02-76SF00515.

Appendix: BES kernel and BFKL sources

The sources κ and $\tilde{\kappa}$ are infinite-dimensional vectors with j -th component given by

$$\kappa_j(u_r, n_r) = - \int_0^\infty \frac{dt}{t} \frac{J_j(2gt)}{e^t - 1} \phi_j(t; u_r, n_r), \quad (32)$$

and similarly for $\tilde{\kappa}$ with ϕ replaced by $\tilde{\phi}$. Here $J_j(x)$ denote Bessel functions of the first kind, and we have

$$\begin{aligned} \phi_j &= \frac{1}{2} \left[e^{\frac{t(1+(-)^j)}{2}} - (-)^j e^{\frac{t(1-(-)^j)}{2}} \right] \cos(u_r t) e^{-\frac{n_r t}{2}} - J_0(2gt), \\ \tilde{\phi}_j &= \frac{1}{2} \left[e^{\frac{t(1+(-)^j)}{2}} + (-)^j e^{\frac{t(1-(-)^j)}{2}} \right] \sin(u_r t) e^{-\frac{n_r t}{2}}. \end{aligned} \quad (33)$$

The matrices \mathbb{Q} and \mathbb{M} in (7) are given by

$$\begin{aligned} \mathbb{Q}_{ij} &= \delta_{ij} (-1)^{i+1} i, & \mathbb{M} &= (1 + \mathbb{K})^{-1}, \\ \mathbb{K}_{ij} &= 2j (-1)^{j(i+1)} \int_0^\infty \frac{dt}{t} \frac{J_i(2gt) J_j(2gt)}{e^t - 1}, \end{aligned} \quad (34)$$

with the latter being simply the kernel of the Beisert-Eden-Staudacher equation [4] as reformulated in [50].

- [1] J. M. Drummond, J. Henn, G. P. Korchemsky, and E. Sokatchev, Nucl. Phys. **B828**, 317 (2010), arXiv:0807.1095 [hep-th].
- [2] J. M. Drummond, J. M. Henn, and J. Plefka, *Strangeness in quark matter. Proceedings, International Conference, SQM 2008, Beijing, P.R. China, October 5-10, 2008*, JHEP **05**, 046 (2009), arXiv:0902.2987 [hep-th].
- [3] Z. Bern, L. J. Dixon, and V. A. Smirnov, Phys. Rev. **D72**, 085001 (2005), arXiv:hep-th/0505205 [hep-th].
- [4] N. Beisert, B. Eden, and M. Staudacher, J. Stat. Mech. **0701**, P01021 (2007), arXiv:hep-th/0610251 [hep-th].
- [5] J. M. Drummond, J. Henn, G. P. Korchemsky, and E. Sokatchev, Nucl. Phys. **B826**, 337 (2010), arXiv:0712.1223 [hep-th].
- [6] V. Del Duca, C. Duhr, and V. A. Smirnov, JHEP **03**, 099 (2010), arXiv:0911.5332 [hep-ph].
- [7] V. Del Duca, C. Duhr, and V. A. Smirnov, JHEP **05**, 084 (2010), arXiv:1003.1702 [hep-th].
- [8] A. B. Goncharov, M. Spradlin, C. Vergu, and A. Volovich, Phys. Rev. Lett. **105**, 151605 (2010), arXiv:1006.5703 [hep-th].
- [9] L. J. Dixon, J. M. Drummond, and J. M. Henn, JHEP **11**, 023 (2011), arXiv:1108.4461 [hep-th].
- [10] L. J. Dixon, J. M. Drummond, and J. M. Henn, JHEP **01**, 024 (2012), arXiv:1111.1704 [hep-th].
- [11] L. J. Dixon, J. M. Drummond, M. von Hippel, and J. Pennington, JHEP **12**, 049 (2013), arXiv:1308.2276 [hep-th].
- [12] L. J. Dixon, J. M. Drummond, C. Duhr, and J. Pennington, JHEP **06**, 116 (2014), arXiv:1402.3300 [hep-th].
- [13] L. J. Dixon, J. M. Drummond, C. Duhr, M. von Hippel, and J. Pennington, *Proceedings, 12th DESY Workshop on Elementary Particle Physics: Loops and Legs in Quantum Field Theory (LL2014): Weimar, Germany, April 27-May 2, 2014*, PoS **LL2014**, 077 (2014), arXiv:1407.4724 [hep-th].
- [14] L. J. Dixon and M. von Hippel, JHEP **10**, 065 (2014), arXiv:1408.1505 [hep-th].
- [15] L. J. Dixon, M. von Hippel, and A. J. McLeod, JHEP **01**, 053 (2016), arXiv:1509.08127 [hep-th].
- [16] S. Caron-Huot, L. J. Dixon, A. McLeod, and M. von Hippel, Phys. Rev. Lett. **117**, 241601 (2016), arXiv:1609.00669 [hep-th].
- [17] S. Caron-Huot, L. J. Dixon, F. Dulat, M. von Hippel, A. J. McLeod, and G. Papathanasiou, JHEP **08**, 016 (2019), arXiv:1903.10890 [hep-th].
- [18] J. Golden and M. Spradlin, JHEP **08**, 154 (2014), arXiv:1406.2055 [hep-th].
- [19] J. Golden, M. F. Paulos, M. Spradlin, and A. Volovich, J. Phys. **A47**, 474005 (2014), arXiv:1401.6446 [hep-th].
- [20] J. M. Drummond, G. Papathanasiou, and M. Spradlin, JHEP **03**, 072 (2015), arXiv:1412.3763 [hep-th].
- [21] L. J. Dixon, J. Drummond, T. Harrington, A. J. McLeod, G. Papathanasiou, and M. Spradlin, JHEP **02**, 137 (2017), arXiv:1612.08976 [hep-th].
- [22] J. Drummond, J. Foster, O. Gürdogan, and G. Papathanasiou, JHEP **03**, 087 (2019), arXiv:1812.04640 [hep-th].
- [23] A. B. Goncharov, (2001), math/0103059v4.

- [24] B. Basso, A. Sever, and P. Vieira, Phys. Rev. Lett. **111**, 091602 (2013), arXiv:1303.1396 [hep-th].
- [25] B. Basso, A. Sever, and P. Vieira, JHEP **01**, 008 (2014), arXiv:1306.2058 [hep-th].
- [26] B. Basso, A. Sever, and P. Vieira, JHEP **08**, 085 (2014), arXiv:1402.3307 [hep-th].
- [27] B. Basso, A. Sever, and P. Vieira, JHEP **09**, 149 (2014), arXiv:1407.1736 [hep-th].
- [28] B. Basso, S. Caron-Huot, and A. Sever, JHEP **01**, 027 (2015), arXiv:1407.3766 [hep-th].
- [29] L. F. Alday and J. M. Maldacena, JHEP **06**, 064 (2007), arXiv:0705.0303 [hep-th].
- [30] J. M. Drummond, G. P. Korchemsky, and E. Sokatchev, Nucl. Phys. **B795**, 385 (2008), arXiv:0707.0243 [hep-th].
- [31] A. Brandhuber, P. Heslop, and G. Travaglini, Nucl. Phys. **B794**, 231 (2008), arXiv:0707.1153 [hep-th].
- [32] Z. Bern, L. J. Dixon, D. A. Kosower, R. Roiban, M. Spradlin, C. Vergu, and A. Volovich, Phys. Rev. **D78**, 045007 (2008), arXiv:0803.1465 [hep-th].
- [33] J. M. Drummond, J. Henn, G. P. Korchemsky, and E. Sokatchev, Nucl. Phys. **B815**, 142 (2009), arXiv:0803.1466 [hep-th].
- [34] R. C. Brower, H. Nastase, H. J. Schnitzer, and C.-I. Tan, Nucl. Phys. **B814**, 293 (2009), arXiv:0801.3891 [hep-th].
- [35] J. Bartels, L. N. Lipatov, and A. Sabio Vera, Phys. Rev. **D80**, 045002 (2009), arXiv:0802.2065 [hep-th].
- [36] V. Del Duca and E. W. N. Glover, JHEP **05**, 056 (2008), arXiv:0802.4445 [hep-th].
- [37] J. Bartels, L. N. Lipatov, and A. Sabio Vera, Eur. Phys. J. **C65**, 587 (2010), arXiv:0807.0894 [hep-th].
- [38] R. C. Brower, H. Nastase, H. J. Schnitzer, and C.-I. Tan, Nucl. Phys. **B822**, 301 (2009), arXiv:0809.1632 [hep-th].
- [39] V. Del Duca, C. Duhr, and E. W. N. Glover, JHEP **12**, 097 (2008), arXiv:0809.1822 [hep-th].
- [40] V. Del Duca, S. Druc, J. Drummond, C. Duhr, F. Dulat, R. Marzucca, G. Papathanasiou, and B. Verbeek, JHEP **08**, 152 (2016), arXiv:1606.08807 [hep-th].
- [41] L. N. Lipatov, Theor. Math. Phys. **170**, 166 (2012), arXiv:1008.1015 [hep-th].
- [42] V. S. Fadin and L. N. Lipatov, Phys. Lett. **B706**, 470 (2012), arXiv:1111.0782 [hep-th].
- [43] J. Bartels, A. Kormilitzin, L. N. Lipatov, and A. Prygarin, Phys. Rev. **D86**, 065026 (2012), arXiv:1112.6366 [hep-th].
- [44] L. Lipatov, A. Prygarin, and H. J. Schnitzer, JHEP **01**, 068 (2013), arXiv:1205.0186 [hep-th].
- [45] J. Bartels, A. Kormilitzin, and L. Lipatov, Phys. Rev. **D89**, 065002 (2014), arXiv:1311.2061 [hep-th].
- [46] V. Del Duca, S. Druc, J. Drummond, C. Duhr, F. Dulat, R. Marzucca, G. Papathanasiou, and B. Verbeek, JHEP **06**, 116 (2018), arXiv:1801.10605 [hep-th].
- [47] R. Marzucca and B. Verbeek, JHEP **07**, 039 (2019), arXiv:1811.10570 [hep-th].
- [48] Explicitly $\delta(z) = \pi \log \frac{|\rho_1|^2}{|1-\rho_1|^2|1-\rho_{N-5}|^2}$ where the ρ_i are defined in terms of the z_i via $z_i = \frac{(1-\rho_{i+1})(\rho_i-\rho_{i-1})}{(1-\rho_{i-1})(\rho_i-\rho_{i+1})}$ together with $\rho_0 = 0$ and $\rho_{N-4} = \infty$, see e.g. [41, 43, 45, 65].
- [49] S. Caron-Huot, JHEP **05**, 093 (2015), arXiv:1309.6521 [hep-th].
- [50] M. K. Benna, S. Benvenuti, I. R. Klebanov, and A. Scardicchio, Phys. Rev. Lett. **98**, 131603 (2007), arXiv:hep-th/0611135 [hep-th].
- [51] Explicitly the helicity-flip kernels are $I_r^+ = 1$ and $I_r^- = x_r^-/x_r^+$, where the Zhukowski variables x_r^\pm are given in (14).
- [52] The measure is similarly obtained and is given by $\frac{dv_r}{2\pi} \tilde{\Phi}_r = \frac{du_r}{2\pi} \frac{g^2(x_r^+x_r^- - g^2) \exp(A - f_{rr} + f_{\tilde{r}\tilde{r}})}{x_r^+x_r^- \sqrt{(x_r^+x_r^- - g^2)(x_r^-x_r^+ - g^2)}}$, see [28].
- [53] J. Bartels, V. Schomerus, and M. Sprenger, JHEP **10**, 67 (2014), arXiv:1405.3658 [hep-th].
- [54] J. M. Drummond and G. Papathanasiou, JHEP **02**, 185 (2016), arXiv:1507.08982 [hep-th].
- [55] L. J. Dixon, C. Duhr, and J. Pennington, JHEP **1210**, 074 (2012), arXiv:1207.0186 [hep-th].
- [56] T. Bargheer, G. Papathanasiou, and V. Schomerus, JHEP **05**, 012 (2016), arXiv:1512.07620 [hep-th].
- [57] F. C. S. Brown, <http://www.ihes.fr/~brown/RHpaper5.pdf>.
- [58] F. C. S. Brown, (2015), 1512.06410 [math.NT].
- [59] G. Papathanasiou, JHEP **11**, 150 (2013), arXiv:1310.5735 [hep-th].
- [60] G. Papathanasiou, Int.J.Mod.Phys. **A29**, 1450154 (2014), arXiv:1406.1123 [hep-th].
- [61] J. Pennington, JHEP **1301**, 059 (2013), arXiv:1209.5357 [hep-th].
- [62] J. Broedel and M. Sprenger, JHEP **05**, 055 (2016), arXiv:1512.04963 [hep-th].
- [63] O. Schnetz, Commun. Num. Theor. Phys. **08**, 589 (2014), arXiv:1302.6445 [math.NT].
- [64] T. Bargheer, (2016), arXiv:1606.07640 [hep-th].
- [65] V. Del Duca, C. Duhr, F. Dulat, and B. Penante, JHEP **01**, 162 (2019), arXiv:1811.10398 [hep-th].
- [66] T. Bargheer, V. Chestnov, and V. Schomerus, (2019), arXiv:1906.00990 [hep-th].