

DESY 85-100  
September 1985



JET PRODUCTION AT HIGH TRANSVERSE MOMENTA BY INTERACTIONS  
OF TWO QUASI-REAL PHOTONS

by

*PLUTO Collaboration*

ISSN 0418-9833

NOTKESTRASSE 85 · 2 HAMBURG 52

**DESY behält sich alle Rechte für den Fall der Schutzrechtserteilung und für die wirtschaftliche Verwertung der in diesem Bericht enthaltenen Informationen vor.**

**DESY reserves all rights for commercial use of information included in this report, especially in case of filing application for or grant of patents.**

**To be sure that your preprints are promptly included in the  
HIGH ENERGY PHYSICS INDEX ,  
send them to the following address ( if possible by air mail ) :**

**DESY  
Bibliothek  
Notkestrasse 85  
2 Hamburg 52  
Germany**

**Jet Production at High Transverse Momenta  
by Interactions of two Quasi-Real Photons**

PLUTO COLLABORATION

CH. BERGER, H. GENZEL, W. LACKAS, J. PIELORZ<sup>a</sup>,  
F. RAUPACH, W. WAGNER<sup>b</sup>,

*I. Phys. Institut der RWTH Aachen<sup>c</sup>, Federal Republic of Germany*

A. KLOVNING, E. LILLESTÖL,  
*University of Bergen<sup>d</sup>, Norway*

J. BÜRGER, L. CRIEGEE, A. DEUTER, F. FERRAROTTO<sup>e</sup>, G. FRANKE,  
M. GASPERO<sup>e</sup>, CH. GERKE, G. KNIES, B. LEWENDEL, J. MEYER,  
U. MICHELSEN, K.H. PAPE, B. STELLA<sup>e</sup>, U. TIMM,  
G.G. WINTER, M. ZAOKHARA<sup>f</sup>, W. ZIMMERMANN,  
*Deutsches Elektronen-Synchrotron (DESY), Hamburg, Federal Republic of Germany*

P.J. BUSSEY, S.L. CARTWRIGHT<sup>g</sup>, J.B. DANTON, B.T. KING<sup>h</sup>, C. RAINE,  
J.M. SCARR, I.O. SKILLICORN, K.M. SMITH, J.C. THOMSON<sup>i</sup>,  
*University of Glasgow<sup>j</sup>, U.K.*

O. ACHTERBERG, V. BLOBEL, D. BURKART, K. DIELMANN, M. FEINDT,  
H. KAPITZA<sup>k</sup>, B. KOPPITZ, M. KRÜGER<sup>l</sup>, M. POPPE,  
H. SPITZER, R. VAN STAA,  
*II. Institut für Experimentalphysik der Universität Hamburg<sup>e</sup>,  
Federal Republic of Germany*

C.Y. CHANG, R.G. GLASSER, R.G. KELLOGG, S.J. MAXFIELD<sup>m</sup>,  
R.O. POLVADO<sup>n</sup>, B. SECHI-ZORN<sup>o</sup>, J.A. SKARD, A. SKUJA,  
A.J. TYLKA<sup>o</sup>, G.E. WELCH, G.T. ZORN,  
*University of Maryland<sup>p</sup>, U.S.A.*

F. ALMEIDA<sup>q</sup>, A. BÄCKER, F. BARREIRO<sup>r</sup>, S. BRANDT, K. DERIKUM<sup>s</sup>,  
C. GRUPEN, H.J. MEYER, H. MÜLLER, B. NEUMANN,  
M. ROST, K. STUPPERICH, G. ZECH,  
*Universität-Gesamthochschule Siegen<sup>c</sup>, Federal Republic of Germany*

G. ALEXANDER, G. BELLA, Y. GNAT, J. GRUNHAUS,  
*University of Tel-Aviv<sup>t</sup>, Israel*

H. JUNGE, K. KRASKI, C. MAXEINER, H. MAXEINER,  
H. MEYER, D. SCHMIDT,  
*Universität-Gesamthochschule Wuppertal<sup>c</sup>, Federal Republic of Germany*

- a) Deceased
- b) Now at University of California at Davis, Davis, Ca., U.S.A.
- c) Supported by the BMFT, Federal Republic of Germany
- d) Partially supported by the Norwegian Council for Science and the Humanities
- e) Rome University, partially supported by I.N.F.N., Sezione di Roma, Italy
- f) Institute of Nuclear Physics, Cracow, Poland
- g) Now at Rutherford Appelton Laboratory, Chilton, U.K.
- h) Now at University of Liverpool, Liverpool, U.K.
- i) Now at Glasgow College of Technology, Glasgow, U.K.
- j) Supported by the U.K. Science and Engineering Research Council
- k) Now at Carleton University, Ottawa, Ontario, Canada
- l) Now at Universität Karlsruhe, Federal Republic of Germany
- m) Now at University of Massachusetts, Amherst, Mass., U.S.A.
- n) Now at Northeastern University, Boston, Mass., U.S.A.
- o) Now at Cosmic Ray Astrophysics Branch, Code 4154.8, U.S. Naval Research Laboratory, Washington, DC 20375, U.S.A.
- p) Partially supported by the Department of Energy, U.S.A.
- q) On leave of absence from Inst. de Fisica, Universidad Federal do Rio de Janeiro, Brasil
- r) On leave of absence at Universidad Autonoma de Madrid, Spain
- s) Now at BESSY, Berlin, Federal Republic of Germany
- t) Partially supported by the Israeli Academy of Sciences and Humanities, Basic Research Foundation

**Abstract.** An experimental study of two jet production by interactions of two quasi-real photons is presented. The data for production of jets with high transverse momentum squared,  $p_T^2 > 10 \text{ GeV}^2$ , are found to be consistent with the fractional charged quark-parton model. If gauge integer charged quark models are considered, then the gluon mass is less than 5 MeV at the 95% confidence level.

The two photon production of quark jets can be studied via the interaction of high energy  $e^+e^-$  beams in the reaction

$$e^+e^- \rightarrow e^+e^-\gamma^*\gamma^*, \quad \gamma^*\gamma^* \rightarrow q\bar{q} \quad (1)$$

where  $\gamma^*$  represents a virtual photon. The final quark-antiquark ( $q\bar{q}$ ) system is detected as a pair of hadronic jets. For small electron scattering angles the photons are quasi-real with  $Q^2$ , the photon mass squared,  $\approx 0$ .

Reaction (1) allows the possibility of distinguishing models with fractional quark charges <sup>(1)</sup> (FCQ) from those with integer charges <sup>(2)</sup> (ICQ) by measurement of the ratio for two photon interactions

$$R_{\gamma\gamma} = \frac{\sigma(e^+e^- \rightarrow e^+e^- + \text{hadrons})}{\sigma(e^+e^- \rightarrow e^+e^- \mu^+\mu^-)} = \frac{1}{3} \sum_{nf} \left( \sum_{nc} q(f,c)^2 \right)^2 \quad (2)$$

where  $nf(nc)$  is the number of flavors (colors (=3)), and  $q(f,c)$  is the quark charge for flavor  $f$  and color  $c$  in units of electron charge. This is to be contrasted with single photon reactions, where, assuming the hadrons are produced as color singlets, such differentiation is not possible. In  $e^+e^-$  annihilation, for example, the ratio

$$R = \frac{\sigma(e^+e^- \rightarrow \text{hadrons})}{\sigma(e^+e^- \rightarrow \mu^+\mu^-)} = \frac{1}{3} \sum_{nf} \left( \sum_{nc} q(f,c)^2 \right)^2 \quad (3)$$

is identical for both FCQ and ICQ models, since for a given flavor the quark charge summed over color is necessarily the same in each case. The ratio (2) must be measured for jets produced at high transverse momentum ( $p_T$ ) so that reaction (1) dominates over hadron production by other processes such as the vector meson dominance mechanism <sup>(3)</sup> (VMD).

One method to study the reaction  $e^+e^- \rightarrow e^+e^- + \text{hadrons}$  is when at least one of the scattered electrons is detected (tagged mode) with consequent reduction of backgrounds. In an earlier study <sup>(4)</sup> results have been reported for single tagged events in which one photon was off-shell ( $Q > 0.1$  GeV), which, in particular at high jet  $p_T$ , agree with the prediction of the FCQ model. However, gauge models with integer charge quarks (gauge ICQ) have been proposed <sup>(5)</sup> in which color SU(3) is broken and gluons acquire both mass and electric charge. In this case equation (2) must be modified <sup>(6)</sup> and such models can be distinguished from the unbroken fractional charge model, in practice, only when quasi-real photons are used. For this reason results on jet production by quasi-real photons, using anti-tagging, are of interest.

The data presented here were obtained at the PETRA storage ring (at DESY, Hamburg) with a beam energy of 17.3 GeV ( $\sqrt{s} = 34.7$  GeV) using the PLUTO detector

which has been described in previous publications <sup>(7)</sup>. The central detector measures charged particles for angles at  $> 25$  degrees to the beam axis with cylindrical proportional chambers in 16.5 Kgauss magnetic field. Photons are detected in lead-scintillator shower counters for angles  $> 15$  degrees. In addition, two forward spectrometers <sup>(8)</sup> measure charged particles between 90 and 260 mr and photons between 23 and 260 mr. Charged particles are measured with resolution  $\sigma_p/p = 3\% \cdot p(\text{GeV})$  and showers with energy resolution of  $(16.5\% - 28\%)/\sqrt{E}$ .

The photon-photon interactions of reaction (1) are characterised by a total hadronic mass,  $W$ , much less than the  $e^+e^-$  center of mass energy,  $\sqrt{s}$ , since energy is carried off by the scattered electrons. In the present analysis predominantly quasi-real photon interactions are selected by using the forward spectrometers to reject events with detected electrons, retaining mainly those events where the scattered electrons are at angles less than 23 mr.

Accepted events are required to satisfy the following conditions:

- a) No electron detected in the forward spectrometers with energy greater than 5 GeV.
- b) Measured hadronic vertex  $< 30$  mm from beam crossing point to reduce beam-gas background.
- c) Number of charged tracks in each jet  $\geq 2$  in order to reduce background due to  $\tau$  production. Since a two jet analysis is performed, the total number of charged tracks observed is  $\geq 4$ .
- d) Absolute value of charge of detected hadrons  $|\Sigma q| \leq 2$ .
- e) Total transverse momentum of the detected hadronic system less than 1.5 GeV/c. This limit preferentially selects interactions of quasi-real photons and reduces background due to badly measured annihilation events.
- f) Only events with total observed hadronic mass,  $W_{vis}$ , less than 12 GeV were used in order to maximise the signal relative to the annihilation background which peaks at  $W \approx \sqrt{s}$ . A cut of  $W_{vis} > 6$  GeV is applied to allow a reliable determination of the jet axis using a thrust algorithm. The value of  $W_{vis}$  is calculated from charged track momenta (assuming pions) and the energies of the showers.

With the above cuts the data sample consists of 3203 events. The corresponding integrated luminosity is  $18.7 \text{ pb}^{-1}$ .

Beam-gas interactions, as determined from the event vertex distribution, contribute a background of 0.25% to the final data sample and can be ignored. Additional

background events arise from the following processes:

$$e^+e^- \rightarrow e^+e^-r^+r^- \quad (4)$$

$$e^+e^- \rightarrow r^+r^- \quad (5)$$

$$e^+e^- \rightarrow \text{hadrons} \quad (6)$$

and have been studied by Monte Carlo simulations and included in the calculated background. Reaction (4) gives a negligible background contribution, while reaction (5) gives a small contribution which is discussed later.

The annihilation process (6) is the most important source of background for high  $p_T$  jets produced in untagged photon-photon interactions, since for some events  $W_{vis}$  has a small value either due to high particle loss (charged or neutral) or due to hard radiation off one or other initial state electron. The contribution of this background has been studied by generating annihilation events using the Lund jet Monte Carlo <sup>(9)</sup>, and including first order initial state radiative corrections according to the method of Berends and Kleiss <sup>(10)</sup>. The resulting events are then processed through the PLUTO detector simulation program, after which they are treated identically to the data and used to perform the background subtractions. Similarly, Monte Carlo events for reaction (1) have been generated for comparison with the high jet transverse momentum ( $p_T$ ) data in accordance with equation (2) and assuming that in the process

$$\gamma^* \gamma^* \rightarrow q \bar{q} \quad (7)$$

the photons couple directly to the fractionally charged quarks of unbroken color SU(3). We will refer to this as the quark-parton model (QPM). Detailed information on this simulation is given in a previous PLUTO report <sup>(4)</sup>.

We present our experimental results as observed distributions, uncorrected for acceptance, which are then compared with simulations which include the acceptance effects.

Each event has been analysed by using a thrust algorithm, in the rest frame of the observed hadrons, to define two hadronic jets. To isolate the reaction (7) we must consider data for jets produced at high  $p_T$ , where we have measured the transverse momentum relative to the electron beam direction. At lower  $p_T$ , multi-jet events and QCD corrections play a role. Jet production by the VMD mechanism dominates at small  $p_T$ . Fig. 1(a) shows the rapidity distribution of the total hadronic system for events with  $p_T^2 > 10 \text{ GeV}^2$  and  $6 < W_{vis} < 12 \text{ GeV}$ . The expected QPM contribution is shown as the lower histogram (an essentially flat distribution is expected for fixed  $W$  before acceptance effects) and the data are compared with the sum of QPM and annihilation simulations. The agreement between data and simulation is good. The figure indicates the importance of the background from annihilation events for high

jet  $p_T$ , which can however be reduced by a cut excluding low rapidity events whose low  $W_{vis}$  is mainly due to track or shower loss. Consequently an additional cut requiring  $|\text{rapidity}| > 0.15$  is used in what follows. At lower  $p_T$  such a peak at low rapidity is less prominent since track loss will also lead to change of observed rapidity. Fig. 1(b) shows the  $W_{vis}$  distribution for the data with  $p_T^2 > 10 \text{ GeV}^2$  and  $|\text{rapidity}| > 0.15$  compared to the Monte Carlo simulation. The agreement is satisfactory. The Monte Carlo also describes the charge balance, charged particle multiplicity distributions and  $W_{vis}$  for  $|\text{rapidity}| < 0.15$ . The neutral multiplicity is slightly underestimated by the Monte Carlo. For  $6 < W_{vis} < 12 \text{ GeV}$  and total hadronic transverse momentum  $> 1.5 \text{ GeV}/c$ , or for  $12 < W_{vis} < 15 \text{ GeV}$ , where the QPM contribution is expected to be small, the annihilation Monte Carlo gives an acceptable description of the data. These checks indicate that systematic uncertainties in the background simulation are no more than 20%.

Background from  $r$  production by reaction (5) is due to undetected neutrals (e.g. neutrinos) resulting in a low value for  $W_{vis}$ . It is strongly suppressed by selection (f). The remaining background events tend to have low rapidity and the applied rapidity cut reduces this source of background to an insignificant level.

Figure 2 shows the observed jet transverse momentum squared ( $p_T^2$ ) distribution for  $6 < W_{vis} < 12 \text{ GeV}$  after subtraction of the annihilation background. The  $p_T$  of the two observed jets may differ by up to  $1.5 \text{ GeV}/c$ , the difference being limited by cut (e) above. The higher  $p_T$  jet has been used as the better approximation to the quark  $p_T$  as indicated by Monte Carlo studies. For comparison the QPM Monte Carlo prediction is also shown. The data for  $p_T^2 > 10 \text{ GeV}^2$  are consistent with the QPM prediction (solid curve) but not with the naive ICQ model (factor 2.65 higher than QPM, dashed curve). The observed thrust distributions for these high  $p_T^2$  events (not shown) are in good agreement with the Monte Carlo predictions. At small  $p_T^2$  the VMD process is expected to dominate the QPM contribution. The sudden rise of the data below  $10 \text{ GeV}$  is a so far unexplained excess in the region  $5 < p_T^2 < 10 \text{ GeV}^2$  over VDM, which was also observed for interactions of low  $Q^2$  photons in our single-tag data <sup>(4)</sup>. This region of excess is associated with lower mean thrust than is expected for the QPM.

Figure 3 shows (upper data points) the charged particle  $p_T^2$  distribution for the events shown in figure 2. For  $p_T^2 > 3 \text{ GeV}^2$  the data exceed the QPM Monte Carlo prediction (upper curve) by a factor of  $\approx 3$ . This result is consistent with that of the TASSO Collaboration <sup>(11)</sup> who find data/QPM  $\approx 4$  for  $p_T^2 > 2.5 \text{ GeV}^2$ . The greater excess at low  $p_T^2$  is due to the VMD contribution. However, if only events with jet  $p_T^2 > 10 \text{ GeV}^2$  are used, the lower set of data points is obtained. For this case, the data are in agreement with the QPM prediction (lower curve) again demonstrating the validity of the QPM model for events with jet  $p_T^2 > 10 \text{ GeV}^2$ .

We now discuss the limits set on the gauge ICQ model by our data. For jet  $p_T^2 > 10 \text{ GeV}^2$  (data of figure 2), the ratio of the 33 observed events (after background subtraction) to the QPM prediction is  $1.1 \pm 0.4$  ( $\pm 0.4$  systematic) and the data are therefore consistent with the fractionally charged quarks of unbroken SU(3) at the limit of high  $p_T$ . For integer charged quarks the expected ratio would be 2.65 for 4 quark flavors, deviating by more than 3 standard deviations from the observed value. This result is consistent with results on other processes involving two photons, such as  $e^+e^- \rightarrow q\bar{q}\gamma$  <sup>(12)</sup> and  $\gamma q \rightarrow \gamma q$  <sup>(13)</sup>. However, if color SU(3) is broken as in the gauge ICQ model <sup>(5,6)</sup>, then equation (2) becomes

$$R_{\gamma\gamma} = \frac{1}{3} \sum_{nf} \left( \sum_{nc} (q(f,c))^2 + 2/9 D(1) D(2) \right)^2 \quad (8)$$

Here,  $q(f,c)$  is the fractional (FCQ) quark charge and

$$D(i) = M_{gluon}^2 / (M_{gluon}^2 + Q(i)^2)$$

$Q(i)^2$  = absolute momentum transfer squared of  $i^{\text{th}}$  photon,

and the factor 2/9 arises from the octet contribution to the effective quark charge. In addition a large contribution to jet production arises from production of charged gluon pairs for which the cross section is given by a lengthy formula <sup>(6)</sup>.

For the high jet  $p_T$  data sample we have computed the ratio,

$$R_{\gamma\gamma} (\text{gauge ICQ model}) / R_{\gamma\gamma} (\text{FCQ model})$$

including acceptance effects, and both quark and gluon production for a range of values of  $M_{gluon}$ . Comparing the calculated ratios with the measured ratio of  $1.1 \pm 0.4$  ( $\pm 0.4$ ) we obtain the following limit (including systematic error) for the gluon mass

$$M_{gluon} < 5 \text{ MeV at } 95\% \text{ C.L.}$$

With  $M_{gluon} < 5 \text{ MeV}$  the gauge ICQ model is indistinguishable from the FCQ model for tagged data where  $Q^2 > 0.1 \text{ GeV}^2$  for one of the photons. Thus single tag data is expected to be consistent with the FCQ model for  $p_T^2 > 10 \text{ GeV}^2$  as observed <sup>(4)</sup>. The single tagged data of the JADE <sup>(14)</sup> and TASSO <sup>(15)</sup> collaboration are also consistent with the FCQ model at high  $p_T$ .

In conclusion, we find that for jets produced by interaction of quasi-real photons the cross section is consistent with the expectation of the fractionally charged quark model for jet  $p_T^2 > 10 \text{ GeV}^2$  and  $6 < W_{vis} < 12 \text{ GeV}$ . If broken SU(3) is considered then  $M_{gluon} < 5 \text{ MeV}$  at the 95% confidence level.

## Acknowledgements

The university groups thank the DESY directorate for their hospitality. We acknowledge the excellent performance of the PETRA machine group, the DESY computer center, and of all the engineers and technicians who participated in construction and maintenance of the apparatus.

## References



- 1) M. Gell-Mann, Phys. Lett. 8, 214 (1964)
- 2) M. Han and Y. Nambu, Phys. Rev. 139b, 1006 (1965)
- 3) V.M. Budnev, I.F. Ginzburg, F.V. Meledin and V.G. Serbo, Phys. Reports 150, 181 (1975)
- 4) PLUTO Collaboration, Ch. Berger et al., Z. Phys. C26, 191 (1984)
- 5) J.C. Pati and A. Salam, Phys. Rev. D8, 1240 (1973)
- 6) R.M. Godbole et al., Phys. Lett. 142, 91 (1984)
- 7) L. Criegee and G. Knies, Phys. Reports 83, 83 (1982)
- 8) PLUTO Collaboration, Ch. Berger et al., Z. Phys. C26, 199 (1984)
- 9) T. Sjöstrand, Comput. Phys. Comm. 27, 243 (1982)
- 10) F.A. Berends and R. Kleiss, Nucl. Phys. B178, 141 (1981)
- 11) TASSO Collaboration, M. Althoff et al., Phys. Lett. 138B, 219 (1984)
- 12) MAC Collaboration, E. Fernandez et al., Phys. Rev. Lett. 54, 95 (1985)
- 13) NA14 Collaboration, P. Astbury et al., Phys. Lett. 152B, 419 (1985)
- 14) JADE Collaboration, W. Bartel et al., Phys. Lett. 107B, 163 (1981)
- 15) TASSO Collaboration, R. Brandelik et al., Phys. Lett. 107B, 1290 (1981)


## Figure Captions

- Figure 1(a) The absolute value of the total hadronic rapidity for events with  $p_T^2 > 10 \text{ GeV}^2$  and  $6 < W_{vis} < 12 \text{ GeV}$ . The lower histogram indicates the QPM Monte Carlo prediction. The upper histogram is the prediction for the sum of QPM and annihilation background. The first bin includes an estimated  $7 \pm 2$  events from the reaction  $e^+e^- \rightarrow \tau\tau$ .
- Figure 1(b) The observed hadronic mass distribution for events with  $p_T^2 > 10 \text{ GeV}^2$  and  $|Rapidity| > 0.15$ . The upper and lower histograms as for 1(a).


Figure 2 The observed jet  $p_T^2$  distribution for events with  $6 < W_{vis} < 12$  GeV after subtraction of the annihilation background. The errors are statistical only. The solid (dashed) curve is drawn through the Monte Carlo prediction for fractional (integer) charged quarks and should be compared with data at the bin centers.

Figure 3 The upper data points show the charged particle  $p_T^2$  distribution for the events of figure 2 (annihilation background subtracted). The lower data points (note shifted scale) show the charged particle  $p_T^2$  distribution for those events with the jet  $p_T^2 > 10$  GeV<sup>2</sup>. The errors are statistical only. The curves are drawn through the QPM Monte Carlo predictions and should be compared with the data at the bin centers.





39154  
 Fig. 2



39155  
 Fig. 3