


DEUTSCHES ELEKTRONEN-SYNCHROTRON **DESY**

DESY 85-106
October 1985



QUARK AND LEPTON MASSES IN A SIX DIMENSIONAL $SO(12)$ MODEL

by

C. Wetterich

Deutsches Elektronen-Synchrotron DESY, Hamburg

ISSN 0418-9833

NOTKESTRASSE 85 · 2 HAMBURG 52

DESY behält sich alle Rechte für den Fall der Schutzrechtserteilung und für die wirtschaftliche Verwertung der in diesem Bericht enthaltenen Informationen vor.

DESY reserves all rights for commercial use of information included in this report, especially in case of filing application for or grant of patents.

**To be sure that your preprints are promptly included in the
HIGH ENERGY PHYSICS INDEX,
send them to the following address (if possible by air mail) :**

**DESY
Bibliothek
Notkestrasse 85
2 Hamburg 52
Germany**

symmetric solutions, one may interpret them as spontaneous symmetry breaking of $SO(10)$.

I will concentrate on an $SU(5)$ symmetric example with $n=3, m=p=1$. Harmonic expansion of the six dimensional spinors leads to three massless chiral fermion generations. Under the generation group $SU(2)_G \times U(1)_q$ the fermions of the type u, d, u^c and e^c in the 10 of $SU(5)$ transform as a doublet with charge $q = \frac{1}{2}$ plus a singlet with $q = -\frac{1}{2}$. In contrast, the fields e, ν and d^c in the $\bar{5}$ of $SU(5)$ transform as a triplet with $q = -\frac{1}{2}$. (In addition the model leads to massless right handed neutrinos at this stage.) One observes that quarks and leptons within the same generation transform differently with respect to the generation group $SU(2)_G \times U(1)_q$. This feature is generic for all $m, p \neq 0$. It has important consequences for predictions on mass ratios, since Yukawa couplings will depend on $SU(2)_G \times U(1)_q$ transformation properties.

For a calculation of Yukawa couplings one needs to identify candidates for the weak Higgs doublet responsible for breaking of $SU(2)_L \times U(1)_Y$. They must be contained in the $SO(12)$ gauge fields, noting that the adjoint of $SO(12)$ contains a complex $SO(10) - 10$ plet with $q = \pm 1$. One therefore has two independent $SU(2)_L$ doublets H_1 and H_2 within the six dimensional gauge fields. The internal components of six dimensional gauge fields transform as four dimensional scalars and their harmonic expansion leads to an infinite tower of scalar doublets with various $SU(2)_G$ quantum numbers. However, only the lowest modes can have Yukawa couplings to the chiral fermions whereas all higher modes decouple due to $SU(2)_G$ symmetry. At this stage, the low energy Higgs field must be a combination

$$\varphi = \gamma_i H_i \quad (6)$$

$$\text{with } \sum_i |\gamma_i|^2 = 1.$$

In my example ($n=3, p=m=1$) the lowest modes transform as a $SU(2)_G$ singlet (H_1) and a triplet (H_2^-, H_2^0, H_2^+). Comparison with the quantum numbers of quarks and leptons immediately implies that H_1 can only couple to the bilinear of the $SU(2)_G$ singlet, charge two third quark and antiquark, which I will identify with the top quark:

$$L_{Yuk,1} \sim t_L t_L^c H_1 + \text{h.c.} \quad (7)$$

A similar analysis for H_2 gives the general form

$$L_{Yuk,2} \sim (b_L b_L^c + \tau_L \tau_L^c) H_2^- + (c_L, u_L) \begin{pmatrix} H_2^- & H_2^0 \\ H_2^0 & H_2^+ \end{pmatrix} \begin{pmatrix} c_L^c \\ u_L^c \end{pmatrix} + \text{h.c.} \quad (8)$$

where the different coupling constants still have to be calculated. No Yukawa couplings are allowed for s, μ, d and e as a consequence of the symmetries of the model.

Since the different doublets have all different $SU(2)_G$ quantum numbers, their mixing will be small if $SU(2)_G$ is broken at a scale M_G somewhat below the scale M_C characteristic for spontaneous compactification⁵⁾. Let me assume that the leading contribution to the low energy doublet d comes from H_1 and that the admixtures from H_2^-, H_2^0 and H_2^+ are suppressed by coefficients r_i involving different powers of the ratio M_G/M_C so that

$$\langle H_1 \rangle \gg \langle H_2^- \rangle \gg \langle H_2^0 \rangle, \langle H_2^+ \rangle \quad (9)$$

Once d gets a vacuum expectation value, this would result in a mass hierarchy

$$m_t \gg m_b, m_\tau, m_c \gg m_s, m_\mu, m_u, m_d, m_e \quad (10)$$

resembling already a little bit the observed pattern.

One may go one step further and explicitly calculate the Yukawa couplings of H_1 and H_2 . One finds that they are all proportional to the four dimensional gauge coupling with ratios being given by ratios of $SU(2)_G$ - Clebsch-Gordon-coefficients. The approximation $\langle H_2^0 \rangle = \langle H_2^+ \rangle = 0$ results in the following predictions for fermion masses (M_W in the W -boson mass)

$$m_t = 2M_W \quad (11)$$

$$m_b = m_\tau \quad (12)$$

$$m_c = \sqrt{\frac{4}{3}} m_b \quad (13)$$

$$m_u = m_s = m_d = m_\mu = m_e = 0 \quad (14)$$

Especially eq.(13) can certainly not be considered a good approximation to a realistic spectrum.

In any case, the monopole solution (4) should not be taken as a realistic ground state (even in the limit of unbroken $SU(2)_L \times U(1)_Y$). Its symmetry is too large - $SU(5)$ and the generation group have to be spontaneously broken - and the spectrum of bosonic fields contains tachyons indicating an instability. To obtain a more realistic model with a possible approximate ground state where all gauge symmetries except $SU(3)_C \times SU(2)_L \times U(1)_Y$ are spontaneously broken, one may introduce²⁾ a six dimensional scalar field in the fifth rank totally antisymmetric tensor representation 792 of $SO(12)$. With respect to the $SO(10) \times U(1)_q$ subgroup this field decomposes

$$792 \rightarrow 126_0 + \overline{126}_0 + 210_{+1} + 120_0 \quad (15)$$

The $SU(3)_C \times SU(2)_L \times U(1)_Y$ singlets in 126 and 210 can be responsible for spontaneous symmetry breaking, including the generation group. In the approximation of an unbroken abelian symmetry $U(1)_{\tilde{q}}$ (which is a combination of $U(1)_{\tilde{q}}$, $U(1)_G$, $U(1)_R$ and $U(1)_{B-L}$ with charge $\tilde{q} = I_{3G} + \frac{1}{2}q + aI_{3R}$) one can again perform a quantum number analysis for the various Higgs doublets. One finds that the only two sorts of fields which can mix with H_1 are H_2^- and h , a combination of scalar doublets from the 792-scalar. The mass matrices for the quarks and charged leptons are

$$M_U = \begin{pmatrix} a_{33}H_1 & a_{32}h & 0 \\ a_{23}h & a_{22}H_2^- & 0 \\ 0 & 0 & 0 \end{pmatrix} \quad (16)$$

$$M_D = \begin{pmatrix} b_{33}H_2^- & 0 & b_{32}h^* \\ 0 & b_{22}h^* & 0 \\ 0 & 0 & 0 \end{pmatrix} \quad (17)$$

$$M_E = \begin{pmatrix} c_{33}H_2^- & 0 & 0 \\ 0 & c_{22}h^* & 0 \\ c_{13}h^* & 0 & 0 \end{pmatrix} \quad (18)$$

The non zero Yukawa couplings a_{ij}, b_{ij}, c_{ij} will be calculable for any given ground state with $SU(3)_C \times SU(2)_L \times U(1)_Y \times U(1)_{\tilde{q}}$ symmetry. This symmetry implies that the first generation remains massless

$$m_d = m_u = m_e = 0 \quad (19)$$

For $H_1 \gg H_2^- \gg h$ one finds the pattern

$$m_t \gg m_b, m_c, m_\tau \gg m_\mu, m_s \gg m_u, m_d, m_e \quad (20)$$

In addition, the admixture of h leads to non vanishing mixing angles and CP violating phase. This scenario predicts mixing angles of order $h/H_2^- \sim m_s/m_b \sim$ a few percent for the third generation. Finally, the fermions of the first generation acquire masses once $U(1)_{\tilde{q}}$ is spontaneously broken. These masses put a lower limit on the corresponding symmetry breaking scale. In a realistic model, corrections from $U(1)_{\tilde{q}}$ breaking would replace the zeros in (16)-(18) by entries of order 1-50 MeV. The qualitative structure of these mass matrices is therefore fixed and it would be interesting to investigate if this is consistent with the observed phenomenology for mixing angles and CP violation.

To conclude, let me resume the general scenario how I imagine that higher dimensional theories could explain the hierarchical structure of fermion masses. One observes four groups for the masses of fermions:

$$\begin{matrix} m_t \\ m_b, m_c, m_\tau \\ m_s, m_\mu \\ m_d, m_u, m_e \end{matrix} \quad (21)$$

The overall scale changes from one group to the next roughly by a factor 20. I emphasize that these groups do not correspond exactly to the generations. For example, the second group contains quarks of the second and the third generation. In four dimensional models where generations appear essentially as repetitions, this feature is difficult to understand. I propose an explanation where these groups are induced by quantum numbers of a generation group. Quarks and leptons within the same generation may transform differently with respect to this generation group, as I have shown above.

In usual four dimensional gauge theories Yukawa couplings are free parameters and one may take them as small as one needs. This gives only little predictivity, but these models can accommodate the observed spectrum. In higher dimensional models, four dimensional Yukawa couplings are not free. Typically, they turn out to be of the same order of magnitude as the gauge coupling g or they vanish due to symmetries. At first sight, this seems to run into conflict with a hierarchical mass pattern. However, higher dimensional models typically contain several or many fields with the quantum numbers of the Higgs doublet. As we have seen above, different doublets may couple to different fermions. The idea is that there are essentially four sorts of fields H_1, H_2, H_3, H_4 where H_1 couples only to top and H_2 to the second group, whereas H_3 does not couple to the first generation. If the low energy doublet is mainly H_1 with small admixtures of H_2, H_3 and H_4 obeying

$$\langle H_1 \rangle : \langle H_2 \rangle : \langle H_3 \rangle : \langle H_4 \rangle \sim 20 \quad (22)$$

the qualitative structure of the mass hierarchies would be obtained. Already at this stage the scenario is very restrictive for higher dimensional model building. It is not easy to find models with appropriate quantum numbers for the doublets H_i so that the above pattern of couplings is realized. However, if electron and top quark are allowed to couple to the same scalar field, one typically expects a prediction $m_e \approx m_t$ since all Yukawa couplings are of the same order. Thus a serious problem arises for models not consistent with the above scenario.

It may seem that I only have replaced the problem of explaining small Yukawa couplings by a similar one for the small mixings between scalar fields. The second one, however, will be easier to solve, at least if the model exhibits somewhat different length scales. If the generation group is spontaneously broken at a scale M_G smaller than the compactification scale M_C ($M_G/M_C \sim 1/20$, for example), the mixing of fields with different generation quantum numbers would be suppressed by different powers of M_G/M_C . The hierarchical structure of fermion masses would then be explained by a fine structure of scales near the compactification scale.

ACKNOWLEDGEMENT

I would like to thank the organizers for their work making this pleasant conference possible and successful.

REFERENCES

- 1) C. Wetterich, Nucl. Phys. B223 (1983) 109;
E. Witten, Proc. 1983 Shelter Island II Conf. (MIT press, 1984).
- 2) C. Wetterich, Nucl. Phys. B244 (1984) 359; Nucl. Phys. B260 (1985) 402;
CERN preprint TH 4154 (1985), to appear in Nucl. Phys. B.
- 3) L. Alvarez-Gaumé and E. Witten, Nucl. Phys. B234 (1984) 269.
- 4) E. Cremmer and J. Scherk, Nucl. Phys. B108 (1976) 409; Nucl. Phys. B118 (1977) 61.
Z. Horvath, L. Palla, E. Cremmer and J. Scherk, Nucl. Phys. B127 (1977).
S. Randjbar-Daemi, A. Salam and J. Strathdee, Nucl. Phys. B214 (1983) 491;
Phys. Lett. 124B (1983) 345.
S. Randjbar-Daemi and C. Wetterich, Phys. Lett. 148B (1984) 48.
- 5) G. Lazarides, Q. Shafi and C. Wetterich, Nucl. Phys. B181 (1981) 287.