

DEUTSCHES ELEKTRONEN-SYNCHROTRON
Ein Forschungszentrum der Helmholtz-Gemeinschaft



DESY 20-186
CP3-20-50
MCNet-20-24
VBSCAN-PUB-11-20
IFJPAN-IV-2021-2
arXiv:2011.02547
November 2020

Majorana Neutrinos in Same-Sign $W^\pm W^\pm$ Scattering at the LHC: Breaking the TeV Barrier

B. Fuks

*Sorbonne Université, CNRS, Laboratoire de Physique Théorique et Hautes Énergies,
LPTHE, Paris, France*

and

Institut Universitaire de France, Paris, France

J. Neundorf, K. Peters

Deutsches Elektronen-Synchrotron DESY, Hamburg

R. Ruiz

Institute of Nuclear Physics, Polish Academy of Sciences, Cracow, Poland

and

*Centre for Cosmology, Particle Physics and Phenomenology (CP3),
Université Catholique de Louvain, Chemin du Cyclotron, Louvain la Neuve, Belgium*

M. Saimpert

CERN, Geneva, Switzerland

ISSN 0418-9833

NOTKESTRASSE 85 - 22607 HAMBURG

DESY behält sich alle Rechte für den Fall der Schutzrechtserteilung und für die wirtschaftliche Verwertung der in diesem Bericht enthaltenen Informationen vor.

DESY reserves all rights for commercial use of information included in this report, especially in case of filing application for or grant of patents.

To be sure that your reports and preprints are promptly included in the
HEP literature database
send them to (if possible by air mail):

DESY Zentralbibliothek Notkestraße 85 22607 Hamburg Germany	DESY Bibliothek Platanenallee 6 15738 Zeuthen Germany
---	---

Majorana Neutrinos in Same-Sign $W^\pm W^\pm$ Scattering at the LHC: Breaking the TeV Barrier

Benjamin Fuks,^{1,2,*} Jonas Neundorf,^{3,†} Krisztian Peters,^{3,‡} Richard Ruiz,^{4,5,§} and Matthias Saimpert^{6,¶}

¹*Sorbonne Université, CNRS, Laboratoire de Physique Théorique et Hautes Énergies, LPTHE, F-75005 Paris, France*

²*Institut Universitaire de France, 103 boulevard Saint-Michel, 75005 Paris, France*

³*Deutsches Elektronen-Synchrotron, Notkestraße 85, 22607 Hamburg, Germany*

⁴*Institute of Nuclear Physics, Polish Academy of Sciences, ul. Radzikowskiego, Cracow 31-342, Poland*

⁵*Centre for Cosmology, Particle Physics and Phenomenology (CP3),*

Université Catholique de Louvain, Chemin du Cyclotron, Louvain la Neuve, B-1348, Belgium

⁶*CERN - 1211 Geneva 23 - Switzerland*

(Dated: March 11, 2021)

We revisit the sensitivity to non-resonant, heavy Majorana neutrinos N in same-sign $W^\pm W^\pm$ scattering at the $\sqrt{s} = 13$ TeV LHC and its high-luminosity upgrade. As a benchmark scenario, we work in the context of the Phenomenological Type I Seesaw model, relying on a simulation up to next-to-leading order in QCD with parton shower matching. After extensively studying the phenomenology of the $pp \rightarrow \mu^\pm \mu^\pm jj$ process at the amplitude and differential levels, we design a simple collider analysis with remarkable signal-background separation power. At 95% confidence level we find that the squared muon-heavy neutrino mixing element $|V_{\mu N}|^2$ can be probed down to about $0.06 - 0.3$ ($0.03 - 0.1$) for $m_N = 1 - 10$ TeV with $\mathcal{L} = 300 \text{ fb}^{-1}$ (3 ab^{-1}). For heavier masses of $m_N = 20$ TeV, we report sensitivity for $|V_{\mu N}|^2 \gtrsim 0.5$ (0.3). The $W^\pm W^\pm$ scattering channel can greatly extend the mass range covered by current LHC searches for heavy Majorana neutrinos and particularly adds invaluable sensitivity above a few hundred GeV. We comment on areas where the analysis can be improved as well as on the applicability to other tests of neutrino mass models.

I. INTRODUCTION

Following the discovery of neutrino oscillations [1, 2], uncovering the origin of neutrinos' tiny masses and their large mixing angles are among the most pressing questions in particle physics today [3, 4]. To address these mysteries, neutrino mass models, collectively known as Seesaw models, do so by hypothesizing a variety of states that couple to the Standard Model's (SM) lepton and Higgs sectors [5]. These states include new charged or gauge-singlet (or sterile) fermions, scalars with exotic gauge quantum numbers, as well as gauge bosons of new symmetries, and mediate the non-conservation of lepton number (LN) and/or charged lepton flavor number over a range of mass scales and coupling strengths. For reviews of Seesaw models and their tests, see Refs. [6–9]. Despite these viable solutions, there remains a lack of clear guidance from both experiment and theory as to what is realized by nature. It is therefore necessary to broadly approach this challenge in complementary aspects.

To this extent, tests of neutrino mass models at the Large Hadron Collider (LHC) are supported by a number of signatures, including searches for dijet resonances [10, 11], many-lepton final states [12–17] and

LN-violating lepton pairs [13, 18–20], but rely mostly on mechanisms mediated by quark-antiquark ($q\bar{q}$) annihilation [21]. However, due to its high center-of-mass energy (\sqrt{s}), the LHC is also effectively an electroweak (EW) boson collider [22–24]. This in turn opens a multitude of complementary channels. For example, in the context of the Phenomenological Type I Seesaw model [6, 25], the $W\gamma$ fusion channel [26–29] has already helped direct searches for heavy neutrinos N with masses beyond a few hundred GeV improve sensitivity to active-sterile neutrino mixing matrix elements $V_{\ell N}$. In fact, with an integrated luminosity of $\mathcal{L} \approx 36 \text{ fb}^{-1}$ of proton-proton collisions at $\sqrt{s} = 13$ TeV, $|V_{\ell N}|^2 \gtrsim \mathcal{O}(0.01 - 1)$ are excluded for lepton flavors $\ell \in \{e, \mu\}$ and sterile neutrino masses in the range $m_N = 100 \text{ GeV} - 1 \text{ TeV}$ [13, 18]. With the full LHC data set, this degree of sensitivity is anticipated to reach masses up to $m_N = 3 - 4 \text{ TeV}$ [30].

Motivated by the recent experimental observations of EW vector boson fusion / scattering (VBF) at the LHC [31–35], we revisit the sensitivity of same-sign $W^\pm W^\pm$ scattering to TeV-scale Majorana neutrinos at $\sqrt{s} = 13$ TeV. As a benchmark scenario, we work in the framework of the Phenomenological Type I Seesaw and focus on the production of same-sign muon pairs ($\mu^\pm \mu^\pm$) without substantial transverse momentum imbalance via the spacelike exchange of an N in $W^\pm W^\pm$ scattering [36],

$$W^\pm W^\pm \rightarrow \mu^\pm \mu^\pm. \quad (1.1)$$

As shown in figure 1, this is essentially a realization of neutrinoless $\beta\beta$ ($0\nu\beta\beta$) decay at large momentum transfer when mediated at dimension $d = 7$ [37–40]. While

* fuks@lpthe.jussieu.fr

† jonas.neundorf@desy.de

‡ krisztian.peters@desy.de

§ rruiz@ifj.edu.pl

¶ matthias.saimpert@cern.ch




FIG. 1. Diagrammatic representation of same-sign $\ell_i^+ \ell_j^+$ production in same-sign W^+W^+ scattering in pp collisions mediated by a Majorana neutrino N . Drawn with JaxoDraw [44].

past studies have investigated the importance of this channel [6, 26, 36, 41–43], most works were restricted to sub-TeV m_N , and therefore subject to signal processes with more dominant cross sections. As the heavy neutrino exchange in equation (1.1) is always non-resonant, the channel is complementary to other processes, such as the $qq' \rightarrow N\ell$ annihilation and $W\gamma \rightarrow N\ell$ fusion mechanisms, which become kinematically inaccessible for sterile neutrinos that are too heavy. For similar reasons, the channel is robust against the impact of long N lifetimes.

To conduct this study, we employ a state-of-the-art simulation tool chain that allows us to model SM backgrounds and, for the first time, the $W^\pm W^\pm \rightarrow \ell_i^\pm \ell_j^\pm$ signal process at next-to-leading order (NLO) in QCD with parton shower (PS) matching. We report remarkable signal-background separation power and attribute this to the signal process exhibiting both VBF and LN-violating topologies. For the $pp \rightarrow \mu^\pm \mu^\pm jj$ collider signature with forward jet-tagging and simple selection cuts, we find that $|V_{\mu N}|^2 \gtrsim 0.06-0.3$ ($0.03-0.1$) can be probed at 95% confidence level (CL) for $m_N = 1-10$ TeV with $\mathcal{L} = 300 \text{ fb}^{-1}$ (3 ab^{-1}). For masses of $m_N = 20$ TeV we find sensitivity for $|V_{\mu N}|^2$ down to 0.5 (0.3).

The remainder of this work is organized in the following manner: First, we describe in section II our theoretical framework and give an overview of current experimental constraints. Next, we summarize our computational setup (section III) and simulation prescriptions (section IV). In section V we explore extensively the phenomenology of the $W^\pm W^\pm \rightarrow \ell_i^\pm \ell_j^\pm$ signal process at the amplitude and differential levels. This is then used in section VI to design our collider analysis and estimate the LHC’s sensitivity to LN violation (LNV) in $W^\pm W^\pm$ scattering. We provide an outlook in section VII on areas where our analysis can be improved as well as its applicability to other tests of neutrino mass models at colliders. Finally, we conclude in section VIII. Where relevant, technical derivations and details on software modifications are reported in appendices A, B, and C.

II. THEORETICAL FRAMEWORK

We describe in this section the theoretical framework in which we work (section II.1) and summarize current experimental constraints on the model (section II.2).

II.1. The Phenomenological Type I Seesaw Model

To study the sensitivity of the LHC to the LN-violating $W^\pm W^\pm \rightarrow \ell_i^\pm \ell_j^\pm$ process, we work in the context of the Phenomenological Type I Seesaw model [6, 25]. Like the eponymous mechanism [45–51] or its low-scale variants [52–57], the model hypothesizes the existence of Majorana neutrinos ($N_{k'}$) that couple to SM particles through mass mixing with light neutrinos (ν_k). More precisely, the renormalizable Lagrangian of the theory,

$$\mathcal{L}_{\text{Type I}} = \mathcal{L}_{\text{SM}} + \mathcal{L}_{\text{Kin.}} + \mathcal{L}_{\text{Y}}, \quad (2.1)$$

extends the SM Lagrangian (\mathcal{L}_{SM}) by kinetic and Majorana mass terms ($\mathcal{L}_{\text{Kin.}}$) for $n_R \geq 2$ right-handed (RH) neutrinos (ν_R^i), as well as by Yukawa couplings (\mathcal{L}_{Y}) between the SM Higgs field, the SM lepton doublets (L^j) and the gauge-singlet fermions ν_R^i .

After EW symmetry breaking, flavor eigenstates of active, left-handed (LH) neutrinos ($\nu_{L\ell}$) can be generically [6] decomposed into light (ν_k) and heavy ($N_{k'}$) mass eigenstates with mass eigenvalues m_{ν_k} and $m_{N_{k'}}$:

$$\nu_{L\ell} = \sum_{k=1}^3 U_{\ell k} \nu_k + \sum_{k'=4}^{n_R+3} V_{\ell k'} N_{k'}. \quad (2.2)$$

Here, the complex-valued matrix elements $U_{\ell k}$ ($V_{\ell k'}$) parameterize the mixing between the LH interaction state $\nu_{L\ell}$ and the light (heavy) mass eigenstate ν_k ($N_{k'}$). Formally, the matrix elements satisfy the relationship $UU^\dagger + VV^\dagger = \mathbb{I}$. For clarity and without loss of generality, we order states such that $m_{N_{k'}} < m_{N_{k'+1}}$.

To leading order in active-sterile mixing $|V_{\ell k'}|$, equation (2.2) gives rise to the following effective, charged current component of the theory’s interaction Lagrangian,

$$\begin{aligned} \Delta\mathcal{L} = & -\frac{g_W}{\sqrt{2}} W_\mu^+ \sum_{k=1}^3 \sum_{\ell}^{\tau} [\bar{\nu}_k U_{\ell k}^* \gamma^\mu P_L \ell] \\ & -\frac{g_W}{\sqrt{2}} W_\mu^+ \sum_{k'=4}^{n_R+3} \sum_{\ell}^{\tau} [\bar{N}_{k'} V_{\ell k'}^* \gamma^\mu P_L \ell] + \text{H.c.} \end{aligned} \quad (2.3)$$

In the above expression, $g_W \approx 0.65$ is the SM weak gauge coupling constant and $P_{L/R} = (1/2)(1 \mp \gamma^5)$ are the standard chiral projection operators in four-component notation. Similar terms can be derived for the neutral current (Z) and Higgs interactions for both heavy Dirac and Majorana neutrinos [6]. For simplicity we solely investigate the phenomenology of the lightest heavy neutrino mass eigenstate, which we relabel as $N \equiv N_{k'=4}$ (so that $V_{\ell N} \equiv V_{\ell k'=4}$), and decouple all other heavy eigenstates.

In short-distance scattering and decay processes involving only a single heavy neutrino, the mixing factors that appear in equation (2.3) act as effective couplings and factor out of amplitudes. For resonant production of a heavy neutrino, this allows one to define a “bare” cross section σ_0 such that [58]

$$\sigma(pp \rightarrow N\ell^\pm + X) \equiv |V_{\ell N}|^2 \times \sigma_0(pp \rightarrow N\ell^\pm + X). \quad (2.4)$$

When the $W^\pm W^\pm \rightarrow \ell_i^\pm \ell_j^\pm$ process involves only a single t -channel exchange of a heavy neutrino, such a factorization is also possible. More specifically, one can define a “bare” t -channel cross section given by

$$\sigma(pp \rightarrow \ell_i^\pm \ell_j^\pm + X) \equiv |V_{\ell_i N} V_{\ell_j N}|^2 \times \sigma_0(pp \rightarrow \ell_i^\pm \ell_j^\pm + X). \quad (2.5)$$

For our purposes, it suffices to add that these expressions hold at NLO in QCD [29, 59]. Under certain assumptions, the above decomposition or a similar one can hold for processes involving multiple interfering heavy neutrino mass eigenstates. Such expressions, however, may not always be tractable due to large interference effects. For further discussions, see Refs. [60–64].

II.2. Model Constraints

In its most general construction, the free parameters of the Phenomenological Type I Seesaw model (beyond those of the SM) consist of the neutrino masses m_{ν_k} , $m_{N'_k}$, and the neutrino mixing elements $U_{\ell k}$, $V_{\ell k'}$. Imposing flavor symmetries on the lepton sector, however, can reduce the number of independent degrees of freedom, as discussed in Refs. [60–62, 65, 66] and references therein. It is also possible to constrain these parameters by tying the lightness of the ν_k neutrinos to beyond the SM physics, such as to dark matter and the baryon asymmetry of the universe, as done for example in Refs. [67, 68]. For our purposes, we take mass and mixing parameters to be phenomenologically independent. We do this to develop a collider analysis in a flavor-model-independent fashion and is motivated by the desire to broaden sensitivity to a range of ultraviolet completions.

Beyond theoretical considerations are the following recent experimental constraints on the model:

- **Direct constraints from $0\nu\beta\beta$ searches:** After an exposure of 127.2 kg-yr, the GERDA experiment reports at 90% CL the following lower limit on the $0\nu\beta\beta$ decay half-life in ^{76}Ge [69]:

$$T_{1/2}^{0\nu} > 1.8 \cdot 10^{26} \text{ yr}. \quad (2.6)$$

Assuming that the nuclear $0\nu\beta\beta$ process is only mediated by heavy neutrinos, this translates into an upper limit on their masses and mixing of

$$\left| \sum_{k'=4}^{n_R+3} \frac{V_{ek'}^2}{m'_{k'}} \right| < (2.33 - 4.12) \cdot 10^{-6} \text{ TeV}^{-1}, \quad (2.7)$$

where the variation stems from the uncertainties in the nuclear matrix element. For further details on the derivation of this constraint, see appendix A.

- **Direct constraints from collider searches:** At $\sqrt{s} = 13 \text{ TeV}$ and with $\mathcal{L} \approx 36 \text{ fb}^{-1}$ of data, searches for the $pp \rightarrow \ell_i \ell_j \ell_k + E_T^{\text{miss}}$ signature with $\ell \in \{e, \mu\}$, by the CMS experiment constrain active-sterile neutrino mixing at 95% CL to be [13],

$$\begin{aligned} |V_{\ell N}|^2 &\lesssim 10^{-5} - 10^{-2} \text{ for } 1 \text{ GeV} < m_N < m_W, \\ |V_{\ell N}|^2 &\lesssim 10^{-2} - 1 \text{ for } m_W < m_N < 1.2 \text{ TeV}. \end{aligned} \quad (2.8)$$

Constraints from the ATLAS experiment with the same integrated luminosity are comparable for $m_N < m_W$ but weaker for $m_N > m_W$ due to the absence of the $W\gamma$ channel in their signal modeling [15]. Searches for the $pp \rightarrow \ell_i^\pm \ell_j^\pm + n_j$ signature by CMS yield only slightly more stringent constraints due to a larger signal-over-background ratio [18].

- **Indirect constraints on $V_{\ell k'}$:** For $n_R = 3$ sterile neutrinos with masses above the EW scale, a global study of precision EW data, searches for non-unitarity in quark mixing, and searches for lepton flavor violation and non-universality of weak decays constrain active-sterile neutrino mixing to be [70],

$$\begin{aligned} \sqrt{2|\eta_{ee}|} &< 0.050, & \sqrt{2|\eta_{e\mu}|} &< 0.026, \\ \sqrt{2|\eta_{\mu\mu}|} &< 0.021, & \sqrt{2|\eta_{e\tau}|} &< 0.052, \\ \sqrt{2|\eta_{\tau\tau}|} &< 0.075, & \sqrt{2|\eta_{\mu\tau}|} &< 0.035, \end{aligned} \quad (2.9)$$

at 95% CL. The parameter $\eta_{\ell\ell'}$ is related to the heavy neutrino mixing matrix $V_{\ell k'}$ by $\sqrt{2|\eta_{\ell\ell'}|} = \sum_{k'=4}^6 \sqrt{V_{\ell k'} V_{\ell' k'}^*}$. For the scenario that we consider, *i.e.*, $\ell = \ell' = \mu$ with only one heavy neutrino species, this translates to an upper limit of

$$|V_{\mu N}|^2 < 4.41 \cdot 10^{-4} \text{ at } 95\% \text{ CL}. \quad (2.10)$$

- **Direct constraints on the absolute mass scale of light neutrinos:** Attempts to measure the light neutrino mass scale directly from the kinematic end point in β decay with the KATRIN experiment [71] constrain the light neutrino masses to satisfy [72]

$$m(\nu_e) < 1.1 \text{ eV} \text{ at } 90\% \text{ CL}. \quad (2.11)$$

- **Constraints on neutrino masses from cosmology:** An analysis of neutrinos’ impact on the cosmic microwave background, supernovae, large scale structure, and big bang nucleosynthesis constrains the sum of light neutrino masses to be [73]

$$\sum m_\nu \lesssim 0.26 \text{ eV} \text{ at } 95\% \text{ CL}. \quad (2.12)$$

- **Neutrino oscillation measurements of $U_{\ell k}$:** In the absence of sterile neutrinos or additional new physics, the elements of the light neutrino mixing matrix $U_{\ell k}$ have been fit to or constrained by long and short baseline neutrino oscillation data under the condition that U is unitary [74]. Relaxing this constraint, however, greatly weakens the goodness of fit, particularly in the τ flavor sector [75, 76].

III. COMPUTATIONAL SETUP

We now summarize the computational setup used in this study. We start with section III.1 where we document our Monte Carlo (MC) simulation chain. In section III.2 we list the numerical values used for SM inputs, and similarly in section III.3 the numerical values used for non-SM inputs. A description on how we model the $W^\pm W^\pm \rightarrow \ell_i^\pm \ell_j^\pm$ signal process and leading background processes in pp collisions is deferred to section IV.

III.1. Monte Carlo Setup

To investigate same-sign $W^\pm W^\pm$ scattering when mediated by a heavy Majorana neutrino at the $\sqrt{s} = 13$ TeV LHC, we employ a state-of-the-art simulation tool chain. For hard, parton-level scattering processes, we use the general-purpose MC event generator MADGRAPH5_AMC@NLO (MG5AMC) (version 2.7.1.2) [77, 78], which enables us to simulate tree-induced processes in the SM up to NLO in QCD [79–82]. Processes involving heavy neutrinos are simulated up to NLO in QCD by importing into MG5AMC the default variant of the HEAVYN [29] FEYNRULES UFO libraries [83–87]. For select backgrounds we perform jet matching up to one additional parton (at the Born level) at NLO in QCD precision using the FxFx matching procedure [88], as further detailed in section IV. To simulate the decay of W bosons, we impose the spin-correlated narrow width approximation as implemented in MADSPIN [89, 90].

All signal and background events are passed through PYTHIA8 (version 243) [91] for QCD and QED parton showering up to leading logarithmic (LL) accuracy, hadronization, and multiparton interaction / underlying event modeling. Decays of heavy-flavored hadrons and τ leptons are handled internally by the PS. Following Refs. [92, 93], we apply the improved color reconnection [94] and dipole recoil [95] models available in PYTHIA8. For simplicity, activity from additional proton-proton interactions occurring during the same bunch crossing, *i.e.*, pileup, is assumed to be subtracted from experimental data by dedicated algorithms and, therefore, are not included in our simulations.

In our analysis, hadron-level events are passed to the fast detector simulator DELPHES (version 3.4.2) [96]. There, particle-level clustering of hadrons is handled according to the anti- k_T algorithm [97–99] as implemented

in FASTJET [100, 101], with a radius parameter $R = 0.4$. To emulate experimental reconstruction with realistic detector resolution and particle identification, detector responses are tuned using the ATLAS configuration card available in the DELPHES repository. Particle-level distributions at LO+PS and NLO+PS were checked using MADANALYSIS5 (version 1.8) [102–104].

III.2. Standard Model Inputs

For SM inputs we work in the $n_f = 4$ active quark flavor scheme with a Cabibbo-Kobayashi-Maskawa mixing matrix that is diagonal with unit entries. Unless specified, we assume the following mass and coupling values:

$$\begin{aligned} m_t(m_t) &= 172.9 \text{ GeV}, & m_b(m_b) &= 4.7 \text{ GeV}, \\ m_\tau &= 1.777 \text{ GeV}, & m_h &= 125.1 \text{ GeV}, \\ M_Z &= 91.188 \text{ GeV}, & \alpha_{\text{QED}}^{-1}(M_Z) &= 132.5070, \\ G_F &= 1.166390 \cdot 10^{-5} \text{ GeV}^2. \end{aligned} \quad (3.1)$$

For scattering computations at both LO and NLO in QCD we employ the NNPDF3.1 NLO+LUXqed parton distribution function (PDF) set with $\alpha_s(M_Z) = 0.118$ (lhaid=324900) [105–107]. For non-perturbative dynamics we tune the shower with the ATLAS A14 central tune (Tune:pp=21) as paired with the NNPDF2.3 LO+QED PDF set with $\alpha_s(M_Z) = 0.130$ (pdfcode=247000) [108]. PDF and α_s scale evolution are handled by LHAPDF (version 6.2.3) [109]. PDF uncertainties are extracted using replicas PDFs [107, 109].

For signal and background processes, we set the nominal ($\zeta = 1.0$) collinear factorization (μ_f) and QCD renormalization (μ_r) scales to be half the sum over each visible, final-state particle’s transverse energy:

$$\mu_f, \mu_r = \zeta \times \mu_0, \text{ with } \mu_0 = \frac{1}{2} \sum_f \sqrt{m_f^2 + p_{T,f}^2}. \quad (3.2)$$

Here m_f and $p_{T,f}$ stand for the mass and transverse momentum of the final-state particle f respectively. The shower factorization scale ($\mu_s = \zeta \times \hat{\mu}_s$) is kept at its default value prescribed in Ref. [78]. To estimate the size of higher-order QCD corrections, we vary discretely and independently μ_f, μ_r, μ_s , over the set $\zeta = \{0.5, 1.0, 2.0\}$ to obtain a nine- or 27-point uncertainty band.

III.3. Heavy Neutrino Inputs

For simulations involving the heavy neutrino N we use the default inputs of the Majorana neutrino variant of the HEAVYN NLO UFO libraries [29]. SM particle masses are updated according to equation (3.1). As we are interested in the benchmark scenario featuring only one heavy mass eigenstate, additional heavy mass eigenstates are decoupled by setting $m_{N_5}, m_{N_6} = 10^{10}$ GeV. To deactivate e

and τ flavor mixing, we set

$$|V_{eN}|^2, |V_{\tau N}|^2 = 0, \quad |V_{\mu N}|^2 = 1.0. \quad (3.3)$$

Sensitivity to smaller values of $|V_{\mu N}|$ are obtained by a naïve rescaling of cross sections, which is permissible by equation (2.5). Notably, as N is never resonantly produced, its total width (Γ_N) and lifetime can be ignored.

IV. SIGNAL AND BACKGROUND MODELING IN MADGRAPH5_AMC@NLO

Generically speaking, collider processes that feature either LNV or VBF exhibit characteristic kinematical and topological properties that enable remarkable background rejection capabilities. Consequentially, background modeling for the $W^\pm W^\pm \rightarrow \ell_i^\pm \ell_j^\pm$ channel is hindered by high background rejection rates, and hence by poor MC efficiencies. In this context, we report the development of efficient MC modeling prescriptions that overcome such difficulties for our signal (section IV.1) and leading backgrounds (section IV.2) at NLO+PS within the MG5AMC simulation framework.

IV.1. Signal Modeling

To model the $W^\pm W^\pm \rightarrow \ell_i^\pm \ell_j^\pm$ process when mediated by the exchange of a t -channel Majorana neutrino N , as shown in figure 1, we consider the gauge-invariant set of $W^\pm W^\pm$ scattering diagrams contributing to the following $2 \rightarrow 4$, Born-level process at $\mathcal{O}(\alpha^4)$

$$q_1 q_2 \rightarrow q'_1 q'_2 \ell_i^\pm \ell_j^\pm. \quad (4.1)$$

Here q denotes any light quark or antiquark. We neglect interference with the s -channel process, $q\bar{q} \rightarrow W^{\pm*} \rightarrow N\ell_i^\pm \rightarrow \ell_i^\pm \ell_j^\pm q\bar{q}$. To justify this, we require that the leading dijet system at the analysis level carries a large invariant mass, which suppresses the $N^{(*)} \rightarrow W^{(*)}\ell \rightarrow q\bar{q}\ell$ splitting chain. For m_N below the TeV scale, neglecting the $q\bar{q}$ annihilation mechanism is also justified by the narrow width approximation. Under this the leading contributions to the s -channel process are factorizable and non-interfering with t -channel diagrams; interfering contributions are $\mathcal{O}(\Gamma_N/m_N) \ll 1$, and hence insignificant.

To carry out this modeling in MG5AMC at NLO in QCD using the HEAVYN libraries, we employ the syntax¹

```
import model SM_HeavyN_NLO
define p = g u c d s u~ c~ d~ s~
define j = p
generate p p > mu+ mu+ j j QED=4 QCD=0 $$
w+ w- / n2 n3 [QCD]
add process p p > mu- mu- j j QED=4 QCD=0 $$
w+ w- / n2 n3 [QCD]
```

¹ For further details on syntax and usage, see Refs. [29, 78].

Formally, the Born-level matrix element for equation (4.1) is finite in the absence of phase space cuts. At $\mathcal{O}(\alpha_s)$, however, an infrared-safe definition for external states is needed. We therefore require at the generator-level that QCD partons are sequentially clustered according to the anti- k_T algorithm with $R = 0.4$ and that the transverse momentum (p_T) and pseudorapidity (η) of these clusters satisfy the following demands:

$$p_T^j > 20 \text{ GeV} \quad \text{and} \quad |\eta^j| < 5.5. \quad (4.2)$$

As a technical remark, we relax checks on infrared pole cancellation in MADFKS. Such checks are automatically raised in MG5AMC for VBF processes due to the possible omission of virtual diagrams that are mixed NLO QCD-EW corrections and not pure QCD contributions at $\mathcal{O}(\alpha_s)$. In our case such diagrams do not exist and therefore bypassing the check only impacts the MC efficiency. To do this we set #IRPoleCheckThreshold=-1.0d0 in the file Cards/FKS_params.dat.

IV.2. Background Modeling

Due to the presence of forward, high- p_T jets, a high-energy, same-sign lepton pair, and an absence of (light) neutrinos in our signal process, several background processes that are traditionally present in collider searches for LNV can be readily suppressed through simple kinematic requirements. Among many examples are: demanding that the leading dijet system carries a large invariant mass, stringent p_T cuts on the same-sign leptons, and vetoing events with three or more charged leptons. As a result, background categories such as associated top quark production, $t\bar{t}B$ with $B \in \{W^\pm/\gamma^*/Z, h\}$; single top quark channels, tB ; and triboson production WWV can be neglected for the purposes of our study.

The leading background channels that remain after such baseline selection criteria include: the mixed EW-QCD channel $W^\pm W^\pm jj$ (section IV.2.1), the pure EW channel $W^\pm W^\pm jj$ (section IV.2.2), and the mixed EW-QCD diboson+jets process $W^\pm V + nj$ with $V \in \{\gamma^*/Z\}$ (section IV.2.3). To model these processes in MG5AMC, we employ the prescriptions described below.

IV.2.1. QCD Production of Same-Sign $W^\pm W^\pm jj$

Due to its similar topology and kinematic scales, the mixed EW-QCD production (henceforth labeled QCD production) of $pp \rightarrow W^\pm W^\pm jj$ is a prominent background for the $pp \rightarrow \ell^\pm \ell^\pm jj$ signal process. However, a defining characteristic of this mode, which at the Born level occurs at $\mathcal{O}(\alpha^2 \alpha_s^2)$, is the t -channel exchange of a gluon. This indicates an intermediate flow of color. As such, the presence of a third, high- p_T jet is significantly more likely in this process than in the signal process. Such radiation can induce recoils in the $(W^\pm W^\pm)$ -system, siphon energy from the two forward jets, or give

TABLE I. Generator-level cross sections [fb] and cuts, μ_f, μ_r scale uncertainty [%], PDF uncertainties [%], and perturbative order for leading backgrounds at $\sqrt{s} = 13$ TeV.

Process	Order	Cuts	$\sigma^{\text{Gen.}}$ [fb]	$\pm\delta_{\mu_f, \mu_r}$	$\pm\delta_{\text{PDF}}$
$W^\pm W^\pm jj$ (QCD)	NLO in QCD	Eq. (4.2)	385	+10% -10%	+1% -1%
$W^\pm W^\pm jj$ (EW)	NLO in QCD	Eq. (4.2) + diagram removal	254	+1% -1%	+1% -1%
Inclusive $W^\pm V$ ($3\ell\nu$)	FxFx (1j)	Eqs. (4.3), (4.4)	2,520	+5% -6%	+1% -1%

rise to excess central hadronic activity [110–114]. One is thus motivated to consider the process at NLO in QCD.

We do this in MG5AMC by using the syntax:

```
import model loop_sm
generate p p > w+ w+ j j QED=2 QCD=2 [QCD]
add process p p > w- w- j j QED=2 QCD=2 [QCD]
```

As in the signal process, we impose the criteria of equation (4.2) on outgoing QCD partons. We report in the first line of table I the corresponding generator-level cross section at $\sqrt{s} = 13$ TeV, which reaches $\sigma \sim 385$ fb, along with residual μ_f, μ_r scale and PDF uncertainties. Before parton showering, resonant W bosons are decayed to muons (see section III.1 for related details).

For simplicity, we neglect contributions from leptonic tau decays. We do so because the presence of additional light neutrinos results in events with characteristically softer muons and larger momentum imbalances. Such features can be tamed by tuning the pre-selection and signal region cuts used in section VI.

IV.2.2. EW Production of Same-Sign $W^\pm W^\pm jj$

Like the previous case, pure EW production of $pp \rightarrow W^\pm W^\pm jj$, which at the Born level occurs at $\mathcal{O}(\alpha^4)$, is a leading background to the $pp \rightarrow \ell^\pm \ell^\pm jj$ signal process. Similarities to the signal process’s topology, kinematic characteristics, and, importantly, color flow are identifiable in the $W^\pm W^\pm \rightarrow W^\pm W^\pm$, VBF sub-process. While featuring a large cross section at the inclusive level, other sub-processes, such as resonant triboson production, are essentially removed through basic selection criteria.

To efficiently model the EW same-sign $W^\pm W^\pm$ channel at NLO in QCD, we make a variant of the so-called “vector boson fusion approximation” and neglect resonant triboson production in a gauge-invariant manner. We do this under the presumption that relevant analysis-level selection cuts are applied. Other non-resonant, interfering diagrams are kept. The syntax we employ is

```
generate p p > w+ w+ j j $$
a z w+ w- QCD=0 QED=4 [QCD]
add process p p > w- w- j j $$
a z w+ w- QCD=0 QED=4 [QCD]
```

We apply the same loose, generator-level cuts as listed in equation (4.2) and report good numerical stability. In the second line of table I, we report the generator-level cross section at $\sqrt{s} = 13$ TeV, which reaches $\sigma \sim 254$ fb, and associated uncertainties. Resonant W bosons are decayed to muons before parton showering.

IV.2.3. Inclusive Diboson Spectrum

Due to its picobarn-scale rate, the inclusive $pp \rightarrow 3\ell\nu + X$ spectrum (which we imprecisely label “diboson spectrum” even though the process includes interference with all non-resonant diagrams) contributes to the same-sign dilepton signature $pp \rightarrow \ell^\pm \ell^\pm jj$ through pathological configurations of the final-state kinematics. Such configurations include, for example, when two or more initial-state QCD emissions both possess large p_T but the odd-sign charged lepton is not successfully identified because it is too soft in p_T or too forward in $|\eta|$.

While a bulk of these phase space configurations are captured by the fixed-order matrix element for the $pp \rightarrow 3\ell\nu jj$ process, which at the Born level occurs at $\mathcal{O}(\alpha^4 \alpha_s^2)$, convergence in perturbative QCD requires that outgoing QCD partons are hard (high p_T) and central (low $|\eta|$). In light of the anticipated scales of the ($3\ell\nu$)-system, the component of phase space where one QCD parton is central and one is forward is better described by the fixed-order matrix element for the $pp \rightarrow 3\ell\nu j$ sub-process, with the second j being populated by the PS. Likewise, for two forward emissions, the phase space is better described best by the $pp \rightarrow 3\ell\nu$ sub-process with two PS emissions.

To model these complications we extend the brute-force prescription of Ref. [30]. This entails starting with the $2 \rightarrow 4$ process, $pp \rightarrow 3\ell\nu$ at NLO in QCD, with all interfering, non-resonant diagrams and without invoking the narrow width approximation for intermediate W, Z bosons. Instead, loose, generator-level cuts on leptons are imposed to regulate s - and t -channel divergences, thereby keeping matrix elements finite and perturbative. Specifically, we require that charged leptons satisfy

$$m_{\text{os}}^{\ell\ell} > 8 \text{ GeV} \quad \text{and} \quad |\eta^\ell| < 4.0, \quad (4.3)$$

where $m_{\text{os}}^{\ell\ell}$ is the invariant mass of any opposite-sign, charged lepton pair, independent of flavor. While smaller $m_{\text{os}}^{\ell\ell}$ thresholds can still regulate $\gamma^* \rightarrow \ell\ell$ splittings, we refrain from doing so to avoid contributions from vector meson resonances. Such states are not modeled with the perturbative event generator MG5AMC.

To account for additional central jet multiplicities, we match the inclusive $pp \rightarrow 3\ell\nu$ spectrum at NLO+PS up to its first jet multiplicity (relative to the Born level) at NLO in QCD using the FxFx prescription [88]. The relevant MG5AMC syntax in this case is

```
define ell = e+ mu+ ta+ e- mu- ta-
define vv = ve vm vt ve~ vm~ vt~
generate p p > ell ell ell vv
```



```

QED=4 QCD=0 [QCD] @0
add process p p > e1l e1l vv j
QED=4 QCD=1 [QCD] @1

```

Explicitly, we use the following jet-matching inputs:

$$p_T^j > 30 \text{ GeV}, \quad |\eta^j| < 5.5, \quad Q_{\text{cut}}^{\text{FxFx}} = 65 \text{ GeV}, \quad (4.4)$$

where $Q_{\text{cut}}^{\text{FxFx}}$ is the FxFx matching scale. With this setup, hadronic observables are accurate to at least LO+PS(LL), with the two leading jets being defined at all momenta and rapidities. We report in the third line of table I the generator-level cross section of the diboson spectrum at NLO in QCD with FxFx matching to the first jet multiplicity (FxFx1j) and its associated uncertainties. At $\sqrt{s} = 13 \text{ TeV}$ the rate is about $\sigma \sim 2.5 \text{ pb}$, which is just slightly larger than the rate at NLO, which we compute to be $\sigma \sim 2.3 \text{ pb}$.

With this setup, we capture configurations where the odd-sign charged lepton in the $3\ell\nu 2j + X$ final state is too forward or too soft to be identified as an analysis-quality charged lepton. A disadvantage of this setup, however, is the limited MC statistics when the two same-sign charged leptons carry $p_T^\ell \gtrsim 100 - 150 \text{ GeV}$ but the odd-sign lepton is much softer. To enrich MC statistics for this region of phase space, we introduce tailored generator-level cuts into the MG5AMC phase space integration routines. Enriched samples are combined with the baseline FxFx1j sample. Overlap is removed through cuts on $p_T^{\ell_2}$. For technical details of this modeling, see appendix B.

V. HEAVY NEUTRINOS IN $W^\pm W^\pm$ SCATTERING AT THE LHC

In this section we investigate the phenomenology of the $W^\pm W^\pm \rightarrow \ell_i^\pm \ell_j^\pm$ process when mediated by a heavy Majorana neutrino at the LHC. To do this, we examine the integrated (section V.1) and differential (section V.2) cross sections of the $W^\pm W^\pm$ channel, and place special focus on the low- (section V.1.1) and high-mass (section V.1.2) limits of the intermediate neutrino, on the impact of QCD corrections (section V.1.3), and on potential violations of partial-wave unitarity (section V.1.4).

V.1. Total Production Rate

As a first step, we present in the upper panel of figure 2 and as a function of heavy neutrino mass m_N , the total cross section for the full $2 \rightarrow 4$, hadron-level process

$$pp \rightarrow \ell^\pm \ell^\pm jj + X. \quad (5.1)$$

More precisely, we evaluate the bare cross section, as defined in equation (2.5), at NLO in QCD and for LHC collisions at $\sqrt{s} = 13 \text{ TeV}$, assuming the exchange of a single heavy neutrino that couples to a single charged lepton flavor. In equation (5.1), X denotes any additional




FIG. 2. Upper: As a function of heavy neutrino mass m_N [GeV], the bare cross section $\sigma/|V_{\ell N} V_{\ell N}|^2$ [fb] for the $W^\pm W^\pm$ signal process at NLO in QCD (purple band), as well as the bare cross sections $\sigma/|V_{\ell N}|^2$ for the CCDY (black band) and $W\gamma$ fusion (green band) processes at NLO in QCD. Band thickness corresponds to the residual scale uncertainty. Lower: The QCD K -factor for each channel.

hadronic and photonic activity present in the inclusive process. The band thickness corresponds to the residual renormalization and collinear factorization scale dependence at NLO, as quantified in section III.2. We assume the generator-level cuts of equation (4.2). To quantify the size of $\mathcal{O}(\alpha_s)$ corrections, we show in the lower panel of figure 2 the NLO in QCD K -factor, defined as the ratio of the NLO and LO cross sections:

$$K^{\text{NLO}} = \sigma^{\text{NLO}} / \sigma^{\text{LO}}. \quad (5.2)$$

For the mass range $m_N = 40 \text{ GeV} - 20 \text{ TeV}$, we report that bare cross sections at NLO in QCD, QCD K -factors, as well as scale and PDF uncertainties roughly span

$$\sigma^{\text{NLO}} : 0.1 - 20 \text{ fb}, \quad (5.3)$$

$$K^{\text{NLO}} : 1.05 - 1.4, \quad (5.4)$$

$$\delta\sigma_{\mu_r, \mu_f} / \sigma : \pm 1\% - \pm 5\%, \quad (5.5)$$

$$\delta\sigma_{\text{PDF}} / \sigma : \pm 1\% - \pm 2\%. \quad (5.6)$$

A summary of bare cross sections and uncertainties for the $W^\pm W^\pm \rightarrow \ell^\pm \ell^\pm$ process at representative heavy neutrino masses is listed in table II.

To compare to other heavy neutrino processes,² we also present in figure 2 the bare cross sections at NLO in

² For these additional channels, we follow the prescription of

TABLE II. For representative heavy neutrino masses (m_N) and active-sterile mixing $V_{\ell N} = 1$, the $pp \rightarrow \ell^\pm \ell^\pm jj + X$ cross section [fb] at NLO in QCD, with residual scale uncertainties [%], PDF uncertainties [%], and NLO K -factor.

m_N	σ^{NLO} [fb]	$\pm\delta_{\mu_f, \mu_r}$	$\pm\delta_{\text{PDF}}$	K^{NLO}
150 GeV	13.3	+1% -2%	+1% -1%	1.09
1.5 TeV	8.45	+4% -4%	+1% -1%	1.26
5.0 TeV	1.52	+5% -5%	+2% -2%	1.32
15 TeV	0.190	+5% -5%	+2% -2%	1.32

QCD, as defined in equation (2.4), the associated scale uncertainties, and QCD K -factor for the $2 \rightarrow 2$, charged current Drell-Yan (CCDY) process (black band),

$$pp \rightarrow W^{\pm(*)} + X \rightarrow N\ell^\pm + X, \quad (5.7)$$

and the $2 \rightarrow 3$, $W\gamma$ fusion process (green band),

$$pp \rightarrow N\ell^\pm j + X. \quad (5.8)$$

We find several notable observations: First is that, quantitatively, the bare, same-sign WW cross section is about 4–6 (1–3) orders of magnitude smaller than the CCDY ($W\gamma$) process for $m_N \sim 50 - 100$ GeV. This is much smaller than the 4 (2) orders of magnitude that one expects from naïve power counting. Second is that while the bare rates of resonant channels fall precipitously for increasing m_N , which is due to suppression in both the matrix element and available phase space, the $W^\pm W^\pm$ rate moderately *increases* before slowly decreasing. For an active-sterile mixing of $|V_{\ell N}|^2 = 1$, this leads to the $W^\pm W^\pm$ rate surpassing the $W\gamma$ rate at $m_N \sim 500 - 600$ GeV and the CCDY rate at $m_N \sim 700$ GeV. Due to the different sensitivities of the three channels to active-sterile mixing, the crossover occurs at higher neutrino masses for smaller values of $|V_{\ell N}|^2$. For example: at $|V_{\ell N}|^2 = 0.1$, the $W^\pm W^\pm$ cross section surpasses the $W\gamma$ (CCDY) rate at $m_N = 1.8 - 1.9$ (1.2–1.3) TeV.

While mixing can alter the precise values of these crossovers, the qualitative picture does not change. For instance: independent of $|V_{\ell N}|^2$, the $W^\pm W^\pm \rightarrow \ell^\pm \ell^\pm$ cross section exhibits a qualitatively different dependence on m_N than in the CCDY and $W\gamma$ channels. This leads to the $W^\pm W^\pm$ rate at $\sqrt{s} = 13$ TeV to be the same at both $m_N \sim 40$ GeV and $m_N \sim 2.5$ TeV. Moreover, unlike resonant production of heavy neutrinos via $q\bar{q}$ annihilation or $W\gamma$ fusion, heavy neutrinos in t -channel

processes like $W^\pm W^\pm \rightarrow \ell_i^\pm \ell_j^\pm$ are non-resonant. So while there is a kinematic suppression in the $W^\pm W^\pm$ matrix element at very large m_N , there is no corresponding phase space suppression. This manifests in the cross section as a milder dependence on increasing sterile neutrino masses. Interestingly, as heavy neutrinos are never on-shell in t -channel exchanges they can never manifest as a long-lived particle. Therefore, search complications associated with displaced vertices are not present.

In comparison to past work, this is the first evaluation of the full $2 \rightarrow 4$, same-sign $W^\pm W^\pm$ scattering process in pp collisions at NLO in QCD. At LO, the literature [6, 36, 39, 41, 42] is admittedly in disagreement with itself. Qualitatively, the dependence on collider energy and heavy neutrino masses in all these works are consistent. Quantitatively, large differences exist. In some cases, differences can be traced to the omission of numerical pre-factors in analytic and/or numerical results, theoretical uncertainties associated with the effective W approximation [22–24], and uncertainties in PDF sets. In other cases, the lack of documented inputs and possible phase space cuts hinder precise comparisons. Support for our numerical results include agreement with analytical expressions for helicity amplitudes. For further details, we refer to sections V.1.1 and V.1.2, and appendix C.

To further understand the dependence on m_N in the $W^\pm W^\pm$ channel, we consider for illustration purposes the matrix element for the $2 \rightarrow 2$, $W^\pm W^\pm \rightarrow \ell^\pm \ell^\pm$ subprocess. For the momentum and helicity assignments

$$W_\mu^+(p_1^W, \lambda_1^W) + W_\nu^+(p_2^W, \lambda_2^W) \rightarrow \ell^+(p_1^\ell, \lambda_1^\ell) + \ell^+(p_2^\ell, \lambda_2^\ell), \quad (5.9)$$

and the invariants $M_{WW}^2 = (p_1^W + p_2^W)^2$, $t = (p_1^W - p_1^\ell)^2$, and $u = (p_1^W - p_2^\ell)^2$, the helicity amplitudes are given by

$$-i\mathcal{M} = \varepsilon_\mu(p_1^W, \lambda_1^W) \varepsilon_\nu(p_2^W, \lambda_2^W) \mathcal{T}^{\mu\nu}(p_1^\ell, p_2^\ell, \lambda_1^\ell, \lambda_2^\ell) + (t \leftrightarrow u). \quad (5.10)$$

Here ε are the usual helicity polarization vectors for massive gauge bosons in the unitary gauge, the $(t \leftrightarrow u)$ term accounts for final-state lepton exchange, and following the Feynman rules of Refs. [115, 116], the LN-violating current (\mathcal{T}) in the HELAS convention [117] is³ [36]

$$\mathcal{T}^{\mu\nu} = -i \left(\frac{-ig_W}{\sqrt{2}} \right)^2 \frac{V_{\ell N} V_{\ell N}}{(t - m_N^2)} \times \quad (5.11)$$

$$\begin{aligned} & [\bar{u}(p_1^\ell, \lambda_1^\ell = R) \gamma^\mu P_R (\not{p}_N + m_N) \gamma^\nu P_L v(p_2^\ell, \lambda_2^\ell = R)] \\ & = -i \left(\frac{-ig_W}{\sqrt{2}} \right)^2 \frac{V_{\ell N} V_{\ell N}}{(t - m_N^2)} \times m_N \times \\ & [\bar{u}(p_1^\ell, \lambda_1^\ell = R) \gamma^\mu \gamma^\nu P_L v(p_2^\ell, \lambda_2^\ell = R)]. \end{aligned} \quad (5.12)$$

Ref. [29] with updated inputs as listed in section IV. To regulate the $W\gamma$ fusion matrix element, we use the phase space cuts of equation (4.2) as well as require $p_T^\ell > 10$ GeV and $|\eta^\ell| < 4.0$. Total widths of SM particles are kept at their SM values.

³ In several instances our analytic results differ from those in Ref. [36]. We do not speculate on their specific origin but note that some omissions are obviously typographical.

In the above we assume a clockwise fermion flow of leptons [115, 116], $p_N = (p_1^W - p_1^\ell)$ is the momentum of the internal sterile neutrino, and $u(p, \lambda)$ and $v(p, \lambda)$ are the standard helicity spinors for massless, spin-1/2 fermions.

Crucially, differences in Feynman rules for LN-violating fermion currents relative to the standard rules for LN-conserving ones give rise to an effective parity inversion in the $W_1^+ - \ell_1^+ - N$ vertex and spinor for ℓ_1^+ [118, 119]. This implies [120, 121] that the successive gauge interactions involving massless particles in \mathcal{T} are *helicity inverting* and not helicity preserving as one usually finds in SM gauge interactions involving massless, external particles. Subsequently, projection operators select for the heavy neutrino's RH helicity state, and hence the factor of $m_N \times \mathbb{I}_4$ in \mathcal{T} . This is in contrast to LN-conserving currents, such as in the process $W^+W^- \rightarrow \ell^+\ell^-$, where projection operators select for the LH helicity state, and hence the \not{p}_N term in \mathcal{T} .

We report that exact, analytic evaluation of equations (5.10) and (5.12) yields somewhat bulky expressions without obvious insights. This is despite being a $2 \rightarrow 2$ process and can be tied to the added algebraic complication of the incoming W bosons being massive and carrying a longitudinal polarization. Instead, we focus on the low- (section V.1.1) and high-mass (section V.1.2) limits of the intermediate heavy neutrino. For technical details and intermediate expressions, see appendix C.

V.1.1. Low-Mass Limit

We consider first the limit where masses of both W and N are small compared to the $W^\pm W^\pm$ scattering scale, *i.e.*, when $m_W, m_N \ll M_{WW}$. In this limit, the LN-violating tensor current in equation (5.12) scales as

$$\mathcal{T}^{\mu\nu} \propto g_W^2 V_{\ell N} V_{\ell N} \frac{m_N M_{WW}}{(t - m_N^2)} \quad (5.13)$$

$$\sim g_W^2 V_{\ell N} V_{\ell N} \frac{m_N}{M_{WW}} + \mathcal{O}\left(\frac{m_N^2}{M_{WW}^2}, \frac{m_W^2}{M_{WW}^2}\right), \quad (5.14)$$

where the M_{WW} factor in the numerator originates from the two lepton spinors, $u(p^\ell), v(p^\ell) \sim \sqrt{M_{WW}}$. In this same limit, the scattering of longitudinally polarized W bosons is enhanced over the scattering of the transverse polarizations. This enhancement can be seen in the polarization vectors themselves, which scale as

$$\varepsilon_\mu(p^W, \lambda^W = \pm) \sim \mathcal{O}(1), \quad (5.15)$$

$$\varepsilon_\mu(p^W, \lambda^W = 0) \sim \frac{p_\mu^W}{m_W} + \mathcal{O}\left(\frac{m_W}{M_{WW}}\right) \quad (5.16)$$

$$\sim \mathcal{O}\left(\frac{M_{WW}}{m_W}\right). \quad (5.17)$$

This shows that in the high-energy limit the $W^\pm W^\pm \rightarrow \ell^\pm \ell^\pm$ process is driven by $W_0^\pm W_0^\pm$ scattering and that

the corresponding matrix element scales as

$$-i\mathcal{M} = \varepsilon_\mu(\lambda_1^W = 0)\varepsilon_\nu(\lambda_2^W = 0)\mathcal{T}^{\mu\nu} + (t \leftrightarrow u) \quad (5.18)$$

$$\sim g_W^2 V_{\ell N} V_{\ell N} \frac{m_N}{m_W^2} M_{WW}. \quad (5.19)$$

Remarkably, after squaring \mathcal{M} , the quadratic dependence on M_{WW} is canceled by the flux factor in the definition of the parton-level cross section $\hat{\sigma}$. This renders the rate independent of M_{WW} but quadratic in m_N ,

$$\hat{\sigma}(W^+W^+ \rightarrow \ell^+\ell^+) \sim g_W^4 |V_{\ell N}|^4 \frac{m_N^2}{m_W^4}. \quad (5.20)$$

Thus, we can attribute the growth in the same-sign WW scattering rate seen in figure 2 for sub-TeV heavy neutrinos to the cancellation of momentum scales in high-energy $W_0^\pm W_0^\pm$ scattering in tandem with helicity inversion in the LN-violating lepton current.

After a more careful computation (see appendix C), the $W^+W^+ \rightarrow \ell_1^+ \ell_2^+$ cross section for n_R heavy neutrinos is

$$\hat{\sigma}(W^+W^+ \rightarrow \ell_1^+ \ell_2^+) = \frac{g_W^4 (2 - \delta_{\ell_1 \ell_2})}{2^5 3^2 \pi m_W^4} \left| \sum_{k=4}^{n_R+3} V_{\ell_1 k} m_{N_k} V_{\ell_2 k} \right|^2 + \mathcal{O}\left(\frac{m_N^2}{M_{WW}^2}, \frac{M_{WW}^2}{m_W^2}\right). \quad (5.21)$$

V.1.2. High-Mass Limit

We consider now the kinematic limit where the W boson's mass and all momentum-transfer scales are small compared to the sterile neutrino's mass, *i.e.*, the decoupling limit [122] where $m_W^2, M_{WW}^2, |t|, |u| \ll m_N^2$. In this limit, N -exchanges can be treated as contact interactions, and the pole structure of its propagator can be systematically expanded. Doing this causes the LN-violating tensor current of equation (5.12) to scale as

$$\mathcal{T}^{\mu\nu} \propto g_W^2 V_{\ell N} V_{\ell N} \frac{m_N M_{WW}}{(t - m_N^2)} \quad (5.22)$$

$$\sim -g_W^2 V_{\ell N} V_{\ell N} \frac{M_{WW}}{m_N} + \mathcal{O}\left(\frac{|t|}{m_N^2}\right). \quad (5.23)$$

An analogous expression holds for the u -channel.

A consequence of this expansion is that the angular dependence that encapsulates forward- and backward-scattering enhancements in gauge interactions becomes a sub-leading contribution in the propagator. This implies that for most $W^\pm W^\pm$ polarization combinations the forward (t -channel) and backward (u -channel) helicity amplitudes are indistinguishable, save for a relative minus sign that triggers an exact destructive interference. As a result, the only non-vanishing amplitudes are those where the incoming $W^\pm W^\pm$ states carry the same helicity.

Noting once more the enhancement of longitudinal-longitudinal scattering over other $W^\pm W^\pm$ helicity configurations, one finds that the leading contribution to the

matrix element for the $W_0^\pm W_0^\pm \rightarrow \ell^\pm \ell^\pm$ process scales as

$$-i\mathcal{M} = \varepsilon_\mu(\lambda_1^W = 0)\varepsilon_\nu(\lambda_2^W = 0)\mathcal{T}^{\mu\nu} + (t \leftrightarrow u) \quad (5.24)$$

$$\sim g_W^2 \frac{V_{\ell N} V_{\ell N}}{m_N} \frac{M_{WW}^3}{m_W^2}. \quad (5.25)$$

After squaring, the dependence on M_{WW} is partially compensated by the flux factor in the parton-level cross section. The result is a total rate that scales as

$$\hat{\sigma}(W^+W^+ \rightarrow \ell^+\ell^+) \sim g_W^4 \frac{|V_{\ell N}|^4}{m_N^2} \frac{M_{WW}^4}{m_W^4}. \quad (5.26)$$

Immediately, we see that the m_N^{-2} factor originating from the heavy neutrino propagator is never fully offset by the LN-violating current or other factors at the cross section level. Ultimately, this is responsible for the drop in cross section that occurs in figure 2 for increasing m_N .

After a more careful computation (see appendix C), one finds that the parton-level, $W^+W^+ \rightarrow \ell_1^+\ell_2^+$ cross section for the exchange of n_R heavy neutrinos is:

$$\begin{aligned} \hat{\sigma}(W^+W^+ \rightarrow \ell_1^+\ell_2^+) &= \frac{g_W^4(2 - \delta_{\ell_1\ell_2})}{2^7 3^2 \pi} \frac{M_{WW}^4}{m_W^4} \quad (5.27) \\ &\times \left| \sum_{k=4}^{n_R+3} \frac{V_{\ell_1 N_k} V_{\ell_2 N_k}}{m_{N_k}} \right|^2 + \mathcal{O}\left(\frac{M_{WW}^2}{m_N^2}, \frac{M_{WW}^2}{m_W^2}\right). \end{aligned}$$

We stress that the transition rate's dependence on masses and mixing elements of heavy neutrinos mirrors the scaling behavior found in nuclear $0\nu\beta\beta$ decay rates [6, 36]. Notably, the coherent summation over V permits complex phases to trigger potentially large cancellations in analogy to the ‘‘funnel behavior’’ in $0\nu\beta\beta$ decay.

V.1.3. QCD Corrections to $W^\pm W^\pm \rightarrow \ell^\pm \ell^\pm$

Returning to figure 2, we recall that past investigations into the $W^\pm W^\pm \rightarrow \ell_i^\pm \ell_j^\pm$ process historically [36, 41, 43] employed the effective W approximation [22–24]. In this approximation, W bosons are treated as constituents of the proton and the $2 \rightarrow 2$, $W^\pm W^\pm \rightarrow \ell_i^\pm \ell_j^\pm$ scattering rate is convolved with W boson PDFs of the proton. Only later in Ref. [6] was the full $2 \rightarrow 4$ process with forward jets evaluated. In all these cases, however, only LO estimates of cross sections were calculated. Therefore, we are motivated to comment on the size of NLO in QCD corrections and residual uncertainties in the full $2 \rightarrow 4$ process, particularly in relation to those of the CCDY and $W\gamma$ modes, which were first reported in Refs. [29, 123].

In the lower panel of figure 2 we show as a function of heavy neutrino mass the QCD K -factors, as defined in equation (5.2), for the same-sign $W^\pm W^\pm$, CCDY, and $W\gamma$ processes. Band thicknesses correspond to the residual μ_r, μ_f dependence at NLO. Over the mass range $m_N \sim 40 \text{ GeV} - 20 \text{ TeV}$, we find that QCD corrections

gradually and uniformly increase the total $W^\pm W^\pm$ rate by +5% to +35%. As in deeply inelastic scattering, one-loop QCD corrections to spacelike EW emissions in VBF do not appreciably alter cross section normalizations [124]. Consequentially, we attribute the increase in cross section at NLO to real, initial-state radiation. This purported reliance on an $\mathcal{O}(\alpha_s(\mu_r))$ splitting is supported by the increased scale dependence at larger m_N and the lack of a scale dependence in the $2 \rightarrow 4$ process. In comparison to the other channels, the K -factor for $W^\pm W^\pm$ sits just below (above) the CCDY ($W\gamma$) curve for $m_N \sim 750 \text{ GeV}$ and overtakes both at $m_N \gtrsim 1 \text{ TeV}$.

For corrections beyond NLO in QCD, one can consider two complementary directions. The first pertains to higher-order QCD corrections while the second pertains to EW corrections. Based on results for VBF production of the SM Higgs boson [125–127], we anticipate that improvements at $\mathcal{O}(\alpha_s^2)$ and $\mathcal{O}(\alpha_s^3)$ have only a modest impact on total cross sections and distributions. At NLO in EW, corrections to the LO rate typically scale as

$$\left| \frac{\delta\sigma^{\text{NLO-EW}}}{\sigma^{\text{LO}}} \right| \sim \frac{g_W^2}{4\pi} \log \frac{M_{WW}^2}{m_W^2}. \quad (5.28)$$

As we show in section V.2, scales for M_{WW} at the LHC range about $M_{WW} \sim 300 \text{ GeV} - 600 \text{ GeV}$ for a large array of heavy neutrino masses. This translates to a modest uncertainty of $\delta\sigma^{\text{NLO-EW}}/\sigma^{\text{LO}} \sim 9\%$ to 13%. We anticipate that in both cases the impact of missing higher-order corrections is negligible for discovery purposes.

V.1.4. Partial-wave unitarity in $W_0^\pm W_0^\pm \rightarrow \ell^\pm \ell^\pm$

As a brief remark, we interestingly note that the scattering amplitude for the $W^\pm W^\pm \rightarrow \ell^\pm \ell^\pm$ process exhibits poor high-energy behavior when initiated by a pair of longitudinally polarized W bosons. As evident in equations (5.19) and (5.25), and more precisely in appendix C, matrix elements for both the low- and high-mass limits scale with some positive power of the (WW)-scattering energy, M_{WW} . This is distinct from the CCDY channel where no such scaling behavior is present. For fixed heavy neutrino masses and mixing, such a dependence on M_{WW} implies [128–133] that the matrix elements violate partial-wave unitarity above some scattering energy threshold E_U , unless additional physics cancels this dependence. As both amplitudes depend on the mass of an internal Majorana neutrino, it is possible that the unitarity violation is actually tied to the explicit breaking of LN symmetry in the Type I Seesaw model. If so, then it can potentially be resolved through the spontaneous breaking of LN symmetry via a Higgs-like mechanism.

While a systematic study of partial-wave unitarity in the Phenomenological Type I Seesaw model lies beyond the scope of this work, we can nevertheless provide a qualitative outlook. Following Ref. [122], the $J = 0$ partial-wave amplitude of the $W_0^\pm W_0^\pm \rightarrow \ell^\pm \ell^\pm$ process

is related to its matrix element \mathcal{M} by the relationship

$$a_{J=0} = \frac{1}{32\pi} \int_{-1}^1 d\cos\theta_1 \mathcal{M}(W_0^\pm W_0^\pm \rightarrow \ell_1^\pm \ell_2^\pm). \quad (5.29)$$

Here θ_1 is the polar angle of ℓ_1 in the frame of the (WW) -system. For the low- and high-mass limits, the partial-wave amplitudes are given to lowest order by

$$\begin{aligned} \text{low-mass : } \quad a_0 &\approx (2 - \delta_{\ell_1 \ell_2}) \frac{g_W^2 M_{WW}}{16\pi m_W^2} \\ &\times \left| \sum_{k=4}^{n_R+3} V_{\ell_1 N_k} m_{N_k} V_{\ell_2 N_k} \right|, \quad (5.30) \end{aligned}$$

$$\begin{aligned} \text{high-mass : } \quad a_0 &\approx (2 - \delta_{\ell_1 \ell_2}) \frac{g_W^2 M_{WW}^3}{32\pi m_W^2} \\ &\times \left| \sum_{k=4}^{n_R+3} \frac{V_{\ell_1 N_k} V_{\ell_2 N_k}}{m_{N_k}} \right|. \quad (5.31) \end{aligned}$$

Requiring $|a_J| < 1$ implies that the $W_0^\pm W_0^\pm$ channel saturates unitarity at $M_{WW} = E_U$, with

$$\text{low-mass : } E_U = \frac{16\pi m_W^2 / [(2 - \delta_{\ell_1 \ell_2}) g_W^2]}{\left| \sum_{k=4}^{n_R+3} V_{\ell_1 N_k} m_{N_k} V_{\ell_2 N_k} \right|}, \quad (5.32)$$

$$\text{high-mass : } E_U^3 = \frac{32\pi m_W^2 / [(2 - \delta_{\ell_1 \ell_2}) g_W^2]}{\left| \sum_{k=4}^{n_R+3} \frac{V_{\ell_1 N_k} V_{\ell_2 N_k}}{m_{N_k}} \right|}. \quad (5.33)$$

In the low-mass limit and assuming a single heavy neutrino of mass $m_N = 1$ TeV with an active-sterile mixing of $|V_{\ell N}|^2 = 10^{-1}$ (10^{-2}) [10^{-3}], partial-wave unitarity saturates at about $E_U \sim 7.6$ TeV (76 TeV) [760 TeV]. Working instead in the high-mass limit, the mass and mixing upper bound from equation (2.7) as derived from $0\nu\beta\beta$ decay searches implies a lower bound on the saturation scale of $E_U \sim 72$ TeV – 87 TeV. This suggests that if a discovery of $0\nu\beta\beta$ decay is made by current-generation experiments, and if the decay is mediated by a heavy Majorana neutrino, then a future pp collider with $\sqrt{s} = 100$ TeV may be able to probe this high-energy behavior. However, we caution that in both cases the saturation scale is acutely sensitive to the values of active-sterile mixing elements and heavy neutrino masses. In order to minimize theoretical bias in estimating the LHC's sensitivity to the $W^\pm W^\pm \rightarrow \ell^\pm \ell^\pm$ process, we do not impose the above constraints on the Phenomenological Type I Seesaw's parameter space, and in fact defer further discussion of partial-wave unitarity to future work.

V.2. Kinematic Properties of $W^\pm W^\pm \rightarrow \ell_i^\pm \ell_j^\pm$

We now turn to exploring the kinematic properties of the $W^\pm W^\pm \rightarrow \ell_i^\pm \ell_j^\pm$ signal process at $\sqrt{s} = 13$ TeV. As

NLO-vs-LO comparisons of VBF kinematics are extensively documented, we restrict ourselves to NLO+PS(LL) distributions where available and neglect comparisons to properties at LO+PS(LL). For concreteness, we fix $\ell_i = \ell_j = \mu$ and set simulation inputs as prescribed in section III. For each of the following observable we assume the representative benchmark masses $m_N = 750$ GeV (darkest), 1.5 TeV (dark), and 5 TeV (light). Events are normalized to $\mathcal{L} = 300$ fb $^{-1}$. Also shown for each distribution is the residual μ_r, μ_f, μ_s uncertainty (band thickness) as obtained from a 27-point variation envelope.

Throughout this section we work with particle-level objects. We do so to emulate detector thresholds (but not detector resolution) according to our analysis in section VI.2 and to ensure the infrared safety of observable definitions. In practice, this means that after parton showering we impose anti- k_T ($R = 0.4$) clustering on all hadronic activity. We also require that electron, muon, hadronic tau, and jet candidates satisfy the following requirements

$$p_T^{e(\mu)} \{ \tau_h \} \{ j \} > 10 \text{ (27) [20] \{25\} GeV}, \quad (5.34)$$

$$|\eta^{e(\mu)} \{ \tau_h \} \{ j \}| < 2.5 \text{ (2.7) [2.5] \{4.5\}}. \quad (5.35)$$

Particle identification efficiencies and mistagging rates are kept at their default values in MADANALYSIS5 [102–104]. That is to say, we consider an ideal setup in which identification efficiencies are set to unity and mistagging rates are set to zero. No kinematic smearing is applied.

Given these stipulations, we define our signal as the

$$pp \rightarrow \mu^\pm \mu^\pm jj + X \quad (5.36)$$

process, where X denotes the possibility of additional hadronic or photonic activity. More precisely, we require events to possess exactly two same-sign μ candidates and at least two j candidates. Events containing any number of e or τ_h candidates are rejected. This setup implies that we remain inclusive with respect to soft and forward objects that fail candidacy requirements. For clarity, objects are ranked by their p_T , with $p_T^k > p_T^{k+1}$.

We start with figure 3 where we plot the (a,b) p_T and (c,d) η distributions of the (a,c) leading (μ_1) and (b,d) sub-leading (μ_2) muon in our signal process at NLO+PS. We foremost note the lack of any resonant structure in both p_T^μ distributions. This follows from the absence of s -channel resonances in the $W^\pm W^\pm \rightarrow \mu^\pm \mu^\pm$ sub-process. Instead, we find a p_T behavior reminiscent of open particle production that plateaus for a few hundred GeV and then falls due to kinematic suppression. There is a steeper falloff for smaller m_N . The p_T spectra indicate that the $(\mu^\pm \mu^\pm)$ -system, or equivalently the $(W^\pm W^\pm)$ -system, possesses a large invariant mass that reaches several hundred GeV. We observe that both muons tend toward smaller values of $|\eta|$, independent of the heavy neutrino mass, indicating an absence of forward scattering.

Moving onto figure 4, we present the same information as in figure 3 but for the (a,c) leading (j_1) and (b,d) sub-leading (j_2) jets. In both p_T spectra we observe peaks at




FIG. 3. Kinematic distributions at $\sqrt{s} = 13$ TeV of the same-sign $W^\pm W^\pm \rightarrow \mu^\pm \mu^\pm$ signal process at NLO+PS with residual μ_r, μ_f, μ_s uncertainty envelope (band thickness), for $m_N = 750$ GeV (darkest), 1.5 TeV (dark), and 5 TeV (light), of the (a,c) leading and (b,d) sub-leading μ^\pm (a,b) transverse momentum (p_T) and (c,d) pseudorapidity (η).

$p_T^j \sim m_W/2$, which is characteristic of the VBF process and is due to the recoil against the t -channel emission of W bosons. Consistently, we observe in the η distributions that the two jets are forward, with maxima in the forward direction near $|\eta^j| \sim 3$ and a suppression of central activity at $|\eta^j| \sim 0$. We find that the shapes of all observables in figure 4 are insensitive to the values of m_N under consideration. This follows from the fact that quarks in the $2 \rightarrow 4$ process do not directly couple to the LN-violating current, and therefore act like spectators.

As the $pp \rightarrow \mu^\pm \mu^\pm jj + X$ process is simulated at NLO+PS, one has access to the $pp \rightarrow \mu^\pm \mu^\pm jjj + X$ process at LO+PS. We are thus able to explore the QCD radiation pattern in the VBF process. In figure 5 we show the (a) p_T and (b) η distribution of the trailing jet j_3 at LO+PS. For all considered values of m_N , we observe an inclination toward lower p_T , with most of the phase space

sitting between the threshold at $p_T^{j_3} \sim p_T = 25$ GeV and $p_T^{j_3} \sim 40$ GeV. This is just below the characteristic p_T of the two leading jets. We note a strong suppression of central ($|\eta^{j_3}| \lesssim 2$) tertiary jets. This does not mean an absence of QCD radiation for $|\eta^{j_3}| \lesssim 2$, only that it is soft. Most of the activity resides in the forward direction, peaking at $|\eta| \sim 3-4$, again independent of m_N . As this is well in the vicinity of the leading jets it is likely that the $q \rightarrow qg$ and $g \rightarrow q\bar{q}$ splittings responsible for j_3 involve smaller momentum transfers, which results in shallower opening angles between j_3 and its companion.

To further investigate the dynamics of the $W^\pm W^\pm \rightarrow \mu^\pm \mu^\pm$ sub-process, we consider in figure 6 observables that are built from the momenta of two or more particles. We start with figure 6(a), where we plot the azimuthal




FIG. 4. Same as figure 3 but for the (a,c) leading and (b,d) sub-leading jet.

separation of the same-sign $\mu^\pm\mu^\pm$ pair, defined as

$$\Delta\varphi(\mu_1, \mu_2) = \vec{p}_T^{\mu_1} \cdot \vec{p}_T^{\mu_2} / |\vec{p}_T^{\mu_1}| |\vec{p}_T^{\mu_2}|. \quad (5.37)$$

We find that the leptons exhibit a strong back-to-back trajectory with the separation peaking (curtailing) at $\Delta\varphi(\mu_1, \mu_2) \approx \pi$ (0). This is despite being a 4-body final state at LO, which would suggest a sizable recoil against the $(jj + X)$ -system. For increasing heavy neutrino masses we observe a higher tendency for back-to-back trajectories. The marginal-to-moderate recoil that is found suggests that modeling the $2 \rightarrow 4$ signal process as a $2 \rightarrow 2$ process within the effective W approximation as done in Refs. [36, 41, 43] is a fair approximation.

In figure 6(b) we focus on the distribution of the missing transverse energy E_T^{miss} , defined per event as the magnitude of the two-momentum recoil against all vis-

ible (vis) objects, regardless of their energy,

$$E_T^{\text{miss}} = |\vec{p}_T^{\text{miss}}|, \quad \vec{p}_T^{\text{miss}} = - \sum_{k \in \{\text{vis}\}} \vec{p}_T^k. \quad (5.38)$$

We find that the distribution strongly peaks at $E_T^{\text{miss}} \lesssim 10$ GeV, in line with expectations of a $2 \rightarrow 4$ process without outgoing light neutrinos. As we are working without any detector resolution effects, the nonzero E_T^{miss} originates from the weak decays to light neutrinos of mesons generated in the parton shower. Aside from differences in the rate normalization, we observe no substantial dependence of E_T^{miss} on the heavy neutrino mass.

In figure 6(c) we show the invariant mass distribution of the two highest p_T (leading) jets, given by

$$M(j_1, j_2) = \sqrt{(p^{j_1} + p^{j_2})^2}. \quad (5.39)$$

For the heavy neutrino masses under consideration, we see that the peaks of the invariant mass spectra occur at




FIG. 5. Same as figure 3 but for the (a) p_T and (b) η of the trailing jet (j_3) at LO+PS.

$M(j_1, j_2) \sim 1000 \text{ GeV} - 1200 \text{ GeV}$, with a peak position at larger $M(j_1, j_2)$ for larger m_N . A narrow collection of events at $M(j_1, j_2) \ll 500 \text{ GeV}$ is also observed. These low-mass events are attributed to instances of one forward jet possessing relatively low p_T while another jet undergoes a hard $q^* \rightarrow qq$ splitting. In such cases the (qq) pair can be identified as the leading jet pair but still return a small $M(j_1, j_2)$ since this corresponds to the (q^*) -system's virtuality, which is favored to be small in massless QCD. The dependence on m_N indicates that the hadronic activity is not completely decoupled from the $W^\pm W^\pm \rightarrow \mu^\pm \mu^\pm$ sub-process, and therefore can potentially offer a handle on determining the value of m_N . This is relevant given the mild scale uncertainty bands.

The $M(j_1, j_2)$ spectra point to the signal process being driven by valence quark-valence quark scattering involving large momentum fractions, *i.e.*, $x_B > M(j_1, j_2)/\sqrt{s} \sim 0.1$. In comparison to the p_T^μ distributions of figure 3, which show charged lepton momenta reaching a few hundred GeV, we see that comparable momentum fractions are propagated into the $W^\pm W^\pm \rightarrow \mu^\pm \mu^\pm$ sub-process. For example: estimating the incoming W boson energies by those of the muons, $E_W \sim E_\mu \sim p_T^\mu \sim 100 \text{ GeV} - 300 \text{ GeV}$, and the outgoing quark energies from the invariant mass of the two leading jets, which are also back-to-back, $E_q^{out} \sim M(j_1, j_2)/2 \gtrsim 500 \text{ GeV} - 1000 \text{ GeV}$, then the typical momentum fractions carried by the W reach

$$x_W \equiv \frac{E_W}{E_q^{in}} = \frac{E_W}{(E_W + E_q^{out})} \lesssim 0.1 - 0.4. \quad (5.40)$$

Moving onto figure 6(d), we show the pseudorapidity difference between the two leading jets, defined as

$$\Delta\eta(j_1, j_2) = \eta^{j_1} - \eta^{j_2}. \quad (5.41)$$

We report several notable features. First is the symmetric behavior around $\Delta\eta = 0$, which stems from having a

symmetric beam configuration. Second is that most of the phase space populates the region where $|\Delta\eta| \gtrsim 2$ and appears independent of heavy neutrino masses. Third is the presence of a modest collection of events with $|\Delta\eta| \lesssim 1$. Such events are consistent with the low-mass distribution in figure 6(c) originating from $q^* \rightarrow qq$ and $g^* \rightarrow q\bar{q}$ splittings with relatively small opening angles.

Beyond one- and two-particle observables are those sensitive to the global activity of the $pp \rightarrow \mu^\pm \mu^\pm jj + X$ process. In particular, we consider in figure 7(a) the scalar sum of p_T over all jets in an event (H_T),

$$H_T = \sum_{k \in \{\text{jets}\}} |\vec{p}_T^k|, \quad (5.42)$$

and in figure 7(b), the scalar sum of the p_T of all reconstructed particle candidates (reco) (X_T),

$$X_T = \sum_{k \in \{\text{reco}\}} |\vec{p}_T^k|. \quad (5.43)$$

In the first case and for all heavy neutrino masses we observe that H_T peaks at $H_T \sim 100 \text{ GeV}$ and uniformly decreases for larger values of H_T . As the net contribution of the two (three) leading jets in the signal scales as

$$p_T^{j_1} + p_T^{j_2} (+p_T^{j_3}) \sim 2 \times \frac{M_W}{2} (+p_T^{j \min}) = 80 \text{ (105) GeV}, \quad (5.44)$$

the H_T distribution suggests the presence of little high- p_T hadronic activity beyond these leading objects. For the X_T case, we observe a slight dependence on N 's mass, with the distributions' maxima occurring at $X_T \sim 600 \text{ GeV} - 750 \text{ GeV}$ and tending towards larger values for larger m_N . We attribute this sensitivity to the dependence of muon p_T , as seen in figure 3, which also peaks at larger values for increasing m_N . In comparing




FIG. 6. Same as figure 3 but for the (a) azimuthal separation of the leading same-sign $\mu^\pm\mu^\pm$ pair $\Delta\varphi(\mu_1, \mu_2)$, (b) missing transverse energy E_T^{miss} , (c) invariant mass of the leading dijet system $M(j_1, j_2)$, and (d) pseudorapidity difference of the same system $\Delta\eta(j_1, j_2)$.

X_T to the $(\mu^\pm\mu^\pm jj(j))$ -system itself, we find that the scalar sum of H_T and the same-sign muon pair p_T ,

$$H_T + p_T^{\mu_1} + p_T^{\mu_2} \sim 100 \text{ GeV} + (200 \text{ GeV} - 400 \text{ GeV}) + (100 \text{ GeV} - 200 \text{ GeV}) \quad (5.45)$$

$$= 400 \text{ GeV} - 700 \text{ GeV}, \quad (5.46)$$

undershoots the peak of X_T by $\Delta X_T \sim 50 \text{ GeV} - 200 \text{ GeV}$. This indicates a sizable presence of electromagnetic activity (photons), which one can anticipate from the presence of jets and muons with TeV-scale momenta.

Finally, we consider observables that measure the relative amounts of hadronic and leptonic activity in a given event. Such quantities are sensitive to the color structure of hard scattering processes [30, 92, 134, 135], and hence employable in dynamic jet vetoes for color-singlet signal processes. While a full exploration of jet vetoes

in $W^\pm W^\pm \rightarrow \mu^\pm\mu^\pm$ is outside our scope, we consider as representative cases in figures 7(c) and 7(d) respectively, the ratio of H_T and the leading charged lepton p_T :

$$r_{\mu_1}^{H_T} = H_T/p_T^{\mu_1}, \quad (5.47)$$

and the ratio of X_T and the leading charged lepton p_T :

$$r_{\mu_1}^{X_T} = X_T/p_T^{\mu_1}. \quad (5.48)$$

In the first case, we observe that a majority of the phase space sits well below $r_{\mu_1}^{H_T} = 1$. This indicates that on an event-by-event basis more transverse momentum is carried by the leading muon than in all jets combined. This is unlike diboson and top quark processes where the situation is reversed [30, 92, 135]. We find that the ratios peak just above $r_{\mu_1}^{H_T} \sim 0.25$ and are largely independent of the heavy neutrino masses that we consider. This is




FIG. 7. Same as figure 3 but for the (a) scalar sum of all jet p_T (H_T), (b) scalar sum of all visible p_T (X_T), (c) ratio $H_T/p_T^{\mu_1}$, (d) ratio $X_T/p_T^{\mu_1}$.

also consistent with naïve estimations from the individual H_T and $p_T^{\mu_1}$ distributions, which suggest

$$\frac{H_T}{p_T^{\mu_1}} \sim \frac{100 \text{ GeV}}{(200 \text{ GeV} - 400 \text{ GeV})} \sim 0.25 - 0.5. \quad (5.49)$$

In the $r_{\mu_1}^{X_T}$ case, we observe a sharp cut-off at $r_{\mu_1}^{X_T} \sim 2$ that is largely independent of the heavy neutrino's mass. This can be tied to the disparity of momentum scales between leptons and jets. In particular, since $p_T^{\mu_1}, p_T^{\mu_2} \gg p_T^j$, one finds that the ratio scales roughly as

$$r_{\mu_1}^{X_T} \sim \frac{H_T + p_T^{\mu_1} + p_T^{\mu_2}}{p_T^{\mu_1}} \sim \frac{p_T^{\mu_1} + p_T^{\mu_2}}{p_T^{\mu_1}} \sim 2. \quad (5.50)$$

The approximation $p_T^{\mu_1} \sim p_T^{\mu_2}$ is again consistent with $2 \rightarrow 2$ scattering and the back-to-back trajectories found in the azimuthal separation distribution in figure 6(a).

VI. SENSITIVITY AT THE LHC AND HL-LHC

In this section we estimate the discovery potential of heavy Majorana neutrinos in same-sign WW scattering at the LHC and its high-luminosity upgrade. After summarizing our simulated detector setup in section VI.1, we build our event selection menu in section VI.2, and present our findings in section VI.3. Our analysis includes signal and background processes that are normalized to an integrated luminosity of $\mathcal{L} = 300 \text{ fb}^{-1}$ for the $\sqrt{s} = 13 \text{ TeV}$ LHC, and to $\mathcal{L} = 3 \text{ ab}^{-1}$ for the HL-LHC.

VI.1. Detector Modeling and Particle Identification

Particle objects considered throughout the analysis are defined using the ATLAS configuration card available

TABLE III. Pre-selection and signal region cuts.

Pre-selection Cuts	
$p_T^{\mu_1 (\mu_2)} > 27$ (10) GeV,	$ \eta^\mu < 2.7, \quad n_\mu = 2,$
$p_T^j > 25$ GeV,	$ \eta^j < 4.5, \quad n_j \geq 2,$
$Q_{\mu_1} \times Q_{\mu_2} = 1,$	$M(j_1, j_2) > 700$ GeV
Signal Region Cuts	
$p_T^{\mu_1}, p_T^{\mu_2} > 300$ GeV	

from the DELPHES repository. In the results that follow the ATLAS card was modified to construct jet candidates from hadronic activity using the anti- k_T sequential clustering algorithm with a distance parameter $R = 0.4$. This value is more widely used in recent ATLAS data analyses than the default value of $R = 0.6$. All the other parameters in the configuration card, which include particle identification and mis-tagging efficiencies as well as the fiducial geometry definition, are left unchanged.

As summarized in section III.1, the impact of pileup is assumed to be subtracted from events as one would do with real data. While we neglect the presence of additional low- p_T , pileup jets in our samples, the impact on particle resolution is at least partly encapsulated in the momentum smearing routines in DELPHES.

VI.2. Event Selection

Our analysis is designed to be as simple as pragmatically possible. We do this to establish a baseline sensitivity and discovery potential at the LHC that broadly covers Majorana masses spanning $m_N = 50$ GeV – 20 TeV. Investigating improvements that target specific mass regimes is left to future work. As discussed in section VII.1, we do not exploit all the kinematic characteristics reported in section V.2. We omit, for example, cuts on E_T^{miss} or hadronic activity that are well-established handles in searches for LNV at colliders [6, 25, 30]. Thus, a more tailored analysis by ATLAS or CMS should yield improvements over the outlook presented here.

To identify our LN-violating collider signature,

$$pp \rightarrow \mu^\pm \mu^\pm jj + X, \quad (6.1)$$

which is characterized by two same-sign muons and at least two jets, we first apply a loose event selection (called pre-selection in the following). This reduces the number of background processes that must be considered while keeping a high selection efficiency for the signal process. The pre-selection also includes requirements needed to ensure the near 100% efficiency of the inclusive, single-muon trigger chains used to record collision data in ATLAS [136] and CMS [137] during Run 2 and the future Run 3 of the LHC. A summary of pre-selection requirements, adapted to Run 2 trigger and acceptance thresholds of ATLAS, is listed in the top of Table III.

TABLE IV. Visible signal cross sections (and efficiencies) after applying different selections to the simulated events.

m_N	$\sigma^{\text{Gen.}}$ [fb]	$\sigma^{\text{Pre.}}$ [fb] (\mathcal{A})	σ^{SR} [fb] (ϵ)
150 GeV	13.3	3.7 (28%)	0.5 (14%)
1.5 TeV	8.45	3.18 (38%)	1.9 (63%)
5 TeV	1.52	0.58 (38%)	0.46 (79%)
15 TeV	0.190	0.072 (38%)	0.056 (78%)

TABLE V. Expected number of SM background events in the Signal Region at the (HL-)LHC with $\mathcal{L} = 300 \text{ fb}^{-1}$ (3 ab^{-1}).

Collider	QCD $W^\pm W^\pm jj$	EW $W^\pm W^\pm jj$	$W^\pm V(3\nu)$	Total
LHC	0.05	0.52	0.14	0.71
HL-LHC	0.49	5.17	1.40	7.10

At pre-selection we require that events have exactly two isolated muon candidates. Muon candidates must have the same electric charge Q and reside within the fiducial volume of $|\eta| < 2.7$. The leading muon must have $p_T > 27$ GeV in order to ensure high efficiency of the muon trigger, whereas the sub-leading muon p_T just needs to be above the reconstruction threshold of $p_T > 10$ GeV. Events with additional lepton candidates, including hadronically decaying τ leptons, are vetoed.

At least two jets with $p_T > 25$ GeV and $|\eta| < 4.5$ must be present in each event. The invariant mass of the system constituted by the two highest p_T jets passing these criteria, $M(j_1, j_2)$, must also exceed 700 GeV. This suppresses interfering sub-processes (see section IV.1) and enriches the signal sample with a topology that corresponds to the scattering of weak vector bosons.

For the heavy neutrino mass range under consideration, we report that about $\mathcal{A} \sim 20\%$ to 40% of signal events survive pre-selection cuts. For representative m_N (first column), we list in Table IV the generator-level cross section $\sigma^{\text{Gen.}}$ (second column) assuming a nominal active-sterile mixing as set in equation (3.3), the pre-selection-level cross section $\sigma^{\text{Pre.}}$ (third column), and the pre-selection acceptance rate $\mathcal{A} = \sigma^{\text{Pre.}}/\sigma^{\text{Gen.}}$.

After pre-selection cuts, we anticipate that the leading background processes consist of mixed EW-QCD production of $W^\pm W^\pm jj$, pure EW production of $W^\pm W^\pm jj$, and the inclusive diboson(+jets) spectrum $W^\pm V + nj$ with $V \in \{\gamma^*/Z^{(*)}\}$. For compactness we label these:

$$W^\pm W^\pm jj \text{ (QCD)} : pp \rightarrow W^\pm W^\pm jj \rightarrow 2\mu^\pm jj + X, \quad (6.2)$$

$$W^\pm W^\pm jj \text{ (EW)} : pp \rightarrow W^\pm W^\pm jj \rightarrow 2\mu^\pm jj + X, \quad (6.3)$$

$$W^\pm V(3\nu) : pp \rightarrow 3\nu + X. \quad (6.4)$$

Our MC modeling of these backgrounds is described in section IV.2 and their generator-level rates are summarized in Table I. We have checked that other processes satisfying the same-sign muon signature do not appre-




FIG. 8. The distribution for (a) $M(j_1, j_2)$ and (b) $p_T^{\mu 2}$ after pre-selection cuts for the signal process assuming representative heavy neutrino masses $m_N = 750$ GeV (darkest cross), 1.5 TeV (dark cross), and 5 TeV (light cross). Also shown are the inclusive $W^\pm V$ (light), EW $W^\pm W^\pm jj$ (dark), and QCD $W^\pm W^\pm jj$ (darkest) backgrounds, as well as their sum (black cross).

ciably survive pre-selection. For example: using the K -factors of Ref. [138], we estimate that the rate for the $t\bar{t}W^\pm \rightarrow 2\mu^\pm + X$ process after pre-selection cuts, but minus the $M(j_1, j_2)$ requirement, is $\sigma_{t\bar{t}W^\pm} \sim 1.2$ ab. The $M(j_1, j_2)$ criterion reduces this an order of magnitude.

We acknowledge that we do not fully account for backgrounds that are difficult to simulate accurately with MC simulations alone. This includes opposite-sign muon pairs in which one muon is reconstructed with the incorrect charge, or events where one of the same-sign muons has a “fake” origin, *e.g.*, a muon candidate originating from a jet. While such backgrounds are sub-dominant in the dimuon final state, this is less so for other lepton flavors [30, 139–141]. Whatever the final state, such backgrounds should be investigated carefully by experiments through dedicated studies based on collision data, as usually done in collider searches for LNV [18, 20].

To help define our analysis’s signal region, we present in figure 8 the distribution of (a) $M(j_1, j_2)$ and (b) $p_T^{\mu 2}$ for the signal process after pre-selection cuts, assuming representative heavy neutrino masses $m_N = 750$ GeV (darkest cross), 1.5 TeV (dark cross), and 5 TeV (light cross). We also plot after pre-selection cuts the $W^\pm V$ (light), EW $W^\pm W^\pm jj$ (dark), and QCD $W^\pm W^\pm jj$ (darkest) backgrounds, as well as their sum (black cross). The curves are normalized to $\mathcal{L} = 300 \text{ fb}^{-1}$.

Qualitatively, we observe that background processes tend towards smaller values of transverse momentum and invariant mass while the signal process tends towards larger values and exhibit broader, wider distributions. More quantitatively, we observe in figure 8(a) that background processes peak at $M(j_1, j_2) \lesssim 800$ GeV and taper off for larger invariant masses. This contrasts with the signal samples, which peak at $M(j_1, j_2) \gtrsim 900$ GeV,

plateau for a couple hundred GeV, and then gradually fall off. While the lightest heavy neutrinos benchmarks stay above the SM background for most all values of $M(j_1, j_2)$, we observe that the heaviest benchmark mass at $m_N = 5$ TeV only exceeds the background for $M(j_1, j_2) \gtrsim 4.5$ TeV. Values of active-sterile mixing below unity will naturally worsen this separation power.

In figure 8(b) we observe that all backgrounds peak at $p_T^{\mu 2} \sim m_N/2 \sim 40$ GeV – 45 GeV, and quickly dissipate at higher p_T . As anticipated, this shows that backgrounds are driven by resonant weak boson production, though not exclusively. Signal rates become more prominent for $p_T \gtrsim 50$ GeV – 100 GeV. For heavy neutrino masses beyond a few hundred GeV we report high selection efficiencies when requiring $p_T^{\mu 2}$ above this range, but less so for lower masses. In this regime, developing an alternative analysis strategy may increase the sensitivity but goes beyond the scope of this work.

For $p_T^{\mu 2} \gtrsim 300$ GeV, we find that the total background rate is strongly suppressed. Subsequently, due to its simplicity, we define our signal region by requiring, in addition to pre-selection requirements, that both same-sign leptons carry p_T above 300 GeV. We summarize this in the bottom of Table III. For the heavy neutrino masses under consideration, we find that about $\varepsilon \sim 15\%$ to 80% of pre-selection signal events survive signal region requirements. For representative masses, we report in last column of table IV the signal rate cross section σ^{SR} and the corresponding selection efficiency $\varepsilon = \sigma^{\text{SR}}/\sigma^{\text{Pre}}$.

After all selection cuts, we find that the total background rate reaches about $\sigma_b^{\text{All cuts}} \approx 2.35$ ab. For each background and their sum, we list in Table V the expected number of background events after full selection for the nominal LHC scenario (LHC) with $\mathcal{L} = 300 \text{ fb}^{-1}$




FIG. 9. Expected 95% CL sensitivity at the $\sqrt{s} = 13$ TeV LHC (300 fb^{-1}) and HL-LHC (3 ab^{-1}) on the squared active-sterile mixing element $|V_{\mu N}|^2$ as a function of heavy neutrino mass following the $W^\pm W^\pm \rightarrow \mu^\pm \mu^\pm$ analysis described in the text. Also shown are the direct limits (35.9 fb^{-1}) set by CMS using the CCDY and $W\gamma$ fusion channels [13], an extrapolation of the CMS to the HL-LHC, and indirect limits [70].

as well as for the high-luminosity scenario (HL-LHC), where our estimate is computed by simply rescaling the luminosity to $\mathcal{L} = 3 \text{ ab}^{-1}$. At the LHC, less than one background event is expected to pass the selection.

VI.3. Results

To quantify the expected excess number of $W^\pm W^\pm \rightarrow \mu^\pm \mu^\pm$ signal events over the number of SM background events, we follow the recommendations of Ref. [142] and employ asymptotic distributions of test statistics. In particular, we define our signal significance Z as [143, 144]:

$$Z = \frac{(n - n_b)}{|n - n_b|} \sqrt{2 \left[n \log x - \frac{n_b^2}{\delta_b^2} \log y \right]}, \quad \text{with} \quad (6.5)$$

$$x = \frac{n(n_b + \delta_b^2)}{n_b^2 + n\delta_b^2}, \quad \text{and} \quad y = 1 + \frac{\delta_b^2(n - n_b)}{n_b(n_b + \delta_b^2)}. \quad (6.6)$$

Here, $n = n_s + n_b$ is the total number of observed events, $n_s = \mathcal{L} \times \sigma_s^{\text{SR}}$ is the number of signal events expected for an integrated luminosity of \mathcal{L} and signal region rate σ_s^{SR} . The quantity $n_b = \mathcal{L} \times \sigma_b^{\text{SR}}$ is the number of background events expected for a signal region rate σ_b^{SR} , and δ_b is the uncertainty in n_b . Based on experimental measurements of the $W^\pm W^\pm$ scattering process and associated control regions at $\sqrt{s} = 13$ TeV [35, 145], we conservatively estimate our background uncertainty to be 20%, *i.e.*, we set $\delta_b = 0.2$.

TABLE VI. Expected exclusion (excl.) and discovery (disc.) limits at the LHC (300 fb^{-1}) and HL-LHC (3 ab^{-1}) on the squared heavy neutrino mixing with the muon $|V_{\mu N}|^2$ following the analysis described in the text, as well as acceptance and efficiency with respect to the generator-level cross section.

m_N [GeV]	$\mathcal{L} = 300 \text{ fb}^{-1}$		$\mathcal{L} = 3 \text{ ab}^{-1}$		$\frac{\sigma^{\text{SR}}}{\sigma^{\text{gen.}}} [\%]$
	$ V_{\mu N}^{\text{excl.}} ^2$	$ V_{\mu N}^{\text{disc.}} ^2$	$ V_{\mu N}^{\text{excl.}} ^2$	$ V_{\mu N}^{\text{disc.}} ^2$	
50	0.55	0.81	0.31	0.53	0.6
150	0.13	0.24	0.072	0.13	3.9
300	0.080	0.15	0.044	0.077	7.8
450	0.064	0.12	0.035	0.062	12.1
600	0.058	0.10	0.032	0.056	15.6
750	0.057	0.10	0.031	0.054	18.2
900	0.056	0.10	0.031	0.054	21.1
1000	0.056	0.10	0.031	0.054	22.2
1250	0.059	0.11	0.033	0.057	24.2
1500	0.063	0.12	0.034	0.060	26.2
1750	0.067	0.12	0.037	0.064	27.1
2000	0.071	0.13	0.039	0.068	28.4
2500	0.081	0.15	0.044	0.078	29.4
5000	0.14	0.25	0.074	0.13	31.4
7500	0.19	0.36	0.11	0.19	32.2
10000	0.25	0.46	0.14	0.24	32.5
15000	0.34	0.62	0.18	0.32	32.6
20000	0.49	0.81	0.27	0.47	32.6

As discussed in Ref. [142], the significance estimate Z is consistent with a Poisson-counting likelihood where the background-rate nuisance parameters are constrained by auxiliary Poisson measurements [143]. This constraint is performed, for example, by using control samples enriched with background events. Defining such control samples, which are ultimately employed in the likelihood fits that constrain the normalization of backgrounds in the signal region, is beyond our scope. They are, however, commonly carried out in LHC analyses by choosing control samples in regions of phase space as close as possible to the signal region, but where no signal is expected.

We report in Table VI that $|V_{\mu N}|^2 \gtrsim 0.06 - 0.6$ ($0.03 - 0.3$) can be probed at 95% CL for $m_N = 50 \text{ GeV} - 20 \text{ TeV}$ with $\mathcal{L} = 300 \text{ fb}^{-1}$ (3 ab^{-1}). Under the assumption that the mass scale of one or more heavy neutrinos is much heavier than collision scales at $\sqrt{s} = 13 \text{ TeV}$, then in analogy to interpretations of searches for $0\nu\beta\beta$ decay, the LHC expected sensitivity at 95% CL can be expressed as

$$\left| \sum_{k'=4}^{n_R+3} \frac{V_{\mu N_k}^2}{m_{N_k}} \right| \gtrsim 2.5 (1.4) \cdot 10^{-2} \text{ TeV}^{-1}. \quad (6.7)$$

Due to higher background rates, we anticipate slightly worse sensitivity for the $e^\pm \mu^\pm jj$ and $e^\pm e^\pm jj$ channels. For final-states involving hadronic taus, we anticipate

even weaker (but still comparable) sensitivity due to tagging efficiencies. Dedicated studies of these complementary signatures are strongly encouraged.

We find that the proposed analysis has a strong potential to significantly extend the current sensitivity of ATLAS and CMS searches for resonant heavy neutrino masses beyond $m_N \sim 750$ GeV and up to masses at the $\mathcal{O}(10 \text{ TeV})$ scale, as shown in figure 9. Similar to the existing ATLAS and CMS searches, the proposed analysis however does not reach a sensitivity comparable to the limits set by indirect precision measurements (see section II.2). The analysis does, however, offer a direct test of the $0\nu\beta\beta$ decay mechanism in lepton flavor configurations that are not accessible at nuclear energy scales.

VII. OUTLOOK

Discovering the $W^\pm W^\pm \rightarrow \ell_i^\pm \ell_j^\pm$ process in LHC collisions would present unambiguous evidence that LN is violated at the TeV scale, and have far-reaching repercussions for both theory and experiment. In light of the encouraging sensitivity reported in section VI.3, we now briefly consider the outlook for further improvements to our proposed experimental analysis (section VII.1), as well as the possible application of our work to other scenarios, including when LN is conserved (section VII.2).

VII.1. Improving the Experimental Analysis

The analysis cuts chosen and outlined in section VI.2 include only a simple set of selection criteria, which were derived to obtain a good significance for a large range of heavy neutrino masses and with the aim of being robust against the effects of finite detector resolution. Obviously, the selection can be optimized for individual neutrino masses. This is especially true for the lower mass range, where our proposed analysis can add sensitivity to current searches for resonant heavy neutrinos. Such improvements can be roughly grouped into those which further suppress the $W^\pm W^\pm$ or diboson backgrounds.

An optimized analysis that also takes into account the resolution and acceptance of specific sub-detectors should considerably improve our sensitivity estimate. An obvious choice would be to use E_T^{miss} in the event selection. An upper cut on E_T^{miss} would especially help reduce the impact of $W^\pm W^\pm$ production, which has otherwise the same topology as the signal, but was omitted since it is particularly sensitive to detector resolution. As examined in section V.2, the heavy neutrino signal is characterized by high muon momenta that can reach the TeV scale. Even for a small fractional mis-measurement of muon momentum this can induce a considerable amount of E_T^{miss} . Hence correlating the E_T^{miss} and muon momentum (or similarly jet momentum) in the selection would be a way to mitigate some of the resolution effects.

WW scattering processes, such as the heavy neutrino signal considered in this paper, commonly feature suppressed central hadronic activity. As a consequence, implementing static jet vetoes [110–113], or their dynamic counterpart [30, 92, 135, 146, 147], can be exploited to further reduce diboson and top quark processes (or in general all non-VBF backgrounds). One can optimize the corresponding selections based on detector efficiency and resolution for low momentum jets as well as take into account LHC pileup conditions. Since the signal process discussed in this paper is modeled at NLO in QCD with parton shower matching, it can be used to study improvements connected to central hadronic activity.

Lastly, our projections for the HL-LHC consisted of only a re-scaling of the results obtained for a smaller data set. However, the planned detector upgrades for ATLAS [148] and CMS [149] will also allow one to refine the analysis’s selection criteria. In particular, the extended coverage of the tracking system will be highly relevant for WW scattering processes due to the use of tracking information for jets in the forward region.

VII.2. Applications to Other Seesaw Searches

In this study we have focused exclusively on the LN-violating process $pp \rightarrow \ell_i^\pm \ell_j^\pm jj$, when mediated by same-sign WW scattering and the t -channel exchange of a heavy Majorana neutrino, as shown in figure 1. In a complementary fashion, the LN-conserving process $pp \rightarrow \ell_i^\pm \ell_j^\mp jj$, which proceeds through opposite-sign WW scattering, is also possible [26, 36]. One could anticipate that the differences in helicity inversion (see section V.1) between the LN-violating $W^\pm W^\pm \rightarrow \ell_i^\pm \ell_j^\pm$ sub-process and the LN-conserving $W^+ W^- \rightarrow \ell_i^\pm \ell_j^\mp$ sub-process results in substantially different kinematic distributions. However, for heavy neutrinos in the range of $m_N = 750$ GeV to 5 TeV, this may not be the case.

As reported in section V.2, we found that the $W^\pm W^\pm \rightarrow \ell_i^\pm \ell_j^\pm$ sub-process in $pp \rightarrow \ell_i^\pm \ell_j^\pm jj$ behaves like a factorizable system with kinematics that are nearly independent of the hadronic environment. This means that much of the charged lepton kinematic properties are driven more by momentum conservation in $2 \rightarrow 2$ scattering than some complex spin correlation. It is arguable that many of the kinematic leverages over backgrounds that we find, *e.g.*, large dijet invariant masses and back-to-back charged lepton trajectories, will also hold for the LN-conserving case. While the $pp \rightarrow \ell_i^\pm \ell_j^\mp jj$ collider signature inherently has a much larger background rate than the LN-violating one, we stress that lepton flavor violation is forbidden in the SM. Therefore, requiring that $\ell_i \neq \ell_j$ and that E_T^{miss} is small, as done for example in Ref. [150] for low-scale Type I Seesaw models [52–57], can significantly reduce SM backgrounds.

As a final remark, we comment on the applicability of our analysis to other neutrino mass models. This include, for example, the Type II Seesaw model [51, 151–154],

wherein the LN-violating, VBF sub-process $W^\pm W^\pm \rightarrow \Delta^{\pm\pm(*)} \rightarrow \ell_i^\pm \ell_j^\pm$ can occur through a possibly non-resonant, s -channel, doubly charged Higgs boson $\Delta^{\pm\pm}$, as well as Left-Right (LR) symmetric models [155–159], wherein the LN-violating, VBF sub-process $W_R^\pm W_R^\pm \rightarrow \ell_i^\pm \ell_j^\pm$ can proceed through two W_R^\pm gauge bosons and a t -channel Majorana neutrino. For LR scenarios this is interesting in the event that resonant production of W_R is not within the kinematic reach of the LHC [160, 161]. As both scenarios can mimic our $pp \rightarrow \ell_i^\pm \ell_j^\pm jj$ collider signature, its discovery does not automatically prove the existence of RH neutrinos. On the other hand, as both processes occur through the scattering of two color-singlet, massive gauge bosons, most of the color and Lorentz structure remains identical to the original case that we study. Therefore, we anticipate again that the VBF sub-processes approximately factorize, resulting in charged lepton kinematics that resemble those presented in section V.2. As a result, the collider analysis that we propose can readily and justifiably be recast in terms of the Type II and LR symmetric models.

VIII. SUMMARY AND CONCLUSIONS

Motivated by the possible non-conservation of LN in nature, we have investigated the potential to search for heavy Majorana neutrinos in same-sign $W^\pm W^\pm$ scattering at the LHC and the HL-LHC. The experimental signature of two forward jets from VBF, two same-sign leptons, and the lack of substantial missing transverse momenta is interesting in its own right as, to our knowledge, it was not yet explored experimentally at the LHC.

As a benchmark scenario we use the Phenomenological Type I Seesaw model with two key aspects to be probed experimentally: the mass m_N of a mostly sterile neutrino N and its mixing with the active neutrinos $|V_{\ell N}|$. Current searches at the LHC target resonant production modes, such as the Drell-Yan and $W\gamma$ fusion mechanisms, which have the advantage of a factor $|V_{\ell N}|^2$ less suppression compared to the $W^\pm W^\pm \rightarrow \ell_i^\pm \ell_j^\pm$ channel. They however suffer from rapidly falling scattering rates at increasing heavy neutrino masses due to matrix element and phase space suppression. For these reasons LHC searches that employ resonant production modes can only probe masses up to $m_N = 3 - 4$ TeV [30].

To conduct this study, we developed in section IV simulation prescriptions at NLO in QCD with parton shower-matching for both the VBF signal process and backgrounds based on the HEAVYN UFO libraries and the MG5AMC simulation suite. We then extensively studied in section V the phenomenology of the $W^\pm W^\pm \rightarrow \ell_i^\pm \ell_j^\pm$ process at the amplitude and differential levels. We find that “bare” cross section for the full, $2 \rightarrow 4$ signal process at NLO peaks for heavy neutrino masses of around 1 TeV and can reach up to $\sigma/|V_{\ell N}|^4 \sim 10$ fb at $\sqrt{s} = 13$ TeV. Apart from the large rapidity gap between the two leading jets and large dijet invariant mass, the signal also

features very high lepton momenta, among other characteristics, which can be exploited for an effective background suppression.

In section VI we designed our collider analysis, employing the DELPHES framework to simulate the response of a typical LHC detector. Our analysis was deliberately kept simple and considers only final states with same-sign muon pairs to obtain reliable and robust results. Accordingly, dedicated analyses exploiting the suppressed QCD radiation in VBF processes, the angular separation of the same-sign lepton pair, or the correlation between the measured missing transverse momenta and very high- p_T leptons should improve on our projected sensitivity.

In section VI.3 we show that with the LHC Run 2 and expected Run 3 data sets, $|V_{\mu N}|^2$ can be probed down to $0.06 - 0.3$ at 95% CL for heavy neutrino masses in the range $m_N = 1 - 10$ TeV, and that masses at $m_N = 20$ TeV can be probed for $|V_{\mu N}|^2$ down to 0.5. At the HL-LHC, this can be improved by a factor of 2. Comparable results are anticipated for other lepton flavors. We find that the $W^\pm W^\pm$ fusion channel extends significantly the current mass reach based on resonant production modes and adds valuable sensitivity to the masses above a few hundred GeV. Finally in section VII, we give an outlook on areas where our proposed analysis can be improved as well as on complementary applications of our results.

ACKNOWLEDGEMENTS

The authors are grateful to Andreas Hoecker, Jordy de Vries, Juergen Reuter, Alishaan Tamarit for helpful discussions. BF and RR acknowledge the (virtual) hospitality of the DESY ATLAS group.

JN and KP acknowledge support by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany’s Excellence Strategy – EXC 2121 “Quantum Universe” – 390833306. RR is supported under the UCLouvain fund “MOVE-IN Louvain” and acknowledge the contribution of the VBSCan COST Action CA16108. This work has received funding from the European Union’s Horizon 2020 research and innovation programme as part of the Marie Skłodowska-Curie Innovative Training Network MCnetITN3 (grant agreement no. 722104), FNRS “Excellence of Science” EOS be.h Project No. 30820817.

Computational resources have been provided by the supercomputing facilities of the Université catholique de Louvain (CISM/UCL) and the Consortium des Équipements de Calcul Intensif en Fédération Wallonie Bruxelles (CÉCI) funded by the Fond de la Recherche Scientifique de Belgique (F.R.S.-FNRS) under convention 2.5020.11 and by the Walloon Region.

Appendix A: Constraints on heavy Majorana neutrinos from $0\nu\beta\beta$ decay searches

In this appendix we derive constraints on heavy Majorana neutrinos that arise from direct searches for nuclear $0\nu\beta\beta$ decay as reported in equation (2.7). To do this, we assume that the decay is solely mediated by the exchange of heavy states N_k of mass m_{N_k} that couple to SM particles according to the Lagrangian of equation (2.3). We work in the standard factorization picture [162–165]. This stipulates that the transition rate for the decay process of nucleus (A, Z) into nucleus $(A, Z + 2)$,

$$(A, Z) \rightarrow (A, Z + 2) + 2e^-, \quad (\text{A1})$$

can be expressed as a product of the two-body phase space factor $G_{0\nu}$ for the (e^-e^-) -system; a nuclear matrix element (NME) \mathcal{A} ; and the propagators for the states N_k . Under the assumption that the $0\nu\beta\beta$ decay process is mediated only by the t -channel exchange of W bosons and N_k , the NME simplifies to $\mathcal{A} \approx \mathcal{A}_N$. Explicitly, these are related to the decay half-life ($T_{1/2}^{0\nu}$) by

$$\frac{1}{T_{1/2}^{0\nu}} = G_{0\nu} m_p^2 |\mathcal{A}_N|^2 \left| \sum_k V_{ek}^2 \frac{m_{N_k}}{(t - m_{N_k}^2)} \right|^2. \quad (\text{A2})$$

The proton mass $m_p \approx 0.938$ GeV is introduced to render \mathcal{A}_N dimensionless. Typical momentum transfers in nuclear $0\nu\beta\beta$ decay are of the order $\sqrt{|t|} \sim \mathcal{O}(0.1 \text{ GeV})$. As we are interested in EW-scale and TeV-scale neutrinos, this allows us to expand equation (A2) and obtain

$$\left| \sum_k V_{ek}^2 \frac{m_{N_k}}{(t - m_{N_k}^2)} \right|^2 = \left| \sum_k V_{ek}^2 \frac{-1}{m_{N_k} \left(1 - \frac{t}{m_{N_k}^2}\right)} \right|^2 \quad (\text{A3})$$

$$= \left| \sum_k \frac{V_{ek}^2}{m_{N_k}} \right|^2 + \mathcal{O}\left(\frac{t}{m_{N_k}^2}\right). \quad (\text{A4})$$

By neglecting $\mathcal{O}(t/m_{N_k}^2)$ contributions we can invert equation (A2) and translate an experimental lower bound on $T_{1/2}^{0\nu}$ into an upper bound on the mixing-over-mass ratio of Majorana neutrinos. In the following we focus on the ${}^{76}\text{Ge} \rightarrow {}^{76}\text{Se} + 2e^-$ transition, and consider the recent experimental limits by the GERDA experiment [69, 166].

Following Ref. [167], we use the NMEs of Refs. [168, 169]. These employ the so-called Self-consistent Renormalized Quasiparticle Random Phase Approximation (SRQRPA), and make use of two potential models to describe the nucleon-nucleon interactions, namely the Argonne and Charge Dependent Bonn (CD-Bonn) models. The calculations moreover rely on intermediate (Intm.) and large (Large) single-particle spacing, *i.e.*, eigenstate multiplicity, an axial-vector coupling constant $g_A = 1.25$, and a nuclear radius $R = 1.1 \text{ fm} \times A^{1/3}$. For ${}^{76}\text{Ge}$, we

TABLE VII. Values of the NMEs for the nucleon-nucleon interaction models under consideration [168, 169] (second column), and the corresponding exclusion limits at 90% CL on heavy neutrino masses and mixing extracted from results by the GERDA experiment [69] (third column).

NME model		\mathcal{A}_N	$ \sum_k V_{ek}^2/m_k $
Argonne	intm.	232.8	$4.12 \times 10^{-6} \text{ TeV}^{-1}$
Argonne	large	264.9	$3.62 \times 10^{-6} \text{ TeV}^{-1}$
CD-Bonn	intm.	351.1	$2.73 \times 10^{-6} \text{ TeV}^{-1}$
CD-Bonn	large	411.5	$2.33 \times 10^{-6} \text{ TeV}^{-1}$

list in Table VII the \mathcal{A}_N for the four nuclear potential configurations. The uncertainty in \mathcal{A}_N is estimated by considering the envelope spanned by the configurations.

We use the phase space factor $G_{0\nu} = G_{0\nu}^{(0)} g_A^4$, as derived by Ref. [170], assuming an axial-vector cutoff of $g_A = 1.25$. While the polarization-dependent component $G_{0\nu}^{(1)}$ is non-zero, its impact on the total $0\nu\beta\beta$ decay rate vanishes after phase space integration. The derivation of the energy-dependent component $G_{0\nu}^0 \propto 1/R^2$ in Ref. [170] uses a nuclear radius of $R = 1.2 \text{ fm} \times A^{1/3}$. Hence, for consistency with our NMEs, we rescale it by

$$G_{0\nu}^{(0)}(r_0 = 1.1 \text{ fm}) = G_{0\nu}^{(0)}(r_0 = 1.2 \text{ fm}) \times \left(\frac{1.2}{1.1}\right)^2. \quad (\text{A5})$$

For ${}^{76}\text{Ge}$ we obtain the phase space factor,

$$G_{0\nu} \approx 6.866 \times 10^{-15} \text{ yr}^{-1}. \quad (\text{A6})$$

When added in quadrature, the total estimated uncertainty in this number spans about $\delta G_{0\nu} \approx 7\% - 9\%$ [170]. This is considerably smaller than the NME uncertainty, and therefore is neglected in our final constraints on N_k .

After an exposure of $\mathcal{E} = 127.2 \text{ kg}\cdot\text{yr}$, GERDA reports a lower limit on the $0\nu\beta\beta$ decay half-life in ${}^{76}\text{Ge}$ of [69]

$$T_{1/2}^{0\nu} > T_{90\% \text{ CL}}^{\text{GERDA}} = 1.8 \times 10^{26} \text{ yr at 90\% CL}. \quad (\text{A7})$$

For the range of NMEs, this translates into the following upper limit on Majorana neutrino masses and mixing:

$$\left| \sum_{k=4}^{n_R+3} \frac{V_{ek}^2}{m_k} \right| < (2.33 - 4.12) \times 10^{-6} \text{ TeV}^{-1}. \quad (\text{A8})$$

For each NME that we consider we report in Table VII the corresponding limit on the mixing-to-mass ratio. For related discussions on $0\nu\beta\beta$ decay, see Refs. [171–173].

Appendix B: Tailored phase space cuts on leading leptons in MADGRAPH5_AMC@NLO

We document here our implementation of tailored phase space cuts in the event generator MG5AMC.

As described in section IV.2.3, our baseline modeling of the diboson spectrum $pp \rightarrow 3\ell\nu + X$, provides limited MC statistics when the two same-sign leptons carry $p_T^\ell \gtrsim 100 - 150$ GeV but the odd-sign lepton is much softer. To populate this phase space region, we introduce into MG5AMC's phase space integration routines tailored generator-level cuts ($p_T^{\ell\text{-cut}}$) on the p_T of the two leading charged leptons, independent of charge. This is in addition to the baseline cuts of equations (4.3) and (4.4); no further cut is applied to the trailing charged lepton.

High-statistics samples with $p_T^{\ell\text{-cut}} = 75$ GeV and 200 GeV are stitched to the baseline FxFx1j sample through cuts on the truth-level p_T of the sub-leading charged lepton. Within statistical uncertainty, we report that the shape and normalization of the high- p_T tails for the leading charged leptons in the stitched sample reproduce those in the baseline FxFx1j sample. To do this, we make several modifications⁴ to the files `cuts.f` and `setcuts.f` in the `SubProcesses` working directory. Working with version 2.7.1.2 of MG5AMC and for the case of $p_T^{\ell\text{-cut}} = 75$ GeV (`pTlXCut = 75 GeV`), we add to the header in `cuts.f` at L72:

```
c define user cuts for pT12
  double precision pTlXCut,pTlXSum
  double precision pTlXMin,pTlXMax
  logical gotLep1
  parameter(pTlXCut = 75.d0)

and at L159 add the lines

  pTlXSum = 0.d0
  pTlXMax = 0.d0
  pTlXMin = 0.d0
  gotLep1 = .false.
c get pT of hardest and softest charged lepton:
  do i=nincoming+1,nexternal
    if (is_a_lp(i).or.is_a_lm(i)) then
c hypothesize that i hardest and softest
      if(.not.gotLep1) then
        pTlXMax = pt_04(p(0,i))
        pTlXMin = pt_04(p(0,i))
        gotLep1 = .true.
      endif
      pTlXSum = pTlXSum + pt_04(p(0,i))
c update if i is harder or softer
      if(pt_04(p(0,i)).gt.pTlXMax) then
        pTlXMax = pt_04(p(0,i))
      endif
      if(pt_04(p(0,i)).lt.pTlXMin)
        pTlXMin = pt_04(p(0,i))
      endif
    endif
  enddo
c check if subleading lepton pT is hard enough
```

```
pTlXSum = pTlXSum - pTlXMax - pTlXMin
if(pTlXSum.lt.pTlXCut.or.
& pTlXMax.lt.pTlXCut) then
  passcuts_user=.false.
  return
endif
```

In practice, the magnitude of the charged leptons' transverse momenta are first added, then the largest and smallest p_T are subtracted to extract the p_T of the sub-leading charged lepton. If either the leading or sub-leading p_T are below the $p_T^{\ell\text{-cut}}$ threshold (`pTlXCut`), then the phase space point is rejected.

To ameliorate inefficient phase space sampling associated with our cuts, we increment the boundary of the PDF convolution integral $\tau_{\min} = \hat{s}/s$, where $\sqrt{\hat{s}}$ (\sqrt{s}) is the partonic (hadronic) c.m. energy, by $1.5 \times p_T^{\ell\text{-cut}}$. To do this we modify `setcuts.f` at about L422 with:

```
double precision pTlXCut,cutFact
parameter (pTlXCut = 75.d0)
parameter (cutFact = 1.5d0)
```

and at L421 add the following

```
c Increment for pTlXCut on charged leptons
  taumin(iFKS,ichan)=
& taumin(iFKS,ichan)+pTlXCut*cutFact
  taumin_j(iFKS,ichan)=
& taumin_j(iFKS,ichan)+pTlXCut*cutFact
  taumin_s(iFKS,ichan)=
& taumin_s(iFKS,ichan)+pTlXCut*cutFact
```

This is inserted just after the `enddo` closure tag at about L421 and just before the line

```
stot = 4d0*ebeam(1)*ebeam(2)
```

For phase space cuts beyond $p_T^{\ell\text{-cut}} \sim 150$ GeV, we observe a severe instability in phase space integration. As documented elsewhere (see footnote 4), this failure is attributed to inefficient phase space sampling for non-resonant diagrams with massive τ leptons. Hence, for the $p_T^{\ell\text{-cut}} = 200$ GeV sample, we import into MG5AMC the model file `loop_sm-no_tau_mass`, which assumes a massless τ lepton. For looser $p_T^{\ell\text{-cut}}$, we find that this results in sub-percent differences in the cross section normalization from the `loop_sm` model file.

Appendix C: Helicity amplitudes for same-sign WW scattering via heavy Majorana neutrinos

Here we document our calculation of helicity amplitudes for same-sign WW scattering to same-sign lepton pairs when mediated by a heavy Majorana neutrino, as shown diagrammatically in figure 1 and discussed in section V.1. To build a succinct picture of the physics, we employ the effective W approximation [22–24]. In this

⁴ An initial version of these modifications was documented in the URL answers.launchpad.net/mg5amcnlo/+question/691233.

formalism, which is akin to collinear factorization in perturbative QCD, the W boson is treated as a parton of the proton. This allows us to focus on the $2 \rightarrow 2$ subprocess

$$W_\mu^+(p_1^W, \lambda_1^W) + W_\nu^+(p_2^W, \lambda_2^W) \rightarrow \ell^+(p_1^\ell, \lambda_1^\ell) + \ell^+(p_2^\ell, \lambda_2^\ell), \quad (\text{C1})$$

where p and λ denote the 4-momenta and helicities of external particles. The amplitudes reported here supplement the analytic results reported in section V. They are also complementary to the numerical results reported throughout the main text, which evaluate precisely the full $2 \rightarrow 4$ helicity amplitudes for the $pp \rightarrow \ell^\pm \ell^\pm jj + X$ process using the HEAVY NLO UFO libraries [29] in conjunction with the MG5AMC [77, 78] MC event generator (see sections III and IV for related details).

For the above process, we work in the hard-scattering frame, which is equivalent to the WW scattering frame. In this frame, we align coordinate axes such that

$$p_1^W = \frac{M_{WW}}{2}(1, 0, 0, +\beta_W), \quad \beta_W = \sqrt{1 - 4r_W}, \quad (\text{C2})$$

$$p_2^W = \frac{M_{WW}}{2}(1, 0, 0, -\beta_W), \quad r_W = \frac{m_W^2}{M_{WW}^2}, \quad (\text{C3})$$

$$p_1^\ell = \frac{M_{WW}}{2}(1, \sin \theta_1 \cos \phi_1, \sin \theta_1 \sin \phi_1, \cos \theta_1), \quad (\text{C4})$$

$$p_2^\ell = p_1^W + p_2^W - p_1^\ell. \quad (\text{C5})$$

Here $M_{WW}^2 = (p_1^W + p_2^W)^2$ is the invariant mass of the (WW) -system and the remaining invariants are given by

$$t = (p_1^W - p_1^\ell)^2 = m_W^2 - \frac{M_{WW}^2}{2}(1 - \beta_W \cos \theta_1), \quad (\text{C6})$$

$$u = (p_1^W - p_2^\ell)^2 = m_W^2 - \frac{M_{WW}^2}{2}(1 + \beta_W \cos \theta_1). \quad (\text{C7})$$

Working in the unitary gauge and assuming a clockwise fermion flow of leptons [115, 116], the helicity amplitudes in the HELAS basis [117] are given by

$$-i\mathcal{M} = \varepsilon_\mu(p_1^W, \lambda_1^W) \varepsilon_\nu(p_2^W, \lambda_2^W) \mathcal{T}^{\mu\nu}(p_1^\ell, p_2^\ell, \lambda_1^\ell, \lambda_2^\ell) + (t \leftrightarrow u), \quad (\text{C8})$$

where the $(t \leftrightarrow u)$ term accounts for final-state lepton exchange, and the LN-violating tensor current $\mathcal{T}^{\mu\nu}$ is

$$\mathcal{T}^{\mu\nu} = -i \left(\frac{-ig_W}{\sqrt{2}} \right)^2 \frac{V_{\ell N} V_{\ell N}}{(t - m_N^2)} \times \quad (\text{C9})$$

$$\begin{aligned} & [\bar{u}(p_1^\ell, \lambda_1^\ell) \gamma^\mu P_R (\not{p}_N + m_N) \gamma^\nu P_L v(p_2^\ell, \lambda_2^\ell)] \\ & = -i \left(\frac{-ig_W}{\sqrt{2}} \right)^2 \frac{V_{\ell N} V_{\ell N}}{(t - m_N^2)} \times m_N \times \quad (\text{C10}) \\ & [\bar{u}(p_1^\ell, \lambda_1^\ell) \gamma^\mu \gamma^\nu P_L v(p_2^\ell, \lambda_2^\ell)]. \end{aligned}$$

We assume the exchange of a single sterile neutrino mass eigenstate N with momentum $p_N = (p_1^W - p_1^\ell)$ and mass

m_N . For the more general case of multiple heavy neutrinos N_k with masses m_{N_k} , one would substitute:

$$\mathcal{T}^{\mu\nu} \rightarrow \sum_{k=3}^{n_R+3} \mathcal{T}_k^{\mu\nu}, \quad (\text{C11})$$

$$\frac{V_{\ell N} V_{\ell N} m_N}{(t - m_N^2)} \rightarrow \sum_{k=3}^{n_R+3} \frac{V_{\ell N_k} V_{\ell N_k} m_{N_k}}{(t - m_{N_k}^2)}, \quad (\text{C12})$$

to capture the interference of multiple propagating mass eigenstates. Similarly, for two final-state lepton flavors, one substitutes, $V_{\ell N_k} V_{\ell N_k} \rightarrow V_{\ell_1 N_k} V_{\ell_2 N_k}$. Importantly, the spinor and Lorentz index contractions are not modified and therefore are the same for any number of t -channel Majorana neutrino exchanges.

Explicit evaluation (and inspection) of $\mathcal{T}^{\mu\nu}$ shows that the tensor is non-vanishing only when both final-state antileptons carry right-handed polarizations, $(\lambda_1^\ell, \lambda_2^\ell) = (R, R)$. Beyond this, the full matrix element also vanishes identically when both incoming W bosons carry opposite transverse polarizations, *i.e.*, when $(\lambda_1^W, \lambda_2^W) = (\pm 1, \mp 1)$, which follows from an orthogonality of d - and p -wave states. For both t -channel (second column) and u -channel (third column) exchanges of a heavy neutrino, we list in table VIII the exact helicity amplitudes as a function of external particle helicity (first column).

We find that t - and u -channel tensor structures for the $(\lambda_1^W, \lambda_2^W) = (0, \pm 1)$ and $(\pm 1, 0)$ configurations differ simply by a global factor of -1 . The t - and u -channel structures for $(\lambda_1^W, \lambda_2^W) = (\pm, \pm 1)$ differ by exchanges of sine and cosine functions, whereas for $(\lambda_1^W, \lambda_2^W) = (0, 0)$ there is a minor difference in the polar angle dependence. For a fixed set of helicity polarizations, the matrix element of equation (C8) is obtained by the standard coherent summation of t - and u -channel terms.

1. Low-mass limit

When the heavy neutrino and W boson masses are both small compared to the (WW) -scattering scale, *i.e.*, $m_N, m_W \ll M_{WW}$, we can expand each of the squared matrix elements in powers of the ratios

$$r_N = \frac{m_N^2}{M_{WW}^2} \quad \text{and} \quad r_W = \frac{m_W^2}{M_{WW}^2}. \quad (\text{C13})$$

Doing so reveals that remaining transverse-transverse permutations, *i.e.*, $(\lambda_1^W, \lambda_2^W) = (\pm 1, \pm 1)$, as well as LH-longitudinal channels, $(\lambda_1^W, \lambda_2^W) = (-1, 0)$ and $(0, -1)$, either vanish or are sub-leading. In the latter cases, we see the emergence of a helicity suppression that can compete or overcome longitudinal polarization enhancements, which scale as $\varepsilon^\mu(k, 0) \sim k^\mu/m_W + \mathcal{O}(m_W/k^0)$.

To lowest order in r_N and r_W , the RH-longitudinal helicity configurations, $(\lambda_1^W, \lambda_2^W) = (+1, 0)$ and $(0, +1)$, are also sub-leading, but at a parametrically milder degree than the previous combinations. This follows from

TABLE VIII. Helicity amplitude components for the $W_\mu^+(p_1^W, \lambda_1^W) + W_\nu^+(p_2^W, \lambda_2^W) \rightarrow \ell^+(p_1^\ell, \lambda_1^\ell) + \ell^+(p_2^\ell, \lambda_2^\ell)$ process, according to external helicities (first column) for the t -channel (second column) and u -channel (third column) heavy neutrino exchange.

$\lambda_1^W \lambda_2^W \lambda_1^\ell \lambda_2^\ell$	$\varepsilon_\mu(p_1^W, \lambda_1^W) \varepsilon_\nu(p_2^W, \lambda_2^W) \mathcal{T}^{\mu\nu}(p_1^\ell, p_2^\ell) / -i \left(\frac{-ig_W}{\sqrt{2}} \right)^2 \frac{V_{LN} V_{LN} m_N}{(t-m_N^2)}$	$\varepsilon_\mu(p_1^W, \lambda_1^W) \varepsilon_\nu(p_2^W, \lambda_2^W) \mathcal{T}^{\mu\nu}(p_2^\ell, p_1^\ell) / -i \left(\frac{-ig_W}{\sqrt{2}} \right)^2 \frac{V_{LN} V_{LN} m_N}{(u-m_N^2)}$
+1 +1 R R	$2e^{-i\phi_1} M_{WW} \cos^2 \left(\frac{\theta_1}{2} \right)$	$2e^{-i\phi_1} M_{WW} \sin^2 \left(\frac{\theta_1}{2} \right)$
+1 -1 R R	0	0
+1 0 R R	$-\frac{1}{2\sqrt{2}r_W} M_{WW} (1 + \beta_W) \sin \theta_1$	$\frac{1}{2\sqrt{2}r_W} M_{WW} (1 + \beta_W) \sin \theta_1$
-1 +1 R R	0	0
-1 -1 R R	$2e^{-i\phi_1} M_{WW} \sin^2 \left(\frac{\theta_1}{2} \right)$	$2e^{-i\phi_1} M_{WW} \cos^2 \left(\frac{\theta_1}{2} \right)$
-1 0 R R	$-\frac{1}{2\sqrt{2}r_W} e^{-i2\phi_1} M_{WW} (1 - \beta_W) \sin \theta_1$	$\frac{1}{2\sqrt{2}r_W} e^{-i2\phi_1} M_{WW} (1 - \beta_W) \sin \theta_1$
0 +1 R R	$\frac{1}{2\sqrt{2}r_W} e^{-i2\phi_1} M_{WW} (1 + \beta_W) \sin \theta_1$	$-\frac{1}{2\sqrt{2}r_W} e^{-i2\phi_1} M_{WW} (1 + \beta_W) \sin \theta_1$
0 -1 R R	$\frac{1}{2\sqrt{2}r_W} M_{WW} (1 - \beta_W) \sin \theta_1$	$-\frac{1}{2\sqrt{2}r_W} M_{WW} (1 - \beta_W) \sin \theta_1$
0 0 R R	$-\frac{1}{2r_W} e^{-i\phi_1} M_{WW} (1 - 2r_W - \beta_W \cos \theta_1)$	$-\frac{1}{2r_W} e^{-i\phi_1} M_{WW} (1 - 2r_W + \beta_W \cos \theta_1)$

the alignment of the (WW) -system's angular momentum with that of the dilepton system and a single longitudinal polarization enhancement. Explicitly, we obtain

$$|\mathcal{M}(W^+W^+ \rightarrow \ell^+\ell^+)|_{(\lambda_1^W, \lambda_2^W)=(+1,0),(0,+1)}^2 = 2g_W^4 |V_{\ell N}|^4 \left(\frac{r_N}{r_W} \right) \cot^2 \theta_1 + \mathcal{O}(r_N^2, r_W^0). \quad (\text{C14})$$

This expression does not account for the $(1/2!)$ multiplicity factor for identical final-state particles.

We find that the leading polarization configuration in this kinematic limit is the longitudinal-longitudinal channel, $(\lambda_1^W, \lambda_2^W) = (0, 0)$. We attribute its survival in the expansion to the double longitudinal enhancement and is

$$|\mathcal{M}(W^+W^+ \rightarrow \ell^+\ell^+)|_{(\lambda_1^W, \lambda_2^W)=(0,0)}^2 = g_W^4 |V_{\ell N}|^4 \left(\frac{r_N}{r_W} \right) + \mathcal{O}(r_N^2, r_W^{-1}). \quad (\text{C15})$$

To build the partonic, $2 \rightarrow 2$ cross section $\hat{\sigma}$ we employ the standard relationship between scattering rates and matrix elements. This is given by the phase space integral

$$\hat{\sigma}(W^+W^+ \rightarrow \ell^+\ell^+) = \int dPS_2 \frac{d\hat{\sigma}}{dPS_2}, \quad (\text{C16})$$

where the 2-body phase space volume measure is

$$dPS_2(p_1^W + p_2^W; p_1^\ell, p_2^\ell) = \frac{d\cos\theta_1 d\phi_1}{2(4\pi)^2} \beta_W, \quad (\text{C17})$$

and the differential scattering rate is

$$\frac{d\hat{\sigma}}{dPS_2} = \frac{1}{2M_{WW}^2 \beta_W} \frac{1}{\mathcal{S}} \sum_{\{\lambda\}} |\mathcal{M}(\{\lambda\})|^2. \quad (\text{C18})$$

The symmetry factor $\mathcal{S} = 3^2 \cdot 2!$ accounts for spin-averaging over initial-state W polarizations and identical, final-state particles, while the incoherent summation

runs over all external helicities $\{\lambda_1^W, \lambda_2^W, \lambda_1^\ell, \lambda_2^\ell\}$. The velocity factor $\beta_W = \sqrt{1 - 4r_W}$ accounts for the masses of incoming beam particles.

After phase integration over the azimuthal direction, the leading contribution to the polar distribution of ℓ_1^+ in $W^+W^+ \rightarrow \ell^+\ell^+$ scattering is given analytically by

$$\frac{d\hat{\sigma}}{d\cos\theta_1} = \frac{g_W^4}{2^6 3^2 \pi m_W^4} |V_{\ell N}|^4 m_N^2 + \mathcal{O}(r_N^2, r_W^{-1}). \quad (\text{C19})$$

After integration over the polar angle, the total rate is

$$\hat{\sigma} = \frac{g_W^4}{2^5 3^2 \pi m_W^4} |V_{\ell N}|^4 m_N^2 + \mathcal{O}(r_N^2, r_W^{-1}). \quad (\text{C20})$$

For this limit, we report agreement between this expression and numerical evaluations of the same $2 \rightarrow 2$ process using the HEAVYN model with MG5aMC. For multiple heavy neutrinos coupling to potentially different charged lepton flavors, the above generalizes to

$$\hat{\sigma}(W^+W^+ \rightarrow \ell_1^+\ell_2^+) = \frac{g_W^4 (2 - \delta_{\ell_1 \ell_2})}{2^5 3^2 \pi m_W^4} \times \left| \sum_{k=4}^{n_R+3} V_{\ell_1 N_k} m_{N_k} V_{\ell_2 N_k} \right|^2 + \mathcal{O}(r_N^2, r_W^{-1}). \quad (\text{C21})$$

2. High-mass limit

When the heavy neutrino mass is large compared to the $W^\pm W^\pm \rightarrow \ell^+\ell^+$ scattering scale, *i.e.*, when $m_N \gg M_{WW}, m_W$, one can work in the decoupling limit [122] and treat the exchange of N as a point-like, contact interaction. Formally, this entails expanding the heavy neutrino propagator such that

$$\frac{1}{t - m_N^2} = \frac{-1}{m_N^2} + \mathcal{O}\left(\frac{|t|^2}{m_N^4}\right), \quad (\text{C22})$$

$$\frac{1}{u - m_N^2} = \frac{-1}{m_N^2} + \mathcal{O}\left(\frac{|u|^2}{m_N^4}\right). \quad (\text{C23})$$

Inserting this expansion into the amplitudes listed in table VIII reveals a strong destructive interference among most of the helicity permutations. In particular, the only non-vanishing channels correspond to those with incoming W bosons carrying identical polarizations, *i.e.* $(\lambda_1^W, \lambda_2^W) = (0, 0)$ and $(\pm 1, \pm 1)$. Explicitly, the matrix elements for these configurations are given by

$$\begin{aligned} \mathcal{M}(W^+W^+ \rightarrow \ell^+\ell^+) \Big|_{(\lambda_1^W, \lambda_2^W) = (\pm 1, \pm 1)} & \quad (\text{C24}) \\ = -ie^{-i\phi_1} g_W^2 \frac{V_{\ell N}^2}{m_N} M_{WW} + \mathcal{O}\left(\frac{|t|^2}{m_N^4}, \frac{|u|^2}{m_N^4}\right), \end{aligned}$$

$$\begin{aligned} \mathcal{M}(W^+W^+ \rightarrow \ell^+\ell^+) \Big|_{(\lambda_1^W, \lambda_2^W) = (0, 0)} & \quad (\text{C25}) \\ = ie^{-i\phi_1} g_W^2 \frac{V_{\ell N}^2}{m_N} M_{WW} \frac{(1 - 2r_W)}{2r_W} + \mathcal{O}\left(\frac{|t|^2}{m_N^4}, \frac{|u|^2}{m_N^4}\right). \end{aligned}$$

In comparing the two expressions one can see the impact of the longitudinal polarization enhancements, which are responsible for the relative factor of $(M_{WW}/m_W)^2$.

After squaring and integrating over the azimuthal angle, we obtain the leading contributions in the decoupling limit to the polar distribution of ℓ_1^+ in $W^+W^+ \rightarrow \ell^+\ell^+$ scattering. For each polarization channel, this is:

$$\begin{aligned} \frac{d\hat{\sigma}}{d \cos \theta_1} \Big|_{(\lambda_1^W, \lambda_2^W) = (\pm 1, \pm 1)} & \quad (\text{C26}) \\ = \frac{g_W^4}{2^6 \pi} \frac{|V_{\ell N}|^4}{m_N^2} + \mathcal{O}(r_N^{-1}), \end{aligned}$$

$$\begin{aligned} \frac{d\hat{\sigma}}{d \cos \theta_1} \Big|_{(\lambda_1^W, \lambda_2^W) = (0, 0)} & \quad (\text{C27}) \\ = \frac{g_W^4}{2^8 \pi} \frac{|V_{\ell N}|^4}{m_N^2} \frac{(1 - 2r_W)^2}{r_W^2} + \mathcal{O}(r_N^{-1}). \end{aligned}$$

We report good agreement between these expressions and numerical evaluations of helicity-polarized cross sections in this kinematic limit using the HEAVYN model [29] with MG5AMC in conjunction with the formalism of Ref. [174].

After integrating over the polar angle and averaging over all W boson helicities, the total scattering rate is

$$\hat{\sigma} = \frac{g_W^4}{2^7 3^2 \pi} \frac{|V_{\ell N}|^4}{m_N^2} \frac{(1 - 4r_W + 12r_W^2)}{r_W^2} + \mathcal{O}(r_N^{-1}). \quad (\text{C28})$$

For n_R heavy neutrinos coupling to potentially different charged lepton flavors, the above generalizes to

$$\begin{aligned} \hat{\sigma}(W^+W^+ \rightarrow \ell_1^+\ell_2^+) & = \frac{g_W^4(2 - \delta_{\ell_1\ell_2})}{2^7 3^2 \pi r_W^2} & (\text{C29}) \\ & \times \left| \sum_{k=4}^{n_R+3} \frac{V_{\ell_1 N_k} V_{\ell_2 N_k}}{m_{N_k}} \right|^2 + \mathcal{O}(r_N^{-1}, r_W^{-1}). \end{aligned}$$

-
- [1] Q. Ahmad *et al.* (SNO), Direct evidence for neutrino flavor transformation from neutral current interactions in the Sudbury Neutrino Observatory, *Phys. Rev. Lett.* **89**, 011301 (2002), [arXiv:nucl-ex/0204008](#).
 - [2] Y. Ashie *et al.* (Super-Kamiokande), A Measurement of atmospheric neutrino oscillation parameters by SUPER-KAMIOKANDE I, *Phys. Rev. D* **71**, 112005 (2005), [arXiv:hep-ex/0501064](#).
 - [3] R. K. Ellis *et al.*, Physics Briefing Book: Input for the European Strategy for Particle Physics Update 2020, (2019), [arXiv:1910.11775 \[hep-ex\]](#).
 - [4] *2020 Update of the European Strategy for Particle Physics* (CERN Council, Geneva, 2020).
 - [5] E. Ma, Pathways to naturally small neutrino masses, *Phys. Rev. Lett.* **81**, 1171 (1998), [arXiv:hep-ph/9805219](#).
 - [6] A. Atre, T. Han, S. Pascoli, and B. Zhang, The Search for Heavy Majorana Neutrinos, *JHEP* **05**, 030, [arXiv:0901.3589 \[hep-ph\]](#).
 - [7] F. F. Deppisch, P. Bhupal Dev, and A. Pilaftsis, Neutrinos and Collider Physics, *New J. Phys.* **17**, 075019 (2015), [arXiv:1502.06541 \[hep-ph\]](#).
 - [8] Y. Cai, J. Herrero-García, M. A. Schmidt, A. Vicente, and R. R. Volkas, From the trees to the forest: a review of radiative neutrino mass models, *Front. in Phys.* **5**, 63 (2017), [arXiv:1706.08524 \[hep-ph\]](#).
 - [9] Y. Cai, T. Han, T. Li, and R. Ruiz, Lepton Number Violation: Seesaw Models and Their Collider Tests, *Front. in Phys.* **6**, 40 (2018), [arXiv:1711.02180 \[hep-ph\]](#).
 - [10] G. Aad *et al.* (ATLAS), Search for new resonances in mass distributions of jet pairs using 139 fb⁻¹ of pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector, *JHEP* **03**, 145, [arXiv:1910.08447 \[hep-ex\]](#).
 - [11] A. M. Sirunyan *et al.* (CMS), Search for high mass dijet resonances with a new background prediction method in proton-proton collisions at $\sqrt{s} = 13$ TeV, *JHEP* **05**, 033, [arXiv:1911.03947 \[hep-ex\]](#).
 - [12] S. Chatrchyan *et al.* (CMS), A Search for a Doubly-Charged Higgs Boson in pp Collisions at $\sqrt{s} = 7$ TeV, *Eur. Phys. J. C* **72**, 2189 (2012), [arXiv:1207.2666 \[hep-ex\]](#).
 - [13] A. M. Sirunyan *et al.* (CMS), Search for heavy neutral leptons in events with three charged leptons in proton-proton collisions at $\sqrt{s} = 13$ TeV, *Phys. Rev. Lett.* **120**, 221801 (2018), [arXiv:1802.02965 \[hep-ex\]](#).
 - [14] M. Aaboud *et al.* (ATLAS), Search for doubly charged scalar bosons decaying into same-sign W boson pairs with the ATLAS detector, *Eur. Phys. J. C* **79**, 58 (2019), [arXiv:1808.01899 \[hep-ex\]](#).
 - [15] G. Aad *et al.* (ATLAS), Search for heavy neutral leptons in decays of W bosons produced in 13 TeV pp collisions using prompt and displaced signatures with the ATLAS

- detector, *JHEP* **10**, 265, [arXiv:1905.09787 \[hep-ex\]](#).
- [16] A. M. Sirunyan *et al.* (CMS), Search for physics beyond the standard model in multilepton final states in proton-proton collisions at $\sqrt{s} = 13$ TeV, *JHEP* **03**, 051, [arXiv:1911.04968 \[hep-ex\]](#).
- [17] G. Aad *et al.* (ATLAS), Search for type-III seesaw heavy leptons in dilepton final states in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector, (2020), [arXiv:2008.07949 \[hep-ex\]](#).
- [18] A. M. Sirunyan *et al.* (CMS), Search for heavy Majorana neutrinos in same-sign dilepton channels in proton-proton collisions at $\sqrt{s} = 13$ TeV, *JHEP* **01**, 122, [arXiv:1806.10905 \[hep-ex\]](#).
- [19] A. M. Sirunyan *et al.* (CMS), Search for a heavy right-handed W boson and a heavy neutrino in events with two same-flavor leptons and two jets at $\sqrt{s} = 13$ TeV, *JHEP* **05**, 148, [arXiv:1803.11116 \[hep-ex\]](#).
- [20] M. Aaboud *et al.* (ATLAS), Search for heavy Majorana or Dirac neutrinos and right-handed W gauge bosons in final states with two charged leptons and two jets at $\sqrt{s} = 13$ TeV with the ATLAS detector, *JHEP* **01**, 016, [arXiv:1809.11105 \[hep-ex\]](#).
- [21] W.-Y. Keung and G. Senjanovic, Majorana Neutrinos and the Production of the Right-handed Charged Gauge Boson, *Phys. Rev. Lett.* **50**, 1427 (1983).
- [22] S. Dawson, The Effective W Approximation, *Nucl. Phys. B* **249**, 42 (1985).
- [23] G. L. Kane, W. Repko, and W. Rolnick, The Effective W +, Z_0 Approximation for High-Energy Collisions, *Phys. Lett. B* **148**, 367 (1984).
- [24] Z. Kunszt and D. E. Soper, On the Validity of the Effective W Approximation, *Nucl. Phys. B* **296**, 253 (1988).
- [25] F. del Aguila and J. Aguilar-Saavedra, Distinguishing seesaw models at LHC with multi-lepton signals, *Nucl. Phys. B* **813**, 22 (2009), [arXiv:0808.2468 \[hep-ph\]](#).
- [26] A. Datta, M. Guchait, and A. Pilaftsis, Probing lepton number violation via majorana neutrinos at hadron supercolliders, *Phys. Rev. D* **50**, 3195 (1994), [arXiv:hep-ph/9311257](#).
- [27] P. Dev, A. Pilaftsis, and U.-k. Yang, New Production Mechanism for Heavy Neutrinos at the LHC, *Phys. Rev. Lett.* **112**, 081801 (2014), [arXiv:1308.2209 \[hep-ph\]](#).
- [28] D. Alva, T. Han, and R. Ruiz, Heavy Majorana neutrinos from $W\gamma$ fusion at hadron colliders, *JHEP* **02**, 072, [arXiv:1411.7305 \[hep-ph\]](#).
- [29] C. Degrande, O. Mattelaer, R. Ruiz, and J. Turner, Fully-Automated Precision Predictions for Heavy Neutrino Production Mechanisms at Hadron Colliders, *Phys. Rev. D* **94**, 053002 (2016), [arXiv:1602.06957 \[hep-ph\]](#).
- [30] S. Pascoli, R. Ruiz, and C. Weiland, Heavy neutrinos with dynamic jet vetoes: multilepton searches at $\sqrt{s} = 14$, 27, and 100 TeV, *JHEP* **06**, 049, [arXiv:1812.08750 \[hep-ph\]](#).
- [31] S. Chatrchyan *et al.* (CMS), Study of Exclusive Two-Photon Production of W^+W^- in pp Collisions at $\sqrt{s} = 7$ TeV and Constraints on Anomalous Quartic Gauge Couplings, *JHEP* **07**, 116, [arXiv:1305.5596 \[hep-ex\]](#).
- [32] G. Aad *et al.* (ATLAS), Evidence for Electroweak Production of $W^\pm W^\pm jj$ in pp Collisions at $\sqrt{s} = 8$ TeV with the ATLAS Detector, *Phys. Rev. Lett.* **113**, 141803 (2014), [arXiv:1405.6241 \[hep-ex\]](#).
- [33] A. M. Sirunyan *et al.* (CMS), Observation of electroweak production of same-sign W boson pairs in the two jet and two same-sign lepton final state in proton-proton collisions at $\sqrt{s} = 13$ TeV, *Phys. Rev. Lett.* **120**, 081801 (2018), [arXiv:1709.05822 \[hep-ex\]](#).
- [34] M. Aaboud *et al.* (ATLAS), Observation of electroweak $W^\pm Z$ boson pair production in association with two jets in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector, *Phys. Lett. B* **793**, 469 (2019), [arXiv:1812.09740 \[hep-ex\]](#).
- [35] A. M. Sirunyan *et al.* (CMS), Measurements of production cross sections of WZ and same-sign WW boson pairs in association with two jets in proton-proton collisions at $\sqrt{s} = 13$ TeV, *Phys. Lett. B* **809**, 135710 (2020), [arXiv:2005.01173 \[hep-ex\]](#).
- [36] D. A. Dicus, D. D. Karatas, and P. Roy, Lepton non-conservation at supercollider energies, *Phys. Rev. D* **44**, 2033 (1991).
- [37] F. del Aguila, A. Aparici, S. Bhattacharya, A. Santamaria, and J. Wudka, Effective Lagrangian approach to neutrinoless double beta decay and neutrino masses, *JHEP* **06**, 146, [arXiv:1204.5986 \[hep-ph\]](#).
- [38] L. Lehman, Extending the Standard Model Effective Field Theory with the Complete Set of Dimension-7 Operators, *Phys. Rev. D* **90**, 125023 (2014), [arXiv:1410.4193 \[hep-ph\]](#).
- [39] M. Aoki, K. Enomoto, and S. Kanemura, Probing charged lepton number violation via $\ell^\pm \ell'^\pm W^\mp W^\mp$, *Phys. Rev. D* **101**, 115019 (2020), [arXiv:2002.12265 \[hep-ph\]](#).
- [40] B. Fuks, J. Neundorff, K. Peters, R. Ruiz, and M. Saimpert, Probing the Weinberg Operator at Colliders, (2020), [arXiv:2012.09882 \[hep-ph\]](#).
- [41] A. Ali, A. Borisov, and N. Zamorin, Majorana neutrinos and same sign dilepton production at LHC and in rare meson decays, *Eur. Phys. J. C* **21**, 123 (2001), [arXiv:hep-ph/0104123](#).
- [42] O. Panella, M. Cannoni, C. Carimalo, and Y. N. Srivastava, Signals of heavy Majorana neutrinos at hadron colliders, *Phys. Rev. D* **65**, 035005 (2002), [arXiv:hep-ph/0107308](#).
- [43] C.-S. Chen, C.-Q. Geng, and D. V. Zhuridov, Same-sign single dilepton productions at the LHC, *Phys. Lett. B* **666**, 340 (2008), [arXiv:0801.2011 \[hep-ph\]](#).
- [44] D. Binosi, J. Collins, C. Kaufhold, and L. Theussl, JaxoDraw: A Graphical user interface for drawing Feynman diagrams. Version 2.0 release notes, *Comput. Phys. Commun.* **180**, 1709 (2009), [arXiv:0811.4113 \[hep-ph\]](#).
- [45] P. Minkowski, $\mu \rightarrow e\gamma$ at a Rate of One Out of 10^9 Muon Decays?, *Phys. Lett. B* **67**, 421 (1977).
- [46] T. Yanagida, Horizontal gauge symmetry and masses of neutrinos, *Conf. Proc. C* **7902131**, 95 (1979).
- [47] M. Gell-Mann, P. Ramond, and R. Slansky, Complex Spinors and Unified Theories, *Conf. Proc. C* **790927**, 315 (1979), [arXiv:1306.4669 \[hep-th\]](#).
- [48] S. Glashow, The Future of Elementary Particle Physics, *NATO Sci. Ser. B* **61**, 687 (1980).
- [49] R. N. Mohapatra and G. Senjanovic, Neutrino Mass and Spontaneous Parity Nonconservation, *Phys. Rev. Lett.* **44**, 912 (1980).
- [50] R. E. Shrock, General Theory of Weak Leptonic and Semileptonic Decays. 1. Leptonic Pseudoscalar Meson Decays, with Associated Tests For, and Bounds on, Neutrino Masses and Lepton Mixing, *Phys. Rev. D* **24**, 1232 (1981).
- [51] J. Schechter and J. Valle, Neutrino Masses in $SU(2) \times$

- U(1) Theories, *Phys. Rev. D* **22**, 2227 (1980).
- [52] R. Mohapatra, Mechanism for Understanding Small Neutrino Mass in Superstring Theories, *Phys. Rev. Lett.* **56**, 561 (1986).
- [53] R. Mohapatra and J. Valle, Neutrino Mass and Baryon Number Nonconservation in Superstring Models, *Phys. Rev. D* **34**, 1642 (1986).
- [54] J. Bernabeu, A. Santamaria, J. Vidal, A. Mendez, and J. Valle, Lepton Flavor Nonconservation at High-Energies in a Superstring Inspired Standard Model, *Phys. Lett. B* **187**, 303 (1987).
- [55] E. K. Akhmedov, M. Lindner, E. Schnapka, and J. Valle, Left-right symmetry breaking in NJL approach, *Phys. Lett. B* **368**, 270 (1996), [arXiv:hep-ph/9507275](#).
- [56] E. K. Akhmedov, M. Lindner, E. Schnapka, and J. Valle, Dynamical left-right symmetry breaking, *Phys. Rev. D* **53**, 2752 (1996), [arXiv:hep-ph/9509255](#).
- [57] M. Gavela, T. Hambye, D. Hernandez, and P. Hernandez, Minimal Flavour Seesaw Models, *JHEP* **09**, 038, [arXiv:0906.1461 \[hep-ph\]](#).
- [58] T. Han and B. Zhang, Signatures for Majorana neutrinos at hadron colliders, *Phys. Rev. Lett.* **97**, 171804 (2006), [arXiv:hep-ph/0604064](#).
- [59] R. Ruiz, QCD Corrections to Pair Production of Type III Seesaw Leptons at Hadron Colliders, *JHEP* **12**, 165, [arXiv:1509.05416 \[hep-ph\]](#).
- [60] A. Pilaftsis, Radiatively induced neutrino masses and large Higgs neutrino couplings in the standard model with Majorana fields, *Z. Phys. C* **55**, 275 (1992), [arXiv:hep-ph/9901206](#).
- [61] J. Kersten and A. Y. Smirnov, Right-Handed Neutrinos at CERN LHC and the Mechanism of Neutrino Mass Generation, *Phys. Rev. D* **76**, 073005 (2007), [arXiv:0705.3221 \[hep-ph\]](#).
- [62] K. Moffat, S. Pascoli, and C. Weiland, Equivalence between massless neutrinos and lepton number conservation in fermionic singlet extensions of the Standard Model, (2017), [arXiv:1712.07611 \[hep-ph\]](#).
- [63] W. Chao, Z.-g. Si, Y.-j. Zheng, and S. Zhou, Testing the Realistic Seesaw Model with Two Heavy Majorana Neutrinos at the CERN Large Hadron Collider, *Phys. Lett. B* **683**, 26 (2010), [arXiv:0907.0935 \[hep-ph\]](#).
- [64] R. M. Godbole, S. P. Maharathy, S. Mandal, M. Mitra, and N. Sinha, Interference Effect in LNV and LNC Meson Decays for Left Right Symmetric Model, (2020), [arXiv:2008.05467 \[hep-ph\]](#).
- [65] S. King, Unified Models of Neutrinos, Flavour and CP Violation, *Prog. Part. Nucl. Phys.* **94**, 217 (2017), [arXiv:1701.04413 \[hep-ph\]](#).
- [66] Z.-z. Xing, Flavor structures of charged fermions and massive neutrinos, *Phys. Rept.* **854**, 1 (2020), [arXiv:1909.09610 \[hep-ph\]](#).
- [67] T. Asaka and M. Shaposhnikov, The ν MSM, dark matter and baryon asymmetry of the universe, *Phys. Lett. B* **620**, 17 (2005), [arXiv:hep-ph/0505013](#).
- [68] T. Asaka, S. Blanchet, and M. Shaposhnikov, The ν MSM, dark matter and neutrino masses, *Phys. Lett. B* **631**, 151 (2005), [arXiv:hep-ph/0503065](#).
- [69] M. Agostini *et al.* (GERDA), Final Results of GERDA on the Search for Neutrinoless Double- β Decay, (2020), [arXiv:2009.06079 \[nucl-ex\]](#).
- [70] E. Fernandez-Martinez, J. Hernandez-Garcia, and J. Lopez-Pavon, Global constraints on heavy neutrino mixing, *JHEP* **08**, 033, [arXiv:1605.08774 \[hep-ph\]](#).
- [71] A. Osipowicz *et al.* (KATRIN), KATRIN: A Next generation tritium beta decay experiment with sub-eV sensitivity for the electron neutrino mass. Letter of intent, (2001), [arXiv:hep-ex/0109033](#).
- [72] M. Aker *et al.* (KATRIN), Improved Upper Limit on the Neutrino Mass from a Direct Kinematic Method by KATRIN, *Phys. Rev. Lett.* **123**, 221802 (2019), [arXiv:1909.06048 \[hep-ex\]](#).
- [73] A. Loureiro *et al.*, On The Upper Bound of Neutrino Masses from Combined Cosmological Observations and Particle Physics Experiments, *Phys. Rev. Lett.* **123**, 081301 (2019), [arXiv:1811.02578 \[astro-ph.CO\]](#).
- [74] I. Esteban, M. Gonzalez-Garcia, A. Hernandez-Cabezudo, M. Maltoni, and T. Schwetz, Global analysis of three-flavour neutrino oscillations: synergies and tensions in the determination of θ_{23} , δ_{CP} , and the mass ordering, *JHEP* **01**, 106, [arXiv:1811.05487 \[hep-ph\]](#).
- [75] S. Parke and M. Ross-Lonergan, Unitarity and the three flavor neutrino mixing matrix, *Phys. Rev. D* **93**, 113009 (2016), [arXiv:1508.05095 \[hep-ph\]](#).
- [76] S. A. R. Ellis, K. J. Kelly, and S. W. Li, Leptonic Unitarity Triangles, (2020), [arXiv:2004.13719 \[hep-ph\]](#).
- [77] T. Stelzer and W. Long, Automatic generation of tree level helicity amplitudes, *Comput. Phys. Commun.* **81**, 357 (1994), [arXiv:hep-ph/9401258](#).
- [78] J. Alwall, R. Frederix, S. Frixione, V. Hirschi, F. Maltoni, O. Mattelaer, H. S. Shao, T. Stelzer, P. Torrielli, and M. Zaro, The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations, *JHEP* **07**, 079, [arXiv:1405.0301 \[hep-ph\]](#).
- [79] S. Frixione and B. R. Webber, Matching NLO QCD computations and parton shower simulations, *JHEP* **06**, 029, [arXiv:hep-ph/0204244](#).
- [80] R. Frederix, S. Frixione, F. Maltoni, and T. Stelzer, Automation of next-to-leading order computations in QCD: The FKS subtraction, *JHEP* **10**, 003, [arXiv:0908.4272 \[hep-ph\]](#).
- [81] V. Hirschi, R. Frederix, S. Frixione, M. V. Garzelli, F. Maltoni, and R. Pittau, Automation of one-loop QCD corrections, *JHEP* **05**, 044, [arXiv:1103.0621 \[hep-ph\]](#).
- [82] V. Hirschi and O. Mattelaer, Automated event generation for loop-induced processes, *JHEP* **10**, 146, [arXiv:1507.00020 \[hep-ph\]](#).
- [83] N. D. Christensen and C. Duhr, FeynRules - Feynman rules made easy, *Comput. Phys. Commun.* **180**, 1614 (2009), [arXiv:0806.4194 \[hep-ph\]](#).
- [84] N. D. Christensen, P. de Aquino, C. Degrande, C. Duhr, B. Fuks, M. Herquet, F. Maltoni, and S. Schumann, A Comprehensive approach to new physics simulations, *Eur. Phys. J. C* **71**, 1541 (2011), [arXiv:0906.2474 \[hep-ph\]](#).
- [85] C. Degrande, C. Duhr, B. Fuks, D. Grellscheid, O. Mattelaer, and T. Reiter, UFO - The Universal FeynRules Output, *Comput. Phys. Commun.* **183**, 1201 (2012), [arXiv:1108.2040 \[hep-ph\]](#).
- [86] A. Alloul, N. D. Christensen, C. Degrande, C. Duhr, and B. Fuks, FeynRules 2.0 - A complete toolbox for tree-level phenomenology, *Comput. Phys. Commun.* **185**, 2250 (2014), [arXiv:1310.1921 \[hep-ph\]](#).
- [87] C. Degrande, Automatic evaluation of UV and R2 terms for beyond the Standard Model Lagrangians: a proof-of-principle, *Comput. Phys. Commun.* **197**, 239 (2015),

- arXiv:1406.3030 [hep-ph].
- [88] R. Frederix and S. Frixione, Merging meets matching in MC@NLO, *JHEP* **12**, 061, arXiv:1209.6215 [hep-ph].
- [89] P. Artoisenet, R. Frederix, O. Mattelaer, and R. Rietkerk, Automatic spin-entangled decays of heavy resonances in Monte Carlo simulations, *JHEP* **03**, 015, arXiv:1212.3460 [hep-ph].
- [90] J. Alwall, C. Duhr, B. Fuks, O. Mattelaer, D. G. Öztürk, and C.-H. Shen, Computing decay rates for new physics theories with FeynRules and MadGraph 5 aMC@NLO, *Comput. Phys. Commun.* **197**, 312 (2015), arXiv:1402.1178 [hep-ph].
- [91] T. Sjöstrand, S. Ask, J. R. Christiansen, R. Corke, N. Desai, P. Ilten, S. Mrenna, S. Prestel, C. O. Rasmussen, and P. Z. Skands, An introduction to PYTHIA 8.2, *Comput. Phys. Commun.* **191**, 159 (2015), arXiv:1410.3012 [hep-ph].
- [92] B. Fuks, K. Nordström, R. Ruiz, and S. L. Williamson, Sleptons without Hadrons, *Phys. Rev. D* **100**, 074010 (2019), arXiv:1901.09937 [hep-ph].
- [93] B. Jäger, A. Karlberg, S. Plätzer, J. Scheller, and M. Zaro, Parton-shower effects in Higgs production via Vector-Boson Fusion, *Eur. Phys. J. C* **80**, 756 (2020), arXiv:2003.12435 [hep-ph].
- [94] J. R. Christiansen and P. Z. Skands, String Formation Beyond Leading Colour, *JHEP* **08**, 003, arXiv:1505.01681 [hep-ph].
- [95] B. Cabouat and T. Sjöstrand, Some Dipole Shower Studies, *Eur. Phys. J. C* **78**, 226 (2018), arXiv:1710.00391 [hep-ph].
- [96] J. de Favereau, C. Delaere, P. Demin, A. Giammanco, V. Lemaitre, A. Mertens, and M. Selvaggi (DELPHES 3), DELPHES 3, A modular framework for fast simulation of a generic collider experiment, *JHEP* **02**, 057, arXiv:1307.6346 [hep-ex].
- [97] S. Catani, Y. L. Dokshitzer, M. Seymour, and B. Webber, Longitudinally invariant K_t clustering algorithms for hadron hadron collisions, *Nucl. Phys. B* **406**, 187 (1993).
- [98] S. D. Ellis and D. E. Soper, Successive combination jet algorithm for hadron collisions, *Phys. Rev. D* **48**, 3160 (1993), arXiv:hep-ph/9305266.
- [99] M. Cacciari, G. P. Salam, and G. Soyez, The anti- k_t jet clustering algorithm, *JHEP* **04**, 063, arXiv:0802.1189 [hep-ph].
- [100] M. Cacciari and G. P. Salam, Dispelling the N^3 myth for the k_t jet-finder, *Phys. Lett. B* **641**, 57 (2006), arXiv:hep-ph/0512210.
- [101] M. Cacciari, G. P. Salam, and G. Soyez, FastJet User Manual, *Eur. Phys. J. C* **72**, 1896 (2012), arXiv:1111.6097 [hep-ph].
- [102] E. Conte, B. Fuks, and G. Serret, MadAnalysis 5, A User-Friendly Framework for Collider Phenomenology, *Comput. Phys. Commun.* **184**, 222 (2013), arXiv:1206.1599 [hep-ph].
- [103] E. Conte, B. Dumont, B. Fuks, and C. Wymant, Designing and recasting LHC analyses with MadAnalysis 5, *Eur. Phys. J. C* **74**, 3103 (2014), arXiv:1405.3982 [hep-ph].
- [104] E. Conte and B. Fuks, Confronting new physics theories to LHC data with MADANALYSIS 5, *Int. J. Mod. Phys. A* **33**, 1830027 (2018), arXiv:1808.00480 [hep-ph].
- [105] A. Manohar, P. Nason, G. P. Salam, and G. Zanderighi, How bright is the proton? A precise determination of the photon parton distribution function, *Phys. Rev. Lett.* **117**, 242002 (2016), arXiv:1607.04266 [hep-ph].
- [106] A. V. Manohar, P. Nason, G. P. Salam, and G. Zanderighi, The Photon Content of the Proton, *JHEP* **12**, 046, arXiv:1708.01256 [hep-ph].
- [107] V. Bertone, S. Carrazza, N. P. Hartland, and J. Rojo (NNPDF), Illuminating the photon content of the proton within a global PDF analysis, *SciPost Phys.* **5**, 008 (2018), arXiv:1712.07053 [hep-ph].
- [108] R. D. Ball, V. Bertone, S. Carrazza, L. Del Debbio, S. Forte, A. Guffanti, N. P. Hartland, and J. Rojo (NNPDF), Parton distributions with QED corrections, *Nucl. Phys. B* **877**, 290 (2013), arXiv:1308.0598 [hep-ph].
- [109] A. Buckley, J. Ferrando, S. Lloyd, K. Nordström, B. Page, M. Rüfenacht, M. Schönherr, and G. Watt, LHAPDF6: parton density access in the LHC precision era, *Eur. Phys. J. C* **75**, 132 (2015), arXiv:1412.7420 [hep-ph].
- [110] V. D. Barger, K.-m. Cheung, T. Han, and R. Phillips, Strong W^+W^+ scattering signals at pp supercolliders, *Phys. Rev. D* **42**, 3052 (1990).
- [111] V. D. Barger, K.-m. Cheung, T. Han, and D. Zeppenfeld, Single forward jet tagging and central jet vetoing to identify the leptonic WW decay mode of a heavy Higgs boson, *Phys. Rev. D* **44**, 2701 (1991), [Erratum: *Phys.Rev.D* 48, 5444 (1993)].
- [112] J. Bjorken, Rapidity gaps and jets as a new physics signature in very high-energy hadron hadron collisions, *Phys. Rev. D* **47**, 101 (1993).
- [113] R. Fletcher and T. Stelzer, Rapidity gap signals in Higgs production at the SSC, *Phys. Rev. D* **48**, 5162 (1993), arXiv:hep-ph/9306253.
- [114] V. D. Barger, R. Phillips, and D. Zeppenfeld, Mini - jet veto: A Tool for the heavy Higgs search at the LHC, *Phys. Lett. B* **346**, 106 (1995), arXiv:hep-ph/9412276.
- [115] A. Denner, H. Eck, O. Hahn, and J. Kublbeck, Feynman rules for fermion number violating interactions, *Nucl. Phys. B* **387**, 467 (1992).
- [116] A. Denner, H. Eck, O. Hahn, and J. Kublbeck, Compact Feynman rules for Majorana fermions, *Phys. Lett. B* **291**, 278 (1992).
- [117] H. Murayama, I. Watanabe, and K. Hagiwara, HELAS: HELicity amplitude subroutines for Feynman diagram evaluations, (1992), KEK-91-11.
- [118] B. Kayser, Majorana Neutrinos and their Electromagnetic Properties, *Phys. Rev. D* **26**, 1662 (1982).
- [119] R. N. Mohapatra and P. B. Pal, *Massive Neutrinos in Physics and Astrophysics; 3rd ed.*, World Scientific lecture notes in physics (World Scientific, Singapore, 2004).
- [120] T. Han, I. Lewis, R. Ruiz, and Z.-g. Si, Lepton Number Violation and W' Chiral Couplings at the LHC, *Phys. Rev. D* **87**, 035011 (2013), arXiv:1211.6447 [hep-ph].
- [121] R. Ruiz, Quantitative study on helicity inversion in Majorana neutrino decays at the LHC, *Phys. Rev. D* **103**, 015022 (2021), arXiv:2008.01092 [hep-ph].
- [122] T. Appelquist and J. Carazzone, Infrared Singularities and Massive Fields, *Phys. Rev. D* **11**, 2856 (1975).
- [123] R. E. Ruiz, *Hadron Collider Tests of Neutrino Mass-Generating Mechanisms*, Ph.D. thesis, Pittsburgh U. (2015), arXiv:1509.06375 [hep-ph].
- [124] T. Han, G. Valencia, and S. Willenbrock, Structure function approach to vector boson scattering in p p

- collisions, *Phys. Rev. Lett.* **69**, 3274 (1992), arXiv:hep-ph/9206246.
- [125] P. Bolzoni, F. Maltoni, S.-O. Moch, and M. Zaro, Higgs production via vector-boson fusion at NNLO in QCD, *Phys. Rev. Lett.* **105**, 011801 (2010), arXiv:1003.4451 [hep-ph].
- [126] P. Bolzoni, F. Maltoni, S.-O. Moch, and M. Zaro, Vector boson fusion at NNLO in QCD: SM Higgs and beyond, *Phys. Rev. D* **85**, 035002 (2012), arXiv:1109.3717 [hep-ph].
- [127] F. A. Dreyer and A. Karlberg, Vector-Boson Fusion Higgs Pair Production at N³LO, *Phys. Rev. D* **98**, 114016 (2018), arXiv:1811.07906 [hep-ph].
- [128] J. M. Cornwall, D. N. Levin, and G. Tiktopoulos, Derivation of Gauge Invariance from High-Energy Unitarity Bounds on the s Matrix, *Phys. Rev. D* **10**, 1145 (1974), [Erratum: *Phys.Rev.D* 11, 972 (1975)].
- [129] B. W. Lee, C. Quigg, and H. Thacker, Weak Interactions at Very High-Energies: The Role of the Higgs Boson Mass, *Phys. Rev. D* **16**, 1519 (1977).
- [130] M. S. Chanowitz and M. K. Gaillard, The TeV Physics of Strongly Interacting W's and Z's, *Nucl. Phys. B* **261**, 379 (1985).
- [131] T. Appelquist and M. S. Chanowitz, Unitarity Bound on the Scale of Fermion Mass Generation, *Phys. Rev. Lett.* **59**, 2405 (1987), [Erratum: *Phys.Rev.Lett.* 60, 1589 (1988)].
- [132] D. A. Dicus and H.-J. He, Scales of fermion mass generation and electroweak symmetry breaking, *Phys. Rev. D* **71**, 093009 (2005), arXiv:hep-ph/0409131.
- [133] D. A. Dicus and H.-J. He, Scales of mass generation for quarks, leptons and majorana neutrinos, *Phys. Rev. Lett.* **94**, 221802 (2005), arXiv:hep-ph/0502178.
- [134] F. Campanario, R. Roth, and D. Zeppenfeld, QCD radiation in *WH* and *WZ* production and anomalous coupling measurements, *Phys. Rev. D* **91**, 054039 (2015), arXiv:1410.4840 [hep-ph].
- [135] S. Pascoli, R. Ruiz, and C. Weiland, Safe Jet Vetoes, *Phys. Lett. B* **786**, 106 (2018), arXiv:1805.09335 [hep-ph].
- [136] G. Aad *et al.* (ATLAS), Muon reconstruction performance of the ATLAS detector in proton-proton collision data at $\sqrt{s} = 13$ TeV, *Eur. Phys. J. C* **76**, 292 (2016), arXiv:1603.05598 [hep-ex].
- [137] A. Sirunyan *et al.* (CMS), Performance of the CMS muon detector and muon reconstruction with proton-proton collisions at $\sqrt{s} = 13$ TeV, *JINST* **13** (06), P06015, arXiv:1804.04528 [physics.ins-det].
- [138] S. von Buddenbrock, R. Ruiz, and B. Mellado, Anatomy of inclusive *t \bar{t} W* production at hadron colliders, (2020), arXiv:2009.00032 [hep-ph].
- [139] G. Aad *et al.* (ATLAS), Search for supersymmetry at $\sqrt{s} = 13$ TeV in final states with jets and two same-sign leptons or three leptons with the ATLAS detector, *Eur. Phys. J. C* **76**, 259 (2016), arXiv:1602.09058 [hep-ex].
- [140] E. Alvarez, D. A. Faroughy, J. F. Kamenik, R. Morales, and A. Szykman, Four Tops for LHC, *Nucl. Phys. B* **915**, 19 (2017), arXiv:1611.05032 [hep-ph].
- [141] CMS (CMS Collaboration), Jet \rightarrow Tau Misidentification Probability with 2016 pp collision data at $\sqrt{s}=13$ TeV, (2017), CMS-DP-2017-036.
- [142] ATLAS (ATLAS Collaboration), Formulae for Estimating Significance, (2020), ATL-PHYS-PUB-2020-025.
- [143] R. D. Cousins, J. T. Linnemann, and J. Tucker, Evaluation of three methods for calculating statistical significance when incorporating a systematic uncertainty into a test of the background-only hypothesis for a Poisson process, *Nucl. Instrum. Meth. A* **595**, 480 (2008), arXiv:physics/0702156.
- [144] G. Cowan, K. Cranmer, E. Gross, and O. Vitells, Asymptotic formulae for likelihood-based tests of new physics, *Eur. Phys. J. C* **71**, 1554 (2011), [Erratum: *Eur.Phys.J.C* 73, 2501 (2013)], arXiv:1007.1727 [physics.data-an].
- [145] M. Aaboud *et al.* (ATLAS), Observation of electroweak production of a same-sign *W* boson pair in association with two jets in *pp* collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector, *Phys. Rev. Lett.* **123**, 161801 (2019), arXiv:1906.03203 [hep-ex].
- [146] J. K. Michel, P. Pietrulewicz, and F. J. Tackmann, Jet Veto Resummation with Jet Rapidity Cuts, *JHEP* **04**, 142, arXiv:1810.12911 [hep-ph].
- [147] J. Michel, *Factorization and Resummation for Precision Physics at the LHC*, Ph.D. thesis, Hamburg U. (2020).
- [148] ATLAS, Letter of Intent for the Phase-II Upgrade of the ATLAS Experiment, (2012), CERN-LHCC-2012-022.
- [149] CMS, Technical Proposal for the Phase-II Upgrade of the CMS Detector, (2015), CMS-TDR-15-02.
- [150] S. Antusch, E. Cazzato, O. Fischer, A. Hammad, and K. Wang, Lepton Flavor Violating Dilepton Dijet Signatures from Sterile Neutrinos at Proton Colliders, *JHEP* **10**, 067, arXiv:1805.11400 [hep-ph].
- [151] M. Magg and C. Wetterich, Neutrino Mass Problem and Gauge Hierarchy, *Phys. Lett. B* **94**, 61 (1980).
- [152] T. Cheng and L.-F. Li, Neutrino Masses, Mixings and Oscillations in SU(2) x U(1) Models of Electroweak Interactions, *Phys. Rev. D* **22**, 2860 (1980).
- [153] R. N. Mohapatra and G. Senjanovic, Neutrino Masses and Mixings in Gauge Models with Spontaneous Parity Violation, *Phys. Rev. D* **23**, 165 (1981).
- [154] G. Lazarides, Q. Shafi, and C. Wetterich, Proton Lifetime and Fermion Masses in an SO(10) Model, *Nucl. Phys. B* **181**, 287 (1981).
- [155] J. C. Pati and A. Salam, Lepton Number as the Fourth Color, *Phys. Rev. D* **10**, 275 (1974), [Erratum: *Phys.Rev.D* 11, 703-703 (1975)].
- [156] R. N. Mohapatra and J. C. Pati, Left-Right Gauge Symmetry and an Isoconjugate Model of CP Violation, *Phys. Rev. D* **11**, 566 (1975).
- [157] R. Mohapatra and J. C. Pati, A Natural Left-Right Symmetry, *Phys. Rev. D* **11**, 2558 (1975).
- [158] G. Senjanovic and R. N. Mohapatra, Exact Left-Right Symmetry and Spontaneous Violation of Parity, *Phys. Rev. D* **12**, 1502 (1975).
- [159] G. Senjanovic, Spontaneous Breakdown of Parity in a Class of Gauge Theories, *Nucl. Phys. B* **153**, 334 (1979).
- [160] P. S. B. Dev, R. N. Mohapatra, and Y. Zhang, Probing the Higgs Sector of the Minimal Left-Right Symmetric Model at Future Hadron Colliders, *JHEP* **05**, 174, arXiv:1602.05947 [hep-ph].
- [161] R. Ruiz, Lepton Number Violation at Colliders from Kinematically Inaccessible Gauge Bosons, *Eur. Phys. J. C* **77**, 375 (2017), arXiv:1703.04669 [hep-ph].
- [162] M. Doi, T. Kotani, H. Nishiura, K. Okuda, and E. Takasugi, Neutrino Mass, the Right-handed Interaction and the Double Beta Decay. 1. Formalism, *Prog. Theor. Phys.* **66**, 1739 (1981), [Erratum: *Prog.Theor.Phys.* 68, 347 (1982)].

- [163] M. Doi, T. Kotani, H. Nishiura, and E. Takasugi, DOUBLE BETA DECAY, *Prog. Theor. Phys.* **69**, 602 (1983).
- [164] H. Primakoff and S. Rosen, Double beta decay, *Rept. Prog. Phys.* **22**, 121 (1959).
- [165] T. Tomoda, Double beta decay, *Rept. Prog. Phys.* **54**, 53 (1991).
- [166] K. Ackermann *et al.* (GERDA), The GERDA experiment for the search of $0\nu\beta\beta$ decay in ^{76}Ge , *Eur. Phys. J. C* **73**, 2330 (2013), [arXiv:1212.4067 \[physics.ins-det\]](#).
- [167] S. Pascoli, M. Mitra, and S. Wong, Effect of cancellation in neutrinoless double beta decay, *Phys. Rev. D* **90**, 093005 (2014), [arXiv:1310.6218 \[hep-ph\]](#).
- [168] A. Faessler, A. Meroni, S. Petcov, F. Simkovic, and J. Vergados, Uncovering Multiple CP -Nonconserving Mechanisms of $(\beta\beta)_{0\nu}$ Decay, *Phys. Rev. D* **83**, 113003 (2011), [arXiv:1103.2434 \[hep-ph\]](#).
- [169] A. Meroni, S. Petcov, and F. Simkovic, Multiple CP non-conserving mechanisms of $(\beta\beta)_{0\nu}$ -decay and nuclei with largely different nuclear matrix elements, *JHEP* **02**, 025, [arXiv:1212.1331 \[hep-ph\]](#).
- [170] J. Kotila and F. Iachello, Phase space factors for double- β decay, *Phys. Rev. C* **85**, 034316 (2012), [arXiv:1209.5722 \[nucl-th\]](#).
- [171] M. Mitra, G. Senjanovic, and F. Vissani, Neutrinoless Double Beta Decay and Heavy Sterile Neutrinos, *Nucl. Phys. B* **856**, 26 (2012), [arXiv:1108.0004 \[hep-ph\]](#).
- [172] V. Cirigliano, W. Dekens, J. de Vries, M. Graesser, and E. Mereghetti, A neutrinoless double beta decay master formula from effective field theory, *JHEP* **12**, 097, [arXiv:1806.02780 \[hep-ph\]](#).
- [173] M. J. Dolinski, A. W. Poon, and W. Rodejohann, Neutrinoless Double-Beta Decay: Status and Prospects, *Ann. Rev. Nucl. Part. Sci.* **69**, 219 (2019), [arXiv:1902.04097 \[nucl-ex\]](#).
- [174] D. Buarque Franzosi, O. Mattelaer, R. Ruiz, and S. Shil, Automated predictions from polarized matrix elements, *JHEP* **04**, 082, [arXiv:1912.01725 \[hep-ph\]](#).