

# DEUTSCHES ELEKTRONEN-SYNCHROTRON **DESY**

DESY 88-023  
February 1988



## MASSIVE NEUTRINOS

by

P. Langacker

*Deutsches Elektronen Synchrotron DESY, Hamburg*

ISSN 0418-9833

NOTKESTRASSE 85 · 2 HAMBURG 52

**DESY behält sich alle Rechte für den Fall der Schutzrechtserteilung und für die wirtschaftliche Verwertung der in diesem Bericht enthaltenen Informationen vor.**

**DESY reserves all rights for commercial use of information included in this report, especially in case of filing application for or grant of patents.**

**To be sure that your preprints are promptly included in the  
HIGH ENERGY PHYSICS INDEX,  
send them to the following address ( if possible by air mail ) :**

**DESY  
Bibliothek  
Notkestrasse 85  
2 Hamburg 52  
Germany**

## Massive Neutrinos\*

Paul Langacker†  
Deutsches Elektronen Synchrotron, DESY  
D-2000 Hamburg 52, Fed. Rep. Germany

February 25, 1988

### Abstract

Most extensions of the standard  $SU_2 \times U_1$  electroweak model predict nonzero neutrino masses. In this talk I summarize the theoretical expectations for neutrino mass in a variety of possible extensions and discuss their implications for various types of experiments.

## 1 Introduction

The question of whether the neutrino has a nonzero mass is one of the most important issues in both particle physics and astrophysics. In the minimal  $SU_2 \times U_1$  model the neutrinos are predicted to be massless. However, extensions of the standard model involving new  $SU_2$ -singlet neutral fermions (the right-handed neutrino partners needed for Dirac mass terms) or new Higgs representations (to generate Majorana masses) allow non-zero masses. [1] In fact, most extensions of the standard model (e.g. most grand unified theories other than  $SU_5$ ) involve one or both of these mechanisms.

Neutrino mass is also of great importance for astrophysics and cosmology. Masses in the 10 eV range could account for the dark matter of the universe, while masses  $\leq 10^{-2}$  eV could resolve the Solar neutrino problem.

In this talk I will describe the complicated subject neutrino mass: the principle theoretical models and their implications will be described, and the experimental situation will be briefly summarized.

## 2 Weyl, Majorana, and Dirac Neutrinos.

For the weak interactions it is convenient to deal with Weyl two-component spinors  $\psi_L$  or  $\psi_R$ , each of which represents two physical degrees of freedom. The field  $\psi_L$  can annihilate

\*To be published in the Proceedings of the X<sup>th</sup> Workshop on Particles and Nuclei: Neutrino Physics, Heidelberg, October 1987.

†Permanent address: Department of Physics, University of Pennsylvania, Philadelphia, PA 19104.

a left-handed ( $L$ ) particle or create a right-handed ( $R$ ) antiparticle, while  $\psi_L^\dagger$  annihilates a  $L$ -particle or creates an  $R$ -antiparticle. For a  $\psi_R$  field the roles of  $L$  and  $R$  are reversed. An ordinary four-component Dirac field  $\psi$  can be written as the sum  $\psi = \psi_L + \psi_R$  of two Weyl fields, where  $\psi_L$  and  $\psi_R$  are just the chiral projections  $\psi_{L,R} = P_{L,R}\psi$ , with  $P_{L,R} = (1 \pm \gamma_5)/2$ .

Alternatively, one can consider Weyl fermions that do not have distinct partners of the opposite chirality. We will see below that such spinors correspond to particles that are either massless or carry no conserved quantum numbers.

In the free field limit a Weyl field  $\psi_L$  can be written as

$$\psi_L(x) = \sum_{\vec{p}} \left[ b_L(\vec{p}) u_L(\vec{p}) e^{-ip \cdot x} + d_R^\dagger(\vec{p}) v_R(\vec{p}) e^{+ip \cdot x} \right], \quad (1)$$

where  $\sum_{\vec{p}}$  represents  $\int d^3\vec{p} / \sqrt{(2\pi)^3 2E}$ . In(1),  $b_L$  and  $d_R$  are annihilation operators for  $L$  particles and  $R$ -antiparticles, respectively, and  $u_L$  and  $v_R$  are the corresponding (4-component) spinors satisfying  $P_L u_L = u_L$ ,  $P_L v_R = v_R$ ,  $P_R u_L = P_R v_R = 0$ . For a  $\psi_R$  spinor one simply interchanges  $L$  and  $R$ . Equation (1) differs from an ordinary (Dirac) free field in that there is no sum over spin.

It is apparent from (1) that each left-handed (right-handed) particle is necessarily associated with a right-handed (left-handed) antiparticle. The right-handed antiparticle [2] field  $\psi_R^c$  is not independent of  $\psi_L$ , but is closely related to  $\psi_L$ . One has  $\psi_R^c = C \bar{\psi}_L^T$ , where  $C$  is the charge conjugation matrix, defined by  $C \gamma_\mu C^{-1} = -\gamma_\mu^T$ . Similarly, for a  $R$ -Weyl spinor,  $\psi_L^c = C \bar{\psi}_R^T$ . In the special case that  $\psi_L$  is the chiral projection  $P_L \psi$  of a Dirac field  $\psi$ ,  $\psi_R^c$  is just the  $R$ -projection  $P_R \psi^c$  of the antiparticle field  $\psi^c = C \bar{\psi}^T$ .

If  $\psi_R$  and  $\psi_R^c$  both exist, they have the opposite values for all additive quantum numbers. Since the quarks and charged leptons carry conserved quantum numbers (e.g. color and electric charge), they must be Dirac fields - i.e.  $\psi_R$  and  $\psi_R^c$  must be distinct. The only quantum number associated with the neutrinos is lepton number, however, and it is conceivable that that is violated in nature. As we will see, that will allow for two very different possibilities for neutrino mass.

The known neutrinos of the first family are the left-handed electron neutrino  $\nu_{eL}$  and its CP partner, the right-handed "antineutrino"  $\nu_{eR}^c = C \bar{\nu}_{eL}^T$ . These are associated with the  $e_L^-$  and  $e_R^+$ , respectively, in ordinary charged current weak interactions.

Mass terms always take left- and right-handed fields into each other. If one introduces a new field  $N_R$  (distinct from  $\nu_R^c$ ) and its CP conjugate  $N_L^c = C \bar{N}_R^T$  into the theory, then one can write a Dirac (lepton number conserving) mass term

$$-L_{Dirac} = m_D \bar{\nu}_L N_R + h.c., \quad (2)$$

which connects  $N_R$  and  $\nu_L$ . In this case  $\nu_L$ ,  $N_R$ ,  $N_L^c$  and  $\nu_R^c$  form a four component Dirac particle - i.e. one can define  $\nu \equiv \nu_L + N_R$ ,  $\nu^c = N_L^c + \nu_R^c = C \bar{\nu}^T$ , so that  $-L_{Dirac} = m_D \bar{\nu} \nu$ . Clearly lepton number is conserved in this case, because there is no transition between  $\nu$  and  $\nu^c$ . In the free field limit the Dirac neutrino field  $\nu$  has the canonical expression

$$\nu_{Dirac}(x) = \sum_{\vec{p}} \sum_{S=L,R} \left[ b_S(\vec{p}) u_S(\vec{p}) e^{-ip \cdot x} + d_S^\dagger(\vec{p}) v_S(\vec{p}) e^{+ip \cdot x} \right], \quad (3)$$

Usually, the  $N_R$  is an  $SU_2 \times U_1$  singlet, with  $m_D$  generated by an ordinary Higgs doublet, and  $L = L_e + L_\mu + L_\tau$  is conserved in the three family generalization. This possibility is

most similar to the way in which masses are generated for the other fermions ( $e^-$ ,  $u$ ,  $d$ , etc.) in the standard model, but it is difficult to understand why  $m_\nu$  is so small in this case.

Another possibility [3] is that  $N_R$  is a known doublet neutrino, such as  $\nu_{eR}^c$ . This is a variation on the Konopinski-Mahmoud model. [4] Then  $\nu_{eL}$ ,  $\nu_{eR}^c$ ,  $\nu_{\tau L}$  and  $\nu_{eR}^c$  can be combined to form a Dirac neutrino with  $L_e - L_\tau$  conserved.

For the generalization of (2) to  $F$  fermion families one has

$$-L_{Dirac} = \bar{n}_L^0 m_D N_R^0 + h.c., \quad (4)$$

where  $m_D$  is an arbitrary [5]  $F \times F$  mass matrix, and  $n_L^0$  and  $N_R^0$  are  $F$ -component vectors; thus  $n_L^0 = (n_{1L}^0 \ n_{2L}^0 \ \dots \ n_{FL}^0)^T$ , where  $n_{iL}^0$  are the "weak eigenstate" neutrinos - i.e.  $n_{iL}^0$  is associated with  $e_{iL}^-$  in weak transitions. The weak eigenstate neutrinos are related to the neutrinos  $n_{iL}$ ,  $N_{iR}$  of definite mass by unitary transformations  $n_L^0 = V_L n_L$ ,  $N_R^0 = V_R N_R$ , where  $V_L$  and  $V_R$  are  $F \times F$  unitary matrices, determined by

$$V_L^\dagger m_D V_R = m_d = \text{diag}(m_1 \ m_2 \ \dots \ m_F) \quad (5)$$

and  $m_d$  is the diagonal matrix of physical neutrino masses.  $V_L$  and  $V_R$  can be determined by

$$V_L^\dagger m_D m_D^\dagger V_L = V_R^\dagger m_D^\dagger m_D V_R = m_d^2 \quad (6)$$

( $m_D m_D^\dagger$  and  $m_D^\dagger m_D$  are Hermitian). In general  $V_L$  and  $V_R$  are unrelated. If there are no degeneracies then  $V_L$  and  $V_R$  are determined uniquely by (6) up to diagonal phase matrices; i.e. if  $V_{L,R}$  satisfy (6) then so do  $V_{L,R} K_{L,R}$ , where  $K_{L,R}$  are diagonal phase matrices associated with the unobservable phases of the  $n_{iL}$  and  $N_{iR}$  fields. Usually one chooses  $K_L$  to put  $V_L$  into a simple conventional form. Then  $K_R$  is determined by the requirement that  $m_d$  be real.

In the presence of neutrino mass the leptonic weak charged current becomes

$$J_W^\mu = (\bar{\nu}_e \ \bar{\nu}_\mu \ \bar{\nu}_\tau) V_L^\dagger \gamma^\mu (1 + \gamma^5) \begin{pmatrix} e^- \\ \mu^- \\ \tau^- \end{pmatrix} \quad (7)$$

so that  $V_L^\dagger$  is just the analogue of the CKM quark mixing matrix. It describes the relative strengths [7] of the weak transition between the various charged leptons and neutrinos of definite mass.

In a Majorana (lepton number violating) mass term one avoids the need for a new fermion field by coupling the  $\nu_L$  to its CP conjugate  $\nu_R^c$ :

$$-L_M = \frac{1}{2} m \bar{\nu}_L \nu_R^c + h.c. = \frac{1}{2} m \bar{\nu}_L C \bar{\nu}_L^T + h.c. \quad (8)$$

$L_M$  can be thought of as creating or annihilating two neutrinos, and violates lepton number by  $\Delta L = \pm 2$ .  $\nu_L$  and  $\nu_R^c$  can be combined to form a two component Majorana neutrino  $\nu = \nu_L + \nu_R^c$ , so that  $-L_M = \frac{1}{2} m \bar{\nu} \nu$ . One has  $\nu = C \bar{\nu}^T$ , i.e. a Majorana neutrino is its own antiparticle. In the free field limit  $\nu$  is just

$$\nu(x) = \sum_{\bar{p}} \sum_{s=L,R} [b_s(\bar{p}) u_s(\bar{p}) e^{-ipx} + b_s^\dagger(\bar{p}) v_s(\bar{p}) e^{+ipx}], \quad (9)$$

i.e. it has the same form as for a free Dirac field (cf (3)) except that there is no distinction between  $b$  and  $d$  annihilation operators.

The Majorana mass  $m$  in (8) can be generated by the vacuum expectation value (VEV) of a new Higgs triplet [8] or as a higher order effective operator. Majorana masses are popular amongst theorists because they are so different from quark and lepton masses, and there is therefore the possibility of explaining why  $m_\nu$  is so small (if it is non-zero).

For  $F$  fermion families, the Majorana mass term is

$$-L_M = \frac{1}{2} \bar{n}_L^0 M n_R^0 + h.c. \quad (10)$$

where  $M$  is an  $F \times F$  Majorana mass matrix and  $n_L^0$  and  $n_R^0$  are  $F$  component vectors: i.e.  $n_L^0 = (n_{1L}^0 \ \dots \ n_{FL}^0)^T$ ,  $n_R^0 = (n_{1R}^0 \ \dots \ n_{FR}^0)^T$ , where  $n_{iL}^0$  and  $n_{iR}^0$  are weak eigenstate neutrinos and "antineutrinos", related by

$$n_{iR}^0 = C \bar{n}_{iL}^{0T} \quad (11)$$

From (11) one can prove the identity  $\bar{n}_{iL}^0 n_{jR}^0 = \bar{n}_{jL}^0 n_{iR}^0$ , from which it follows that the Majorana mass matrix  $M$  must be symmetric:  $M = M^T$ . Proceeding in analogy to the Dirac case, one can relate the  $n_{iL}^0$  and  $n_{jR}^0$  to mass eigenstate neutrino fields by  $n_L^0 = U_L n_L$ ,  $n_R^0 = U_R n_R^c$ , where  $U_L$  and  $U_R$  are  $F \times F$  unitary matrices chosen so that

$$U_L^\dagger M U_R = M_d = \text{diag}(m_1 \ m_2 \ \dots \ m_F), \quad (12)$$

where  $M_d$  is a diagonal matrix of Majorana mass eigenvalues. Unlike the Dirac case (for which  $m_D$  was an arbitrary matrix and  $V_L$  and  $V_R$  unrelated), the symmetry of  $M$  implies a relation between  $U_L$  and  $U_R$ , viz

$$U_L = U_R^* K^\dagger, \quad (13)$$

where  $K$  is unitary and symmetric. That is, just as in the Dirac case,  $U_L$  is determined from  $U_L^\dagger M M^\dagger U_L = M_d^2$  to be of the form  $U_L = \bar{U}_L K_L$ , where  $K_L$  is a matrix of phases that can be chosen for convenience.  $U_R$  is then determined from (13), where  $K$  is chosen so that  $M_d$  is real and positive. If there are no degeneracies then  $K$  is just a matrix of phases. [9] One can always pick  $K_L$  such that  $K = I$ , but it is not always convenient to do so.

In terms of the mass eigenstates, (10) reduces to

$$-L_M = \frac{1}{2} \sum_{i=1}^F m_i \bar{n}_i n_i + h.c. = \frac{1}{2} \sum_{i=1}^F m_i \bar{n}_i n_i, \quad (14)$$

where  $n_i = n_{iL} + n_{iR}^c$  is the  $i^{\text{th}}$  Majorana mass eigenstate. [10] Written in terms of the  $n_{iL}$ , the weak charged current assumes a form analogous to (7), with  $U_L^\dagger$  replacing  $V_L^\dagger$  to describe the leptonic mixing. [11]

There are several physical distinctions between Dirac and Majorana neutrinos. If the  $\nu_e$  is Majorana, for example, one could have the sequence  $\pi^+ \rightarrow e^+ \nu_e$  followed by  $\nu_e p \rightarrow e^+ n$ . The combined process violates lepton number by two units and is allowed for Majorana but not Dirac neutrinos. Similarly, a hypothetical heavy neutrino  $N$  would undergo the decays  $N \rightarrow e^+ q_1 \bar{q}_2$  and  $N \rightarrow e^- \bar{q}_1 q_2$  with equal rates if it is Majorana, while for a Dirac particle one would have  $N \rightarrow e^+ \bar{q}_1 q_2$ ,  $N^c \rightarrow e^+ q_1 \bar{q}_2$  only [12]. There are differences due

to Fermi statistics in the production of  $\nu\nu$  (Majorana) or  $\nu\nu^c$  (Dirac) pairs near threshold [13], and finally Majorana neutrinos cannot have electromagnetic form factors, such as magnetic moments [14].

It is important to keep in mind, however, that these distinctions must all disappear in the limit that the neutrino mass can be neglected. For  $m_\nu \rightarrow 0$  the  $\nu_R$  component of a Dirac neutrino decouples, and both Majorana and Dirac neutrinos reduce to Weyl two-component neutrinos - there is no difference between them. [15] In particular, lepton number conservation is reestablished smoothly as  $m_\nu \rightarrow 0$  for a Majorana neutrino, because in that limit helicity - which is conserved up to corrections of order  $m_\nu/E_\nu$  - plays the role of an approximate lepton number. For example, the  $\nu_e$  produced in  $\pi^+ \rightarrow e^+\nu_e$  has  $h_\nu = -1$  up to corrections of order  $(m_\nu/E_\nu)^2$  (in rate), while the reaction  $\nu_e p \rightarrow e^+ n$  has a cross section that is suppressed by  $(m_\nu/E_\nu)^2$  for the wrong (negative) helicity.

In many models Dirac and Majorana mass terms are both present. For one doublet neutrino  $\nu_L^0$  (with  $\nu_R^{0c} = C\bar{\nu}_L^{0T}$ ) and one new singlet  $N_R^0$  (with  $N_L^0 = C\bar{N}_R^{0T}$ ), for example, one could have the general mass term

$$-L = \frac{1}{2} \begin{pmatrix} \bar{\nu}_L^0 & \bar{N}_L^0 \end{pmatrix} \begin{pmatrix} m_t & m_D \\ m_D^T & m_S \end{pmatrix} \begin{pmatrix} \nu_R^0 \\ N_R^0 \end{pmatrix} + h.c., \quad (15)$$

where  $m_D = m_D^T$  is a Dirac mass generated by a Higgs doublet (analogous to (2)),  $m_t$  is a Majorana mass for  $\nu_L^0$  generated by a Higgs triplet or effective interaction (cf. (8)), and  $m_S$  is a Majorana mass for  $N_R^0$ , generated by a Higgs singlet or bare mass. Similarly, for  $F$  families (15) still holds provided one interprets  $\nu_L^0$ ,  $N_L^0$ ,  $\nu_R^0$ , and  $N_R^0$  as  $F$  component vectors, and  $m_t$ ,  $m_D$ , and  $m_S$  as  $F \times F$  matrices (with  $m_t = m_t^T$ ,  $m_S = m_S^T$ ). Then, (15) becomes simply

$$-L = \frac{1}{2} \bar{n}_L^0 M n_R^0 + h.c., \quad (16)$$

where  $n_L^0 \equiv (\nu_L^0, N_L^0)^T$  and  $n_R^0 \equiv (\nu_R^0, N_R^0)^T$  are  $2F$  component vectors and  $M$  is the symmetric  $2F \times 2F$  Majorana mass matrix in (15). Equation (16) can be diagonalized in exact analogy with (10-14), yielding finally

$$-L = \frac{1}{2} \sum_{i=1}^{2F} m_i \bar{n}_{iL} n_{iR}^c + h.c. \quad (17)$$

i.e. there are in general  $2F$  Majorana neutrinos, related to  $n_L^0, n_R^0$  by unitary transformations. Unlike the pure Majorana case, however, there is now mixing between particles with different weak interaction properties (e.g.  $n_{iL} = (U_L^\dagger)_{ij} n_{jL}^0$  is a mixture of  $SU_2$  doublets and singlets), which can have important consequences for neutrino oscillations [16] and decays.

It is instructive to see how the Dirac case ( $m_t = m_S = 0$ ) emerges as a limiting case of (15). For a single family one has  $M = m_D \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ . Since  $M$  is Hermitian (for  $m_D$  real) one can diagonalize it by a unitary transformation  $U_L$ . One finds

$$U_L^\dagger M U_L = m_D \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad (18)$$

with  $U_L = \frac{1}{\sqrt{2}} \text{diag}(1, -1)$ ; i.e. the mass eigenstates are

$$\begin{aligned} n_{1L} &= \frac{1}{\sqrt{2}}(\nu_L^0 + N_L^{0c}) \\ n_{2L} &= \frac{1}{\sqrt{2}}(\nu_L^0 - N_L^{0c}) \\ n_{1R}^c &= \frac{1}{\sqrt{2}}(\nu_R^{0c} + N_R) \\ n_{2R}^c &= \frac{1}{\sqrt{2}}(\nu_R^{0c} - N_R). \end{aligned} \quad (19)$$

The negative mass eigenvalue in (18) can be removed by redefining [17] the right-handed fields  $n_{1R} = n_{1R}^c$ ,  $n_{2R} = -n_{2R}^c$ . This is nothing more than taking  $U_R^\dagger M U_R = m_d = m_D \text{diag}(1, 1)$ , where  $U_R$  is given by (13) with  $K = \text{diag}(1, -1)$ . Finally, the two Majorana states  $n_1 = n_{1L} + n_{1R}^c$  and  $n_2 = n_{2L} + n_{2R}^c$  are degenerate. We can therefore reexpress  $L$  in the new basis

$$\begin{aligned} \nu &\equiv \frac{1}{\sqrt{2}}(n_1 + n_2) = \nu_L^0 + N_R^0 \\ \nu^c &\equiv \frac{1}{\sqrt{2}}(n_1 - n_2) = N_L^{0c} + \nu_R^{0c}, \end{aligned} \quad (20)$$

yielding

$$\begin{aligned} -L &= \frac{1}{2} m_D (\bar{n}_{1L} n_{1R}^c + \bar{n}_{2L} n_{2R}^c) + h.c. \\ &= m_D \bar{\nu}_L^0 N_R^0 + h.c. = m_D \bar{\nu} \nu. \end{aligned} \quad (21)$$

This is just a standard Dirac mass term, with a conserved lepton number (i.e. no transition between  $\nu$  and  $\nu^c$ ). A Dirac neutrino is therefore nothing but a pair of degenerate two-component Majorana neutrinos ( $n_1$  and  $n_2$ ), combined to form a 4-component neutrino with a conserved lepton number.

One sometimes refers to a pseudo-Dirac neutrino, which is just a Dirac neutrino to which is added as small lepton number-violating perturbation. For example, one could modify the Dirac mass to

$$M = \begin{pmatrix} \epsilon & m_D \\ m_D & 0 \end{pmatrix}, \quad (22)$$

with  $\epsilon \ll m_D$ . One then finds two Majorana mass eigenstates  $n_\pm$ , with

$$\begin{aligned} n_{+L} &= n_{1L} + \frac{\epsilon}{4} n_{2L} \\ n_{-L} &= -\frac{\epsilon}{4} n_{1L} + n_{2L}, \end{aligned} \quad (23)$$

( $n_{1L}$  and  $N_{2L}$  are defined in (19)), with masses  $m_D \pm \frac{\epsilon}{2}$ .

Other important special cases of (15) are considered below.

### 3 Models of Neutrino Mass

There are many models for neutrino mass [1], all of which have good and bad features. The major classes of models are listed in Table 1, along with the most natural scales for the neutrino masses and for  $\langle m_\nu \rangle$ , an effective mass relevant to neutrinoless double  $\beta$  decay.

Table 1: Models of neutrino mass, along with their most natural scales for the light neutrino masses.

Model	$m_{\nu_e}$	$\langle m_{\nu_e} \rangle$	$m_{\nu_\mu}$	$m_{\nu_\tau}$
Dirac	1 – 10 MeV	0	100 MeV – 1 GeV	1 – 100 GeV
pure Majorana [8] (Higgs triplet)	arbitrary	$m_{\nu_e}$	arbitrary	arbitrary
GUT seesaw [18,19] ( $M \sim 10^{14}$ GeV)	$10^{-11}$ eV	$m_{\nu_e}$	$10^{-6}$ eV	$10^{-3}$ eV
intermediate seesaw [20] ( $M \sim 10^9$ GeV)	$10^{-7}$ eV	$m_{\nu_e}$	$10^{-2}$ eV	10 eV
$SU_{2L} \times SU_{2R} \times U_1$ seesaw [21] ( $M \sim 1$ TeV)	$10^{-1}$ eV	$m_{\nu_e}$	10 KeV	1 MeV
light seesaw [22] ( $M \ll 1$ GeV)	1 – 10 MeV	$\ll m_{\nu_e}$	-	-
charged Higgs [23]	$< 1$ eV	$\ll m_{\nu_e}$	-	-

Dirac neutrinos are exactly like other fermions. They involve a conserved total lepton number (though the individual  $L_e$ ,  $L_\mu$ , and  $L_\tau$  lepton numbers are violated by mixing in general) and therefore do not lead to neutrinoless double beta decay. The problem with Dirac neutrinos is that it is hard to understand why the neutrinos are so much lighter than the other fermions. In the standard model Dirac mass are generated by the vacuum expectation value (VEV)  $v = \sqrt{2}\langle\varphi^0\rangle \simeq (\sqrt{2}G_F)^{-1/2} \simeq 246$  GeV of the neutral component of a doublet of Higgs scalar fields. One has  $m_D = h_\nu v$ , where  $h_\nu$  is the Yukawa coupling




$$L = -\sqrt{2}h_\nu(\bar{\nu}_L \bar{e}_L) \begin{pmatrix} \varphi^0 \\ \varphi^- \end{pmatrix} N_R + h.c. \quad (24)$$

of the neutrino to  $\varphi^0$ .

A  $\nu_e$  mass in the 20 eV range would require an anomalously small Yukawa coupling  $h_{\nu_e} \leq 10^{-10}$ . Moreover,  $h_{\nu_e}$  would have to be smaller by  $m_{\nu_e}/m_e \leq 10^{-4}$  than the analogous Yukawa coupling for the electron. Of course, we do not understand the masses of the other fermions either (or why they range over at least five orders of magnitude), so it is hard to totally exclude the possibility that  $h_{\nu_e}$  is simply small. Nevertheless, the possibility seems sufficiently ugly that it is hard to take seriously unless some mechanism (other than fine-tuning) for the smallness is proposed.

One possibility is that  $h_\nu$  is actually zero to lowest order (tree level) due to some new symmetry, and that  $h_\nu$  is only generated as a higher order correction (i.e. so that  $m_\nu/m_e$  is some power of  $\alpha$ .) This is a very attractive possibility, but no particularly compelling models to implement it have emerged. The idea has recently been resurrected in some superstring inspired models [24], which have difficulty incorporating the seesaw type ideas described below.

Majorana mass terms for the ordinary  $SU_2$ -doublet neutrinos involve a transition from



41175

Figure 1: Dirac, pure Majorana, induced, and charged Higgs generated neutrino masses.

$\nu_R^c$  ( $t_3 = -\frac{1}{2}$ ) into  $\nu_L$  ( $t_3 = +\frac{1}{2}$ ), and therefore must be generated by an operator transforming as a triplet under weak  $SU_2$ .

The simplest possibility is the Gelmini-Roncadelli model [8], in which one introduces a triplet of Higgs fields  $\vec{\varphi}_i = (\varphi_i^0, \varphi_i^-, \varphi_i^{--})$  into the theory. The Yukawa coupling

$$L = \frac{1}{2}h_i(\bar{\nu}_L \bar{e}_L)\vec{\tau} \cdot \vec{\varphi}_i \begin{pmatrix} e_R^c \\ -\nu_R^c \end{pmatrix} = \frac{1}{2}h_i(\bar{\nu}_L \bar{e}_L) \begin{pmatrix} \varphi_i^- & \sqrt{2}\varphi_i^0 \\ \sqrt{2}\varphi_i^{--} & -\varphi_i^- \end{pmatrix} \begin{pmatrix} e_R^c \\ -\nu_R^c \end{pmatrix} \quad (25)$$

then generates a Majorana mass  $m_i = h_i v_i$  for the  $\nu$ , where  $v_i = \sqrt{2}\langle\varphi_i^0\rangle$  is the VEV of the Higgs triplet. Since both  $h_i$  and  $v_i$  are unknown the neutrino mass is unrelated to the other fermions and can in principle be arbitrarily small, at least at tree level.

However, small  $m_{\nu_e}$  is not explained in such models - it is merely parametrized and in fact is almost as problematic as a Dirac mass. The weak neutral current (and  $W$  and  $Z$  masses) require [25]  $v_i \leq 0.08v \sim 20$  GeV. For  $v_i$  close to this limit one requires  $h_i \leq 10^{-9}$ , i.e. almost as bad a fine-tuning as the Dirac case. For  $v_i \ll v$  one can tolerate more reasonable values for  $h_i$ , but then it is difficult to understand the large hierarchy in vacuum expectation values. One generally expects all non-zero VEV's to be comparable

in magnitude unless fine-tunings are performed on the parameters in the Higgs potential. Even if one does this, higher order corrections are likely to upset the hierarchy. [26]

The VEV  $\langle \varphi_t^0 \rangle \neq 0$  necessarily violates lepton number conservation by two units (the Yukawa coupling in (25) does not by itself violate  $L$  because  $\varphi_t$  can be regarded as carrying two units of  $L$ ). If the rest of the Lagrangian conserves  $L$  then lepton-number is spontaneously broken, and there will be an associated massless Goldstone boson, the triplet-Majoron. (This is the version of the model that is usually considered [8].) In this case limits based on stellar energy loss (carried off by Majorons) require [27]  $v_t \leq 2 - 10 \text{ KeV}$ . Implications of the Majoron for neutrino decay and annihilation, cosmology, and neutrinoless double beta decay will be mentioned below.

It is also possible introduce other couplings into the Higgs triplet model which explicitly break lepton number conservation, such as a cubic interaction between  $\tilde{\varphi}_t$  and two Higgs doublets. (This violates  $L$  since  $\tilde{\varphi}_t$  was assigned  $L = 2$  to make (25) invariant). In that case all of the new scalar particles associated with  $\varphi_t$  become massive - i.e. there is no Majoron.

Another mechanism for introducing a Majorana mass is to consider the induced interaction (Fig. 1).

$$L_{eff} = \frac{1}{2} \frac{C}{M} (\bar{\nu}_L \ \bar{e}_L) \bar{\tau} \begin{pmatrix} e_R^c \\ -\nu_R^c \end{pmatrix} \cdot (\varphi^- - \varphi^0) \bar{\tau} \begin{pmatrix} \varphi^0 \\ \varphi^- \end{pmatrix} \quad (26)$$

between two leptons and two Higgs doublets. The Higgs fields in (26) are arranged to transform as an  $SU_2$  triplet, so  $L_{eff}$  is  $SU_2 \times U_1$  invariant; however,  $L_{eff}$  is non-renormalizable, as is evidenced by the dimensional coupling  $C/M$ , where  $M$  is a mass.  $L_{eff}$  cannot therefore be an elementary coupling, but it could be an effective four-particle interaction induced [28] by new physics at some large mass scale  $M$  (just as the four-fermion weak interaction is a nonrenormalizable effective interaction that is really generated by  $W$  and  $Z$  exchange). When  $\varphi^0$  is replaced by its vacuum expectation value, (26) yields an effective Majorana mass  $m \sim Cv^2/M$ , which is naturally small for  $M \gg v$ . For example, if (26) were somehow induced by quantum gravity one would expect  $M \sim 10^{19} \text{ GeV}$  (the Planck scale). Then for  $C \sim 1$  one would have  $m_\nu \sim 10^{-5} \text{ eV}$ .

The most popular realisation of this idea is the seesaw model, [18] in which the underlying physics is the exchange of a very heavy  $SU_2$ -singlet Majorana neutrino  $N_R^0$ , as indicated in Fig. 1. The seesaw model for one family is a special case of the general mass matrix in (15), in which  $m_D$  is a typical Dirac mass (typically assumed to be comparable to  $m_\nu$  or  $m_e$  for the first family) connecting  $\nu_L^0$  to a new  $SU_2$ -singlet  $N_R^0$  and  $m_S \gg m_D$  is a Majorana mass for  $N_R^0$ , presumably comparable to some new (large) physics scale. One typically assumes that  $m_t = 0$  in the seesaw model, i.e. that there is not a Higgs triplet as well. [29] In that case, (15) yields two Majorana mass eigenstates  $n_1$  and  $n_2$  with

$$\begin{aligned} \nu_L^0 &= n_{1L} \cos \theta + n_{2L} \sin \theta \\ N_L^{0c} &= -n_{1L} \sin \theta + n_{2L} \cos \theta \\ \nu_R^{0c} &= -(n_{1R}^c \cos \theta + n_{2R}^c \sin \theta) \\ N_R^0 &= -n_{1R}^c \sin \theta + n_{2R}^c \cos \theta. \end{aligned} \quad (27)$$

The physical masses [30] are

$$\begin{aligned} m_1 &\simeq \frac{m_D^2}{m_S} \ll m_D \\ m_2 &\simeq m_S \end{aligned} \quad (28)$$

and the mixing angle is

$$\tan \theta = \left( \frac{m_1}{m_2} \right)^{1/2} \simeq \frac{m_D}{m_S} \ll 1. \quad (29)$$

Hence, one naturally obtains one very light neutrino, which is mainly the ordinary  $SU_2$  doublet  $(\nu_L^0, \nu_R^{0c})$ , and one very heavy neutrino, which is mainly the singlet  $(N_L^{0c}, N_R^0)$ .

If one does allow  $m_t \neq 0$  (but  $\ll m_S$ ) then there are still two Majorana neutrinos with masses  $|a - \frac{m_D^2}{m_S}|$  and  $m_S$ , respectively, while  $\theta \sim m_D/m_S \ll 1$  still holds. (The minus sign in  $\nu_R^{0c}$  is removed if  $a - \frac{m_D^2}{m_S}$  is positive). In this case, however, one loses the natural explanation of why  $m_1$  is so small, unless  $m_t$  is itself induced by the underlying physics and is of the same order as  $m_D^2/m_S$ .

The seesaw model is easily generalized to  $F$  families. One then has the general  $2F \times 2F$  Majorana mass matrix in (15). Assuming that the eigenvalues of  $m_S$  are all much larger than any of the components of  $m_D$  or  $m_t$  (if it is non-zero) one can calculate the eigenvalues and mixing matrices to leading order in  $m_S^{-1}$ . One finds that there are  $F$  light Majorana neutrinos (consisting of the  $F$  doublets  $(\nu_L^0, \nu_R^{0c})$ , up to corrections of order  $m_D m_S^{-1}$  and  $F$  heavy Majorana neutrinos (consisting of the singlets  $(N_L^{0c}, N_R^0)$ , to  $O(m_D m_S^{-1})$ ). That is, one can write

$$\begin{aligned} \begin{pmatrix} \nu_L^0 \\ N_L^{0c} \end{pmatrix} &= U_L \begin{pmatrix} n_{lL} \\ n_{hL} \end{pmatrix} \\ \begin{pmatrix} \nu_R^{0c} \\ N_R^0 \end{pmatrix} &= U_R \begin{pmatrix} n_{lR}^c \\ n_{hR}^c \end{pmatrix}, \end{aligned} \quad (30)$$

where  $n_{lL}$  and  $n_{hL}$  are  $F$  component vectors of light and heavy Majorana mass eigenstates, respectively, and similarly for  $n_{lR}^c, n_{hR}^c$ . As usual,  $U_L$  and  $U_R$  are  $2F \times 2F$  unitary matrices which diagonalize  $M$  in (15), viz

$$U_L^\dagger \begin{pmatrix} m_t & m_D \\ m_D^T & m_S \end{pmatrix} U_R = m_d = \begin{pmatrix} m_l & 0 \\ 0 & m_h \end{pmatrix}, \quad (31)$$

where  $m_l$  and  $m_h$  are diagonal  $F \times F$  matrices of the  $F$  light and  $F$  heavy eigenvalues, respectively. To leading order in  $m_S^{-1}$  one can write  $U_L^\dagger$  and  $U_R$  in block diagonal form

$$U_L^\dagger = K U_R^T = \begin{pmatrix} K_1 & 0 \\ 0 & K_2 \end{pmatrix} \begin{pmatrix} A^T & -A^T m_D m_S^{-1} \\ D^T m_S^{-1} m_D^\dagger & D^T \end{pmatrix}, \quad (32)$$

where  $A^T$  and  $D^T$  are unitary (to leading order)  $F \times F$  matrices defined by

$$\begin{aligned} m_l &= K_1 A^T (m_t - m_D m_S^{-1} m_D^T) A \\ m_h &= K_2 D^T m_S D \end{aligned} \quad (33)$$

i.e. the mass matrix for the light neutrinos is  $m_l - m_D m_S^{-1} m_D^T$ , which is diagonalized by  $A$ , while that for the heavy neutrinos is  $m_S$ , diagonalized by  $D$ .  $K_1$  and  $K_2$  are diagonal phase

matrices which ensure that  $m_t$  and  $m_b$  are real and positive. We see from (30-33) that indeed there are  $F$  heavy states with masses of  $\mathcal{O}(m_S)$ , and in the simplest case  $m_t = 0$  there are  $F$  states which are naturally very light ( $\mathcal{O}(m_D^2 m_S^{-1})$ ). (For  $m_t \neq 0$  one must separately assume  $m_t$  is small). Furthermore, the mixing between the light and heavy sectors is very small (of  $\mathcal{O}(m_D m_S^{-1})$ ), while the matrices  $A$  and  $D$ , which describe mixings within the two sectors, are in general arbitrary.

There are several classes of seesaw models [18], depending on the scale of  $m_S$ . In simple grand unified models one assumes that the scale is a typical GUT unification scale of around  $10^{14}$  GeV. In many such models (e.g.  $SO_{10}$ ) one has that the neutrino Dirac mass matrix  $m_D$  is the same as  $m_u/k$  where  $m_u$  is the  $u$ -quark mass matrix and  $k \simeq 4.7$  represents the running of the Yukawa couplings between the GUT scale and low energies. If one makes the somewhat ad-hoc assumption that the matrix  $m_S$  is just  $M_X I$ , where  $M_X \sim 10^{14}$  GeV is the unification scale and  $I$  is the identity matrix, one has (for  $m_t = 0$ ) the light eigenvalues

$$m_{\nu_i} \sim \frac{m_{u_i}^2}{M_X k^2} \quad (34)$$

$\sim 10^{-11}$  eV,  $10^{-6}$  eV,  $10^{-3}$  eV, i.e. the neutrino masses are naturally expected to be extremely tiny, and to scale like the squares of the  $u$ ,  $c$ , and  $t$  quark masses. (Equation (34) was computed for  $m_{top} \sim 50$  GeV). Several caveats are in order: the assumption of  $m_S \sim M_X I$  was quite arbitrary. One could easily imagine that the eigenvalues of  $m_S$  are smaller than  $M_X$  due to small Yukawa coupling couplings (increasing  $m_{\nu_i}$ ). Also, they need not be the same. For example, if the  $m_S$  eigenvalues followed the same family hierarchy as the ordinary fermions (i.e.  $m_S \propto m_{u_i}$ ) then one would have  $m_{\nu_i}$  scaling as  $m_{u_i}$  rather than  $m_{u_i}^2$ . (A similar linear hierarchy ensues in some variant GUTs in which  $m_S$  is zero at tree level but is generated at higher orders [31]. Of course, more complicated patterns for  $m_S$  and  $m_D$  (in (33)) are also possible. Furthermore, in many cases loop corrections to the (GUT) Higgs potential may induce [29] VEV's for Higgs representations that can yield a non-zero triplet terms  $m_t$  in (31). These are most likely to affect the smallest masses (e.g.  $m_{\nu_e}$ ). Equation (34) should therefore be regarded only as a typical order of magnitude.

If one does assume that  $m_S = M_X I$ , however, then  $m_u^2/M_X$  is diagonalized by the same transformations that diagonalize  $m_u$ . Since one also has equal electron and  $d$ -quark mass matrices (i.e.  $m_e = m_d/k$ ) in most simple GUTs the final result is that flavor mixing in the lepton sector (analogous to (7)) is described by the same mixing matrix as the CKM quark mixing matrix. This result continues to hold [19] approximately for a far wider class of  $m_S$  than does the simple mass prediction in (34).

Lower mass scales for  $m_S$  imply larger values for the light neutrino masses (and generally less predictive power for  $m_D$ ). Several authors [20] have suggested that the heavy Majorana scale could be the intermediate range  $10^8 - 10^{12}$  GeV associated with invisible axions. For  $m_D \sim m_e$ , and  $m_S \sim 10^9$  GeV, for example, one obtains the values  $\sim 10^{-7}$  eV,  $10^{-2}$  eV,  $10$  eV for  $m_{\nu_e}$ ,  $m_{\nu_\mu}$ ,  $m_{\nu_\tau}$ , respectively.

If  $m_S$  is in the several TeV range (as expected in some left-right symmetric [32]  $SU_{2L} \times SU_{2R} \times U_1$ , models [21], for example) one typically expects (for  $m_D \sim m_e, m_S \propto I$ )  $m_{\nu_e}, m_{\nu_\mu}, m_{\nu_\tau}$  to have relatively large values  $10^{-1}$  eV,  $10$  KeV, and  $1$  MeV, respectively. As we will see, such models run into severe cosmological difficulties unless the mass hierarchy is somehow modified or a fast decay mechanism is found for the  $\nu_\mu$  and  $\nu_\tau$ . Of course,

one could also have  $m_S$  much smaller than the  $SU_{2L} \times SU_{2R} \times U_1$  scale (e.g. in the  $10$  GeV  $- 100$  GeV range), with corresponding larger masses for the light neutrinos. Similar statements apply to models with extra  $Z$  bosons in the  $100$  GeV  $- 10$  TeV range, which usually also have heavy Majorana neutrinos.

Finally, one can consider light seesaw models, in which typically  $m_S \ll 1$  GeV. Such models are very artificial and abandon the principal advantages of the seesaw, because both  $m_D$  and  $m_S$  must be taken unnaturally small to obtain an acceptable  $\nu_e$  mass. Their only virtue is that they yield strongly suppressed neutrinoless double beta decay rates, even though the neutrinos are Majorana.

Seesaw models were first introduced in GUT type models in which lepton number is explicitly violated by the gauge interactions. One can also consider non-gauge seesaw models [33] in which lepton number is spontaneously broken by the VEV of the Higgs field which generates  $m_S$ . Such models imply the existence of a massless Goldstone boson, the singlet-Majoron. [34] Unlike the triplet-Majoron in the Gelmini-Roncadelli model,[8] which can couple strongly to the ordinary neutrinos (coupling  $\sim h_t$ ), the singlet-Majoron effectively decouples from ordinary particles. That is, it couples strongly to the heavy neutrino, with a coupling of order  $m_D m_S^{-1}$  to off-diagonal  $n_{lR}$  vertices, and with strength  $(m_d m_S^{-1})^2$  to light neutrinos.

It is difficult to implement the seesaw model in most superstring inspired models, because there is no Higgs field available to generate a large  $m_S$ . It has been suggested [35] that  $m_S$  could be generated by a higher order effective operator, but such model may run into serious cosmological problems [36].

There have also been variant seesaw models constructed [37] in which the light neutrinos occur in degenerate pairs which can be combined from Dirac neutrinos with a conserved  $L$ .

Finally, I mention the charged Higgs models [23], in which small Majorana masses are generated by loop diagrams involving new charged Higgs bosons with explicit  $L$ -violating couplings (Fig. 1). Viable versions often lead to pseudo-Dirac neutrinos. The approximately conserved lepton number is typically  $L_e - L_\mu + L_\tau$ , for example, rather than  $L$ . The actual mass scale depends on unknown Yukawa couplings and masses.

## 4 Experimental Constraints

There are a number of excellent reviews [1] of the experimental status of neutrino mass. My major purpose in this section is to comment on the implications of the various theoretical models for the different types of experiments.

### 4.1 Kinematic Tests

Direct kinematic limits on the masses of the  $\nu_e$ ,  $\nu_\mu$ , and  $\nu_\tau$  are given in Table 2. The ITEP group [38] has long claimed evidence for a non-zero  $\nu_e$  mass in the  $20$  eV range from tritium  $\beta$  decay, but this has not been confirmed by other groups, and in fact the Zurich-SIN measurement is on the verge of conflicting with the ITEP result. In addition the neutrinos from supernova 1987A observed by the Kamiokande [45] and IMB [46] experiments place upper limits in the  $20$  eV range on the  $\nu_e$  mass (otherwise the arrival times of the detected

Table 2: Kinematic limits/values on neutrino masses.

$17 \text{ eV} < m_{\nu_e} < 40 \text{ eV}$	ITEP [38]
$m_{\nu_e} < 18 \text{ eV}$	Zurich [39]
$m_{\nu_e} < 27 \text{ eV}$	LANL [40]
$m_{\nu_e} < 32 \text{ eV}$	INS-Tokyo [41]
$m_{\nu_e} < O(20 \text{ eV})$	SN1987A [42]
$m_{\nu_\mu} < 0.25 \text{ MeV}$	SIN [43]
$m_{\nu_\tau} < 50 \text{ MeV}$	ARGUS [44]

neutrinos would be spread out more than is observed), but it is hard to make this limit precise because it depends on the details of the neutrino emission [42].

A 20 eV neutrino mass is just in the range that would be most interesting cosmologically, so clearly it is essentially to resolve the situation. Hopefully, the current and next generation of tritium  $\beta$  decay experiments will be sensitive down to a few eV, but it is doubtful whether experiments of this type will ever be able to probe to much lower scales. As can be seen in Table 1, none of the models really predict  $m_{\nu_e}$  in the 20 eV range (the  $SU_{2L} \times SU_{2R} \times U_1$  models come closest), but most can accommodate masses in this range by fine-tuning parameters.

As can be seen in Table 2, the direct kinematic limits on  $m_{\nu_\mu}$  (from  $\pi_{\mu 2}$  decay) and on  $m_{\nu_\tau}$  (from  $\tau \rightarrow \nu_\tau + 5\pi$ ) are relatively weak. The experiments are extremely difficult (the mass scales being probed are very much smaller than the energies released in the decays), so it is unlikely that these measurements will improve by much more than a factor of two.

## 4.2 Heavy Neutrinos

There are many limits [1,47] on possible small admixtures of heavy neutrino states in the  $\nu_e$  or  $\nu_\mu$ , including universality tests in nuclear  $\beta$  decay, searches for secondary peaks or distortions of the lepton spectra in  $\beta$ ,  $\pi$ , and  $K$  decay, searches for the decay products of heavy neutrinos (e.g.  $\nu_h \rightarrow \nu_e e^+ e^-$ ) produced in beam dumps,  $e^+ e^-$  annihilation, or neutrino scattering. The limits on the mixing angle  $U_{ai}$ ,  $a = e$  or  $\mu$ , where  $\nu_a^0 = \sum_i U_{ai} \nu_i$  are quite impressive, especially for  $m_i$  in the range 10 MeV – 10 GeV, where they are comparable to the expectations in (29) of a seesaw model with  $m_1 \sim 10 \text{ eV}$  and  $m_2 = m_i$ . The lower part of this range corresponds to the masses expected in the “light-seesaw” model (Table 1), while the 1 GeV – 1 TeV range is consistent with  $SU_{2L} \times SU_{2R} \times U_1$  models. [21]

As has already been mentioned, heavy neutrinos in the GeV – TeV range are likely to give too large  $m_{\nu_e}$  and  $m_{\nu_\mu}$ , unless the typical seesaw hierarchy  $m_{\nu_i} \propto m_i^n$  or  $m_{\nu_i}^n$ ,  $n = 1$  or 2, for the light neutrinos is avoided or new physics is invoked to ensure fast decays or annihilations for the  $\nu_\mu$  and  $\nu_\tau$ . On the other hand, if such new physics is present some of the limits (those based on decays) may no longer be valid, because in many cases the heavy neutrinos will decay rapidly into unobservable channels (e.g.  $\nu_h \rightarrow \nu_l + \text{Majoron}$ ) before reaching the detector.

## 4.3 Neutrino Oscillations

Neutrino oscillations are a beautiful example of a common quantum phenomenon: viz that if one starts at time  $t = 0$  in a state that is not an energy eigenstate [48] then at later times it can oscillate into another (orthogonal) state. For example, suppose that the  $\nu_e^0$  and a second neutrino  $\nu_a^0$  (e.g.  $\nu_a^0 = \nu_\mu^0$  or  $\nu_\tau^0$ ) are mixtures of two mass eigenstates  $\nu_1$  and  $\nu_2$  with mixing angle  $\theta$ . If at time  $t = 0$  the weak eigenstate  $\nu_e^0 = \cos\theta \nu_1 + \sin\theta \nu_2$  is produced (e.g. in the process  $\pi^+ \rightarrow \pi^0 e^+ \nu_e^0$ ) then at time  $t$  it will have evolved into the state

$$\begin{aligned} \nu_e^0(t) &= \cos\theta \nu_1 e^{-iE_1 t} + \sin\theta \nu_2 e^{-iE_2 t} \\ &\simeq \cos\theta \nu_1 e^{-\frac{-im_1^2 t}{2p}} + \sin\theta \nu_2 e^{-\frac{-im_2^2 t}{2p}}. \end{aligned} \quad (35)$$

In the second form I have assumed relativistic neutrinos  $E_i = \sqrt{p^2 + m_i^2} \sim p + m_i^2/2p$  with definite momentum [49]  $p \gg m_i$ , and have neglected an irrelevant overall phase  $\exp(-ipt)$ . The state  $\nu_e^0(t)$  has a non-trivial overlap with  $\nu_a^0$ . After traveling a distance  $L \sim t$ , there will be a probability

$$\begin{aligned} P(\nu_e \rightarrow \nu_a) &= |\langle \nu_a^0 | \nu_e^0(t) \rangle|^2 \\ &= \sin^2 2\theta \sin^2 \left( \frac{\Delta m^2 L}{4p} \right) \\ &= \sin^2 2\theta \sin^2 \left( \frac{1.27 \Delta m^2 (\text{eV}^2) L (m)}{p (\text{MeV})} \right) \end{aligned} \quad (36)$$

that the state will have evolved into  $\nu_a^0$  (as can be observed in the process  $\nu_a N \rightarrow e_a N'$ , for example), and a probability  $P(\nu_e \rightarrow \nu_e) = 1 - P(\nu_e \rightarrow \nu_a)$  that the state will remain a  $\nu_e^0$ . In (36),  $\Delta m^2 = m_1^2 - m_2^2$ , and the last form is valid for  $\Delta m^2$  in  $\text{eV}^2$ ,  $L$  in  $m$ , and  $p$  in  $\text{MeV}$ . It is seen that the  $\nu_e \rightarrow \nu_a$  probability depends on both the mixing angle  $\theta$  and on  $\Delta m^2 L/p$ . For moderate values of the latter quantity the probability oscillates as a function of  $L$  and  $p$ , while for very large values the oscillations are averaged by a finite-sized detector or non-monochromatic source, (the second factor in (36) averages to 1/2). It is easy to generalize [1] (36) to the case that the initial neutrino is a mixture of more than two mass eigenstates,  $\nu_a^0 = \sum_i U_{ai} \nu_i$ . One obtains

$$P(\nu_e \rightarrow \nu_a) = \sum_i |U_{ei} U_{ai}^*|^2 + \text{Re} \sum_{i \neq j} U_{ei} U_{ai}^* U_{ej}^* U_{aj} e^{-\frac{i(m_i^2 - m_j^2)L}{2p}} \quad (37)$$

Neutrino oscillations can be searched for in (a) appearance experiments, in which one looks for the interactions of  $\nu_a$  in a detector, and (b) disappearance experiments, in which one looks for a reduced  $\nu_e$  flux. In both cases one can compare the observed counting rate with the expectation from known backgrounds (appearance) or from the expected flux (disappearance) as determined, for example, by measuring the electron spectrum from  $n \rightarrow pe^- \bar{\nu}_e$  in reactor  $\bar{\nu}_e$  oscillation experiments. A much cleaner technique is to search for actual oscillations in the appearance or disappearance probabilities as a function of  $L$  or  $p$ , such as by using two detectors at different distances from the source.

There are many limits on neutrino oscillations from accelerator experiments [1] (e.g. counter and emulsion experiments and beam dumps, searching for  $\nu_\mu \rightarrow \nu_e$ ,  $\nu_\mu \rightarrow \nu_\tau$  and




Figure 2: 90% *c.l.* limits on neutrino oscillations, from [55]. (a)  $\nu_\mu \rightarrow \nu_e$  (BNL, CHARM, BEBC, Los Alamos, PS-191),  $\nu_e \rightarrow \nu_\tau$  (E531), and  $\bar{\nu}_e \rightarrow \nu_X$  (Bugey, Gösigen). (b)  $\nu_\mu \rightarrow \nu_\tau$ ,  $\nu_X$ ,  $\nu_\mu$ . The Bugey [52] and PS-191 [54] regions are allowed by positive results. The other contours are exclusion plots (the regions to the right are excluded).

$\nu_e \rightarrow \nu_\tau$ , as well as  $\nu_\mu$  disappearance), and reactors [1] ( $\bar{\nu}_e$  disappearance), as well as on the oscillations of  $\nu_\mu$  produced in cosmic ray interactions in the atmosphere [50]. (Implications for the Solar neutrino problem are discussed below). The results of these searches [51] are summarized in Fig. 2. The Bugey reactor experiment [52] reports a positive signal for  $\bar{\nu}_e$  disappearance, but their results are contradicted by the Gösigen experiment [53]. Similarly, the CERN PS-191 counter experiment [54] reports an excess of  $\nu_e$  events in a  $\nu_\mu$  beam, but their signal is in conflict with several other  $\nu_\mu \rightarrow \nu_e$  experiments [55]. Clearly, a clarification of the situation is essential.

From Fig. 2 it is clear that there are stringent limits on neutrino mixings for  $|\Delta m^2|$  above  $\simeq 1 \text{ eV}^2$ . This should be contrasted with the suggested value  $m_{\nu_e} \sim 17 - 40 \text{ eV}$  by the ITEP experiment [38]. If the ITEP result is correct then most likely the  $\nu_e$  could not have any significant mixing with other neutrinos (the alternative possibility, that the  $\nu_e$  is almost degenerate with another neutrino flavor so that  $|\Delta m^2| \ll m_{\nu_e}^2$ , seems rather contrived but cannot be excluded). A comparison of Fig. 2 with the expectations of various models (Table 1) suggests that  $\nu_\mu \rightarrow \nu_\tau$  oscillations may be the most optimistic possibility for the future. Many of the seesaw-type models predict that the lepton mixing angles are roughly correlated with the corresponding quark mixing angles. This would suggest  $\sin^2 2\theta \sim 10^{-4}, 10^{-2}, 10^{-1}$  for  $\nu_e \leftrightarrow \nu_\tau$ ,  $\nu_\mu \leftrightarrow \nu_\tau$ , and  $\nu_e \leftrightarrow \nu_\mu$ , respectively.

Oscillations between ordinary  $SU_2$  doublet neutrinos ( $\nu_e^0, \nu_\mu^0, \nu_\tau^0$ , and possible fourth

family  $\nu$ 's), known as first class or flavor oscillations, occur for pure Dirac and pure Majorana neutrinos, as well as in the multi-family seesaw models. In models involving both Dirac and Majorana mass terms of comparable magnitude, however, there can be additional light neutrinos, and the mass eigenstates can have significant admixtures of both  $SU_2$  doublets and singlets. In this case second class oscillations [16] can occur, in which the ordinary neutrinos oscillate into  $SU_2$  singlets with negligible interactions. These "sterile" neutrinos are essentially undetectable, so second class oscillations can be observed [56] only in disappearance experiments. Of course, first and second class oscillations can occur simultaneously. For three families, for example, there could be oscillations between six Majorana neutrinos (3 doublets and 3 singlets).

Yet another possibility [57] are models in which the ordinary neutrinos have small mixings with heavy neutrinos. In that case the neutrinos actually produced in weak processes are the projections of the weak eigenstates onto the subspace of light or massless neutrinos. It can easily occur that the projections of the  $\nu_e^0$  and  $\nu_\mu^0$ , for example, are not orthogonal. The result is that a  $\nu_e^0$  could produce an  $e^-$  in a subsequent reaction. Such a non-orthogonality would mimic the effects of oscillation appearance experiments, even if the masses of the light neutrinos are zero or negligible.

#### 4.4 Cosmology

There are many limits on neutrino mass and decays from cosmology [58]. Ordinary light or massless neutrinos would have been produced by such weak processes as  $e^+e^- \leftrightarrow \nu\nu^c$  in the early universe. As long as the weak reaction rate [59]

$$\Gamma_{weak} \sim (\sigma v) n_T \sim G_F^2 T^5 \quad (38)$$

( $(\sigma v) \sim G_F^2 T^2$  is the thermally averaged cross section times relative velocity, and  $n_T \sim T^3$  is the density of target particles, where  $T$  is the temperature) was large compared to the expansion rate  $H \sim T^2/m_p$  (where  $m_p = G_N^{-1/2} \sim 10^{19} \text{ GeV}$  is the Planck scale) the number of neutrinos stayed in equilibrium. However, as soon as  $T$  dropped below the temperature  $T_D \sim (G_F^2 m_p)^{-1/3} \simeq 3 \text{ MeV}$  for which  $\Gamma_{weak} \sim H$ , the weak rate became negligible and the neutrinos decoupled, i.e. effectively stopped interacting. According to most models these neutrinos should remain in the present universe, undisturbed from the first second of the big bang except for a redshifting of their momenta by the expansion of the universe. They are analogous to the  $2.7^\circ\text{K}$  microwave radiation (which decoupled later). If the neutrino masses are much less than  $1 \text{ eV}$  there should be  $\simeq 50$  neutrinos/ $\text{cm}^3$  of each type ( $\nu_{eL}, \nu_{eR}^c$ , etc) with momenta characterized by a thermal spectrum with temperature  $\simeq 1.9^\circ\text{K}$  ( $10^{-4} \text{ eV}$ ). Despite the large number of neutrinos ( $\simeq 10^{10}$  per baryon) they are essentially impossible to detect [60] - [62] because their cross section  $\sim G_F^2 E_\nu^2 \sim 10^{-62} \text{ cm}^2$  is so low. [63]

The major cosmological bound is based on the energy density of the present universe. There are predicted to be so many relic neutrinos that even for a small mass in the  $10 \text{ eV}$  range they would be important. Limits on the energy density imply

$$\sum_i m_{\nu_i} < 40 \text{ eV} \quad (39)$$

where the sum extends over the light, stable (at least compared to the age of the universe) doublet neutrinos. Conversely, a neutrino with mass in this range would dominate the energy density and could account for the dark (missing) matter in galaxies and clusters [64]. In particular, for the ITEP value  $m_{\nu_e} \sim (17 - 40) \text{ eV}$ , the  $\nu_e$  would be an ideal candidate for the dark matter, but one would probably then have to find a mechanism to explain why the  $\nu_e$  is the heaviest neutrino.

• Similarly, the energy density associated with light or massless neutrinos for  $T \sim T_D$  affects nucleosynthesis and leads to the limit  $N_\nu \leq 4$  on the number of neutrino flavors with  $m_\nu \leq 1 \text{ MeV}$ . [65]

There are also a variety of constraints on unstable neutrinos. An ordinary doublet mass eigenstate neutrino  $\nu_2$  (with  $m_{\nu_2} > m_{\nu_1}$ ) is expected to decay into

$$\begin{aligned} \nu_2 &\rightarrow \nu_1 \gamma, & (m_{\nu_2} < 2m_e) \\ \nu_2 &\rightarrow \nu_1 e^+ e^-, & (2m_e < m_{\nu_2} < m_\mu + m_e). \end{aligned} \quad (40)$$

The first decay occurs at one loop, while the second occurs at tree level. Both decays are very slow for small  $m_{\nu_2}$  and the decay products are detectable. There are a large variety of cosmological and astrophysical constraints [66] on  $m_{\nu_2}$  and  $\tau_{\nu_2}$  from the present energy density, the growth of galaxies, the distortion of the  $2.7^\circ\text{K}$  background radiation, the non-observation of the decay photons, supernovae, and nucleosynthesis and breakup. For reasonable mixing angles these limits exclude the range  $40 \text{ eV} - (20 - 40) \text{ MeV}$  for ordinary neutrinos [67] decaying according to (40). Combined with laboratory limits this implies [66,68] that the  $\nu_\mu$  and  $\nu_\tau$  (i.e. their dominant mass eigenstate components) should be lighter than  $40 \text{ eV}$ . In particular, this poses serious problems for the  $\text{TeV}$  scale seesaw model.

Most of the cosmological limits can be evaded if new physics is invoked to allow fast and invisible (except for the relativistic energy of the decay products) decays or annihilation for the heavy neutrinos. One possibility is the decay  $\nu_2 \rightarrow 3\nu_1$ . However, the rate for this mode from off-diagonal  $Z$  couplings [69] is too slow, while models in which the couplings of a Higgs triplet [70] (present in  $SU_{2L} \times SU_{2R} \times U_1$ ) are arranged to allow a fast decay generally run into problems [71] with  $\mu \rightarrow 3e$ .

More promising are models in which  $\nu_2 \rightarrow \nu_1 G$ , where  $G$  is a Goldstone boson [72]-[74] associated with a spontaneously broken global symmetry. Likely examples are the case that  $G$  is a familon [72] (a Goldstone boson associated with a broken family symmetry) or a triplet-Majoron [73]. In fact, for triplet-Majorons one expects the annihilation process  $\nu\nu \rightarrow MM$  (which begins when  $T$  drops below  $v_t$ ) to have removed any relic neutrinos from the present universe [8]. In familon models some care must be taken to avoid unacceptably large flavor changing neutral current effects. The decay  $\nu_2 \rightarrow \nu_1 M$  is too slow in the simpler versions of the singlet-Majoron model [74] to avoid cosmological problems.

The role of spontaneous  $L$  violation in Majoron models in reducing possible initial large lepton asymmetries to cosmologically interesting values at the time of nucleosynthesis is discussed in [75].

#### 4.5 Double Beta Decay

Another important source of information on the  $\nu_e$  mass (if it is Majorana) is neutrinoless double beta decay ( $\beta\beta_{0\nu}$ ).




Figure 3: Diagrams for two neutrino ( $\beta\beta_{2\nu}$ ) and neutrinoless ( $\beta\beta_{0\nu}$ ) double beta decay.

First consider the lepton-number conserving two-neutrino ( $\beta\beta_{2\nu}$ ) process  $(Z, N) \rightarrow (Z + 2, N - 2)e^-e^- \nu_e \bar{\nu}_e$ , which can be thought of as two ordinary beta decays occurring in the same nucleus (Fig. 3). In the context of neutrino mass this process is mainly of interest as a calibration of the calculated nuclear matrix elements that are needed for the neutrinoless case. There has long been a two order of magnitude discrepancy between the predicted rates [76], e.g. for  $^{130}\text{Te} \rightarrow ^{130}\text{Xe}$ , and indirect measurements by geochemical techniques [77]. Within the last year, however, this discrepancy has gone away. The geochemical measurements were confirmed by the first laboratory observation of double beta decay (at Irvine [78].) In addition, several groups [76] have found that previously neglected ground state correlation effects could suppress the matrix element by the required order of magnitude. Furthermore, there is no analogous uncertainty in the  $\beta\beta_{0\nu}$  case.

The neutrinoless double beta decay process  $(Z, N) \rightarrow (Z + 2, N - 2)e^-e^-$ , which violates lepton number by two units, can proceed through the second diagram [79] in Fig. 3. In the absence of mixing the quantity  $\langle m_{\nu_e} \rangle$ , the effective Majorana neutrino mass, is

$$\langle m_{\nu_e} \rangle = \begin{cases} 0, & \text{Dirac neutrino} \\ m_{\nu_e}, & \text{unmixed Majorana neutrino} \end{cases} \quad (41)$$

Although the matrix element is proportional to  $\langle m_{\nu_e} \rangle$ , which is necessarily very small,  $\beta\beta_{0\nu}$  has an enormous advantage in phase space over  $\beta\beta_{2\nu}$ , and could be observable for  $\langle m_{\nu_e} \rangle$  in the  $\text{eV}$  range. Of course, the sum of the electron energies should be a sharp peak in  $\beta\beta_{0\nu}$  (and a continuum for  $\beta\beta_{2\nu}$ ), so the principal difficulty is controlling the background. [80,1] Currently, the most sensitive experiments are for  $^{76}\text{Ge} \rightarrow ^{76}\text{Se} e^-e^-$ . No evidence for  $\beta\beta_{0\nu}$  has been observed, [81] and the lower limit on the lifetime is [55]  $\tau_{1/2} > 9 \times 10^{23} \text{ yr}$  (68% c.l.). According to several calculations of the nuclear matrix elements [82] this implies  $\langle m_{\nu_e} \rangle \leq 1 \text{ eV}$ . However, a recent estimate by Engel et al. [83] yielded a much weaker limit  $\langle m_{\nu_e} \rangle \leq 11 \text{ eV}$ , so caution is advisable.

Even the largest value  $\langle m_{\nu_e} \rangle \leq 11 \text{ eV}$  is smaller than the range  $m_{\nu_e} \sim (17 - 40) \text{ eV}$  suggested by the ITEP experiment. If the latter is correct the simplest possibility is that

the  $\nu_e$  is Dirac. Another possibility [84] is that the  $\nu_e^0$  is a mixture of Majorana mass eigenstate neutrinos. Then,  $\langle m_{\nu_e} \rangle$  becomes

$$\langle m_{\nu_e} \rangle = \sum_i m_i U_{Lei}^2 \xi_i F(m_i, A), \quad (42)$$

where  $m_i \geq 0$  is the physical mass of the  $i^{\text{th}}$  mass eigenstate,  $U_{Lei}$  is the mixing matrix element ( $\nu_{eL}^0 = \sum U_{Lei} \nu_{iL}$ ) and  $\xi_i = \pm 1$  is the CP parity of  $\nu_{iL}$ .  $\xi_i$  is just  $K_{ii}$  in (13), and a negative value  $\xi_i = -1$  means simply that the eigenvalue of  $M$  in (10) was negative before choosing  $K$  to redefine  $\nu_R^c$ . In (42),  $F(m_i, A)$  is a nucleus dependent propagator correction, [85] defined by

$$F(m_i, A) = \frac{\langle e^{-m_i r} / r \rangle}{\langle 1/r \rangle}. \quad (43)$$

It is  $\sim 1$  for  $m_i \ll 10 \text{ MeV}$ . For  $m_i \gg 10 \text{ MeV}$ ,  $F(m_i, A) \ll 1$  (it falls as  $m_i^{-2}$ ) and allows the possibility [86] of  $A$  dependence of  $\langle m_{\nu_e} \rangle$ .

Because of the possibility of negative contributions to  $\langle m_{\nu_e} \rangle$  it is conceivable that there are cancellations so that  $\langle m_{\nu_e} \rangle$  is much smaller than the mass of the dominant Majorana component of  $\nu_e$  (e.g.  $m_1 \sim (17-40) \text{ eV}$ ). Such a cancellation is actually not so contrived as it might first appear. If all of the  $m_i$  are small enough that  $F(m_i, A) = 1$  then from (12)  $\langle m_{\nu_e} \rangle$  is just the  $M_{ee}$  component of the original Majorana mass matrix in (10). As we have seen,  $M_{ee}$  must be generated by a Higgs triplet and vanishes in many models. In fact, the light seesaw model of Table 1 automatically leads to  $\langle m_{\nu_e} \rangle = 0$  for sufficiently small  $m_i$ . For two neutrinos, for example,  $\langle m_{\nu_e} \rangle = m_1 \cos^2 \theta - m_2 \sin^2 \theta$ , which vanishes by (28) and (29). However, the light seesaw model was devised just in order to give  $\langle m_{\nu_e} \rangle = 0$ . For seesaw models with more natural scales  $m_2 \gg 10 \text{ MeV}$  one has that  $F(m_i, A) \ll 1$  and  $U_{e1} \sim 1$ , so that  $\langle m_{\nu_e} \rangle \sim m_{\nu_e}$ . In most Majorana models, therefore, one expects  $\langle m_{\nu_e} \rangle \sim m_{\nu_e}$  unless fine-tuned deviations from the seesaw formula are invoked.

Whether or not the cancellation of the terms in (42) is natural, one can consider whether it is phenomenologically viable. For two neutrinos, for example, the conditions  $m_1 \sim 20 \text{ eV}$ , and  $\langle m_{\nu_e} \rangle \ll m_1$  imply

$$\tan^2 \theta = \frac{m_1}{m_2 F(m_2, A)}, \quad (44)$$

where  $m_1 \leq m_2$ ,  $\tan^2 \theta \leq 1$  since the ITEP experiment presumably measures the dominant component of  $\nu_e$ . However, the reactor oscillation limits in Fig. 2 allow only two possibilities. One is that  $m_1 \simeq m_2$ ,  $0 \sim 45^\circ$ . In that case  $\nu_1$  and  $\nu_2$  can be combined to form a Dirac neutrino (or pseudo-Dirac if the degeneracy is not exact), possibly with a non-canonical lepton number (such as  $L_e - L_\mu + L_\tau$ ) conserved. Alternatively, one can have  $m_2 \geq 450 \text{ eV}$ . However, the various laboratory and cosmological limits exclude [22] almost all values of  $m_2$  except for small windows around  $40 \text{ MeV}$  and  $2 \text{ GeV}$ . Hence, if the ITEP results turn out to be correct they would almost certainly imply either (a) the  $\nu_e$  is Dirac, or (b) there is new physics (such as a Majoron) that evades the cosmological bounds.

There are additional contributions to neutrinoless double beta decay in  $SU_{2L} \times SU_{2R} \times U_1$  models [87]. Typically, such models contain additional charged  $W_R^\pm$  bosons which couple to right-handed currents  $\bar{e}_R \gamma^\mu N_R$ , where  $N_R$  is a heavy Majorana neutrino. The exchange

of a  $N_R$  (rather than a  $\nu_L$  in Fig. 3) yields a new contribution  $M_N F(M_N, A) (M_{WL}/M_{WR})^4$  to  $\langle m_{\nu_e} \rangle$ , which sets non-trivial constraints [21] on  $M_N$  and  $M_{WR}$ . Furthermore, mixed contributions involving one ordinary left-handed current  $\bar{e}_L \gamma^\mu \nu_L$  and one right-handed current  $\bar{e}_R \gamma^\mu N_R$  can yield contributions to  $0^+ \rightarrow 2^+$  decay amplitudes that are not directly proportional to a neutrino mass [88] However, the relevant amplitudes are of order [1,87]

$$\left( \frac{M_{WL}}{M_{WR}} \right)^2 \theta, \quad \zeta \theta, \quad (45)$$

where  $\theta$  is a light-heavy neutrino mixing angle and  $\zeta$  is the  $W_L - W_R$  mixing angle. One typically expects  $(M_{WL}/M_{WR})^2$  and  $\zeta$  to be less than  $10^{-3}$ . Since we expect  $\theta \sim m_D/M_{WR} \leq 10^{-4} - 10^{-5}$  in a typical  $T \text{ eV}$ -seesaw, the expected values for the quantities in (45) are smaller than the experimental limits (of  $\sim 10^{-6}$ ).

One typically has  $\langle m_{\nu_e} \rangle \ll m_{\nu_e}$  for the charged Higgs models [23] because the antisymmetry of the relevant Yukawa coupling forces  $M_{ee}$  to vanish.

## 4.6 The Solar Neutrino Problem

For some years the event rate in the  $^{37}\text{Cl} \rightarrow ^{37}\text{Ar}$  Solar neutrino experiment [89] ( $2.0 \pm 0.3 \text{ SNU}$  [90]) has been considerably below the prediction [91]  $5.8 \pm 2.2 \text{ SNU}$  of the standard Solar model. The discrepancy has recently been confirmed by the Kamiokande group which reports [92] an upper limit on the  $\nu_e$  flux (from  $\nu_e e$  elastic scattering) that is less than half the expected event rate. One explanation for the discrepancy is the existence of vacuum oscillations of the  $\nu_e$  into other neutrinos. These could be important for neutrino mass-squared differences [93]  $\Delta m^2 \equiv m_1^2 - m_2^2$  as small as  $\Delta m^2 \sim (10^{-11} - 10^{-10}) \text{ eV}^2$ , but only if the mixing angles are large.

Another possibility [94] is that the  $\nu_e$  is a Dirac particle with a magnetic moment in the range  $\mu_{\nu_e} \sim (0.6 - 10) \times 10^{-10} \mu_B$ . The  $\nu_e$  spin could then precess in the Solar magnetic field into a sterile right-handed  $\nu_e$ , thus reducing the observed flux by a factor  $\simeq 2$ . The necessary value of  $\mu_{\nu_e}$  is barely consistent with laboratory limits [95] but is probably excluded by astrophysical constraints from nucleosynthesis and stellar cooling [96] (Table 3). The worst objection, however, is that the necessary  $\mu_{\nu_e}$  is unnaturally high. In the standard model with a Dirac mass one expects [97]

$$\mu_\nu = \frac{3G_F m_\nu m_e}{4\pi^2 \sqrt{2}} \mu_B \sim 3 \times 10^{-19} \left( \frac{m_\nu}{1 \text{ eV}} \right) \mu_B \quad (46)$$

which is many orders of magnitude too small. Non-standard models [98] can yield larger  $\mu_\nu$ , but to obtain a sufficiently large value appears highly contrived.

Other canonical explanations involve non-standard Solar models. The existing experiments are mainly sensitive to the relatively high energy (from  $0.81 \text{ MeV}$  up to  $14 \text{ MeV}$ ) neutrinos from  $^8\text{B}$  decay. The flux of these  $^8\text{B}$  neutrinos depends very sensitively on the temperature of the Solar core and could be changed significantly by modifications of the standard Solar model. Recently, there has been much attention to the possibility that weakly interacting massive particles (WIMPs), which could form the dark matter, could carry energy out of the Solar core and lower the central temperature slightly. [58] Less exotic modifications of the standard model are also possible.

Table 3: Limits on the neutrino magnetic moments. A value  $\mu_{\nu_e} \sim (0.6 - 10) \times 10^{-10} \mu_B$  would be needed to resolve the Solar  $\nu$  problem.

laboratory [95]	$\mu_{\bar{\nu}_e} < 1.5 \times 10^{-10} \mu_B$
	$\mu_{\nu_\mu} < 9.5 \times 10^{-10} \mu_B$
Stellar cooling [96] ( $\gamma \rightarrow \nu \bar{\nu}$ )	$\mu_\nu < 0.8 \times 10^{-11} \mu_B$
Nucleosynthesis [96] ( $\nu_R$ produced by spin precession)	$\mu_\nu < 0.5 \times 10^{-10} \mu_B$
Standard model [97] (Dirac mass)	$\mu_\nu \sim 3 \times 10^{-19} \left( \frac{m_\nu}{1 \text{ eV}} \right) \mu_B$

A  ${}^{71}\text{Ga} \rightarrow {}^{71}\text{Ge}$  experiment could distinguish the nonstandard Solar model from the first two possibilities. Most of the expected  ${}^{71}\text{Ga}$  event rate is from the low energy  $pp$  neutrinos, the flux of which can be inferred from the over-all Solar luminosity and is relatively insensitive to the temperature of the Solar core. The predicted  ${}^{71}\text{Ga}$  event rate of  $\simeq 107 \text{ SNU}$  can be reduced at most to around  $78 \text{ SNU}$  in most non-standard Solar models [91,99]. The traditional view has been that a flux lower than this would imply large vacuum oscillations, which would reduce the  ${}^{71}\text{Ga}$  rate by a factor comparable to the  ${}^{37}\text{Cl}$  event rate reduction for most oscillation parameters (e.g. to around  $40 \text{ SNU}$ ).

Yet another possibility, i.e. that neutrinos decay between the Sun and the Earth, is all but excluded by the survival of neutrinos from supernova 1987A, except in some two-component models with large mixing angles. [100]

Recently, Mikheyev and Smirnov [101] have proposed an elegant new solution to the Solar neutrino problem, in which even tiny vacuum mixing angles can be amplified by the coherent interactions of  $\nu_e$  with matter.

Considering  $\nu_e \leftrightarrow \nu_\mu$  oscillations for definiteness, the vacuum oscillation equation in (35) can be described in terms of the weak basis states  $|\nu_e\rangle$  and  $|\nu_\mu\rangle$  by

$$|\nu(t)\rangle = \nu_e(t)|\nu_e\rangle + \nu_\mu(t)|\nu_\mu\rangle, \quad (47)$$

where the coefficients satisfy the Schrödinger-like equation

$$i \frac{d}{dt} \begin{pmatrix} \nu_e(t) \\ \nu_\mu(t) \end{pmatrix} = M_0 \begin{pmatrix} \nu_e(t) \\ \nu_\mu(t) \end{pmatrix}, \quad (48)$$

with

$$M_0 = \begin{pmatrix} \frac{\Delta m^2 \cos 2\theta}{2p} & \frac{-\Delta m^2 \sin 2\theta}{4p} \\ \frac{-\Delta m^2 \sin 2\theta}{4p} & 0 \end{pmatrix}, \quad (49)$$

where an irrelevant term proportional to the identity (which only affects the overall phase) has been dropped. Wolfenstein pointed out [102] that in the presence of matter,  $M_0$  is replaced by the  $M'$ , where

$$M' = M_0 + \begin{pmatrix} \sqrt{2}G_F n_e & 0 \\ 0 & 0 \end{pmatrix} \quad (50)$$




Figure 4: The energy eigenvalues of  $M'$  as a function of  $d$ , the distance from the center of the Sun.

and  $n_e$  is the density of electrons. The new term [102] - [104] is the effect of the coherent forward scattering amplitude for  $\nu_e e^- \rightarrow \nu_e e^-$  via the charged current. The effects of neutral current scattering from  $e^-$ ,  $p$ , and  $n$  have been neglected because they are the same for  $\nu_e$  and  $\nu_\mu$  and only contribute to the overall phase. For  $\Delta m^2 < 0$  (i.e.  $m_{\nu_e} < m_{\nu_\mu}$  [104]) there is a critical density [105]  $n_e^{\text{crit}} = -\Delta m^2 \cos 2\theta / (2\sqrt{2}G_F p)$  for which the diagonal elements of  $M'$  are equal (i.e. zero). At that density a resonance occurs, i.e. even a tiny off diagonal mixing term leads to large mixing effects.

In particular, if  $n_e$  in the Sun varies slowly an adiabatic approximation applies [101, 106].  $\nu_e$ 's produced in the core of the Sun (where  $n_e > n_e^{\text{crit}}$ ) correspond to the larger mass eigenstate  $\nu_2$  of  $M'$  (Fig. 4). Outside the Sun, on the other hand, the higher energy state  $\nu_2$  corresponds to  $\nu_\mu$  for  $\Delta m^2 < 0$ . Hence, if the variation of  $n_e$  with the distance from the center of the Sun is sufficiently slow, the initial  $\nu_e$  will be adiabatically converted to  $\nu_\mu$  as they pass through the resonance.

A number of authors [99],[101],[106] - [108] have analyzed the implications of the Mikheyev-Smirnov-Wolfenstein (MSW) effect for the Solar neutrinos quantitatively. It is found that there are three classes of parameters which can explain the reduction of  ${}^8\text{B}$  neutrinos observed in the  ${}^{37}\text{Cl}$  experiment. These roughly form the sides of a triangle, as is illustrated schematically in Fig. 5. For solution (a) corresponding to  $|\Delta m^2| \sim 5 \times 10^{-5} \text{ eV}^2$ ,  $\sin^2 2\theta \geq 4 \times 10^{-4}$ , the adiabatic hypothesis is valid and  $\simeq 100\%$  conversion occurs. However, only the high energy  ${}^8\text{B}$  neutrinos actually encounter a resonance layer (the central density is too low for the low energy neutrinos) and are converted. For this parameter range one expects little reduction in the counting rate for the gallium experiment (i.e. the effect is similar to non-standard Solar models in that respect).

For solution (b), extending down to  $|\Delta m^2| \sim 10^{-8} \text{ eV}^2$  one has [107]  $|\Delta m^2| \sin^2 2\theta \sim 10^{-7.5} \text{ eV}^2$ . Here the adiabatic approximation starts to break down. All neutrino energies are affected, but the conversion probability is less than unity. For these solutions one expects a significant reduction in the gallium counting rate, similar to large vacuum oscillations or magnetic moments. Solution (c), corresponding to large vacuum mixing angles,




Figure 5: A schematic view of the regions in the  $\Delta m^2 - \sin^2 2\theta$  plane which can explain the Solar neutrino problem via the Mikheyev-Smirnov-Wolfenstein (MSW) effect.

is an extension of the vacuum oscillation solution. In the middle of region (c) one expects a large day-night asymmetry in the  $\nu_e$  counting rate due to MSW regeneration in the Earth at night [99].

The MSW solution is very elegant, but it severely complicates the task of sorting out which if any of the proposed solutions to the Solar  $\nu$  problem is correct. It will take an ambitious program of experiments [99,109] to clarify the matter.

The first two events from SN1987A observed by the Kamiokande experiment [45] point back towards the supernova. They are consistent with  $\nu_e$  from the initial neutronization burst, scattering via  $\nu_e e^- \rightarrow \nu_e e^-$ . However, they could also be  $\bar{\nu}_e$  from the subsequent thermal burst, scattering via  $\bar{\nu}_e p \rightarrow e^+ n$ , which has a much larger cross section (and which produces an isotropic distribution of positrons). If they are indeed  $\nu_e$  they are problematic for the MSW mechanism because one expects  $\nu_e \rightarrow \nu_\mu$  conversions on the way out of the supernova. However, there are two parameter regions (shown in Fig. 5), which would still be consistent [110], corresponding respectively to incomplete conversion in the supernova and reconversion in the Earth. Unfortunately there is no way to determine whether the two events are really  $\nu_e$ 's.

The MSW mechanism is consistent with the expectation of GUT [19] and intermediate scale [20] seesaws. As is illustrated in Fig. 5 the predictions of the GUT seesaw are consistent with  $\nu_e \leftrightarrow \nu_\tau$  conversions in the Sun. In this case, the mass ranges are too small to ever see any direct laboratory effects of neutrino mass. The intermediate scale seesaw could account for the Solar  $\nu$  problem via  $\nu_e \leftrightarrow \nu_\mu$  conversions. In that case,  $\nu_\mu \leftrightarrow \nu_\tau$  oscillations could well be observable in the laboratory.

It is also possible that the Solar  $\nu$  problem could be explained by small neutrino masses, but that neutrino appearance experiments might nevertheless yield positive signals due to

non-orthogonal neutrino states (induced by mixings between very light and very heavy neutrinos. [57])

## 5 Summary

- The question of whether the neutrinos have mass is vital for both particle physics and cosmology. However, there is at present no compelling evidence for neutrino mass. The ITEP result  $m_{\nu_e} \sim (17 - 40) eV$  has not been confirmed by other experiments and is on the verge of being excluded. Although there are two positive indications of neutrino oscillations (with different parameters), these are contradicted by other experiments. The negative results suggest  $m_{\nu_i} \simeq O(1 eV)$  unless there are very small mixings or degeneracies. There are also stringent limits on incoherent mixing with heavy neutrinos.
- The MSW solution to the Solar neutrino problem would work for  $\nu_\mu$  or  $\mu_\tau$  in the  $10^{-2} eV$  range, consistent with intermediate mass or GUT seesaws.
- The nonobservation of neutrinoless double beta decay implies  $\langle m_{\nu_e} \rangle < 1 - 11 eV$ . If the ITEP result is correct this would most likely imply a Dirac neutrino or new physics to evade cosmological bounds.
- A  $\nu$  mass in the 5 - 40 eV range would dominate the energy density of the universe and would be an excellent candidate for the dark matter, though other mechanisms would have to be invoked to explain the initial formation of galaxies. Conversely, the light stable neutrinos must be lighter than  $\simeq 40 eV$ . A variety of astrophysical, laboratory, and cosmological bounds exclude unstable neutrinos up to  $\sim 40 MeV$  (unless new physics is invoked for fast, invisible decays or annihilations), implying that  $m_{\nu_\mu}, m_{\nu_\tau} < 40 eV$ .

## References

- [1] There are several recent reviews of neutrino mass. See, for example, F. Boehm and P. Vogel, *Physics of Massive Neutrinos*, (Cambridge Univ. Press, Cambridge, 1987); J.D. Vergados, *Phys. Rev.* 133, 1 (1986); S.M. Bilenky and S.T. Petcov, *Rev. Mod. Phys.* 59, 671 (1987); R. Eichler, *1987 Int. Symp. on Lepton and Photon Interactions at High Energies*.
- [2] Under a CP transformation  $\psi_L(\vec{x}, t) \rightarrow \gamma^0 \psi_R^c(-\vec{x}, t)$ , so  $\psi_L$  and  $\psi_R^c$  are essentially CP conjugates.
- [3] L. Wolfenstein, *Nucl. Phys.* B186, 147 (1981).
- [4] E.S. Konopinski and M. Mahmoud, *Phys. Rev.* 92, 1045 (1953).
- [5] Fermion mass matrices need not be Hermitian. One could generalize still further and introduce  $K(\neq F)$   $N_R^0$  fields, in which case  $m_D$  would be  $F \times K$  dimensional.

- [6] We have chosen a basis for the charged leptons so that the analogue of  $V_L$  for the electrons is the identity matrix.
- [7]  $V_L^1$  contains  $F^2$  real parameters, of which  $F(F-1)/2$  are angles, analogous to the Cabibbo angle, which describe transitions between families and lead to such effects as neutrino oscillations.  $2F-1$  parameters are unobservable phases, which can be removed by appropriate choices of phases for the  $n_{iL}$  and  $e_{iL}^-$  fields (corresponding phases must be chosen for the  $n_{iR}$  and  $e_{iR}$  to ensure real masses). The remaining  $(F-1)(F-2)/2$  parameters are phases which could lead to CP violation in the leptonic sector.
- [8] G.B. Gelmini and M. Roncadelli, Phys. Lett. 99B, 411 (1981); H. Georgi et al., Nucl. Phys. B193, 297 (1983).
- [9] More generally,  $K$  is block diagonal, the blocks being unitary symmetric matrices in the subspaces of degenerate eigenvalues.
- [10] One has  $n_{iR}^c = K_{ji} C \bar{n}_{jL}^T$ , so that  $n_i$  is self-conjugate under the generalized C transformation  $n_i \rightarrow K_{ji} C \bar{n}_j^T$ .
- [11] The counting of angles in  $U_L^1$  is the same as for the Dirac case. However, there are an additional  $F-1$  CP violating phases associated with the fact that  $U_L$  and  $U_R$  are related by (13). For example, if one chooses  $K_L$  so that  $K=I$ , then the phases of the  $n_{iL}$  are fixed and cannot be redefined to remove phases from  $U_L$ . See S.M. Bilenky, J. Hosek, and S.T. Petcov, Phys. Lett. 94B, 495 (1980); M. Doi et al., Phys. Lett. 102B, 323 (1981).
- [12] W.Y. Keung and G. Senjanovic, Phys. Rev. Lett. 50, 1427 (1983); J.F. Gunion and B. Kayser, Proc. 1984 Summer Study on the SSC, eds R. Donaldson and J.G. Morfin (APS, 1984) p. 153.
- [13] H. Goldberg, Phys. Rev. Lett. 50, 1419 (1983); L. M. Krauss, Phys. Lett. 128B, 37 (1983), Nucl. Phys. B227, 556 (1983).
- [14] For a review, see B. Kayser, Comm. Nucl. Part. Phys. 14, 69 (1985).
- [15] New interactions could in principle distinguish the two cases even for  $m_\nu = 0$ , depending on whether they conserve lepton number.
- [16] S.M. Bilenky and B. Pontecorvo, Lett. Nuovo Cimento 17, 569 (1976). The implications of the existence of Dirac and Majorana neutrino mass terms were also studied in: V. Barger et al., Phys. Rev. Lett. 45, 692 (1976); S.M. Bilenky, J. Hosek and S.T. Petcov, Phys. Lett. 94B, 495 (1980); J. Schechter and J.W. F. Valle, Phys. Rev. D22, 2227 (1980); T.P. Cheng and L.F. Li, Phys. Rev. D22, 2860 (1980); T. Yanagida and M. Yoshimura, Progr. Th. Phys. 64, 1870 (1980).
- [17] The relative sign of  $n_{1R}$  and  $n_{2R}$  will reemerge, however, in neutrinoless double beta decay, where it will be seen to be observable.
- [18] M. Gell-Mann, P. Ramond, and R. Slansky, in *Supergravity*, eds. F. van Nieuwenhuizen and D. Freedman, (North Holland, Amsterdam, 1979) p. 315; T. Yanagida, Progr. Th. Physics B135, 66 (1978).
- [19] P. Langacker, S. T. Petcov, G. Steigman, and S. Toshev, Nucl. Phys. B282, 589 (1987).
- [20] R.N. Mohapatra and G. Senjanovic, Z. Phys. C17, 53 (1983); Q. Shafi and F.W. Stecker, Phys. Rev. Lett. 53, 1292 (1984); K. Kang and M. Shin, Nucl. Phys. B287, 687 (1987); K. Kang and A. Pantziris, Phys. Lett. 193B, 467 (1987); P. Langacker, R.D. Peccei, and T. Yanagida, Mod. Phys. Lett. A1, 541 (1986).
- [21] R.N. Mohapatra and G. Senjanovic, Phys. Rev. D23, 165 (1981). For constraints on the heavy neutrino mass, see R.N. Mohapatra; Phys. Rev. D34, 909 (1986).
- [22] P. Langacker, B. Sathiapalan, and G. Steigman, Nucl. Phys. B266, 669 (1986).
- [23] A. Zee, Phys. Lett. 93B, 389 (1980); 161B, 141 (1985); Nucl. Phys. B264, 99 (1986); K. Tamvakis and J. Vergados, Phys. Lett. 155B, 373 (1985); G.K. Leontaris and K. Tamvakis, Phys. Lett. 191B, 421 (1987); M. Fukugita and T. Yanagida, Phys. Rev. Lett. 58, 1807 (1987).
- [24] A. Masiero, D.V. Nanopoulos, and A. Sanda, Phys. Rev. Lett. 57, 663 (1986).
- [25] U. Amaldi et al., Phys. Rev. D36, 1385 (1987).
- [26] Unlike similar hierarchy problems in comparing the weak and unification scale, we know that there is no supersymmetry to control the radiative corrections.
- [27] D.S.P. Dearborn, D.N. Schramm, and G. Steigman, Phys. Rev. Lett. 56, 26 (1986); H.Y. Cheng, Phys. Rev. D36, 1649 (1987); and references therein.
- [28] R. Barbieri, J. Ellis, and M.K. Gaillard, Phys. Lett. 90B, 249 (1980).
- [29] In many models which implement the seesaw mechanism Higgs triplet fields are also present. Even if these have zero VEV at tree level they may obtain VEV's due to higher order corrections to the Higgs potential. See M. Magg and C. Wetterich, Phys. Lett. 94B, 61 (1980); D. Chang and R.N. Mohapatra, Phys. Rev. D32, 1248 (1985); J. Bijnens and C. Wetterich, DESY 87-042 (1987).
- [30] A simple diagonalization yields eigenvalues  $-m_D^2/m_S$  and  $m_S$ . The sign of the smaller eigenvalue is reversed by choosing  $K = \text{diag}(-1, 1)$ , which accounts for the minus sign in  $\nu_R^{0c}$ .
- [31] E. Witten, Phys. Lett. 91B, 81 (1980).
- [32] Some versions of these models [29] are expected to have induced values for  $m_i$  in (31).
- [33] Y. Chikashige, R.N. Mohapatra, and R.D. Peccei, Phys. Lett. 98B, 265 (1981).

- [34] In many of the intermediate mass seesaw models [20] the singlet-Majoron and the invisible axion are the same particle.
- [35] S. Nandi and U. Sarkar, Phys. Rev. Lett. 56, 564 (1986); J. Derendinger, L. Ibanez, and H.P. Nilles, Nucl. Phys. B267, 365 (1986).
- [36] J. Ellis et al., Phys. Lett. 188B, 415 (1987).
- [37] M. Roncadelli and D. Wyler, Phys. Lett. 133B, 325 (1983); A.S. Joshipura et al., Phys. Lett. 156B, 353 (1985); P. Roy et al, Phys. Rev. D30, 2385 (1984); O. Shanker, Nucl. Phys. B250, 351 (1985); J. Oliensis and C.H. Albright, Phys. Lett. 160B, 121 (1985).
- [38] S. Boris et al., Phys. Rev. Lett. 58, 2019 (1987).
- [39] M. Fritschi et al., Phys. Lett. 173B, 485 (1986).
- [40] J.F. Wilkerson et al, Phys. Rev. Lett. 58, 2023 (1987).
- [41] H. Kawakami et al., '86 *Massive Neutrinos in Physics and Astrophysics*, ed. O. Fackler and J. Tran Than Van.
- [42] See, for example, D. Schramm, *1987 Int. Symp. on Lepton and Photon Interactions at High Energies*; H. Meyer, this volume; E.W. Kolb et al., Phys. Rev. D35, 3598 (1987).
- [43] R. Abela et al., Phys. Lett. B146, 431 (1984).
- [44] ARGUS collaboration, quoted by R. Eichler, [1].
- [45] K. Hirakata et al., Phys. Rev. Lett. 58, 1490 (1987).
- [46] R.M. Bionta et al., Phys. Rev. Lett. 58, 1494 (1987).
- [47] For reviews and a general discussion, see M. Gronau, C.N. Leung, and J.L. Rosner, Phys. Rev. D29, 2539 (1984); F.J. Gilman and S.H. Rhie, Phys. Rev. D32, 324 (1985). F.J. Gilman, Comm. Nucl. Part. Phys. 16, 231 (1986); R.E. Shrock, Phys. Rev. D24, 1232, 1275 (1981).
- [48] A similar phenomenon occurs for a system of coupled classical oscillators.
- [49] For a discussion of quantum mechanical subtleties, such as the uncertainty in  $p$ , see F. Boehm and P. Vogel, [1].
- [50] IMB, J.M. Lo Secco et al., Phys. Lett. 184B, 305 (1987).
- [51] The limits in Fig. 2 are obtained by analyzing the data in terms of the formula in (36) for oscillations between two types of neutrino only. Slightly weaker limits are obtained if one allows mixings between 3 neutrino flavors ( $\nu_e, \nu_\mu, \nu_\tau$ ). See K. Kleinknecht, Comm. Nucl. Part. Phys. 16, 267 (1986).
- [52] J.F. Cavaignac et al., Phys. Lett. 148B, 387 (1984).
- [53] G. Zacek et al., Phys. Rev. D34, 2621 (1986).
- [54] G. Bernardi et al., Phys. Lett. 181B, 173 (1986).
- [55] R. Eichler, [1].
- [56] Another manifestation of the underlying physics would be off-diagonal neutral currents, so that a heavy neutrino could decay into three light neutrinos.
- [57] D. Wyler and L. Wolfenstein, Nucl. Phys. B218, 205 (1983); B.W. Lee and R.E. Shrock, Phys. Rev. D16, 1444 (1977); P. Langacker and D. London, to be published.
- [58] For a review, see G. Gelmini, this volume.
- [59] Both  $\Gamma_{weak}$  and  $H$  have additional factors involving the number of types of relativistic particles at temperature  $T$ .
- [60] P. Langacker, J.P. Leveille, and J. Sheiman, Phys. Rev. D27, 1228 (1983).
- [61] N. Cabibbo and L. Maiani, Phys. Lett. 114B, 115 (1982).
- [62] For a recent review, see B. Müller, this volume.
- [63] For masses in the 10 eV range the  $\nu$ 's would be clustered in the galaxy and their cross sections would be increased by  $10^8$ , but direct detection still appears virtually impossible.
- [64] Massive neutrinos could easily account for the scales of galactic clusters. However, it is hard to see how such structures could have fragmented into galaxies in the time available, unless another ingredient, such as cosmic strings, provided the seed.
- [65] G. Steigman et al., Phys. Lett. 176B, 35 (1986).
- [66] For a recent review, see M. Roos, this volume; M. Turner, AIP Conf. Proc. 72, 266 (1980).
- [67] Most of these limits apply to light  $SU_2$ -singlet neutrinos as well, because in many cases their small mixing with ordinary doublet neutrinos is sufficient to produce them prolifically in the early universe. See [22].
- [68] S. Sarker and A.M. Cooper, Phys. Lett. B148, 347 (1984); L. Krauss, Phys. Rev. Lett. 53, 1976 (1984); R. Cowsik, Phys. Lett. 151B, 52 (1985); H. Harari and Y. Nir, Phys. Lett. 188B, 163 (1987), Nucl. Phys. B292, 251 (1987).
- [69] Y. Hosotani, Nucl. Phys. B191, 411 (1981); M. Davis et al., AP.J. 250, 423 (1981); P. Hut and S. White, Nature 310, 637 (1984); P. Binetruy et al., Phys. Lett. 134B, 147 (1984); A.A. Natale, Phys. Lett. 141B, 323 (1984).
- [70] M. Roncadelli and G. Senjanovic, Phys. Lett. 107B, 59 (1983).
- [71] P.B. Pal, Nucl. Phys. B227, 237 (1983); H. Harari and Y. Nir, [68].

- [72] D.B. Reiss, Phys. Lett. 115B, 217 (1982); F. Wilczek, Phys. Rev. Lett. 49, 1549 (1982); G. Gelmini, S. Nussinov, and T. Yanagida, Nucl. Phys. B219, 31 (1983).
- [73] See [8].
- [74] [33] and J. Schechter and J.W.F. Valle, Phys. Rev. D25, 774 (1982); G.B. Gelmini and J.W.F. Valle, Phys. Lett. 142B, 181 (1984); G.B. Gelmini et al., Phys. Lett. 146B, 311 (1984); A. Kumar and R. N. Mohapatra, Phys. Lett. 150B, 191 (1985).
- [75] P. Langacker, G. Segre, and S. Soni, Phys. Rev. D26, 3425 (1982).
- [76] For a recent review, see A. Faessler, this volume.
- [77] T. Kirsten et al., Phys. Rev. Lett. 50, 474 (1983), Z. Phys. C16, 189 (1983).
- [78] M.K. Moe, this volume.
- [79] Neutrinoless double beta decay could conceivably occur through other mechanisms, such as Higgs exchange, or could be accompanied by the emission of a triplet-Majoron. [81] See G.K. Leontaris and J.D. Vergados, CERN-TH.4835/1987. J.D. Vergados, [1].
- [80] F.T. Avignone, this volume.
- [81] The PNL/USC group [80] has reported evidence for  $^{76}\text{Ge} \rightarrow ^{76}\text{Se} e^-e^- + \text{Majoron}$ , but their results are in contradiction with other experiments.
- [82] W.C. Haxton and G.J. Stephenson, Prog. Part. Nucl. Phys. 12, 409 (1984); T. Tomoda et al., Nucl. Phys. A452, 591 (1986); K. Grotz and H.V. Klapdor, Nucl. Phys. A460, 395 (1986); T. Tomoda and A. Faessler, to be published.
- [83] J. Engel, P. Vogel, and M.R. Zirnbauer, to be published.
- [84] L. Wolfenstein, Phys. Lett. 107B, 77 (1981); M. Doi et al., Phys. Lett. 102B, 323 (1981).
- [85] See W.C. Haxton and G.J. Stephenson, [82], J.D. Vergados, [1].
- [86] A. Halprin, S.T. Petcov, and S.P. Rosen, Phys. Lett. 125B, 335 (1983); C.N. Leung and S.T. Petcov, Phys. Lett. 145B, 416 (1984).
- [87] M. Doi, T. Kotani, and E. Takasugi, Prog. Th. Phys. (Supp) 83, 1 (1985).
- [88] However, in a gauge theory some neutrino must be massive. Otherwise the diagrams cancel. See B. Kayser, *7<sup>th</sup> Moriond Workshop on Searches for New and Exotic Phenomena*, Les Arcs, Jan. 1987.
- [89] J.K. Rowley, B.T. Cleveland and R. Davis, AIP Conference Proc. 126, 1 (1985).
- [90] A Solar neutrino unit (*SNU*) is defined as  $10^{-36}$  interactions per target nucleus per second.
- [91] J.N. Bahcall et al., Rev. Mod. Phys. 54, 767 (1982); J.N. Bahcall, AIP Conference Proc. 126, 60 (1985).
- [92] H. Hirata et al, to be published.
- [93] For recent discussions, see V. Barger et al., Phys. Rev. D24, 538, (1981); S.L. Glashow and L.M. Krauss, Phys. Lett. 190B, 199 (1987); L. Krauss and F. Wilczek, Phys. Rev. Lett. 55, 122 (1985).
- [94] M.B. Voloshin and M.I Vysotsky, ITEP 86-1(1986); L.B. Okun et al., Zh. Eksp. Teor. Fiz. 91, 754 (1986) (JETP 64, 446 (1986)). For earlier references, see [19].
- [95] A.V. Kyuldjiev, Nucl. Phys. B243, 387 (1987); K. Abe et al., Phys. Rev. Lett. 58, 636 (1987).
- [96] For a recent discussion, see M. Fukujita and S. Yazaki, Kyoto preprint. RIFP-709.
- [97] B.W. Lee and R. Shrock, Phys. Rev. D16, 1444 (1977); K. Fujikawa and R. Shrock, Phys. Rev. Lett. 45, 963 (1980).
- [98] M.J. Duncan et al., Phys. Rev. Lett. 191B, 304 (1987); J. Liu, Phys. Rev. D35, 3447 (1987); M. Fukugita and T. Yanagida, [23].
- [99] For a review, see W. Hampel, this volume.
- [100] F.A. Frieman, H.E. Haber, and K. Freese, Phys. Lett. 200B, 115 (1988).
- [101] S.P. Mikheyev and A. Yu Smirnov, Yad Fiz. 42, 1441 (1985) (Sov. Jour. Nucl. Phys. 42, 1441 (1985)); Nuo. Cim. 9C, 17 (1986).
- [102] L. Wolfenstein, Phys. Rev. D17, 2369 (1968); D20, 2634 (1979).
- [103] V. Barger et al., Phys. Rev. D22, 2718 (1980).
- [104] A crucial sign error was corrected in [60].
- [105] The scattering amplitude in  $M'$  changes sign for antineutrinos. Hence, resonance can occur for either  $\nu_e$  or  $\bar{\nu}_e$ , depending on the sign of  $\Delta m^2$ .
- [106] H.A. Bethe, Phys. Rev. Lett. 56, 1305 (1986).
- [107] S.P. Rosen and S.M. Gelb, Phys. Rev. D34, 969 (1986).
- [108] P. Langacker, S.T. Petcov, G. Steigman, and S. Toshev, Nucl. Phys. B282, 589 (1987).
- [109] E.W. Beier et al., 1986 Snowmass Workshop, Pennsylvania UPR-0140E.
- [110] J. Arafune et al., Phys. Rev. Lett. 59, 1864 (1987); P.O. Lagage et al., Phys. Lett. 193B, 127 (1987); L. Wolfenstein, Phys. Lett. 194B, 197 (1987); D. Nötzgold, Phys. Lett. 196B, 315 (1987); S. P. Rosen, Los Alamos LA-UR-87-1296.