

DEUTSCHES ELEKTRONEN-SYNCHROTRON
Ein Forschungszentrum der Helmholtz-Gemeinschaft



DESY 21-133
HU-EP-21/29-RTG
arXiv:2108.13432
August 2021

Bootstrapping Holographic Defect Correlators in $\mathcal{N} = 4$ Super Yang-Mills

J. Barrat

*Institut für Physik und IRIS Adlershof,
Humboldt-Universität, Berlin*

A. Gimenez-Grau, P. Liendo

Deutsches Elektronen-Synchrotron DESY, Hamburg

ISSN 0418-9833

NOTKESTRASSE 85 - 22607 HAMBURG

DESY behält sich alle Rechte für den Fall der Schutzrechtserteilung und für die wirtschaftliche Verwertung der in diesem Bericht enthaltenen Informationen vor.

DESY reserves all rights for commercial use of information included in this report, especially in case of filing application for or grant of patents.

To be sure that your reports and preprints are promptly included in the
HEP literature database
send them to (if possible by air mail):

DESY Zentralbibliothek Notkestraße 85 22607 Hamburg Germany	DESY Bibliothek Platanenallee 6 15738 Zeuthen Germany
---	---

Bootstrapping holographic defect correlators in $\mathcal{N} = 4$ super Yang-Mills

Julien Barrat,¹ Aleix Gimenez-Grau,² Pedro Liendo.²

¹*Institut für Physik und IRIS Adlershof, Humboldt-Universität zu Berlin, Zum Großen Windkanal 2, 12489 Berlin, Germany*

²*DESY Hamburg, Theory Group, Notkestraße 85, D-22607 Hamburg, Germany*

E-mail: julien.barrat@hu-berlin.de, aleix.gimenez@desy.de,
pedro.liendo@desy.de

ABSTRACT: We study two-point functions of single-trace half-BPS operators in the presence of a supersymmetric Wilson line in $\mathcal{N} = 4$ SYM. We use inversion formula technology in order to reconstruct the CFT data starting from a single discontinuity of the correlator. In the planar strong coupling limit only a finite number of conformal blocks contributes to the discontinuity, which allows us to obtain elegant closed-form expressions for two-point functions of single-trace operators \mathcal{O}_J of weight $J = 2, 3, 4$. Our final result passes a number of non-trivial consistency checks: it has the correct discontinuity, it satisfies the superconformal Ward identities, it has a sensible expansion in both defect and bulk OPEs, and is consistent with available results coming from localization. The method is completely algorithmic and can be implemented to calculate correlators of arbitrary weight.

Contents

1	Introduction	2
2	Preliminaries	3
2.1	Setup	3
2.2	Supergravity interpretation	4
2.3	Superconformal kinematics	5
2.4	Topological subsector	8
3	$\langle\langle \mathcal{O}_2 \mathcal{O}_2 \rangle\rangle$ at strong coupling	10
3.1	Lorentzian inversion formula	10
3.2	Computation of the discontinuity	11
3.3	Inversion of $\mathcal{B}_{[0,2,0]}$	12
3.4	Supersymmetrization of the correlator	15
3.5	Final result and comparison to localization	16
4	General identical operators	18
4.1	General discussion	18
4.2	Example 1: $\langle\langle \mathcal{O}_3 \mathcal{O}_3 \rangle\rangle$	19
4.3	Example 2: $\langle\langle \mathcal{O}_4 \mathcal{O}_4 \rangle\rangle$	22
4.4	A conjecture	23
5	Conclusions	24
A	Appendix	25
A.1	Singular part of bulk blocks	25
A.2	Conformal block normalization	25

1 Introduction

The analytic conformal bootstrap is a powerful tool that has seen significant progress in recent years. The basic proposal made in the original papers [1, 2], is that singular terms in a CFT correlator imply the existence of families of operators at large spin. This observation was later systematized and allowed the calculation of CFT data as a series expansion in inverse powers of the spin variable [3]. This line of thinking led to the Lorentzian inversion formula, that neatly captures the dependence of the CFT data as an analytic function of the spin variable [4]. This formula is a fully non-perturbative result valid for any CFT in any dimension.

Apart from giving us an improved understanding of the structure of CFT, the inversion formula is also a powerful calculational tool. Its power lies on the fact that correlators can be reconstructed from a certain discontinuity, which is in general a simpler object. Inversion formula technology is particularly useful for planar theories at strong coupling, where the discontinuity is captured by only a finite number of conformal blocks, simplifying computations considerably. This approach was carried out successfully for four-point functions of half-BPS operators in $\mathcal{N} = 4$ SYM [5, 6], and confirmed a previous conjecture made in Mellin space [7, 8].

Because the presence of a defect does not change the local physics, the main properties that enable the bootstrap for strong coupling correlators are still present if we add a line defect to the configuration. More precisely, in this paper we consider correlators between single trace half-BPS operators and a supersymmetric Wilson line. These are canonical examples of local and non-local operators in $\mathcal{N} = 4$ SYM. One point-function of half-BPS operators in the presence of a line are fixed by kinematics and the overall coefficient can be calculated exactly using matrix-model techniques [9–12]. Less work has been done on two-point correlators in the presence of a line. This is the first case in which correlators depend non-trivially on conformal invariants and therefore capture an infinite amount of CFT data. The only results on the literature so far are an explicit weak-coupling calculation [13], the extreme strong coupling limit where the correlator becomes the square of a one-point function [14, 15], and special kinematical points where the correlator does not depend on invariants and can be calculated using localization [14, 16]. The goal of this paper is to calculate the full two-point correlator as a function of its cross-ratios, to the first non-trivial order at strong coupling, relying only on analytical bootstrap techniques.

The outline of the paper is as follows. In section 2 we review the basic kinematics of our two-point functions, including an interpretation of our bootstrap problem in terms of Witten diagrams. In section 3 we compute the two-point function for the stress-tensor multiplet, which is the half-BPS operator of weight $J = 2$, using Lorentzian inversion technology, and present an elegant closed-form expression for the full correlator. In section 4 we generalize the analysis for operators of higher weight and present explicit solutions for the cases $J = 3, 4$ as a demonstration of our method. Possible future directions are discussed in section 5, while in appendix A we give the bulk and defect conformal blocks needed for the computations.

2 Preliminaries

In this section we review the basic properties of the setup under study, before jumping into the technicalities of the computation in section 3.

2.1 Setup

Consider $\mathcal{N} = 4$ SYM theory with gauge group $U(N)$ in the planar $N \rightarrow \infty$ limit. The most important local operators in our discussion are single-trace chiral-primary operators:

$$\mathcal{O}_J(x, u) := (2\pi)^J \frac{2^{J/2}}{\sqrt{J\lambda^J}} \text{tr} (u \cdot \phi(x))^J. \quad (2.1)$$

Here u is a six-component null polarization vector $u \cdot u = 0$, such that $\mathcal{O}_J(x, u)$ transforms in a symmetric-traceless representation of the R -symmetry group. These are protected operators that preserve half of the supercharges of the theory. The other important observable in our analysis is the supersymmetric Wilson line (sometimes called the Maldacena-Wilson line):

$$\mathcal{W}_\ell := \frac{1}{N} \text{tr} \mathcal{P} \exp \int_{-\infty}^{\infty} d\tau (i\dot{x}^\mu A_\mu + |\dot{x}| \theta \cdot \phi). \quad (2.2)$$

This extended operator also preserves half of the supercharges, and has been studied extensively in the literature. Here θ is a unit-normalized six-component vector $\theta \cdot \theta = 1$ that parametrizes a direction in R -symmetry space.

In this work we study correlators of local operators in the presence of the supersymmetric Wilson line:¹

$$\langle\langle \mathcal{O}_{J_1}(x_1, u_1) \mathcal{O}_{J_2}(x_2, u_2) \dots \rangle\rangle := \frac{1}{\langle \mathcal{W}_\ell \rangle} \langle \mathcal{W}_\ell \mathcal{O}_{J_1}(x_1, u_1) \mathcal{O}_{J_2}(x_2, u_2) \dots \rangle. \quad (2.3)$$

Even though our configuration breaks some of the $PSU(2, 2|4)$ symmetry of $\mathcal{N} = 4$ SYM, these correlation functions are still restricted by the remaining defect (super)conformal symmetry. It is well understood that in defect CFT one-point functions are kinematically fixed, see for example [17]. The simplest observables in which the coordinate dependence is dynamical and not fixed by symmetry are two-point functions of bulk operators. The case $J_1 = J_2 = 2$ was studied recently at weak 't Hooft coupling [13] using standard Feynman diagrams. Here we consider instead a perturbative expansion at large 't Hooft coupling:

$$\langle\langle \mathcal{O}_J(x_1, u_1) \mathcal{O}_J(x_2, u_2) \rangle\rangle \quad \text{as} \quad N \rightarrow \infty, \lambda = g^2 N \gg 1. \quad (2.4)$$

To be more precise, our correlator admits a double expansion in powers of $\frac{\lambda}{N^2}$ and $\frac{1}{\sqrt{\lambda}}$ of the form

$$\langle\langle \mathcal{O}_J \mathcal{O}_J \rangle\rangle = \langle\langle \mathcal{O}_J \mathcal{O}_J \rangle\rangle^{(0)} + \frac{\lambda}{N^2} \left(\langle\langle \mathcal{O}_J \mathcal{O}_J \rangle\rangle^{(1)} + \frac{1}{\sqrt{\lambda}} \langle\langle \mathcal{O}_J \mathcal{O}_J \rangle\rangle^{(2)} + O(\lambda^{-1}) \right) + \dots, \quad (2.5)$$

where \dots stands for terms starting at $\frac{\lambda^2}{N^4}$. As we will discuss shortly, the first two terms in the expansion are somewhat trivial. Our goal then is to use modern bootstrap methods to calculate the $\langle\langle \mathcal{O}_J \mathcal{O}_J \rangle\rangle^{(2)}$ contribution. The above perturbative expansion has a natural interpretation in terms of the holographic dual of $\mathcal{N} = 4$ SYM, to which we now turn.


¹The overall normalization is redundant in the present case because $\langle \mathcal{W}_\ell \rangle = 1$, but it would be important for circular Wilson loops.

two-point function contains an interaction vertex in the bulk [14]. In the exchanged line one sums over all KK modes that can couple to two \mathcal{O}_J 's. Schematically we have



(2.10)

This is the diagram we calculate in the present work.³

Instead of an explicit calculation using the effective action in $AdS_5 \times S^5$, we will bootstrap the result using the bulk-to-defect inversion formula obtained in [27]. The inversion formula reconstructs a correlator from its singular part, which is mathematically captured by a discontinuity:


(2.11)

The crucial property of holographic CFT's is that the discontinuity is dramatically simpler than the correlator. In particular, we show below that the discontinuity receives corrections only from a finite number of single-trace operators exchanged in the bulk. Each of these contributions is schematically a product of a tree-level one-point function and a three-point function:


(2.12)

The one- and three-point functions are known from localization, so all is left is to compute a certain integral and sum over finitely many single-trace contributions. In the rest of the paper, we translate this pictorial representation into a concrete bootstrap algorithm, that fully fixes two-point correlators with minimal external input.

2.3 Superconformal kinematics

The defect CFT associated with the Wilson line (2.2) preserves an $OSp(4^*|4)$ subgroup of the full $PSU(2,2|4)$ symmetry. From the spacetime perspective, the defect preserves an $SO(2,1) \subset SO(4,2)$ subgroup of the full conformal algebra. Furthermore, the presence of a preferred direction θ preserves only $SO(5)_R \subset SO(6)_R$ of the R -symmetry. The kinematics of non-supersymmetric defects have been thoroughly studied in [17]. Below we only give a brief review, highlighting the new features due to supersymmetry, such as superconformal Ward identities and superconformal blocks.

³In the case of holographic four-point functions, there exist contact Witten diagrams which are correctly reconstructed from the inversion formula [5, 6]. If such diagrams are present in the defect case, our bootstrap result should capture them. We defer a more detailed study of the explicit holographic calculation for future work.

2.3.1 Two-point function and cross-ratios

A two-point function in a defect CFT depends on two spacetime cross-ratios, while in our setup there is an extra R -symmetry cross-ratio σ :

$$\langle\langle \mathcal{O}_J(x_1, u_1) \mathcal{O}_J(x_2, u_2) \rangle\rangle = \frac{(u_1 \cdot \theta)^J (u_2 \cdot \theta)^J}{|x_1^\perp|^J |x_2^\perp|^J} \mathcal{F}^{(J)}(z, \bar{z}, \sigma). \quad (2.13)$$

For the spacetime part, we use the cross-ratios z, \bar{z} defined in [27]

$$\frac{z + \bar{z}}{2\sqrt{z\bar{z}}} = \frac{x_1^\perp \cdot x_2^\perp}{|x_1^\perp| |x_2^\perp|}, \quad \frac{(1-z)(1-\bar{z})}{\sqrt{z\bar{z}}} = \frac{x_{12}^2}{|x_1^\perp| |x_2^\perp|}. \quad (2.14)$$

Geometrically z, \bar{z} are coordinates in a plane orthogonal to the defect. Indeed, placing the first operator at $x_1 = (1, 0, 0, 0)$ and the second one in a xy -plane $x_2 = (x, y, 0, 0)$, then $z = x + iy$ and $\bar{z} = x - iy$. In Lorentzian signature, one would instead find that z and \bar{z} are real and independent. On the other hand, the R -symmetry cross-ratio is defined as

$$\sigma := \frac{(u_1 \cdot u_2)}{(u_1 \cdot \theta)(u_2 \cdot \theta)}. \quad (2.15)$$

The correlator $\mathcal{F}^{(J)}(z, \bar{z}, \sigma)$ is a polynomial of order J in the σ cross-ratio. This reflects the number of different ways to contract u_1, u_2 with each other and with θ , the polarization of the Wilson-line defect. Before moving on, let us point out that one can also consider operators restricted to the line and study the corresponding $1d$ CFT, this configuration has been studied recently by a variety of means [28–33]⁴. In this work we consider bulk operators outside the line, which probes the interplay between local physics and the defect.

2.3.2 Structure of the correlator

As discussed above, the leading contributions at strong coupling to $\langle\langle \mathcal{O}_J \mathcal{O}_J \rangle\rangle$ take the simple form (2.8) and (2.9). Comparing with the form of the two-point function (2.13) we find

$$\mathcal{F}^{(J)}(z, \bar{z}, \sigma) = \left(\frac{\sigma \sqrt{z\bar{z}}}{(1-z)(1-\bar{z})} \right)^J + a_J^2 + O\left(\frac{\sqrt{\lambda}}{N^2}\right). \quad (2.16)$$

Our goal is to compute the correlation function at the next order $O(\frac{\sqrt{\lambda}}{N^2})$. For simplicity we decompose the correlator at this order in powers of the R -symmetry cross-ratio σ :⁵

$$\mathcal{F}^{(J)}(z, \bar{z}, \sigma) \Big|_{O(\frac{\sqrt{\lambda}}{N^2})} = \sum_{j=0}^J \sigma^j F_{J-j}^{(J)}(z, \bar{z}). \quad (2.17)$$

Below we use bootstrap methods to reconstruct the functions $F_j^{(J)}(z, \bar{z})$, leading to the final results in equations (3.33), (4.14) and (4.18).

⁴Similar setups have also been considered for $3d$ ABJM theories [34–36].

⁵Note that the definition of $F_j^{(J)}$ in this decomposition is slightly different to the one used in [13]: here the terms are ordered in terms of powers of σ and not Ω . Furthermore, here we keep outside the definition the two leading contributions which are trivial.

2.3.3 Superconformal Ward identities

An important property of our correlator is that it satisfies *superconformal Ward identities*. These were studied in detail for half-BPS boundaries in [37], where it was observed that with a suitable identification of the cross-ratios, they also apply to the half-BPS line defect.⁶ In our conventions the Ward identities read

$$\left(\partial_z + \frac{1}{2}\partial_\omega\right)\mathcal{F}^{(J)}(z, \bar{z}, \sigma)\Big|_{z=\omega} = 0, \quad \left(\partial_{\bar{z}} + \frac{1}{2}\partial_\omega\right)\mathcal{F}^{(J)}(z, \bar{z}, \sigma)\Big|_{\bar{z}=\omega} = 0, \quad (2.18)$$

where the natural variable to use is ω , defined by

$$\sigma = -\frac{(1-\omega)^2}{2\omega}. \quad (2.19)$$

It is not hard to check that the two leading terms (2.16) satisfy these equations. Note that the above Ward identities take a form very similar to other setups in the literature [38–40]. In the case of $\mathcal{N} = 4$ SYM without defects, the Ward identities admit a simple closed-form solution in which the different R -symmetry channels are related by algebraic relations. This drastically simplifies the analysis because in the end one can just work with the independent channels. Sadly, the defect setup of this paper is closer to the Ward identities in three dimensions, where such an algebraic relation does not exist [38]. It is possible however to relate the different R -symmetry channels by the action of a non-local operator. This is obviously more cumbersome, but one can still implement it in order to focus only on the independent portions of the correlator [41]. In this paper we work explicitly with all the R -symmetry channels, however finding a better parameterization for our correlators is an interesting problem that should be studied in more detail.

2.3.4 Crossing equation

The last important ingredient for our calculation are the defect CFT crossing equations. Here we only summarize them, and refer the reader to [17] for further details. In any CFT it is possible to fuse two bulk local operators as a sum of bulk operators. This *bulk OPE* is denoted schematically as $\mathcal{O}_J\mathcal{O}_J \sim \sum_{\mathcal{O}} \lambda_{JJ\mathcal{O}}\mathcal{O}$. On the other hand, the *defect OPE* expands a bulk operator as a sum of defect operators $\mathcal{O}_J \sim \sum_{\hat{\mathcal{O}}} b_{J\hat{\mathcal{O}}}\hat{\mathcal{O}}$. Note that for a bulk one-point function $\langle\langle \mathcal{O} \rangle\rangle \sim a_{\mathcal{O}}$, the defect OPE implies $a_{\mathcal{O}} = b_{\mathcal{O}\hat{\mathbb{1}}}$ where $\hat{\mathbb{1}}$ is the defect identity operator. These two expansions can be inserted in a two-point function, resulting in a *crossing equation*

$$\mathcal{F}^{(J)}(z, \bar{z}, \sigma) = \left(\frac{\sqrt{z\bar{z}}\sigma}{(1-z)(1-\bar{z})}\right)^J \sum_{\mathcal{O}} \lambda_{JJ\mathcal{O}} a_{\mathcal{O}} \mathcal{G}_{\mathcal{O}}^{(J)}(z, \bar{z}, \sigma) = \sum_{\hat{\mathcal{O}}} b_{J\hat{\mathcal{O}}}^2 \hat{\mathcal{G}}_{\hat{\mathcal{O}}}^{(J)}(z, \bar{z}, \sigma). \quad (2.20)$$

It is important to keep in mind that operators in superconformal theories belong to superconformal multiplets. For example, in the crossing equation, \mathcal{O} refers to the superconformal primary operator of a $PSU(2, 2|4)$ representation, which is labeled by the dimension Δ ,

⁶To be precise, z, \bar{z} are mapped to the boundary R -symmetry cross-ratios w_1, w_2 , while σ is mapped to the boundary spacetime cross-ratio -2ξ (this factor -2 ensures the correctness of this map).

the spin ℓ , and an $SO(6)_R$ label K . Similarly, $\hat{\mathcal{O}}$ is the superconformal primary of an $OSp(4^*|4)$ representation labeled by the defect dimension $\hat{\Delta}$, the transverse-spin s , and an $SO(5)_R$ label \hat{K} .⁷ Supersymmetry fixes the contributions of superdescendant operators, so the expansion (2.20) is organized in *superconformal blocks*. These are linear combination of non-supersymmetric conformal blocks:

$$\begin{aligned}\mathcal{G}_{\mathcal{O}}^{(J)}(z, \bar{z}, \sigma) &= \sum_{\Delta, \ell, K} c_{\Delta, \ell, K}^{(J)} h_K(\sigma) f_{\Delta, \ell}(z, \bar{z}), \\ \hat{\mathcal{G}}_{\hat{\mathcal{O}}}^{(J)}(z, \bar{z}, \sigma) &= \sum_{\hat{\Delta}, s, \hat{K}} c_{\hat{\Delta}, s, \hat{K}}^{(J)} \hat{h}_{\hat{K}}(\sigma) \hat{f}_{\hat{\Delta}, s}(z, \bar{z}).\end{aligned}\tag{2.21}$$

The sums range over all operators in the supermultiplets of $\mathcal{O}/\hat{\mathcal{O}}$ that can appear in the bulk/defect OPE of \mathcal{O}_J . The function $f_{\Delta, \ell}(z, \bar{z})$ is a bulk-channel conformal block, see (A.1) for a useful series representation. Similarly, $\hat{f}_{\hat{\Delta}, s}(z, \bar{z})$ is a defect-channel conformal block given in (A.4). Finally, $h_K(\sigma)$ and $\hat{h}_{\hat{K}}(\sigma)$ are bulk and defect R -symmetry blocks respectively. They are polynomials in σ with a simple hypergeometric closed form expression (A.5). The precise relative coefficients $c_{\Delta, \ell, K}^{(J)}$ and $c_{\hat{\Delta}, s, \hat{K}}^{(J)}$ have been presented in the appendix of [13].

2.4 Topological subsector

An important property of our correlators is that they contain a topological subsector. In terms of the cross-ratios, the topological subsector can be obtained by setting $z = \bar{z} = \omega$. This projects out all the non-protected operators from the bulk and defect OPE and only half-BPS protected operators remain. This topological subsector goes beyond our two-point correlators, and it actually describes a closed-subsector of the operator spectrum of the theory. The CFT data of the topological subsector can be obtained by solving Gaussian multi-matrix models [11, 14, 29, 45]. Some of the explicit results that we present below are necessary to fix overall coefficients in our bootstrap analysis, while others provide non-trivial consistency checks of our calculation.

We start by looking at bulk single-trace half-BPS operators \mathcal{O}_J defined in (2.1). We follow the usual normalization conventions such that their two-point function is unit normalized, and the dynamical information is captured by three-point functions. In our conventions

$$\begin{aligned}\langle \mathcal{O}_J(x_1, u_1) \mathcal{O}_J(x_2, u_2) \rangle &= \frac{(u_1 \cdot u_2)^J}{(x_{12}^2)^J}, \\ \langle \mathcal{O}_{J_1}(x_1, u_1) \mathcal{O}_{J_2}(x_2, u_2) \mathcal{O}_{J_3}(x_3, u_3) \rangle &= \lambda_{J_1 J_2 J_3} \frac{(u_1 \cdot u_2)^{J_{123}} (u_2 \cdot u_3)^{J_{231}} (u_3 \cdot u_1)^{J_{312}}}{|x_{12}|^{2J_{123}} |x_{23}|^{2J_{231}} |x_{31}|^{2J_{312}}},\end{aligned}\tag{2.22}$$

with $J_{ijk} := (J_i + J_j - J_k)/2$. The OPE coefficients $\lambda_{J_1 J_2 J_3}$ were originally computed in [46], and they are independent of the coupling:

$$\lambda_{J_1 J_2 J_3} = \frac{\sqrt{J_1 J_2 J_3}}{N}.\tag{2.23}$$

⁷Detailed analysis of the representation theory of these supergroups can be found in the literature [42–44].

As already discussed, one-point functions of single-trace operators in the presence of the supersymmetric Wilson line are kinematically fixed:

$$\langle\langle \mathcal{O}_J(x, u) \rangle\rangle = a_J \frac{(u \cdot \theta)^J}{|x^\perp|^J}. \quad (2.24)$$

Since the normalization of \mathcal{O}_J is determined by the two point function, the coefficient a_J contains dynamical information about the defect CFT.⁸ The precise value can be obtained from a perturbative calculation [9] or from a matrix-model calculation [10–12]

$$a_J = \frac{\sqrt{\lambda J}}{2^{J/2+1} N} \frac{I_J(\sqrt{\lambda})}{I_1(\sqrt{\lambda})} \stackrel{\lambda \gg 1}{\cong} \frac{\sqrt{\lambda J}}{2^{J/2+1} N} \left(1 - \frac{(J+1)(J-1)}{2\sqrt{\lambda}} + O(\lambda^{-1}) \right). \quad (2.25)$$

Note that it is also possible to construct half-BPS multi-trace operators. For example, the OPE between the two single-trace operators $\mathcal{O}_J \times \mathcal{O}_J$ must contain the following *double-trace* operator:

$$\mathcal{O}_{(J,J)}(x, u) := (2\pi)^{2J} \frac{2^{J-1/2}}{J\lambda^J} \text{tr}(u \cdot \phi(x))^J \text{tr}(u \cdot \phi(x))^J. \quad (2.26)$$

The prefactor is chosen to have a unit normalized two-point function. By using the localization techniques of [11, 12, 16], we derived the one-point function of $\mathcal{O}_{(J,J)}$ and its three-point function with single-trace operators. Their product reads

$$a_{(J,J)} \lambda_{JJ(J,J)} = \frac{J\lambda}{2^{J+2} N^2} \frac{I_{2J-1}(\sqrt{\lambda})}{I_1(\sqrt{\lambda})} \stackrel{\lambda \gg 1}{\cong} \frac{J\lambda}{2^{J+2} N^2} \left(1 - \frac{2J(J-1)}{\sqrt{\lambda}} + O(\lambda^{-1}) \right). \quad (2.27)$$

This is all the information we need about bulk protected operators.

The topological sector also captures information involving protected operators localized on the defect. There is a large degeneracy of protected defect operators which has been discussed in [45]. For our purposes, there are two defect operators that play an important role. On the one hand, there are defect operators inserted inside the path-ordering of the Wilson line, which we write as

$$\hat{\mathcal{O}}_{\hat{K}}(\tau, \hat{u}) = \mathcal{W} \left[(\hat{u} \cdot \phi(\tau))^{\hat{K}} \right]. \quad (2.28)$$

These operators are in symmetric traceless representations of the $SO(5)_R$ symmetry preserved by the defect. Therefore, \hat{u} is a null polarization vector orthogonal to the Wilson line polarization $\hat{u} \cdot \theta = \hat{u} \cdot \hat{u} = 0$. On the other hand, it is possible to define a similar operator which lives outside the path-ordering

$$\hat{\mathcal{O}}_{(\hat{K})}(\tau, \hat{u}) = \mathcal{W} \text{tr}(\hat{u} \cdot \phi(\tau))^{\hat{K}}. \quad (2.29)$$

For both of these operators, one should choose a normalization such that their defect two-point functions are unit normalized. The non-trivial dynamical information is then encoded

⁸For the particular case $J = 2$ the coefficient a_2 is proportional to the Bremsstrahlung function [47].

in the bulk-defect two-point function:

$$\begin{aligned}\langle\langle \hat{\mathcal{O}}_{\hat{K}}(\tau_1, \hat{u}_1) \hat{\mathcal{O}}_{\hat{K}}(\tau_2, \hat{u}_2) \rangle\rangle &= \frac{(\hat{u}_1 \cdot \hat{u}_2)^{\hat{K}}}{\tau_{12}^{2\hat{K}}}, \\ \langle\langle \mathcal{O}_J(x_1, u_1) \hat{\mathcal{O}}_{\hat{K}}(\tau_2, \hat{u}_2) \rangle\rangle &= b_{J\hat{K}} \frac{(u_1 \cdot \hat{u}_2)^{\hat{K}} (u_1 \cdot \theta)^{J-\hat{K}}}{((x_1^\perp)^2 + \tau_{12}^2)^{\hat{K}} |x_1^\perp|^{J-\hat{K}}}.\end{aligned}\tag{2.30}$$

For operators of the first type $\hat{\mathcal{O}}_{\hat{K}}$, the bulk-defect coefficient has been calculated at strong coupling in the planar limit [45]

$$b_{J\hat{K}} = \frac{\lambda^{\frac{2-\hat{K}}{4}}}{N} \frac{2^{\frac{3\hat{K}-J-2}{2}} \sqrt{J} \Gamma(\frac{J+\hat{K}+1}{2})}{\sqrt{\hat{K}!} \Gamma(\frac{J-\hat{K}+1}{2})} \left(1 + \frac{4 - 4J^2 + 5\hat{K} + \hat{K}^2}{8\sqrt{\lambda}} + O(\lambda^{-1}) \right).\tag{2.31}$$

On the other hand, the coefficients $b_{J(\hat{K})}$ have not appeared in the literature. An interesting outcome of our analysis is a prediction for the value of $b_{J(j)}$. However, let us stress that our correlators also contain information about infinitely many non-protected operators that are not captured by the topological subsector.

Note that the crossing equation given in (2.20) truncate on both sides in the topological sector [37]. This is known as *microbootstrap*, and in some cases the system of equations can be solved exactly [48].

3 $\langle\langle \mathcal{O}_2 \mathcal{O}_2 \rangle\rangle$ at strong coupling

In this section we compute the correlator $\langle\langle \mathcal{O}_2 \mathcal{O}_2 \rangle\rangle$ in the strong coupling limit. As explained in section 2.2, the two leading contributions are somewhat trivial and take the form given in equations (2.8) and (2.9). As in equation (2.17), we decompose the next correction into three R -symmetry channels:

$$\mathcal{F}^{(2)}(z, \bar{z}, \sigma) \Big|_{O(\frac{\sqrt{\lambda}}{N^2})} = \sigma^2 F_0(z, \bar{z}) + \sigma F_1(z, \bar{z}) + F_2(z, \bar{z}).\tag{3.1}$$

For compactness, here and for the rest of this section we drop the superscripts in the functions $F_j^{(J)}(z, \bar{z})$. In what follows we derive this correlator using the Lorentzian inversion formula presented in [27].

3.1 Lorentzian inversion formula

The idea of the Lorentzian inversion formula is that the *discontinuity* of the correlator is sufficient to extract the full defect CFT data, which in turn can be used for reconstructing the full correlator. For now we consider general single-trace operators $\mathcal{O}_J(x, u)$, and later we focus on the $J = 2$ case. For a codimension-three defect, such as a Wilson line in four dimensions, the inversion formula reads

$$\begin{aligned}b_j(\hat{\Delta}, s) &= \int_0^1 \frac{dz}{2z} z^{-(\hat{\Delta}-s)/2} \int_1^{1/z} \frac{d\bar{z}}{2\pi i} (1 - z\bar{z})(\bar{z} - z) \bar{z}^{-(\hat{\Delta}+s)/2-2} \\ &\times {}_2F_1\left(\frac{1}{2}, 1+s, \frac{3}{2}+s; \frac{z}{\bar{z}}\right) {}_2F_1\left(\frac{1}{2}, 1-\hat{\Delta}, \frac{3}{2}-\hat{\Delta}; z\bar{z}\right) \text{Disc } F_j(z, \bar{z}),\end{aligned}\tag{3.2}$$

where the discontinuity is computed around the branch cut in $\bar{z} \in [1, \infty)$:

$$\text{Disc } F_j(z, \bar{z}) = F_j(z, \bar{z} + i\epsilon) - F_j(z, \bar{z} - i\epsilon), \quad \bar{z} \geq 1. \quad (3.3)$$

The inversion formula is *bosonic* and thus it should be applied to each R -symmetry channel $F_j(z, \bar{z})$ independently. The defect conformal dimensions are encoded in the poles of $b_j(\hat{\Delta}, s)$, while the residues are OPE coefficients:

$$b_j(\hat{\Delta}, s) = - \sum_{n \geq 0} \frac{(b_j^2)_{n,s}}{\hat{\Delta} - (J + s + 2n + \gamma_{n,s})}. \quad (3.4)$$

The coefficients $(b_j)_{n,s}$ capture the normalization of the bulk-defect two-point function $\langle\langle \mathcal{O}_J \hat{\mathcal{O}}_{n,s} \rangle\rangle$, where the exchanged operators $\hat{\mathcal{O}}_{n,s}$ have dimension $\hat{\Delta}_{n,s} = J + s + 2n + \gamma_{n,s}$ and transverse-spin s .⁹ In the case where (small) anomalous dimensions are relevant, second-order poles are also present in the formula when Taylor expanding:

$$b_j(\hat{\Delta}, s) = - \sum_{n \geq 0} \left(\frac{(b_j^2)_{n,s}}{\hat{\Delta} - (J + s + 2n)} + \frac{(b_j^2 \gamma_j)_{n,s}}{(\hat{\Delta} - (J + s + 2n))^2} + \dots \right). \quad (3.5)$$

Once the OPE coefficients and the anomalous dimensions have been obtained, the correlator can be expanded in defect spacetime blocks:

$$\tilde{F}_j(z, \bar{z}) = \sum_{s=0}^{\infty} \sum_{n=0}^{\infty} \left((b_j^2)_{n,s} \hat{f}_{J+s+2n,s}(z, \bar{z}) + (b_j^2 \gamma_j)_{n,s} \partial_{\hat{\Delta}} \hat{f}_{J+s+2n,s}(z, \bar{z}) \right). \quad (3.6)$$

Note that we have introduced the notation \tilde{F}_j instead of F_j , since the inversion formula might miss contributions from low spins $s < s_*$ [27]. In that case, we must add extra terms with spins $s = 0, 1, \dots, s_*$ to \tilde{F}_j in order to recover the full correlator F_j . This procedure will be described in detail in section 3.4. Except for these subtleties, equations (3.2)-(3.6) reconstruct the function F_j using only information in its discontinuity $\text{Disc } F_j$.

3.2 Computation of the discontinuity

The first step in order to apply the inversion formula (3.2) is to compute the discontinuity. As we now show, the discontinuity can be computed even though the full correlator is not known.

The idea is to expand the correlation functions in the bulk channel. The conformal blocks in (A.1) can be written as $f_{\Delta,\ell}(z, \bar{z}) = [(1-z)(1-\bar{z})]^{(\Delta-\ell)/2} \tilde{f}_{\Delta,\ell}(z, \bar{z})$, where the function $\tilde{f}_{\Delta,\ell}(z, \bar{z})$ has an expansion around $z, \bar{z} = 1$ in positive integer powers. As a result, only the prefactor can have non-vanishing discontinuity. Therefore, the contribution of a single bulk operator \mathcal{O} to the discontinuity is

$$\text{Disc } F_j(z, \bar{z})|_{\mathcal{O}} \propto (z\bar{z})^{J/2} (1-z)^{\frac{\Delta-(2J+\ell)}{2}} \tilde{f}_{\Delta,\ell}(z, \bar{z}) \text{Disc} \left[(1-\bar{z})^{\frac{\Delta-(2J+\ell)}{2}} \right]. \quad (3.7)$$

There are two situations when this discontinuity does not vanish:

⁹In general the defect spectrum contains degeneracies, in which case the CFT data has to be understood as a sum over degenerate operators.

1. If Δ is non-integer. This corresponds to \mathcal{O} having an anomalous dimension correcting its tree-level dimension $\Delta = 2J + \ell + 2n + \gamma$.
2. If Δ is integer but $\Delta < 2J + \ell$. This corresponds to \mathcal{O} being a protected single-trace operator, whose dimension is below the double-trace threshold. Note that even though $\text{Disc}(1 - \bar{z})^{-n}$ naively vanishes, for $n > 0$ the singularity at $\bar{z} = 1$ gives a finite contribution to the inversion formula.¹⁰

In our setup the discontinuity only receives contributions of the second type. This claim can be proved by studying the superconformal bulk OPE in detail. From now on we focus on the $J = 2$ case. It was shown in [37] that in the presence of the line defect the OPE $\mathcal{O}_2 \times \mathcal{O}_2$ truncates in the following way:

$$\mathcal{O}_2 \times \mathcal{O}_2 \rightarrow \mathbb{1} + \mathcal{B}_{[0,2,0]} + \mathcal{B}_{[0,4,0]} + \sum_{\ell} \mathcal{C}_{[0,2,0],\ell} + \sum_{\Delta,\ell} \mathcal{A}_{[0,0,0],\ell}^{\Delta}. \quad (3.8)$$

These representations correspond to the operators acquiring a non-vanishing one-point function in the presence of a half-BPS line defect (like the supersymmetric Wilson line we study in this work). The operators in the $\mathcal{B}_{[0,2,0]}$ multiplet have integer dimension $\Delta < 4 + \ell$, so they have non-vanishing discontinuity. The operators in the $\mathcal{B}_{[0,4,0]}$, $\mathcal{C}_{[0,2,0],\ell}$ multiplets have integer dimension $\Delta \geq 4 + \ell$, so they cannot contribute to the discontinuity. Only the unprotected multiplets $\mathcal{A}_{[0,0,0],\ell}^{\Delta}$ can have anomalous dimensions. The scaling dimensions of these multiplets have the following schematic structure [49]

$$\Delta = 2J + 2n + \ell + \frac{1}{N^2} \left(a + \frac{b}{\lambda^{3/2}} + \dots \right) + O(N^{-4}), \quad (3.9)$$

which means anomalous dimensions do not contribute at the order in perturbation theory we are working. This implies all the operators in the bulk OPE (3.8) do not have anomalous dimensions at this order, so the correlator must admit an expansion in integer powers in the limit $z, \bar{z} \rightarrow 1$.

The main consequence of these observations is that only the superblock $\mathcal{G}_{[0,2,0]}$ corresponding to $\mathcal{B}_{[0,2,0]}$ has non-vanishing discontinuity:

$$\text{Disc } \mathcal{F}^{(2)}(z, \bar{z}, \sigma) \Big|_{O(\frac{\sqrt{\lambda}}{N^2})} = \lambda_{222} a_2 \text{Disc} \left(\frac{\sqrt{z\bar{z}} \sigma}{(1-z)(1-\bar{z})} \right)^2 \mathcal{G}_{[0,2,0]}(z, \bar{z}, \sigma). \quad (3.10)$$

In the rest of this section we reconstruct the full correlator from the single superblock $\mathcal{G}_{[0,2,0]}$.

3.3 Inversion of $\mathcal{B}_{[0,2,0]}$

We now invert the superblock $\mathcal{G}_{[0,2,0]}$ in order to extract the defect CFT data, and by resumming the defect expansions, we obtain the correlators given in (3.23), (3.25) and (3.26).

¹⁰This will be proved concretely by deriving equation (3.16).

The superblocks $\mathcal{G}_{[0,K,0]}$ are known and given in equation (4.4). For $K = 2$ they take the form

$$\mathcal{G}_{[0,2,0]}(z, \bar{z}, \sigma) = h_2(\sigma) f_{2,0}(z, \bar{z}) + \frac{1}{180} h_0(\sigma) f_{4,2}(z, \bar{z}), \quad (3.11)$$

where $h_K(\sigma)$ and $f_{\Delta,\ell}(z, \bar{z})$ correspond respectively to R -symmetry and spacetime conformal blocks, which can be found in appendix A. We see that we need to invert two bosonic blocks, namely a scalar block $f_{2,0}$ as well as the stress-tensor block $f_{4,2}$. Using the definition (A.5) for the R -symmetry blocks, we can extract the discontinuities in the three channels:

$$\begin{aligned} \text{Disc } F_0(z, \bar{z}) &= -\lambda_{222} a_2 \text{Disc} \frac{z\bar{z}}{180(1-z)^2(1-\bar{z})^2} (30f_{2,0}(z, \bar{z}) - f_{4,2}(z, \bar{z})), \\ \text{Disc } F_1(z, \bar{z}) &= \lambda_{222} a_2 \text{Disc} \frac{z\bar{z}}{(1-z)^2(1-\bar{z})^2} f_{2,0}(z, \bar{z}), \\ \text{Disc } F_2(z, \bar{z}) &= 0. \end{aligned} \quad (3.12)$$

It is convenient to express the blocks using the following variable:

$$\bar{y} := \frac{1 - \bar{z}}{\sqrt{\bar{z}}}. \quad (3.13)$$

As discussed in the previous section, only negative powers of \bar{y} are relevant for the discontinuity. Since a factor \bar{y}^{-2} comes from the prefactor in (3.12), we have to expand the bulk blocks to order $O(\bar{y})$ as $\bar{y} \rightarrow 0$. Using the methods in appendix A.1 we find

$$\begin{aligned} f_{2,0}(z, \bar{z}) &= -\bar{y} \log z + O(\bar{y}^3), \\ f_{4,2}(z, \bar{z}) &= \bar{y} \left(\frac{90(z+1)}{z-1} - \frac{30(z^2+4z+1) \log z}{(z-1)^2} \right) + O(\bar{y}^3). \end{aligned} \quad (3.14)$$

Let us now invert an arbitrary power \bar{y}^{-p} . In this section we only need the $p = 1$ case, but the case $p \geq 2$ is relevant for section 4. For arbitrary powers of \bar{y} the discontinuity (3.3) results in:

$$\text{Disc } \bar{y}^{-p} = 2i \sin(p\pi) (-\bar{y})^{-p}. \quad (3.15)$$

In principle now we should compute the inversion integral (3.2). In practice this is too hard, so we expand the integrand as $z \rightarrow 0$.¹¹ Each new power of z will give information of a new defect family with higher transverse-twist $\hat{\Delta} - s$. In the $z \rightarrow 0$ limit the integral over \bar{z} is standard and gives the following result:

$$B_p(\beta) = 2i \sin(\pi p) \int_1^\infty \frac{d\bar{z}}{2\pi i} \bar{z}^{-\beta/2-1} (-\bar{y})^{-p} = \frac{\Gamma\left(\frac{\beta+p}{2}\right)}{\Gamma(p)\Gamma\left(\frac{\beta-p+2}{2}\right)}. \quad (3.16)$$

Although for integer $p > 0$ the discontinuity (3.15) naively vanishes, note that the final result is perfectly finite. This is expected, because the correlator is singular at $\bar{z} \rightarrow 1$, and the inversion formula reconstructs the CFT data from this singularity. Note that for $p = 1$ then $B_1(\beta) = 1$, which simplifies our calculations below.

¹¹As a side effect, the expansion as $z \rightarrow 0$ makes the inversion integral convergent order by order in z .

In general the integral over z has the following structure:

$$\begin{aligned}
b(\hat{\Delta}, s) &= \sum_{n \geq 0} \int_0^1 \frac{dz}{2z} z^{-\frac{\hat{\Delta} - (2+s+2n)}{2}} \left[b^{(0,n)}(\hat{\Delta}, s) + b^{(1,n)}(\hat{\Delta}, s) \log z \right] \\
&= - \sum_{n \geq 0} \left(\frac{b^{(0,n)}(\hat{\Delta}, s) + 2\partial_{\hat{\Delta}} b^{(1,n)}(\hat{\Delta}, s)}{\hat{\Delta} - (2 + s + 2n)} + \frac{2b^{(1,n)}(\hat{\Delta}, s)}{(\hat{\Delta} - (2 + s + 2n))^2} + \dots \right)_{\hat{\Delta}=2+s+2n}.
\end{aligned} \tag{3.17}$$

This has to be compared to equation (3.5) in order to obtain the OPE coefficients as well as the product of anomalous dimensions with tree-level OPE coefficients. The presence of logs in equation (3.14) reveals that the scaling dimensions of the defect operators receive anomalous corrections at this order.

In principle the results above are sufficient for extracting the defect CFT data in an algorithmic way and for resumming the correlator using equation (3.6). However, for the $p = 1$ case we can derive a closed-form formula for the defect CFT data corresponding to the bosonic blocks of equation (3.11). In the following we denote by $b_{f_{\Delta,\ell}}(\hat{\Delta}, s)$ the result of the inversion formula performed for individual bosonic spacetime blocks $f_{\Delta,\ell}$. We begin with the scalar block $f_{2,0}$. Using the inversion formula as well as the integrals (3.16) and (3.17) we find

$$\begin{aligned}
b_{f_{2,0}}(\hat{\Delta}, s) &= - \sum_{j,k \geq 0} \frac{(s+1)_j (1/2)_j (1-\hat{\Delta})_k (1/2)_k}{j!(s+3/2)_j k!(3/2-\hat{\Delta})_k} \int_0^1 \frac{dz}{2z} z^{-\frac{\hat{\Delta}-s-2}{2}} z^{j+k} \log z \\
&= - \sum_{j,k \geq 0} \frac{(s+1)_j (1/2)_j (1-\hat{\Delta})_k (1/2)_k}{j!(s+3/2)_j k!(3/2-\hat{\Delta})_k} \frac{-2}{(\hat{\Delta} - s - 2(j+k+1))^2},
\end{aligned} \tag{3.18}$$

where we have expanded the hypergeometric functions of equation (3.2). There are only second-order poles present, thus only the coefficients $b^{(1,n)}$ are non-trivial in equation (3.17). The infinite sum can be obtained in closed form

$$\begin{aligned}
b_{f_{2,0}}^{(0,n)}(\hat{\Delta}, s) &= 0, \\
b_{f_{2,0}}^{(1,n)}(\hat{\Delta}, s) &= \mathcal{C}^{(n)}(\hat{\Delta}, s),
\end{aligned} \tag{3.19}$$

where

$$\mathcal{C}^{(n)}(\hat{\Delta}, s) = - \frac{\Gamma(n + \frac{1}{2}) (\hat{\Delta} - n)_n}{n! \sqrt{\pi} \left(\hat{\Delta} - n - \frac{1}{2} \right)_n} {}_4F_3 \left(\frac{1}{2}, -n, s+1, \hat{\Delta} - n - \frac{1}{2}; 1 \right). \tag{3.20}$$

The calculation of the stress-tensor block proceeds in an analogous way. Using the integrals given above we obtain

$$\begin{aligned}
b_{f_{4,2}}^{(0,n)}(\hat{\Delta}, s) &= 90\mathcal{C}^{(n)}(\hat{\Delta}, s) + 180 \sum_{m=1}^n \mathcal{C}^{(n-m)}(\hat{\Delta}, s), \\
b_{f_{4,2}}^{(1,n)}(\hat{\Delta}, s) &= 30\mathcal{C}^{(n)}(\hat{\Delta}, s) + 180 \sum_{m=1}^n m \mathcal{C}^{(n-m)}(\hat{\Delta}, s).
\end{aligned} \tag{3.21}$$

Putting everything together, with the relative coefficients given by equation (3.12), the bosonic CFT data for $\tilde{F}_0(z, \bar{z})$ reads

$$\begin{aligned} (b_0^2)_{n,s} &= \frac{1}{180} \lambda_{222} a_2 \left(b_{f_{4,2}}^{(0,n)}(\hat{\Delta}, s) - 60 \partial_{\hat{\Delta}} b_{f_{2,0}}^{(1,n)}(\hat{\Delta}, s) + 2 \partial_{\hat{\Delta}} b_{f_{4,2}}^{(1,n)}(\hat{\Delta}, s) \right)_{\hat{\Delta}=J+s+2n}, \\ (b_0^2 \gamma_0)_{n,s} &= -\frac{1}{90} \lambda_{222} a_2 \left(30 b_{f_{2,0}}^{(1,n)}(\hat{\Delta}, s) - b_{f_{4,2}}^{(1,n)}(\hat{\Delta}, s) \right)_{\hat{\Delta}=J+s+2n}. \end{aligned} \quad (3.22)$$

This can be used for resumming the correlator as in equation (3.6). The result takes a very simple form:

$$\tilde{F}_0(z, \bar{z}) = -\lambda_{222} a_2 \frac{z\bar{z}}{2(1-z)(1-\bar{z})} \left[\frac{1+z\bar{z}}{(1-z\bar{z})^2} + \frac{2z\bar{z} \log z\bar{z}}{(1-z\bar{z})^3} \right]. \quad (3.23)$$

The same analysis can be performed for $\tilde{F}_1(z, \bar{z})$, for which we find the following bosonic CFT data

$$\begin{aligned} (b_1^2)_{n,s} &= 2\lambda_{222} a_2 \partial_{\hat{\Delta}} b_{f_{2,0}}^{(1,n)}(\hat{\Delta}, s), \\ (b_1^2 \gamma_1)_{n,s} &= 2\lambda_{222} a_2 b_{f_{2,0}}^{(1,n)}(\hat{\Delta}, s), \end{aligned} \quad (3.24)$$

and the resummation gives a compact expression:

$$\tilde{F}_1(z, \bar{z}) = -\lambda_{222} a_2 \frac{z\bar{z} \log z\bar{z}}{(1-z)(1-\bar{z})(1-z\bar{z})}. \quad (3.25)$$

Finally, since the discontinuity of $F_2(z, \bar{z})$ vanishes (3.12), we simply find

$$\tilde{F}_2(z, \bar{z}) = 0. \quad (3.26)$$

3.4 Supersymmetrization of the correlator

The correlation function obtained in the previous section is *not* supersymmetric, i.e. the three R -symmetry channels given in (3.23), (3.25) and (3.26) do not respect the Ward identities given in (2.18). This happens because the inversion formula misses contributions from low-lying spins $s \leq s_*$ as anticipated in section 3.1.¹² The value of s_* is related to the behavior of the two-point function in the Regge limit $z/\bar{z} \rightarrow 0$ [27], and in principle s_* can be determined by careful analysis of the corresponding Witten diagrams. Instead, in the present work we use the heuristic that s_* should take the minimal value that generates a supersymmetric correlator. As we show below, the resulting correlators make predictions which are in perfect agreement with the expectations from the topological sector.

As we just argued, in order to obtain a supersymmetric correlator, we add defect families with operators of dimensions $\hat{\Delta} = 0, 1, 2, \dots$ and low spin $s \leq s_*$. The OPE coefficients of these operators are unknowns that we fix by imposing the Ward identities (2.18). We have found experimentally that the minimal ansatz consists on taking $s_* = 0$

¹²Such a phenomenon has already been observed for bulk correlators. For example, in the bootstrap of the Wilson-Fisher fixed point there is an ambiguity captured by a single $\ell = 0$ block [50, 51]. In supersymmetric theories, one expects the inversion formula to converge better than in non-supersymmetric ones, see [52] for a recent discussion.

for $\tilde{F}_1(z, \bar{z})$ and $s_* = 1$ for $\tilde{F}_2(z, \bar{z})$. To be precise, we define the final correlators $F_j(z, \bar{z})$ as

$$\begin{aligned} F_0(z, \bar{z}) &= \tilde{F}_0(z, \bar{z}), \\ F_1(z, \bar{z}) &= \tilde{F}_1(z, \bar{z}) + \sum_{n=0}^{\infty} \left(k_n \hat{f}_{n,0}(z, \bar{z}) + p_n \partial_{\hat{\Delta}} \hat{f}_{n,0}(z, \bar{z}) \right), \\ F_2(z, \bar{z}) &= \tilde{F}_2(z, \bar{z}) + \sum_{s=0,1} \sum_{n=0}^{\infty} \left(q_{n,s} \hat{f}_{n+s,s}(z, \bar{z}) + r_{n,s} \partial_{\hat{\Delta}} \hat{f}_{n+s,s}(z, \bar{z}) \right). \end{aligned} \quad (3.27)$$

As mentioned before, the free coefficients k_n , p_n , $q_{n,s}$ and $r_{n,s}$, can be fixed by requiring that the Ward identities are satisfied. In fact this fixes all the coefficients in terms of $q_{0,0}$ and k_1 . Note that $q_{0,0}$ corresponds to the ambiguity $f_{0,0}(z, \bar{z}) = 1$, i.e. the defect identity. However, we know from the Witten diagrams analysis of section 2.2 that the defect identity is given by the constant contribution a_2^2 , and thus

$$q_{0,0} = a_2^2 \Big|_{O(\frac{\sqrt{\lambda}}{N^2})}. \quad (3.28)$$

On the other hand, the unknown k_1 can be determined by demanding a bulk expansion that is consistent with the observations made in section 3.2, i.e. there should not appear anomalous dimensions for bulk operators. This means that the expansion of (3.27) in the limit $z, \bar{z} \rightarrow 1$ should take the form of a power series, without spurious $\log(1 - \bar{z})$ terms. Since the defect expansion (3.27) is natural around $z, \bar{z} \sim 0$, this is only possible after fixing the free coefficients and resumming the correlator. We were able to do so, and the remaining spurious term reads:

$$F_1(z, \bar{z}) \sim \frac{1}{2} (\lambda_{222} a_2 - k_1) \log(1 - \bar{z}) + \dots \quad (3.29)$$

This fixes the coefficient k_1 to be:

$$k_1 = \lambda_{222} a_2. \quad (3.30)$$

3.5 Final result and comparison to localization

We will now present the final result for the correlator $\langle\langle \mathcal{O}_2 \mathcal{O}_2 \rangle\rangle$ using the input of localization for the two remaining free coefficients, namely a_2^2 and $\lambda_{222} a_2$. We can then obtain OPE coefficients of other protected operators which in turn can be checked against the localization data.

The constant contribution a_2^2 from the defect identity can be fixed using equation (2.25):

$$a_2^2 = \frac{\lambda}{N^2} \left(\frac{1}{8} - \frac{3}{8\sqrt{\lambda}} + \dots \right) \quad (3.31)$$

while for the OPE coefficient $\lambda_{222} a_2$ we use the localization results given in equation (2.23) and (2.25):

$$\lambda_{222} a_2 = \frac{\lambda}{N^2} \left(\frac{1}{\sqrt{\lambda}} + \dots \right). \quad (3.32)$$

We thus obtain a correlator without any free coefficient left:

$$\begin{aligned}
F_0(z, \bar{z}) &= -\frac{\sqrt{\lambda}}{2N^2} \frac{z\bar{z}}{(1-z)(1-\bar{z})} \left[\frac{1+z\bar{z}}{(1-z\bar{z})^2} + \frac{2z\bar{z} \log z\bar{z}}{(1-z\bar{z})^3} \right], \\
F_1(z, \bar{z}) &= \frac{\sqrt{\lambda}}{N^2} \left[\log(1+\sqrt{z\bar{z}}) + \frac{z\bar{z}}{(1-z\bar{z})^2} \right. \\
&\quad \left. + \frac{z\bar{z}(5z\bar{z} - 2z^2\bar{z}^2 + z^3\bar{z}^3 - (z+\bar{z})(2-z\bar{z} + z^2\bar{z}^2)) \log z\bar{z}}{2(1-z)(1-\bar{z})(1-z\bar{z})^3} \right], \\
F_2(z, \bar{z}) &= \frac{\sqrt{\lambda}}{8N^2} \left[-3 - \frac{2(z+\bar{z})}{\sqrt{z\bar{z}}} + \frac{(z+\bar{z})(1+z\bar{z}) - 4z\bar{z}}{(1-z\bar{z})^2} \right. \\
&\quad \left. + \frac{2((z+\bar{z})(1+z\bar{z}) - 4z\bar{z}) \log(1+\sqrt{z\bar{z}})}{z\bar{z}} \right. \\
&\quad \left. + \frac{z\bar{z}((z+\bar{z})(3-2z\bar{z} + z^2\bar{z}^2) - 6 + 6z\bar{z} - 4z^2\bar{z}^2) \log z\bar{z}}{(1-z\bar{z})^3} \right]. \tag{3.33}
\end{aligned}$$

Comparing to (A.8), this correlator predicts the OPE coefficient of the double-trace operator $\mathcal{O}_{(2,2)}$

$$\lambda_{22(2,2)} a_{(2,2)} = \frac{\lambda}{N^2} \left(\frac{1}{8} - \frac{1}{2\sqrt{\lambda}} + \dots \right), \tag{3.34}$$

which matches the localization results given in equation (2.27). We can also extract the defect CFT data for the protected operators:

$$\begin{aligned}
b_{21}^2 &= \frac{\lambda}{N^2} \left(\frac{1}{\sqrt{\lambda}} + \dots \right), \\
b_{2(2)}^2 &= 1 + \frac{\lambda}{N^2} \left(-\frac{1}{2\sqrt{\lambda}} + \dots \right). \tag{3.35}
\end{aligned}$$

The OPE coefficient b_{21}^2 can be compared to the direct computation (see equation (2.31)), and we find a perfect match. The OPE coefficient $b_{2(2)}^2$ corresponds to the operator $\hat{\mathcal{O}}_{(2)}$ introduced in equation (2.29) and is a prediction from our result.¹³

Moreover, using the superblocks described in [13, 37], the correlator above can also be used for extracting the supersymmetric CFT data for unprotected operators. The resulting CFT data has to be interpreted as a sum over degenerate operators, and one would need to solve a mixing problem similar to the case of $\mathcal{N} = 4$ SYM without defects [6, 53–55]. Below we provide a few examples, while we postpone the full analysis of the CFT data and the mixing problem to future work. In particular, the product of tree-level coefficients and anomalous dimensions for the unprotected operators at lowest twist $\hat{\Delta} = s + 2$ reads

$$\hat{F}_{s+2,s} \gamma_{s+2,s} \Big|_{\mathcal{O}(\frac{\sqrt{\lambda}}{N^2})} = -\frac{3+2s}{2(1+s)} \frac{\sqrt{\lambda}}{N^2}. \tag{3.36}$$

¹³The observation that $\hat{\mathcal{O}}_{(2)}$ should appear in this type of correlator was first discussed in appendix A of [45]. In principle the operator $\hat{\mathcal{O}}_2$ should also appear, but it can be seen from equation (2.31) that it is not relevant at the present order.

Note that here we use the notation of [13]. It is also possible to obtain a closed form for the OPE coefficients of the semishort operators $(B, 1)_{[1,s]}$:

$$\hat{E}_s \Big|_{O(\frac{\sqrt{\lambda}}{N^2})} = -\frac{1+s}{4(1+2s)} \frac{\sqrt{\lambda}}{N^2}. \quad (3.37)$$

4 General identical operators

In this section we extend the analysis of the previous section for general identical operators $\langle\langle \mathcal{O}_J \mathcal{O}_J \rangle\rangle$. The calculation of $\langle\langle \mathcal{O}_2 \mathcal{O}_2 \rangle\rangle$ carries through almost unchanged, as will be described shortly. As a concrete application we obtain closed-form expressions for the $J = 3, 4$ correlators.

4.1 General discussion

As discussed in section 2, the correlator of interest has the form

$$\mathcal{F}^{(J)}(z, \bar{z}, \sigma) = \left(\frac{\sigma \sqrt{z\bar{z}}}{(1-z)(1-\bar{z})} \right)^J + \frac{J\lambda}{2^{J+2}N^2} + \sum_{j=0}^J \sigma^j F_{J-j}^{(J)}(z, \bar{z}) + O\left(\frac{1}{N^2}\right), \quad (4.1)$$

where we used the leading-order result for the one-point function (2.25). Here we give a general prescription to obtain the functions $F_j^{(J)}(z, \bar{z})$ that contribute at order $\frac{\sqrt{\lambda}}{N^2}$.

The central idea is to reconstruct these functions using the Lorentzian inversion formula (3.2). It was discussed in section 3.2 that only operators with single-trace dimension can contribute to the discontinuity of the correlator. The bulk OPE of \mathcal{O}_J takes the form [37]

$$\mathcal{O}_J \times \mathcal{O}_J \sim \mathbf{1} + \sum_{k=1}^J \mathcal{B}_{[0,2k,0]} + \dots, \quad (4.2)$$

where \dots contains unprotected multiplets that do not contribute to the discontinuity. Furthermore, $\mathcal{B}_{[0,2J,0]}$ has double-twist dimension and does not contribute. From this it follows that, at the order we are working, the discontinuity of the correlator reads

$$\text{Disc } \mathcal{F}^{(J)}(z, \bar{z}, \sigma) \Big|_{O(\frac{\sqrt{\lambda}}{N^2})} = \text{Disc} \left[\left(\frac{\sigma \sqrt{z\bar{z}}}{(1-z)(1-\bar{z})} \right)^J \sum_{k=1}^{J-1} \lambda_{JJ2k} a_{2k} \mathcal{G}_{[0,2k,0]}(z, \bar{z}, \sigma) \right]. \quad (4.3)$$

The superconformal blocks capture the information of half-BPS operators exchanged in the bulk OPE $\mathcal{O}_J \times \mathcal{O}_J$. They were obtained in [13], and we reproduce them here for simplicity:

$$\begin{aligned} \mathcal{G}_{[0,K,0]}(z, \bar{z}, \sigma) &= h_K(\sigma) f_{K,0}(z, \bar{z}) + \frac{(K+2)^2 K}{128(K+1)^2(K+3)} h_{K-2}(\sigma) f_{K+2,2}(z, \bar{z}) \\ &+ \frac{(K-2)(K+2)K^2}{16384(K-1)^2(K+1)(K+3)} h_{K-4}(\sigma) f_{K+4,0}(z, \bar{z}). \end{aligned} \quad (4.4)$$

Using (4.3) as input to the inversion formula, it is possible to generate a series representation of the correlators $\tilde{F}_j^{(J)}(z, \bar{z})$. In concrete examples, these series expansions do not satisfy

the Ward identities, just as we saw for the $\langle\langle \mathcal{O}_2 \mathcal{O}_2 \rangle\rangle$ case. This is a result of the inversion formula not converging for low values of the transverse spin s . Empirically we have found that the minimal set of additions is

$$\begin{aligned}
F_j^{(J)}(z, \bar{z}) &= \tilde{F}_j^{(J)}(z, \bar{z}) \quad \text{for } j = 0, \dots, J-2, \\
F_{J-1}^{(J)}(z, \bar{z}) &= \tilde{F}_{J-1}^{(J)}(z, \bar{z}) + \sum_{n=0}^{\infty} \left(k_n \hat{f}_{n,0}(z, \bar{z}) + p_n \partial_{\hat{\Delta}} \hat{f}_{n,0}(z, \bar{z}) \right), \\
F_J^{(J)}(z, \bar{z}) &= \tilde{F}_J^{(J)}(z, \bar{z}) + \sum_{s=0,1} \sum_{n=0}^{\infty} \left(q_{n,s} \hat{f}_{n+s,s}(z, \bar{z}) + r_{n,s} \partial_{\hat{\Delta}} \hat{f}_{n+s,s}(z, \bar{z}) \right).
\end{aligned} \tag{4.5}$$

Namely, we must add $s = 0$ ambiguities to $F_{J-1}^{(J)}(z, \bar{z})$, while we must add $s = 0, 1$ ambiguities to $F_J^{(J)}(z, \bar{z})$. Using the Ward identities (2.18), it is possible to fix all the free coefficients $k_n, p_n, q_{n,s}, r_{n,s}$ except for $q_{0,0}$ and k_1 . As discussed around equation (3.28), $q_{0,0}$ can be identified with the defect identity a_J^2 and is therefore fixed from localization. Furthermore, if one keeps the bulk OPE coefficients $\lambda_{JK} a_K$ arbitrary, the Ward identities also fix their relative values. For example, the Ward identities together with (4.5) imply

$$\lambda_{JK} a_K \Big|_{O(\frac{\sqrt{\lambda}}{N^2})} = \frac{K}{2^{K/2}} \lambda_{JJ} a_2 \Big|_{O(\frac{\sqrt{\lambda}}{N^2})} \quad \text{for } 2 \leq K \leq J-2. \tag{4.6}$$

These results are in perfect agreement with the known values (2.25) at leading order at large λ . This provides evidence that (4.5) is the correct prescription for the low-spin additions.

At this point, it is possible to resum the series expansion representations for $F_j^{(J)}$. These resummed correlators can be expanded in a series around $z, \bar{z} = 1$, which corresponds to the bulk conformal block decomposition. The expansion contains spurious $\log(1 - \bar{z})$ terms, which would imply that bulk operators get anomalous dimensions. These anomalous dimensions should not be present at the order we are working, so canceling the spurious logarithms fixes the remaining free parameter k_1 . Finally, one uses a_J^2 and $\lambda_{JJ} a_2$ coming from localization to write down the final correlator. This correlation function produces all other OPE coefficients in the protected sector, and we find perfect agreement with the literature.

4.2 Example 1: $\langle\langle \mathcal{O}_3 \mathcal{O}_3 \rangle\rangle$

Let us turn our attention to the case of $\langle\langle \mathcal{O}_3 \mathcal{O}_3 \rangle\rangle$. The purpose of this example is to illustrate the general formalism that we just discussed. Furthermore, we present concrete intermediate results to help the readers interested in reproducing our results.

4.2.1 Inversion of single traces

The first step is to obtain the discontinuity of the correlator, by combining (4.3) and (4.4) with (A.2). For concreteness we look at the coefficient of σ^3 , or equivalently we focus on

the $F_0^{(3)}(z, \bar{z})$ correlator:

$$\begin{aligned} \text{Disc } F_0^{(3)}(z, \bar{z}) &= \lambda_{332} a_2 \frac{z^{3/2} (z^2 - 2z \log z - 1)}{2(1-z)^5} \text{Disc } \frac{1}{\bar{y}^2} \\ &\quad - \lambda_{334} a_4 \frac{z^{3/2} (z^3 + 9z^2 - 9z - 1 - 6z(z+1) \log z)}{4(1-z)^6} \text{Disc } \frac{1}{\bar{y}}. \end{aligned} \quad (4.7)$$

Similar expressions can be easily obtained for the other $F_j^{(3)}(z, \bar{z})$. The next step is to insert the discontinuity in the inversion formula (3.2), and carry out the inversion order by order as $z \rightarrow 0$. Each power z^p induces a defect family with blocks $\hat{f}_{2p+s+2n,s}(z, \bar{z})$ and OPE coefficients given by the poles of $b(\hat{\Delta}, s)$. Similarly, each power $z^p \log z$ induces defect anomalous dimensions $\partial_{\hat{\Delta}} \hat{f}_{2p+s+2n,s}(z, \bar{z})$ given by the double poles of $b(\hat{\Delta}, s)$. All the integrals that are needed are of the form (3.16) with $p = 1, 2$. The resulting defect expansion for the lowest-lying operators is

$$\begin{aligned} \tilde{F}_0^{(3)}(z, \bar{z}) &= \frac{1}{2} (3\lambda_{332} a_2 - \lambda_{334} a_4) \hat{f}_{3,0}(z, \bar{z}) + \frac{1}{2} (5\lambda_{332} a_2 - \lambda_{334} a_4) \hat{f}_{4,1}(z, \bar{z}) \\ &\quad + \frac{2}{21} (121\lambda_{332} a_2 - 73\lambda_{334} a_4) \hat{f}_{5,0}(z, \bar{z}) + 2(5\lambda_{332} a_2 - 3\lambda_{334} a_4) \partial_{\hat{\Delta}} \hat{f}_{5,0}(z, \bar{z}) \\ &\quad + \frac{1}{2} (7\lambda_{332} a_2 - \lambda_{334} a_4) \hat{f}_{5,2}(z, \bar{z}) + \dots \end{aligned} \quad (4.8)$$

Once again, the expansions for other $\tilde{F}_j^{(3)}(z, \bar{z})$ are obtained in an identical manner. Unlike in the $\langle\langle \mathcal{O}_2 \mathcal{O}_2 \rangle\rangle$ case, we were not able to express the defect CFT data in closed forms. However, the previous calculation can be automatized with a computer, and the expansion can be generated efficiently up to high orders.

4.2.2 Supersymmetrization of the correlator

As for the $\langle\langle \mathcal{O}_2 \mathcal{O}_2 \rangle\rangle$ case, the $\tilde{F}_j^{(3)}(z, \bar{z})$ do not give a supersymmetric correlator because the Lorentzian inversion formula can miss low transverse-spin contributions. In order for $\tilde{F}_j^{(3)}(z, \bar{z})$ to satisfy the Ward identities (2.18), we add the following $s = 0, 1$ contributions to our correlators

$$\begin{aligned} F_0^{(3)}(z, \bar{z}) &= \tilde{F}_0^{(3)}(z, \bar{z}), \quad F_1^{(3)}(z, \bar{z}) = \tilde{F}_1^{(3)}(z, \bar{z}), \\ F_2^{(3)}(z, \bar{z}) &= \tilde{F}_2^{(3)}(z, \bar{z}) + \sum_{n=0}^{\infty} \left(k_n \hat{f}_{n,0}(z, \bar{z}) + p_n \partial_{\hat{\Delta}} \hat{f}_{n,0}(z, \bar{z}) \right), \\ F_3^{(3)}(z, \bar{z}) &= \tilde{F}_3^{(3)}(z, \bar{z}) + \sum_{s=0,1} \sum_{n=0}^{\infty} \left(q_{n,s} \hat{f}_{n+s,s}(z, \bar{z}) + r_{n,s} \partial_{\hat{\Delta}} \hat{f}_{n+s,s}(z, \bar{z}) \right). \end{aligned} \quad (4.9)$$

After this addition, the functions $F_j^{(3)}(z, \bar{z})$ should satisfy the Ward identities (2.18), which are highly constraining.

In fact, the only coefficients that remain unfixed are $\lambda_{332} a_2$, k_1 and $q_{0,0}$. In particular, notice that the precise relation between single-trace OPE coefficients is fixed:

$$\lambda_{334} a_4 \Big|_{O(\frac{\sqrt{\lambda}}{N^2})} = \lambda_{332} a_2 \Big|_{O(\frac{\sqrt{\lambda}}{N^2})}. \quad (4.10)$$

Reassuringly, this relation is consistent with localization (2.25). Again, we believe this is strong evidence for the ansatz (4.9) for low transverse-spin ambiguities to be correct. In the case of $\langle\langle \mathcal{O}_4 \mathcal{O}_4 \rangle\rangle$ we find a similar result involving $K = 4, 6$, and for general J we expect equation (4.6) to hold.

4.2.3 Fixing free parameters and final result

At this point, we have a supersymmetric correlator depending on three free parameters. Using the series representation of the correlator (4.8) it is possible to find a closed form expression in terms of rational functions and logarithms. In particular, we find

$$F_2^{(3)}(z, \bar{z}) = -\frac{3}{8}(3\lambda_{332}a_2 + 32k_1) \tanh^{-1} \sqrt{z\bar{z}} + \dots, \quad (4.11)$$

where \dots stand for terms which have an expansion around $z, \bar{z} = 1$ involving only integer powers. On the other hand, $\tanh^{-1} \sqrt{z\bar{z}}$ has an expansion with $\log(1 - \bar{z})$ terms. These logarithms correspond to anomalous dimensions in the bulk, which should be absent by our assumptions. We thus conclude

$$k_1 = -\frac{3}{32}\lambda_{332}a_2. \quad (4.12)$$

In order to fix the last two coefficients we use input from the localization results of the topological sector. Indeed, from (2.25) we have that

$$\lambda_{332}a_2 = \frac{\lambda}{N^2} \left(\frac{3}{2\sqrt{\lambda}} + \dots \right), \quad a_3^2 = \frac{\lambda}{N^2} \left(\frac{3}{32} - \frac{3}{4\sqrt{\lambda}} + \dots \right). \quad (4.13)$$

As discussed around equation (3.28), the coefficient $q_{0,0}$ has a natural interpretation as the defect identity contribution given by a_3^2 . We thus conclude $q_{0,0} = -\frac{3\sqrt{\lambda}}{4N^2}$.

The final result with no free parameters takes a reasonably simple form:

$$\begin{aligned} F_0^{(3)}(z, \bar{z}) &= -\frac{6\sqrt{\lambda}}{8N^2} \frac{(z\bar{z})^{3/2}}{[(1-z)(1-\bar{z})]^2} \left[\frac{1+z\bar{z}}{(1-z\bar{z})^2} + \frac{2z\bar{z} \log z\bar{z}}{(1-z\bar{z})^3} \right], \\ F_1^{(3)}(z, \bar{z}) &= \frac{3\sqrt{\lambda}}{4N^2} \frac{(z\bar{z})^{3/2}}{(1-z)(1-\bar{z})(1-z\bar{z})^4} \left(z^2\bar{z}^2 - 38z\bar{z} + 1 \right. \\ &\quad \left. - \frac{2((z+\bar{z})(z\bar{z}+1)(z^2\bar{z}^2 - 11z\bar{z} + 1) + z\bar{z}(z\bar{z}+5)(5z\bar{z}+1)) \log z\bar{z}}{(1-z)(1-\bar{z})(1-z\bar{z})} \right), \\ F_2^{(3)}(z, \bar{z}) &= -\frac{18\sqrt{\lambda}}{N^2} \frac{(z\bar{z})^{1/2}}{(1-z\bar{z})^2} \left(\frac{z\bar{z}}{(1-z)(1-\bar{z})} - \frac{3(1+z\bar{z})(z^2\bar{z}^2 + 6z\bar{z} + 1)}{32(1-z\bar{z})^2} \right. \\ &\quad \left. - \frac{2z\bar{z}(7+10z\bar{z}+7z^2\bar{z}^2) \log z\bar{z}}{32(1-z\bar{z})^3} + \frac{z\bar{z}(1+z\bar{z}) \log z\bar{z}}{2(1-z)(1-\bar{z})(1-z\bar{z})} \right), \\ F_3^{(3)}(z, \bar{z}) &= -\frac{3\sqrt{\lambda}}{4N^2} \left(1 - \frac{3(z\bar{z})^{1/2}((z+\bar{z})(z^2\bar{z}^2 + 10z\bar{z} + 1) - 3(z\bar{z}+1)^3)}{4(1-z\bar{z})^4} \right. \\ &\quad \left. + \frac{3(z\bar{z})^{3/2}(5z^2\bar{z}^2 + 2z\bar{z} + 5 - 3(z+\bar{z})(z\bar{z}+1)) \log z\bar{z}}{2(1-z\bar{z})^5} \right). \end{aligned} \quad (4.14)$$

In principle, this correlation function contains information of infinitely many unprotected operators, but as for the $\langle\langle \mathcal{O}_2 \mathcal{O}_2 \rangle\rangle$ case we leave a detailed analysis of the OPE for future work. Instead, we focus on the CFT data captured by the topological sector. Comparing the full correlator (4.1) to equation (A.8) gives the bulk data

$$\lambda_{334}a_4 = \frac{\lambda}{N^2} \left(\frac{3}{2\sqrt{\lambda}} + \dots \right), \quad \lambda_{33(3,3)}a_{(3,3)} = \frac{\lambda}{N^2} \left(\frac{3}{32} - \frac{9}{8\sqrt{\lambda}} + \dots \right), \quad (4.15)$$

while comparing to (A.10) gives the defect data

$$b_{31}^2 = \frac{\lambda}{N^2} \left(\frac{27}{16\sqrt{\lambda}} + \dots \right), \quad b_{32}^2 = O\left(\frac{\lambda^0}{N^2}\right), \quad b_{3(3)}^2 = 1 + \frac{\lambda}{N^2} \left(-\frac{3}{4\sqrt{\lambda}} + \dots \right). \quad (4.16)$$

These results are in perfect agreement with the literature, see (2.25), (2.27) and (2.31). Remember that $b_{3(3)}$ cannot be compared to (2.31) because the results of [45] apply to the defect operator $\hat{\mathcal{O}}_3$, while our result applies to the defect operator $\hat{\mathcal{O}}_{(3)}$, see the discussion in footnote 13.

4.3 Example 2: $\langle\langle \mathcal{O}_4 \mathcal{O}_4 \rangle\rangle$

Let us finally consider the $\langle\langle \mathcal{O}_4 \mathcal{O}_4 \rangle\rangle$ two-point function. Since the calculation is essentially identical to the previous one, we skip most of the details. However, it is important to note that the following input from localization is necessary:

$$\lambda_{442}a_2 = \frac{\lambda}{N^2} \left(\frac{2}{\sqrt{\lambda}} + \dots \right), \quad a_4^2 = \frac{\lambda}{N^2} \left(\frac{1}{16} - \frac{15}{16\sqrt{\lambda}} + \dots \right). \quad (4.17)$$

Following all the steps, which can be automatized with the help of a computer, we obtain the correlation function

$$\begin{aligned} F_0^{(4)}(z, \bar{z}) &= -\frac{\sqrt{\lambda}}{N^2} \frac{(z\bar{z})^2}{[(1-z)(1-\bar{z})]^3} \left[\frac{1+z\bar{z}}{(1-z\bar{z})^2} + \frac{2z\bar{z} \log z\bar{z}}{(1-z\bar{z})^3} \right], \\ F_1^{(4)}(z, \bar{z}) &= \frac{\sqrt{\lambda}}{N^2} \frac{(z\bar{z})^2}{(1-z)^2(1-\bar{z})^2(1-z\bar{z})^4} \left(z^2\bar{z}^2 - 38z\bar{z} + 1 \right. \\ &\quad \left. - \frac{2((z+\bar{z})(z\bar{z}+1)(z^2\bar{z}^2 - 11z\bar{z} + 1) + z\bar{z}(z\bar{z}+5)(5z\bar{z}+1)) \log z\bar{z}}{(1-z)(1-\bar{z})(1-z\bar{z})} \right), \\ F_2^{(4)}(z, \bar{z}) &= -\frac{3\sqrt{\lambda}}{2N^2} \frac{(z\bar{z})^2}{(1-z)^2(1-\bar{z})^2(1-z\bar{z})^6} \left(\right. \\ &\quad + 2(z+\bar{z})(z\bar{z}+1)(8z^2\bar{z}^2 - 91z\bar{z} + 8) + 2z\bar{z}(43z^2\bar{z}^2 + 214z\bar{z} + 43) \\ &\quad + \frac{(z\bar{z}+1)(z^4\bar{z}^4 + 14z^3\bar{z}^3 + 270z^2\bar{z}^2 + 14z\bar{z} + 1) \log z\bar{z}}{1-z\bar{z}} \\ &\quad \left. + \frac{(z+\bar{z})(7z^4\bar{z}^4 - 46z^3\bar{z}^3 - 222z^2\bar{z}^2 - 46z\bar{z} + 7) \log z\bar{z}}{1-z\bar{z}} \right), \\ F_3^{(4)}(z, \bar{z}) &= -\frac{\sqrt{\lambda}}{2N^2} \left(\frac{270(z\bar{z})^2(z\bar{z}+1)}{(1-z)(1-\bar{z})(1-z\bar{z})^4} - \frac{z\bar{z}(2z^4\bar{z}^4 + 229z^3\bar{z}^3 + 438z^2\bar{z}^2 + 229z\bar{z} + 2)}{(1-z\bar{z})^6} \right. \\ &\quad + \frac{90(z\bar{z})^2(z^2\bar{z}^2 + 4z\bar{z} + 1) \log z\bar{z}}{(1-z)(1-\bar{z})(1-z\bar{z})^5} - 4 \log(1 + \sqrt{z\bar{z}}) \\ &\quad \left. - \frac{(z\bar{z})^2(2z^5\bar{z}^5 - 14z^4\bar{z}^4 + 117z^3\bar{z}^3 + 325z^2\bar{z}^2 + 395z\bar{z} + 75) \log z\bar{z}}{(1-z\bar{z})^7} \right), \end{aligned}$$

$$\begin{aligned}
F_4^{(4)}(z, \bar{z}) = & \frac{\sqrt{\lambda}}{N^2} \left(-\frac{z + \bar{z}}{2\sqrt{z\bar{z}}} + \frac{(z + \bar{z})(z\bar{z} + 1) - 4z\bar{z}}{2z\bar{z}} \log(1 + \sqrt{z\bar{z}}) \right) \\
& + \frac{2(z + \bar{z})(z\bar{z} + 1)(2z^4\bar{z}^4 - 11z^3\bar{z}^3 + 468z^2\bar{z}^2 - 11z\bar{z} + 2)}{16(1 - z\bar{z})^6} \\
& - \frac{15z^6\bar{z}^6 - 74z^5\bar{z}^5 + 1397z^4\bar{z}^4 + 924z^3\bar{z}^3 + 1397z^2\bar{z}^2 - 74z\bar{z} + 15}{16(1 - z\bar{z})^6} \\
& + \frac{(z\bar{z})^2(z + \bar{z})(z^5\bar{z}^5 - 6z^4\bar{z}^4 + 14z^3\bar{z}^3 + 56z^2\bar{z}^2 + 315z\bar{z} + 70) \log z\bar{z}}{4(1 - z\bar{z})^7} \\
& - \frac{(z\bar{z})^2(4z^5\bar{z}^5 - 28z^4\bar{z}^4 + 189z^3\bar{z}^3 + 245z^2\bar{z}^2 + 385z\bar{z} + 105) \log z\bar{z}}{4(1 - z\bar{z})^7}.
\end{aligned} \tag{4.18}$$

Once again, let us compare the predictions of this correlator with the topological sector. We find the bulk data

$$\begin{aligned}
\lambda_{444}a_4 &= \frac{\lambda}{N^2} \left(\frac{2}{\sqrt{\lambda}} + \dots \right), & \lambda_{446}a_6 &= \frac{\lambda}{N^2} \left(\frac{3}{2\sqrt{\lambda}} + \dots \right), \\
\lambda_{44(4,4)}a_{(4,4)} &= \frac{\lambda}{N^2} \left(\frac{1}{16} - \frac{3}{2\sqrt{\lambda}} + \dots \right),
\end{aligned} \tag{4.19}$$

while the defect data is given by

$$\begin{aligned}
b_{41}^2 &= \frac{\lambda}{N^2} \left(\frac{2}{\sqrt{\lambda}} + \dots \right), & b_{42}^2 &= O\left(\frac{\lambda^0}{N^2}\right), \\
b_{43}^2 &= O\left(\frac{\lambda^0}{N^2}\right), & b_{4(4)}^2 &= 1 + \frac{\lambda}{N^2} \left(-\frac{1}{\sqrt{\lambda}} + \dots \right).
\end{aligned} \tag{4.20}$$

These results are in perfect agreement with the literature, except for $b_{4(4)}$, which is a new prediction from our calculation.

4.4 A conjecture

In the three examples considered in the present paper, the function $F_0^{(J)}(z, \bar{z})$ is very simple:

$$F_0^{(J)}(z, \bar{z}) = -\frac{J\sqrt{\lambda}}{4N^2} \frac{(z\bar{z})^{J/2}}{[(1-z)(1-\bar{z})]^{J-1}} \left[\frac{1+z\bar{z}}{(1-z\bar{z})^2} + \frac{2z\bar{z} \log z\bar{z}}{(1-z\bar{z})^3} \right]. \tag{4.21}$$

It is tempting to conjecture that this relation also holds for $J > 4$. From our bootstrap calculation, (4.21) is a result of a precise combination of the spacetime bulk blocks (A.2), the superblocks (4.4) and the OPE coefficients in the topological sector (2.25). Perhaps from the point of view of the explicit holographic calculation the origin of (4.21) will be more transparent.

Let us also note the similarity between $F_1^{(3)}(z, \bar{z})$ and $F_1^{(4)}(z, \bar{z})$

$$\begin{aligned}
F_1^{(J)}(z, \bar{z}) = & \frac{J\sqrt{\lambda}}{4N^2} \frac{(z\bar{z})^{J/2}}{(1-z)^{J-2}(1-\bar{z})^{J-2}(1-z\bar{z})^4} \left(z^2\bar{z}^2 - 38z\bar{z} + 1 \right. \\
& \left. - \frac{2((z + \bar{z})(z\bar{z} + 1)(z^2\bar{z}^2 - 11z\bar{z} + 1) + z\bar{z}(z\bar{z} + 5)(5z\bar{z} + 1)) \log z\bar{z}}{(1-z)(1-\bar{z})(1-z\bar{z})} \right).
\end{aligned} \tag{4.22}$$

We do not have enough data points to propose a full analytic formula for any J , however our current results look promising. In the discussions below we speculate on what might be the best possible strategy for the future.

5 Conclusions

In this work we have studied the structure of two-point functions of single-trace half-BPS operators in the presence of a supersymmetric Wilson line in $\mathcal{N} = 4$ SYM theory. We used analytical bootstrap techniques in order to reconstruct the correlator at strong coupling. For operators of weight $J = 2, 3, 4$ we obtained fairly simple results presented in (3.33), (4.14), (4.18) which only involve logarithms and rational functions of the cross-ratios.

A natural continuation of our work is the analysis of two-point functions for arbitrary weight $\langle\langle \mathcal{O}_{J_1} \mathcal{O}_{J_2} \rangle\rangle$. One obvious approach is to keep pushing the algorithm presented in this paper to higher values of J . However there might be better strategies. In $\mathcal{N} = 4$ SYM without defects, explicit closed form expressions for half-BPS operators take a particularly simple form in Mellin space [7, 8, 56, 57]. It would be interesting to transform the explicit formulas presented in this paper to Mellin variables. The Mellin space approach for defect CFT was explored in [58], the hope being that this might be the natural language to write the most general correlator $\langle\langle \mathcal{O}_{J_1} \mathcal{O}_{J_2} \rangle\rangle$.

Another interesting line of research is to reproduce our results by an explicit holographic calculation using Witten diagrams. Some observables in holographic defect CFTs have been studied perturbatively at strong coupling, for example bulk one-point functions [11, 26], bulk-defect correlators [45] and correlators localized on the defect [28, 29, 36, 59, 60]. The two-point function of bulk operators was studied to order $O(\frac{\lambda}{N^2})$ in [14], and the next order will involve the calculation of the Witten diagram (2.10). The structural understanding presented in this work might give valuable input for this type of calculation.

The idea of reconstructing correlators starting from their discontinuity is powerful. In defect CFT it seems to be as powerful as in the case for homogeneous CFTs. For monodromy defects in the ε -expansion this method was already used to fully bootstrap two-point correlators of chiral fields [61]. In this work we developed an algorithm that in principle can be used to bootstrap an infinite family of half-BPS correlators. We expect that the same method also works in related $\mathcal{N} = 4$ SYM setups, such as the non-supersymmetric Wilson line [59], Wilson lines in more general representations of the gauge group [60], or even higher codimension defects. Furthermore, many half-BPS defects are known to exist in maximally supersymmetric theories in $d = 3, 4, 6$, and all of them might be prime targets for the analytical bootstrap techniques used here.

Acknowledgments

We thank S. Giombi and M. Preti for useful correspondence. We also thank V. Forini, A. Kaviraj, Z. Komargodski, Y. Linke, G. Peveri, J. Plefka, L. Rastelli, J. Rong, and V. Schomerus for discussions and comments. JB's research is funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) - Projektnummer 417533893/GRK2575 "Rethinking Quantum Field Theory". AG and PL are supported by the DFG through the Emmy Noether research group "The Conformal Bootstrap Program" project number 400570283.

A Appendix

A.1 Singular part of bulk blocks

In this appendix, bulk blocks are studied in the limit $\bar{y} = (1 - \bar{z})/\sqrt{\bar{z}} \rightarrow 0$. These results provide the necessary input for the inversion formula in sections 3 and 4. The starting point are the explicit formulas for bulk blocks derived in [62, 63], which we reproduce here for convenience:

$$\begin{aligned}
f_{\Delta,\ell}(z, \bar{z}) &= \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} \frac{4^{m-n} \left(-\frac{\ell}{2}\right)_m^2 \left(\frac{2-\ell-\Delta}{2}\right)_m}{m!n! \left(-\ell\right)_m \left(\frac{3-\ell-\Delta}{2}\right)_m} \frac{\left(\frac{\Delta-1}{2}\right)_n^2 \left(\frac{\Delta+\ell}{2}\right)_n}{(\Delta-1)_n \left(\frac{\Delta+\ell+1}{2}\right)_n} \frac{\left(\frac{\Delta+\ell}{2}\right)_{n-m}}{\left(\frac{\Delta+\ell-1}{2}\right)_{n-m}} \\
&\times (1 - z\bar{z})^{\ell-2m} {}_4F_3 \left(\begin{matrix} -n, -m, \frac{1}{2}, \frac{\Delta-\ell-2}{2} \\ \frac{2-\Delta-\ell-2n}{2}, \frac{\Delta+\ell-2m}{2}, \frac{\Delta-\ell-1}{2} \end{matrix}; 1 \right) \\
&\times [(1-z)(1-\bar{z})]^{\frac{\Delta-\ell}{2}+m+n} {}_2F_1 \left(\begin{matrix} \frac{\Delta+\ell}{2} - m + n, \frac{\Delta+\ell}{2} - m + n \\ \Delta + \ell - 2(m-n) \end{matrix}; 1 - z\bar{z} \right).
\end{aligned} \tag{A.1}$$

After changing variables from $\bar{z} \rightarrow \bar{y}$ and expanding up to order $O(\bar{y}^3)$, we obtain the following formulas:

$$\begin{aligned}
f_{2,0}(z, \bar{z}) &\sim -\bar{y} \log z + \bar{y}^3 \left(\frac{z+1}{8(z-1)} - \frac{z \log z}{4(z-1)^2} \right), \\
f_{4,0}(z, \bar{z}) &\sim \bar{y}^2 \left(-12 + \frac{6(z+1) \log z}{z-1} \right), \\
f_{4,2}(z, \bar{z}) &\sim \bar{y} \left(\frac{90(z+1)}{z-1} - \frac{30(z^2+4z+1) \log z}{(z-1)^2} \right) \\
&\quad + \bar{y}^3 \left(-\frac{15(z+1)(z^2-20z+1)}{8(z-1)^3} - \frac{15z(z^2+7z+1) \log z}{2(z-1)^4} \right), \\
f_{6,0}(z, \bar{z}) &\sim \bar{y}^3 \left(\frac{90(z+1)}{z-1} - \frac{30(z^2+4z+1) \log z}{(z-1)^2} \right), \\
f_{6,2}(z, \bar{z}) &\sim \bar{y}^2 \left(-\frac{140(11z^2+38z+11)}{3(z-1)^2} + \frac{140(z+1)(z^2+8z+1) \log z}{(z-1)^3} \right), \\
f_{8,2}(z, \bar{z}) &\sim \bar{y}^3 \left(\frac{525(z+1)(5z^2+32z+5)}{(z-1)^3} - \frac{630(z^4+16z^3+36z^2+16z+1) \log z}{(z-1)^4} \right).
\end{aligned} \tag{A.2}$$

The symbol \sim is a reminder that these expressions are valid up to corrections of order $O(\bar{y}^4)$. The same calculation can be carried out to higher orders in \bar{y} , as will be necessary to extend the present work to correlators with $J > 4$.

A.2 Conformal block normalization

In this appendix we review our normalization conventions for conformal blocks. Furthermore, we study the contribution of half-BPS operators to the bulk and defect OPEs. This allows to extract topological subsector data from correlation functions, and compare to the predictions from section 2.4.

The bulk-channel conformal block is given explicitly in (A.1). In the bulk OPE limit $z, \bar{z} \rightarrow 1$ it goes like

$$\lim_{z, \bar{z} \rightarrow 1} f_{\Delta, \ell}(z, \bar{z}) = [(1-z)(1-\bar{z})]^{(\Delta-\ell)/2} (1-z\bar{z})^\ell. \quad (\text{A.3})$$

For the defect-channel conformal block we use the same normalization as [27]

$$\hat{f}_{\hat{\Delta}, s}(z, \bar{z}) = z^{\frac{\hat{\Delta}-s}{2}} \bar{z}^{\frac{\hat{\Delta}+s}{2}} {}_2F_1\left(\frac{1}{2}, -s; \frac{1}{2} - s; \frac{z}{\bar{z}}\right) {}_2F_1\left(\frac{1}{2}, \hat{\Delta}; \hat{\Delta} + \frac{1}{2}; z\bar{z}\right). \quad (\text{A.4})$$

The asymptotics in the defect OPE limit $z, \bar{z} \rightarrow 0$ can be easily extracted. Finally, we use the following form of the R -symmetry blocks:

$$\begin{aligned} h_K(\sigma) &= \sigma^{-K/2} {}_2F_1\left(-\frac{K}{2}, -\frac{K}{2}; -K-1; \frac{\sigma}{2}\right), \\ \hat{h}_{\hat{K}}(\sigma) &= \sigma^{\hat{K}} {}_2F_1\left(-\hat{K}-1, -\hat{K}; -2(\hat{K}+1); \frac{2}{\sigma}\right). \end{aligned} \quad (\text{A.5})$$

Using (2.13) and (2.15), it is possible to check that these blocks satisfy the appropriate Casimir equations.

In the discussion of the main text, it is crucial to compare to the predictions in the topological subsector of section 2.4. We start showing how to extract the topological bulk CFT data from our correlators. The choice of normalization in (2.22) fixes uniquely the bulk OPE

$$\mathcal{O}_J(x_1, u_1) \mathcal{O}_J(x_2, u_2) \Big|_{\mathcal{O}_K} = \lambda_{JJK} \frac{(u_1 \cdot u_2)^{J-\frac{K}{2}} (u_1 \cdot D_u^{(6)})^{\frac{K}{2}} (u_2 \cdot D_u^{(6)})^{\frac{K}{2}} \mathcal{O}_K(x_2, u)}{(x_{12}^2)^{J-\frac{K}{2}} K!(K+1)!} + \dots \quad (\text{A.6})$$

This is the contribution of a single half-BPS operator $\mathcal{O}_K(x, u)$ to the bulk OPE, and we suppress terms subleading as $x_{12}^2 \rightarrow 0$. In order to deal with the R -symmetry polarization, we use the $SO(r)$ Todorov operator

$$(D_u^{(r)})_\mu = \left(\frac{r}{2} - 1 + u \cdot \frac{\partial}{\partial u}\right) \frac{\partial}{\partial u^\mu} - \frac{1}{2} u_\mu \frac{\partial^2}{\partial u \cdot \partial u}. \quad (\text{A.7})$$

One can insert the bulk OPE (A.6) combined with the one-point function (2.24) in the two-point function. Keeping only the leading term as $z, \bar{z} \rightarrow 1$ gives

$$\mathcal{F}^{(J)}(z, \bar{z}, \sigma) \Big|_{\mathcal{O}_K} = \left(\frac{\sqrt{z\bar{z}} \sigma}{(1-z)(1-\bar{z})}\right)^J \lambda_{JJK} a_K h_K(\sigma) [(1-z)(1-\bar{z})]^{\frac{K}{2}} + \dots \quad (\text{A.8})$$

This is the leading contribution of a protected half-BPS operator to the bulk OPE, and it is equally valid for single- and multi-trace operators. It is reassuring that the contribution is proportional to the R -symmetry block (A.5). This result also justifies our choice of the overall normalization for the bulk superblock (4.4).

The story for the defect OPE works in an identical way. The form of the correlators (2.30) fixes uniquely the bulk-defect expansion

$$\mathcal{O}_J(x, u)|_{\hat{\mathcal{O}}_{\hat{K}}} = b_{J\hat{K}} \frac{(u \cdot \theta)^{J-\hat{K}} (u \cdot D_{\hat{u}}^{(5)})^{\hat{K}} \hat{\mathcal{O}}_{\hat{K}}(\tau, \hat{u})}{|x^\perp|^{J-\hat{K}} \hat{K}!(3/2)_{\hat{K}}} + \dots \quad (\text{A.9})$$

As before, we focus on the contribution of a protected defect operator and keep only the leading-order term as $x^\perp \rightarrow 0$. Since the defect operator transforms as an $SO(5)$ symmetric traceless tensor, we use the $r = 5$ version of (A.7). Inserting the defect OPE in the two-point function gives

$$\mathcal{F}^{(J)}(z, \bar{z}, \sigma)|_{\hat{\mathcal{O}}_{\hat{K}}} \sim b_{J\hat{K}}^2 \hat{h}_{\hat{K}}(\sigma) (z\bar{z})^{\hat{K}/2} + \dots \quad (\text{A.10})$$

This is the leading contribution as $z, \bar{z} \rightarrow 0$ of a protected defect operator to the defect OPE.

References

- [1] A. L. Fitzpatrick, J. Kaplan, D. Poland and D. Simmons-Duffin, *The Analytic Bootstrap and AdS Superhorizon Locality*, *JHEP* **12** (2013) 004 [[1212.3616](#)].
- [2] Z. Komargodski and A. Zhiboedov, *Convexity and Liberation at Large Spin*, *JHEP* **1311** (2013) 140 [[1212.4103](#)].
- [3] L. F. Alday, *Large Spin Perturbation Theory for Conformal Field Theories*, *Phys. Rev. Lett.* **119** (2017) 111601 [[1611.01500](#)].
- [4] S. Caron-Huot, *Analyticity in Spin in Conformal Theories*, *JHEP* **09** (2017) 078 [[1703.00278](#)].
- [5] L. F. Alday and S. Caron-Huot, *Gravitational S-matrix from CFT dispersion relations*, *JHEP* **12** (2018) 017 [[1711.02031](#)].
- [6] S. Caron-Huot and A.-K. Trinh, *All tree-level correlators in AdS₅ × S⁵ supergravity: hidden ten-dimensional conformal symmetry*, *JHEP* **01** (2019) 196 [[1809.09173](#)].
- [7] L. Rastelli and X. Zhou, *Mellin amplitudes for AdS₅ × S⁵*, *Phys. Rev. Lett.* **118** (2017) 091602 [[1608.06624](#)].
- [8] L. Rastelli and X. Zhou, *How to Succeed at Holographic Correlators Without Really Trying*, *JHEP* **04** (2018) 014 [[1710.05923](#)].
- [9] G. W. Semenoff and K. Zarembo, *More exact predictions of SUSYM for string theory*, *Nucl. Phys. B* **616** (2001) 34 [[hep-th/0106015](#)].
- [10] K. Okuyama and G. W. Semenoff, *Wilson loops in N=4 SYM and fermion droplets*, *JHEP* **06** (2006) 057 [[hep-th/0604209](#)].
- [11] S. Giombi and V. Pestun, *Correlators of local operators and 1/8 BPS Wilson loops on S² × S² from 2d YM and matrix models*, *JHEP* **10** (2010) 033 [[0906.1572](#)].
- [12] M. Billo, F. Galvagno, P. Gregori and A. Lerda, *Correlators between Wilson loop and chiral operators in N = 2 conformal gauge theories*, *JHEP* **03** (2018) 193 [[1802.09813](#)].
- [13] J. Barrat, P. Liendo and J. Plefka, *Two-point correlator of chiral primary operators with a Wilson line defect in N = 4 SYM*, *JHEP* **05** (2021) 195 [[2011.04678](#)].

- [14] S. Giombi and V. Pestun, *Correlators of Wilson Loops and Local Operators from Multi-Matrix Models and Strings in AdS*, *JHEP* **01** (2013) 101 [[1207.7083](#)].
- [15] E. Buchbinder and A. Tseytlin, *Correlation function of circular Wilson loop with two local operators and conformal invariance*, *Phys. Rev. D* **87** (2013) 026006 [[1208.5138](#)].
- [16] M. Beccaria and A. A. Tseytlin, *On the structure of non-planar strong coupling corrections to correlators of BPS Wilson loops and chiral primary operators*, [2011.02885](#).
- [17] M. Billò, V. Goncalves, E. Lauria and M. Meineri, *Defects in conformal field theory*, *JHEP* **04** (2016) 091 [[1601.02883](#)].
- [18] J. M. Maldacena, *The Large N limit of superconformal field theories and supergravity*, *Int. J. Theor. Phys.* **38** (1999) 1113 [[hep-th/9711200](#)].
- [19] E. Witten, *Anti-de Sitter space and holography*, *Adv. Theor. Math. Phys.* **2** (1998) 253 [[hep-th/9802150](#)].
- [20] S. Gubser, I. R. Klebanov and A. M. Polyakov, *Gauge theory correlators from noncritical string theory*, *Phys. Lett. B* **428** (1998) 105 [[hep-th/9802109](#)].
- [21] J. M. Maldacena, *Wilson loops in large N field theories*, *Phys. Rev. Lett.* **80** (1998) 4859 [[hep-th/9803002](#)].
- [22] S.-J. Rey and J.-T. Yee, *Macroscopic strings as heavy quarks in large N gauge theory and anti-de Sitter supergravity*, *Eur. Phys. J. C* **22** (2001) 379 [[hep-th/9803001](#)].
- [23] J. Erickson, G. Semenoff and K. Zarembo, *Wilson loops in $N=4$ supersymmetric Yang-Mills theory*, *Nucl. Phys. B* **582** (2000) 155 [[hep-th/0003055](#)].
- [24] N. Drukker and D. J. Gross, *An Exact prediction of $N=4$ SUSYM theory for string theory*, *J. Math. Phys.* **42** (2001) 2896 [[hep-th/0010274](#)].
- [25] V. Pestun, *Localization of gauge theory on a four-sphere and supersymmetric Wilson loops*, *Commun. Math. Phys.* **313** (2012) 71 [[0712.2824](#)].
- [26] D. E. Berenstein, R. Corrado, W. Fischler and J. M. Maldacena, *The Operator product expansion for Wilson loops and surfaces in the large N limit*, *Phys. Rev. D* **59** (1999) 105023 [[hep-th/9809188](#)].
- [27] M. Lemos, P. Liendo, M. Meineri and S. Sarkar, *Universality at large transverse spin in defect CFT*, *JHEP* **09** (2018) 091 [[1712.08185](#)].
- [28] S. Giombi, R. Roiban and A. A. Tseytlin, *Half-BPS Wilson loop and AdS_2/CFT_1* , *Nucl. Phys. B* **922** (2017) 499 [[1706.00756](#)].
- [29] S. Giombi and S. Komatsu, *Exact Correlators on the Wilson Loop in $\mathcal{N} = 4$ SYM: Localization, Defect CFT, and Integrability*, *JHEP* **05** (2018) 109 [[1802.05201](#)].
- [30] P. Liendo, C. Meneghelli and V. Mitev, *Bootstrapping the half-BPS line defect*, *JHEP* **10** (2018) 077 [[1806.01862](#)].
- [31] P. Ferrero and C. Meneghelli, *Bootstrapping the half-BPS line defect CFT in $\mathcal{N} = 4$ SYM at strong coupling*, [2103.10440](#).
- [32] D. Grabner, N. Gromov and J. Julius, *Excited States of One-Dimensional Defect CFTs from the Quantum Spectral Curve*, *JHEP* **07** (2020) 042 [[2001.11039](#)].
- [33] A. Cavaglià, N. Gromov, J. Julius and M. Preti, *Integrability and Conformal Bootstrap: One Dimensional Defect CFT*, [2107.08510](#).

- [34] L. Bianchi, L. Griguolo, M. Preti and D. Seminara, *Wilson lines as superconformal defects in ABJM theory: a formula for the emitted radiation*, *JHEP* **10** (2017) 050 [[1706.06590](#)].
- [35] L. Bianchi, M. Preti and E. Vescovi, *Exact Bremsstrahlung functions in ABJM theory*, *JHEP* **07** (2018) 060 [[1802.07726](#)].
- [36] L. Bianchi, G. Bliard, V. Forini, L. Griguolo and D. Seminara, *Analytic bootstrap and Witten diagrams for the ABJM Wilson line as defect CFT₁*, *JHEP* **08** (2020) 143 [[2004.07849](#)].
- [37] P. Liendo and C. Meneghelli, *Bootstrap equations for $\mathcal{N} = 4$ SYM with defects*, *JHEP* **01** (2017) 122 [[1608.05126](#)].
- [38] F. A. Dolan, L. Gallot and E. Sokatchev, *On four-point functions of 1/2-BPS operators in general dimensions*, *JHEP* **09** (2004) 056 [[hep-th/0405180](#)].
- [39] M. Nirschl and H. Osborn, *Superconformal Ward identities and their solution*, *Nucl. Phys.* **B711** (2005) 409 [[hep-th/0407060](#)].
- [40] P. Liendo, C. Meneghelli and V. Mitev, *On Correlation Functions of BPS Operators in 3d $\mathcal{N} = 6$ Superconformal Theories*, *Commun. Math. Phys.* **350** (2017) 387 [[1512.06072](#)].
- [41] S. M. Chester, J. Lee, S. S. Pufu and R. Yacoby, *The $\mathcal{N} = 8$ superconformal bootstrap in three dimensions*, *JHEP* **09** (2014) 143 [[1406.4814](#)].
- [42] F. A. Dolan and H. Osborn, *On short and semi-short representations for four-dimensional superconformal symmetry*, *Annals Phys.* **307** (2003) 41 [[hep-th/0209056](#)].
- [43] C. Cordova, T. T. Dumitrescu and K. Intriligator, *Multiplets of Superconformal Symmetry in Diverse Dimensions*, *JHEP* **03** (2019) 163 [[1612.00809](#)].
- [44] N. B. Agmon and Y. Wang, *Classifying Superconformal Defects in Diverse Dimensions Part I: Superconformal Lines*, [2009.06650](#).
- [45] S. Giombi and S. Komatsu, *More Exact Results in the Wilson Loop Defect CFT: Bulk-Defect OPE, Nonplanar Corrections and Quantum Spectral Curve*, *J. Phys.* **A52** (2019) 125401 [[1811.02369](#)].
- [46] S. Lee, S. Minwalla, M. Rangamani and N. Seiberg, *Three point functions of chiral operators in $D = 4$, $N=4$ SYM at large N* , *Adv. Theor. Math. Phys.* **2** (1998) 697 [[hep-th/9806074](#)].
- [47] D. Correa, J. Henn, J. Maldacena and A. Sever, *An exact formula for the radiation of a moving quark in $N=4$ super Yang Mills*, *JHEP* **1206** (2012) 048 [[1202.4455](#)].
- [48] J. Barrat, A. Gimenez-Grau and P. Liendo, *Unpublished notes*.
- [49] V. Goncalves, *Four point function of $\mathcal{N} = 4$ stress-tensor multiplet at strong coupling*, *JHEP* **04** (2015) 150 [[1411.1675](#)].
- [50] L. F. Alday, *Solving CFTs with Weakly Broken Higher Spin Symmetry*, *JHEP* **10** (2017) 161 [[1612.00696](#)].
- [51] L. F. Alday, J. Henriksson and M. van Loon, *Taming the ϵ -expansion with large spin perturbation theory*, *JHEP* **07** (2018) 131 [[1712.02314](#)].
- [52] M. Lemos, B. C. van Rees and X. Zhao, *Regge trajectories for the $(2,0)$ theories*, [2105.13361](#).
- [53] F. Aprile, J. M. Drummond, P. Heslop and H. Paul, *Quantum Gravity from Conformal Field Theory*, *JHEP* **01** (2018) 035 [[1706.02822](#)].
- [54] F. Aprile, J. M. Drummond, P. Heslop and H. Paul, *Unmixing Supergravity*, *JHEP* **02** (2018) 133 [[1706.08456](#)].

- [55] F. Aprile, J. M. Drummond, P. Heslop and H. Paul, *Loop corrections for Kaluza-Klein AdS amplitudes*, *JHEP* **05** (2018) 056 [[1711.03903](#)].
- [56] L. F. Alday and X. Zhou, *All Holographic Four-Point Functions in All Maximally Supersymmetric CFTs*, *Phys. Rev. X* **11** (2021) 011056 [[2006.12505](#)].
- [57] L. F. Alday and X. Zhou, *All Tree-Level Correlators for M-theory on $AdS_7 \times S^4$* , *Phys. Rev. Lett.* **125** (2020) 131604 [[2006.06653](#)].
- [58] V. Goncalves and G. Itsios, *A note on defect Mellin amplitudes*, [1803.06721](#).
- [59] M. Beccaria, S. Giombi and A. A. Tseytlin, *Correlators on non-supersymmetric Wilson line in $N=4$ SYM and AdS_2/CFT_1* , [1903.04365](#).
- [60] S. Giombi, J. Jiang and S. Komatsu, *Giant Wilson loops and $AdS_2/dCFT_1$* , *JHEP* **11** (2020) 064 [[2005.08890](#)].
- [61] A. Gimenez-Grau and P. Liendo, *Bootstrapping Monodromy Defects in the Wess-Zumino Model*, [2108.05107](#).
- [62] M. Isachenkov, P. Liendo, Y. Linke and V. Schomerus, *Calogero-Sutherland Approach to Defect Blocks*, *JHEP* **10** (2018) 204 [[1806.09703](#)].
- [63] P. Liendo, Y. Linke and V. Schomerus, *A Lorentzian inversion formula for defect CFT*, [1903.05222](#).