

DEUTSCHES ELEKTRONEN-SYNCHROTRON
Ein Forschungszentrum der Helmholtz-Gemeinschaft



DESY 22-002
arXiv:2201.01606
January 2022

Uncertainties on the Theoretical Input for $t\bar{t}j$ Experimental Analyses at the LHC

K. Voss

II. Institut für Theoretische Physik, Universität Hamburg

and

*Zentrum für Teilchenphysik Siegen, Department Physik,
Universität Siegen*

M. V. Garzelli, S.-O. Moch

II. Institut für Theoretische Physik, Universität Hamburg

ISSN 0418-9833

NOTKESTRASSE 85 - 22607 HAMBURG

DESY behält sich alle Rechte für den Fall der Schutzrechtserteilung und für die wirtschaftliche Verwertung der in diesem Bericht enthaltenen Informationen vor.

DESY reserves all rights for commercial use of information included in this report, especially in case of filing application for or grant of patents.

Herausgeber und Vertrieb:

Verlag Deutsches Elektronen-Synchrotron DESY

DESY Bibliothek
Notkestr. 85
22607 Hamburg
Germany

Uncertainties on the theoretical input for $t\bar{t}j$ experimental analyses at the LHC

KATHARINA VOSS^{a,b}, MARIA VITTORIA GARZELLI^a, SVEN-OLAF MOCH^a

^a *II. Institut für Theoretische Physik, Universität Hamburg,
Luruper Chaussee 149, D – 22761 Hamburg, Germany*

^b *Center For Particle Physics Siegen, Department Physik, Universität Siegen,
Walter Flex Str. 3, D – 57068 Siegen, Germany*

The precise measurement of the top-quark mass constitutes one of the main goals of the LHC top-quark physics program. One possibility to extract this parameter uses the ρ_s distribution, which depends on the invariant mass of the $t\bar{t}j$ system. To fully take advantage of the experimental accuracy achievable in measuring top quark production cross sections at the LHC, the theory uncertainties need to be well under control. We present a study of the effect of varying the input parameters of the theoretical calculation on the predicted ρ_s distribution. Thereby we studied the influence of the R parameter in the jet reconstruction procedure, as well as the effect of various renormalization and factorization scale definitions and different PDF sets. A behaviour similar to the one presented here for the ρ_s -distribution was also found for other differential distributions.

PRESENTED AT

14th International Workshop on Top Quark Physics
(videoconference), 13–17 September, 2021

1 Introduction

A possibility to extract the top-quark mass m_t in a well defined mass renormalization scheme is given by the analysis of the normalized ρ_s distribution, as first discussed in [1], which is defined as $(1/\sigma_{t\bar{t}j})d\sigma_{t\bar{t}j}/d\rho_s$, with $\rho_s = 2m_0/m_{t\bar{t}j}$ and $m_0 = 170$ GeV.

A recent extraction of the ATLAS collaboration [2] of m_t in $t\bar{t}j$ hadroproduction using the ρ_s distribution led to an on-shell value of $m_t^{\text{pole}} = 171.1 \pm 0.4(\text{st}) \pm 0.9(\text{sy})_{-0.3}^{+0.7}(\text{th})$ GeV, in which the theory uncertainty is sizeable. Thereby, the scale uncertainty dominates the theory uncertainty and contributes to an uncertainty on m_t^{pole} with $_{-0.2}^{+0.6}$ GeV, while the PDF and α_S uncertainty amounts to ± 0.2 GeV.

In [3] a better description of the high energy tails of differential NLO distributions of the process $pp \rightarrow t\bar{t}j$ with fully leptonic top-quark decay was found when using dynamical renormalization and factorization scales (μ_R, μ_F) instead of a static scale $\mu_0 = m_t$. We therefore present a study of the scale and parton distribution function (PDF) uncertainty using a static and two dynamical scales, with more inclusive cuts in contrast to [3], similar to those recently adopted by the experimental collaborations, and a special focus on the discussed ρ_s distribution.

2 Theory uncertainty in the ρ_s distribution

The NLO differential cross sections of the $pp \rightarrow t\bar{t}j$ process at $\sqrt{s} = 13$ TeV for $m_t^{\text{pole}} = 172$ GeV presented in the following were obtained with the `ttbarj V2` implementation in the POWHEG-BOX [4]. In contrast to the previous `ttbarj V1` implementation [5], in `ttbarj V2` all hard scattering amplitudes are calculated with `OpenLoops2` and the calculation can be parallelized. During the analysis at least one jet satisfying the kinematic cuts $p_T^j > 30$ GeV and $|\eta_j| < 2.4$ was required, with jets reconstructed using the anti- k_T jet clustering algorithm with $R = 0.4$. The default PDFs and $\alpha_S(M_Z)$ value were taken from the CT18NLO PDF set. Besides the static scale $\mu_0 = m_t^{\text{pole}}$ also the two dynamical scales $\mu_0 = H_T^B/2$ and $H_T^B/4$ were investigated defined through $H_T^B = \sqrt{p_{T,t}^B{}^2 + (m_t^{\text{pole}})^2} + \sqrt{p_{T,\bar{t}}^B{}^2 + (m_t^{\text{pole}})^2} + p_{T,j}^B$. The superscript B means that the kinematic variables are evaluated at the underlying Born level in the POWHEG-BOX.

In Fig. 1 the ρ_s distribution and the corresponding seven point scale variation graphs, using the three scale definitions described above, are shown explicitly. The graphs are obtained from $(\mu_R, \mu_F) = (K_R, K_F)\mu_0$ by varying $K_R, K_F \in \{0.5, 1, 2\}$, leaving out the extreme combinations (0.5, 2) and (2, 0.5). In fact, in case of the static scale, for low values of ρ_s , which correspond to large values of $m_{t\bar{t}j}$ and as such to the high-energy region, the description of the ρ_s distribution seems to be unreliable and the scale uncertainty increases rapidly. Further, a crossing of the graphs obtained with different values of K_R, K_F is seen in the interval $0.1 \lesssim \rho_s \lesssim 0.3$. These distinctive

features are not observed using either dynamical scale and the scale variation does not induce such a large shape variation as seen in the static scale case. This leads to




Figure 1: NLO prediction of the ρ_s distribution and the seven point scale variation using as nominal μ_R, μ_F the scales $\mu_0 = m_t, H_T^B/2$ and $H_T^B/4$ (from left to right).

the observation that the scale uncertainty in the normalized ρ_s distribution is clearly reduced using a dynamical instead of the static scale, as shown in Fig. 2. Further




Figure 2: Same as in Fig. 1, but for the normalized ρ_s distribution.

investigation of the scale uncertainty showed that using a dynamical scale with either $R = 0.4$ or $R = 0.8$ leads to a similar scale uncertainty, while in the static scale case the scale uncertainty can be reduced in the region of low ρ_s using $R = 0.8$. Comparing the NLO and LO scale variation uncertainty bands, these overlap over the whole region of the ρ_s distribution using either dynamical scale, while in the predictions obtained with the static scale, the NLO and LO scale variation bands start to depart

from each other in the high-energy region.

Furthermore, we investigated the approximate NLO PDF uncertainty in the ρ_s distribution using four different modern NLO PDF sets CT18NLO, ABMP16, MSHT20 and NNPDF3.1 and the dynamical scale $\mu_0 = H_T^B/4$. In the bulk of the ρ_s distribution the predictions obtained with the different PDF sets agree well, while differences are observed in the high-energy tail, which are not covered by the PDF uncertainty. This was found to stem from differences in the large x -gluon distributions of the corresponding PDF fits.

3 Conclusions

The ρ_s distribution seems to be better described, especially at low ρ_s , when using the dynamical scale $H_T^B/4$ instead of the static scale $\mu_0 = m_t$. This is concluded from the observation that the graphs obtained with different (K_R, K_F) values do not cross using either described dynamical scale and the NLO and LO scale variation bands overlap over the whole ρ_s distribution, while using the static scale these only barely overlap in the high-energy region. The shape variation of the ρ_s distribution induced by the scale variation is smaller when adopting the dynamical scales instead of the static scale. This leads to a strongly reduced scale variation uncertainty in the normalized ρ_s distribution, that can be used for the experimental extraction of the top quark mass. In fact, using the dynamical scales the PDF uncertainty is comparable in size to the scale variation uncertainty in the normalized ρ_s distribution. It was additionally found that while the size of the scale uncertainty does not show dependence on the R -parameter in the anti- k_T jet clustering algorithm using either of the two dynamical scales, the statistics can be increased by using a larger R -value.

References

- [1] S. Alioli et al. A new observable to measure the top-quark mass at hadron colliders. *Eur. Phys. J. C*, 73:2438, 2013.
- [2] G. Aad et al. Measurement of the top-quark mass in $t\bar{t} + 1$ -jet events collected with the ATLAS detector in pp collisions at $\sqrt{s} = 8$ TeV. *JHEP*, 11:150, 2019.
- [3] G. Bevilacqua et al. Off-shell Top Quarks with One Jet at the LHC: A comprehensive analysis at NLO QCD. *JHEP*, 11:098, 2016.
- [4] S. Alioli et al. A general framework for implementing NLO calculations in shower Monte Carlo programs: the POWHEG BOX. *JHEP*, 06:043, 2010.
- [5] S. Alioli et al. Hadronic top-quark pair-production with one jet and parton showering. *JHEP*, 01:137, 2012.