


# DEUTSCHES ELEKTRONEN-SYNCHROTRON **DESY**

DESY 87-031  
April 1987



SEARCH FOR LEPTOQUARKS AND OTHER NEW PARTICLES WITH  
LEPTON-HADRON SIGNATURE IN  $e^+e^-$  INTERACTIONS

by

*JADE Collaboration*

ISSN 0418-9833

**NOTKESTRASSE 85 · 2 HAMBURG 52**

**DESY behält sich alle Rechte für den Fall der Schutzrechtserteilung und für die wirtschaftliche Verwertung der in diesem Bericht enthaltenen Informationen vor.**

**DESY reserves all rights for commercial use of information included in this report, especially in case of filing application for or grant of patents.**

**To be sure that your preprints are promptly included in the  
HIGH ENERGY PHYSICS INDEX,  
send them to the following address ( if possible by air mail ) :**

**DESY  
Bibliothek  
Notkestrasse 85  
2 Hamburg 52  
Germany**

# Search for Leptoquarks and other New Particles with Lepton - Hadron Signature in $e^+e^-$ Interactions

JADE Collaboration

W. Bartel, L. Becker, R. Felst, D. Haidt, G. Knies, H. Krehbiel, P. Laurikainen<sup>1</sup>,  
N. Magnussen<sup>2</sup>, R. Meinke, B. Naroska, J. Olsson, D. Schmidt<sup>2</sup>, P. Steffen  
*Deutsches Elektronen-Synchrotron DESY, Hamburg, Germany*

T. Greenshaw, J. Hagemann, G. Heinzlmann, H. Kado, C. Kleinwort,  
M. Kuhlen, A. Petersen<sup>3</sup>, R. Ramcke, U. Schneekloth<sup>4</sup>, G. Weber  
*II. Institut für Experimentalphysik der Universität Hamburg, Hamburg, Germany*

K. Ambrus, S. Bethke<sup>5</sup>, A. Dieckmann, G. Eckerlin, E. Elsen, J. Heintze,  
K.-H. Hellenbrandt, S. Komamiya<sup>3</sup>, J. von Krogh, H. Rieseberg, H. v. d. Schmitt,  
L. Smolik, J. Spitzer, A. Wagner, M. Zimmer  
*Physikalisches Institut der Universität Heidelberg, Heidelberg, Germany*

C.K. Bowdery, A.J. Finch, F. Foster, G. Hughes, J.M. Nye, I.W. Walker  
*University of Lancaster, Lancaster, England*

J. Allison, R.J. Barlow, J. Chrin, I.P. Duerdoth, F.K. Loebinger, A.A. Macbeth,  
H.E. Mills<sup>6</sup>, P.G. Murphy, K. Stephens, P. Warming<sup>7</sup>  
*University of Manchester, Manchester, England*

R.G. Glasser, P. Hill, J.A.J. Skard, S. R. Wagner<sup>8</sup>, G.T. Zorn  
*University of Maryland, College Park, MD, USA*

S.L. Cartwright<sup>4</sup>, D. Clarke, R. Marshall, R.P. Middleton  
*Rutherford Appleton Laboratory, Chilton, Didcot, England*

T. Kawamoto, T. Kobayashi, M. Minowa, H. Takeda, S. Yamada  
*International Center for Elementary Particle Physics, University of Tokyo, Tokyo, Japan*

<sup>1</sup> University of Helsinki, Helsinki, Finland

<sup>2</sup> Universität-Gesamthochschule Wuppertal, Wuppertal, Germany

<sup>3</sup> Now at SLAC, Stanford, California, USA

<sup>4</sup> Now at MIT, Cambridge, Massachusetts, USA

<sup>5</sup> Now at LBL, Berkeley, California, USA

<sup>6</sup> Now at SCS, Hamburg, Germany

<sup>7</sup> Now at Digitec, Hamburg, Germany

<sup>8</sup> Now at University of Colorado, Boulder, Colorado, USA

Using the JADE detector at PETRA, a search for hadronic events with isolated leptons in  $e^+e^-$  annihilation was performed. No evidence was found for the production of leptoquarks or color octet leptons and mass limits were obtained. Three hadronic events, each of them with two isolated muons, were detected and found to be in agreement with predictions of the  $\alpha^4$  QED process  $e^+e^- \rightarrow q\bar{q}\mu^+\mu^-$ . At the highest PETRA energies,  $46.30 \text{ GeV} \leq \sqrt{s} \leq 46.78 \text{ GeV}$ , five events with low thrust,  $T < 0.8$ , and a muon at a large angle with respect to the thrust axis,  $|\cos\delta| < 0.7$ , were observed. The number of such events expected from an extrapolation of control data at lower energies is  $0.56 \pm 0.18$ . No satisfactory explanation for the excess in terms of background could be found. A corresponding excess of electron events was not seen.

## 1 Introduction

In this paper we report on a study of the reaction  $e^+e^- \rightarrow \mu^+\mu^- + \text{hadrons}$  and on a search for leptoquarks and color octet leptons in  $e^+e^-$  annihilation. Three different event topologies are discussed: two acoplanar jets, a muon + jets + missing energy and two muons + jets. Finally we describe an investigation of hadronic events with low thrust containing isolated leptons, for which the Mark J group observed an unexpected behaviour at the highest PETRA energies [1].

## 2 Event selection and lepton identification

### 2.1 Event selection

The data were collected with the JADE detector [2] at the PETRA  $e^+e^-$  storage ring. Multiparticle events were selected using the criteria described in [3], which are the basis for the selection of multihadronic events in JADE. The most important cuts are:

1. The total lead glass energy in the barrel part of the detector is required to exceed 3 GeV or the energy deposition in each of the endcaps has to be larger than 0.4 GeV.
2. At least four charged particles are demanded to originate from the event vertex.
3. Remaining cosmic ray events,  $\tau$  - pairs and purely leptonic QED events are removed by visual inspection.

Further cuts, in visible energy  $E_{vis} = \sum_i p_i$ , and longitudinal momentum balance  $p_{bal} = \sum_i p_i^z / E_{vis}$ , extract the multihadronic annihilation events from this sample. The sums run over all particle momenta  $p_i$ ,  $p_i^z$  denoting the components along the beam axis.

4.  $E_{vis} > E_b$ , where  $E_b$  is the beam energy.
5.  $|p_{bal}| < 0.4$

### 2.2 Muon identification

Muons are identified as penetrating particles in the muon filter [4], which consists of four successive blocks of absorber material interleaved with layers of drift chambers. It covers 92 % of the full solid angle and presents in total at least 6.4 nuclear absorption lengths to particles with incidence perpendicular to the beam. All tracks recorded in the jet chamber are extrapolated through the

muon filter, and their likelihood of being muons is tested. The following samples of tracks are considered as muon candidates in the analysis [5]:

#### “Muon candidates”

At least two hits in the muon chambers outside the magnet yoke are asked for within a  $2\sigma$  range of the extrapolation of a track taking into account fitting errors and multiple scattering. One missing hit in the  $2\sigma$  range is permitted, except in the case of the outermost intersected layer of chambers, where a hit has to be present.

#### “Good muons”

Further requirements define the “good muons” used in the analysis. The segmented muon filter enables different selection criteria to be chosen, depending on the goals of the physics analysis. Usually loose cuts are used for new particle searches, where high detection efficiency is essential (see for example [6]), while tight cuts give the cleanest muon sample and are used where the properties to be studied are obscured by a large background, for example for the determination of the semi-muonic branching ratios and fragmentation functions of heavy quarks [7]. The “loose” cuts and the “tight” cuts are defined as follows<sup>1</sup>:

“Loose” cuts: In addition to the above criteria, the track momentum has to exceed 1.8 GeV, the track is required to penetrate more than 4.8 absorption lengths to the outermost associated hit in the muon filter and the  $\chi^2$  probability that the jet chamber track matches the  $\mu$  filter hits has to be greater than 1%.

“Tight” cuts: As a further condition there must be no expected hit missing in the muon chambers, and more than 5.8 absorption lengths have to be traversed. At least 3 layers of muon chambers outside the yoke are required to have registered hits which can be associated with the jet chamber track.

### 2.3 Electron identification

Electrons are identified by their energy loss in the gas of the drift chamber and by their electromagnetic shower detected in the lead glass calorimeter. In order to confine the shower to the central part of the lead glass system which has the highest resolution, an acceptance cut of  $|\cos\theta| < 0.76$  is imposed,  $\theta$  being the angle of the particle with the beam axis.

For a particle to be considered an electron candidate the track momentum has to be greater than 1 GeV, and its energy loss as measured in the jet chamber is demanded to agree with the expectation for electrons within  $-1.5\sigma < (dE/dx)_{measured} - (dE/dx)_{electron} < 2.0\sigma$ . The shape of the lead glass cluster associated with the track is required to be consistent with an electromagnetic shower. An electron candidate is removed if it can be combined with another track to form a photon conversion pair. Finally, the deposited energy  $E$  in the leadglass has to agree with the track momentum  $p$ , it being required that  $0.8 < E/p < 3.0$ . Fig. 1 shows the  $E/p$  distribution after all except the last of these cuts. There is a distinct electron peak at  $E/p = 1$ .

### 3 The reaction $e^+e^- \rightarrow q\bar{q}\mu^+\mu^-$

The process  $e^+e^- \rightarrow q\bar{q}\mu^+\mu^-$  is an  $\alpha^4$  QED process. It is a source of events with isolated energetic muons, and can thus be used to check the sensitivity of the detector to such processes

<sup>1</sup>The “loose” cuts correspond to muons of “quality A” and the “tight” cuts to muons of “quality C” in ref. [5].




Figure 1:  $E/p$  distribution after  $dE/dx$  cut

and to study the background for the event topologies to be searched for in the following chapters.




Figure 2: Feynman graphs and corresponding event topologies for  $e^+e^- \rightarrow q\bar{q}\mu^+\mu^-$

Three classes of Feynman graphs, which are displayed in fig. 2, contribute to this process. The cross section was calculated using the formalism and computer code developed by Berends, Daverfeldt and Kleiss [8]. The  $q\bar{q}\mu^+\mu^-$  events generated by this program were fragmented according to the Lund scheme [9] and passed through the detector simulation and analysis chain in order to determine the detection efficiency. The selection criteria, optimized with respect to the 2-photon conversion diagrams (fig. 2 A) which give the main contribution to the total cross section, were as follows:

1. The event had to fulfill the multihadron selection requirements.

2. 2 "muon candidates" with  $p_\mu > 3$  GeV, at least one of them being a "good muon" ("loose" cuts) were asked for.
3. No additional charged tracks, with the exception of other "muon candidates", were allowed in cones with a half opening angle of  $60^\circ$  surrounding each of the two muons.
4. It was required that there be less than 1 GeV lead glass energy in these cones.
5. The remaining events were scanned by a physicist, and it was required that there be two clearly separated tracks visible within the muon filter.

Data corresponding to an integrated luminosity of  $94.2 \text{ pb}^{-1}$  at centre of mass energies ranging from 29.5 to 46.78 GeV were analysed. Three events were found after all cuts, as compared to 1.2 events expected from the QED simulation of  $e^+e^- \rightarrow q\bar{q}\mu^+\mu^-$ . The background, mainly from  $\mu\mu\tau\tau$  and  $q\bar{q}$  events with fake muons, was estimated to be less than 0.3 events. In fig. 3 the muon opening angles and hadronic invariant masses are compared to the QED distributions. Within the limited statistics, event rate and kinematic configurations agree with the QED expectation. The observation of these events at the expected rate gives confidence that the JADE detector is capable of detecting rare events with isolated muons.

## 4 Leptoquarks

### 4.1 Introduction

Light leptoquark bosons with charge  $1/3, 2/3, \dots$ , which decay into a quark and a lepton, appear in several technicolor and composite models [10], [11]. In the present analysis emphasis is placed on a specific model due to B. and F. Schrempp [10] with second generation leptoquarks of charge  $2/3$ . In this model generation quantum number is conserved. Thus a second generation leptoquark  $\chi_\mu$  can only decay into either  $c\bar{\nu}_\mu$  or  $s\mu^+$ .

The cross section for pair production of charged color triplet spinless bosons in  $e^+e^-$  annihilation is

$$\frac{d\sigma}{d\Omega}(e^+e^- \rightarrow \chi\bar{\chi}) = 3 \cdot \frac{\alpha^2}{8s} \cdot Q^2 \cdot \beta^3 \cdot \sin^2\theta \quad (1)$$

where  $\beta = \sqrt{1 - 4m_\chi^2/s}$ ,  $m_\chi$  is the leptoquark mass,  $Q = 2/3$  and  $\theta$  is the production angle with respect to the beam direction. The cross section rises slowly from threshold with increasing energy, due to the  $\beta^3$  factor. Near threshold the contribution to the total hadronic cross section is too small to be measurable, but leptoquarks can be searched for by their distinctive event topologies (see fig. 4):

- A)  $e^+e^- \rightarrow \chi_\mu\bar{\chi}_\mu \rightarrow c\bar{\nu}_\mu\bar{\nu}_\mu$  : two acoplanar jets
- B)  $e^+e^- \rightarrow \chi_\mu\bar{\chi}_\mu \rightarrow c\bar{\nu}_\mu s\bar{\mu}$  or *c.c.* : two jets, a muon and missing momentum
- C)  $e^+e^- \rightarrow \chi_\mu\bar{\chi}_\mu \rightarrow s\mu^+ s\bar{\mu}^-$  : two jets and two muons

Using a Monte Carlo simulation of leptoquark events, methods of isolating them from ordinary events by applying certain cuts were developed. The detection efficiencies after the cuts were also obtained from the simulation. The leptoquarks were fragmented like heavy quarks using the Lund model [9] and the Peterson fragmentation function [12].

Events of classes A), B) and C) were searched for in the high energy data sample corresponding to an integrated luminosity of  $24.5 \text{ pb}^{-1}$  between  $\sqrt{s} = 38.66$  and 46.78 GeV.




Figure 3: Muon opening angle and hadronic invariant mass for simulated  $q\bar{q}\mu^+\mu^-$  events after cuts. The measured values of the three observed events are indicated.




Figure 4: Event topologies for leptoquarks

## 4.2 Analysis

### A) Acoplanar jets

Cuts identical to those used in the searches for the supersymmetric partner of the  $Z^0$  [13] and monojet events [14] previously performed by JADE were applied to the multiparticle events defined in section 2.1. These were:

1.  $2/5 < E_{vis}/\sqrt{s} < 1$
2.  $|\cos \theta_T| < 0.65$ , with  $\theta_T$  the angle between thrust axis and the positron beam direction.
3.  $\phi_{acop} > 40^\circ \cdot (1 + |\cos \theta_T|)$

The acoplanarity angle  $\phi_{acop}$  is given by

$$\phi_{acop} = \arccos \left( -\frac{(\vec{p}_+ \times \vec{n}_z) \cdot (\vec{p}_- \times \vec{n}_z)}{|\vec{p}_+ \times \vec{n}_z| \cdot |\vec{p}_- \times \vec{n}_z|} \right) \quad (2)$$

where  $\vec{p}_+$  and  $\vec{p}_-$  are defined as follows: Each event is split into two halves using a plane perpendicular to the thrust axis.  $\vec{p}_+$  and  $\vec{p}_-$  are then the sums of the momenta of the particles in each of these event halves.  $\vec{n}_z$  is a unit vector in the direction of the positron beam. If there are no particles in one of the event halves then  $\phi_{acop}$  is set to  $180^\circ$ .

No event survived the cuts. The detection efficiency for  $m_\chi = 21$  GeV was found to be 28 %.

### B) Two jets, a muon + missing momentum

The multihadron sample was used for this analysis.

1. A "good muon" selected with "loose" cuts and with momentum  $p_\mu > 5$  GeV was required. In fig. 5 the observed muon momentum spectrum is compared to that of a Monte Carlo simulation of multihadronic events [9]. Also shown is the expected distribution for leptoquark events, which is significantly different from that for ordinary multihadronic events.




Figure 5: Muon momenta  $p_\mu$

2.  $E_{vis} < \sqrt{s}$ , as the events required contain neutrinos which escape detection.
3.  $\phi_{acol} > 40^\circ \cdot (1 + 0.5 |\cos \theta_T|)$   
The acolinearity angle

$$\phi_{acol} = \arccos \left( -\frac{\vec{p}_+ \cdot \vec{p}_-}{|\vec{p}_+| \cdot |\vec{p}_-|} \right) \quad (3)$$

is  $180^\circ$  minus the angle between the two momentum sum vectors defined above. Muons are not included in the momentum sums in order to treat them on an equal footing with the neutrinos in case A). As for the acoplanarity angle, the acolinearity angle is defined to be  $180^\circ$  if all particles are in the same event half.


One event with an isolated muon in one event half opposite to a hadronic system in the other half survived these cuts. The event is shown in fig. 6. It is consistent with background from  $e^+e^- \rightarrow q\bar{q}\mu^-\mu^+$  in which one muon is not detected, or from the decay  $D^\pm \rightarrow K^0\mu^\pm\nu$  with an escaping  $K_L^0$ . The detection efficiency for leptoquark events with  $m_\chi = 21$  GeV is 50 % after the above cuts have been applied.

### C) Two jets + two muons

The search for these events was performed on the multihadron sample, and the selection criteria were:


1. Two "good muons" ("loose" cuts) with momentum  $p_\mu > 3$  GeV
2. The transverse momentum with respect to the thrust axis ( $p_\mu^T$ ) was required to be larger than 5 GeV for at least one of the muons.

After these cuts the detection efficiency for  $m_\chi = 21$  GeV is 44 %. One event passes these cuts; it is event D from the low thrust - isolated muon analysis of chapter 6 and is shown in fig. 7: As



40525

Figure 6: Leptoquark candidate 1. The view along the beam axis shows the tracks of charged particles in the drift chamber, lead glass energies and hits in the muon filter. The centre of mass energy is 43.21 GeV and the muon has a momentum of 6.0 GeV. There is a nuclear interaction in the pressure vessel of the drift chamber, emitting a particle into the muon half. The charged particles alone, not counting those from the nuclear interaction, constitute an invariant mass of 3.9 GeV, thus excluding the possibility of this event being a  $\tau$  - pair event.



40524

Figure 7: Leptoquark candidate 2 (and event D from chapter 6) at  $\sqrt{s} = 46.57$  GeV. The momentum of the  $\mu^+$  is  $21.8^{+18.2}_{-6.8}$  GeV, and that of the  $\mu^-$  is  $3.2 \pm 0.3$  GeV. The isolated converted photon has an energy of 14.4 GeV.

the momentum of the second  $\mu$  is too low to be associated with a high mass leptoquark decay, leptoquarks with  $m_\chi > 16$  GeV cannot be the source of this event.

### 4.3 Results

The detection efficiencies quoted are valid for leptoquark masses of 21 GeV. For lower masses they generally decrease and approach zero for masses below about 5 GeV.

Comparison of the number of candidate events to the number of expected leptoquark events, given the luminosity, cross section and detection efficiency, yields limits for the branching ratios and masses. The accepted events from cases B and C were taken as candidates; a background subtraction was not attempted.

Assuming second generation leptoquarks of charge 2/3, which decay exclusively into second generation fermions, we obtain the bounds (95 % C.L.) shown in fig. 8 for the leptoquark mass and branching ratio  $BR(\chi_\mu \rightarrow c\bar{\nu}_\mu) = 1 - BR(\chi_\mu \rightarrow s\mu^+)$ . The combined limit from the analyses A), B) and C) excludes these leptoquarks from 5 GeV up to 20.8 GeV. The mass region below 5 GeV is not accessible to this analysis.

To account for leptoquarks of other charges and other decay channels, we also give limits in terms of  $Q^2 \cdot BR(\chi \rightarrow q\nu) \cdot BR(\chi \rightarrow q\nu)$ ,  $Q^2 \cdot BR(\chi \rightarrow q\mu) \cdot BR(\chi \rightarrow q\nu)$  and  $Q^2 \cdot BR(\chi \rightarrow q\mu) \cdot BR(\chi \rightarrow q\mu)$  in fig. 9. These have been calculated assuming that the detection efficiencies are independent of quark flavor.

The lower mass limit of 20.8 GeV (95 % C.L.) is not high enough to completely exclude a leptoquark interpretation of a  $2\mu + 2\text{jet}$  event observed by the CELLO collaboration [15], which would suggest a leptoquark mass of about 20.5 GeV. Similar limits as in fig. 8 were obtained by the CELLO collaboration [16].

## 5 Chromoleptons

In some composite models [17],[18] color octet leptons  $l_8$  may be sufficiently light to be produced at present accelerators. They decay into a gluon and their color singlet partner, the ordinary lepton. In  $e^+e^-$  annihilation, colored neutrinos, pair produced via a virtual  $Z^0$ , would result in acoplanar jets, and colored muon pairs, mainly produced via a virtual photon at PETRA energies, in  $2\mu + 2\text{jet}$  events. The analysis of acoplanar jet events and dimuon events in the leptoquark search also puts limits on  $\nu_8$  and  $\mu_8$ .

For the pair production cross section  $\sigma_8$  of colored leptons we assumed

$$\sigma_8 = 8 \cdot F(q^2) \cdot \sigma_1 \quad (4)$$

where 8 is the color factor, and  $F(q^2)$  is a form factor to account for the compositeness of these objects;  $\sigma_1$  is the cross section for the production of color singlet leptons.

Differences in quark and gluon fragmentation were neglected and it was assumed that the lifetime of the color octet particles is too short for these to be visible in the detector. With these assumptions the  $\nu_8\bar{\nu}_8$  and  $\mu_8^+\mu_8^-$  events look the same as the  $c\bar{\nu}_\mu\bar{c}\nu_\mu$  and  $s\mu^+\bar{s}\mu^-$  events, with only the angular distributions for fermions ( $l_8$ ) and bosons ( $\chi$ ) being different.

The resulting limits (95 % C.L.) for the form factors and masses are given in fig. 10. Charged chromoleptons would also contribute to the total hadronic cross section, and the limits derived from the  $R$ -value analysis [3],[6] are also shown. When  $F = 1$ , which is expected since




Figure 8: Limits (95 % C.L.) on mass and branching ratio  $BR(\chi_\mu \rightarrow c\bar{\nu}_\mu)$  for second generation leptoquarks with charge 2/3




Figure 9: Lower mass limits (95 % C.L.) for leptoquarks of charge  $Q$  decaying into a quark and a muon or neutrino.  $BR_1$  and  $BR_2$  are the branching ratios of the leptoquarks.




Figure 10: Mass limits for chromoleptons. The limit from the measurement of the total hadronic cross section ( $R$ -value) is equally valid for  $e_8^\pm$ ,  $\mu_8^\pm$  und  $\tau_8^\pm$ .

$q^2 = s \ll \Lambda_C^2$  ( $\Lambda_C$  is the compositeness scale) for PETRA energies (here  $\langle s \rangle \approx 1900 \text{ GeV}^2$ ), the mass regions given in table 1 can be excluded. A limit on color octet electrons previously published by JADE [6], where the effective coupling  $\lambda$  at the  $ee_g$  vertex enters, is quoted for completeness.

Table 1: Summary of excluded mass regions for color octet leptons ( $F(q^2) = 1$ )

	lower limit	upper limit	analysis
$\nu_8^\pm$	0 GeV	19.8 GeV	$R$ -value
$e_8^\pm$	0 GeV	$173 \text{ GeV} \cdot \lambda^{2/3}$	$R$ -value
$\mu_8^\pm$	5 GeV	23.2 GeV	hadronic dimuon events
$\nu_8$	9 GeV	21.9 GeV	acoplanar jet events

## 6 Hadronic events with low thrust containing isolated leptons

### 6.1 Introduction

All PETRA experiments have set bounds on new heavy quarks [19], from which open production of new charged 2/3 quarks is excluded up to  $m = 23.3 \text{ GeV}$ , and for charged 1/3 quarks up to  $m = 22.7 \text{ GeV}$ . Near threshold, events from heavy quarks would give rise to a spherical event topology, and leptons therein are expected to be produced by the semileptonic decays of the quarks.

Recently the Mark J group [1] reported the observation of an excess of muon inclusive hadronic events with low thrust

$$T = \max \frac{\sum_i |\vec{p}_i \cdot \vec{n}_T|}{\sum_i |\vec{p}_i|} \quad (5)$$

at the highest PETRA energies,  $\sqrt{s} \geq 46.3 \text{ GeV}$ , where the muon had a large isolation angle  $\delta$  with respect to the thrust axis  $\vec{n}_T$ , as shown in fig. 11. A search for this type of events within




Figure 11: Definition of isolation angle  $\delta$

the JADE data sample of  $22.8 \text{ pb}^{-1}$  in the energy range  $38.66 \text{ GeV}$  to  $46.3 \text{ GeV}$  and  $1.8 \text{ pb}^{-1}$  between  $46.3 \text{ GeV}$  and  $46.78 \text{ GeV}$  is described in this chapter.

### 6.2 Inclusive muons

The analysed events were demanded to fulfill the multihadron conditions. Muons used in this chapter were required to satisfy the "loose" criteria as described in section 2.2. In order to allow comparison of results, cuts identical to those used by Mark J were chosen to define the signal region, i. e.  $T < 0.8$  and  $|\cos \delta| < 0.7$ . The results for  $\sqrt{s} \geq 46.3 \text{ GeV}$  were compared to the expectation calculated from data with  $\sqrt{s} < 46.3 \text{ GeV}$ . In fig. 12 the inclusive muon events are shown in the  $T - |\cos \delta|$  plane for both of the centre of mass energy intervals, with the cuts indicated. The signal region for  $\sqrt{s} \geq 46.3 \text{ GeV}$  contains five events compared to an expectation of  $(0.56 \pm 0.18)$  events from an extrapolation of the lower energy data<sup>2</sup>. This expectation is in agreement with a Monte Carlo simulation of multihadron production with five flavors which predicts  $(0.74 \pm 0.11)$  events in the signal region. The probability of a fluctuation of the observed magnitude (or larger) is less than 0.4 %.

<sup>2</sup>Recently we learnt that Mark J [20] did not include the muon when calculating the thrust, and that they did not balance the event before the calculation, as we do. Applying the Mark J procedure, we find a sixth event at  $\sqrt{s} \geq 46.3 \text{ GeV}$ , and the expectation changes slightly to  $0.69 \pm 0.21$ .




Figure 12:  $T$  and  $|\cos \delta|$  for muon inclusive events

The energy flow of the five events, labelled A - E, at  $\sqrt{s} \geq 46.3$  GeV with  $T < 0.8$  and  $|\cos \delta| < 0.7$ , is shown in the plots of fig. 13. In these diagrams the energy flow is rolled out in the  $\cos \theta - \phi$  plane, where  $\theta$  is the polar angle with respect to the beam axis, and  $\phi$  the azimuthal angle. Muons and electromagnetic showers with energies  $E_{e,m.} > 5$  GeV are marked. Event D contains two muons and has already been shown in fig. 7. We note the high muon momenta in events A and D and the energetic electromagnetic clusters, from either an electron or a photon in events A, D and E. Large electromagnetic clusters were also observed by Mark J in their excess events [21]. Except for event B, which is the candidate event with high transverse jet mass and a high  $p^T$  muon found previously in a heavy quark search [6], the events are planar, as can be seen from the  $Q$  - plots [22] attached to the legoplots. The visible energy and missing transverse momentum show no peculiarities when compared to ordinary multihadron events.

In this analysis the standard muon criteria for new particle searches ("loose" cuts) were used, but only two of the five muons fulfill the stricter requirements of the "tight" selection<sup>3</sup>. Two of the others hit regions of the muon filter with fewer chambers, where hadron rejection is reduced.

The observed excess might indicate the threshold of some new particle production above  $\sqrt{s} = 46.3$  GeV, and hence careful checks of the background were performed.

<sup>3</sup>The second muon in event D and the muon in the sixth event, which appears when muons are not included in the thrust calculation, satisfy also the "tight" criteria.




Figure 13: Energy flow diagrams of the five  $\mu$  - events A - E with  $T < 0.8$  and  $|\cos \delta| < 0.7$  at  $\sqrt{s} \geq 46.3$  GeV. The corresponding  $Q$  - plots are also shown.  $Q_1, Q_2, Q_3$  are the eigenvalues of the sphericity tensor,  $(Q_3 - Q_2)/\sqrt{3}$  being plotted as abscissa and  $Q_1$  as ordinate in the  $Q$  - plots.

### 6.3 Background

Various tests have been carried out to check whether the observed events are due to an enhanced background of hadrons misidentified as muons at  $\sqrt{s} \geq 46.3$  GeV as compared to  $\sqrt{s} < 46.3$  GeV. A possible source of background is the increased emission of synchrotron radiation during the high energy scan, which caused spurious hits in the muon filter, especially in the forward region.

The fraction of events containing a muon for the two energy intervals agrees within errors, and also the angular distribution of the muons shows no significant excess over the entire polar range, as shown in fig. 14.




Figure 14: Muon angular distribution:  $\theta$  is the angle between the muon and the beam axis. The arrows mark the polar position of the five signal muons.

The probability that a given hadron track with  $p > 1.8$  GeV is classified as a muon can be determined from the data in the following way. The sample of observed tracks satisfying the muon selection criteria consists of prompt muons and fake muons. The prompt contribution is known from a Monte Carlo simulation, assuming that there are no new processes involved. The fake contribution is then the difference between the number of tracks satisfying the muon selection criteria and the prompt contribution. At the energy of interest,  $\sqrt{s} \geq 46.3$  GeV, the fake probability was calculated to be  $P_{h \rightarrow \mu} = (1.8 \pm 0.5)\%$ . This number is in agreement with a measurement of the  $\pi$  punch-through + decay probability of  $(1.5 \pm 0.3)\%$  using  $\tau^+ \tau^-$  pairs as a source of pion tracks.

In order to estimate the contribution of fake muons to the signal region, events with  $T < 0.8$  and at least 1 track, irrespective of particle type, with  $p > 1.8$  GeV and  $|\cos \delta| < 0.7$  were selected at  $\sqrt{s} \geq 46.3$  GeV. 30 events were found, containing in total 50 tracks fulfilling the above requirements. Assuming a probability of  $P_{h \rightarrow \mu} = (1.8 \pm 0.5)\%$  as determined above to fake a muon, we estimate a background contribution of  $0.90 \pm 0.25$  events to the observed signal of five events. This number is considered to be an upper limit, since most of the hadron tracks lie within jets, where the fake probability is larger than for the isolated tracks we are considering. From prompt muons, an additional  $0.22 \pm 0.03$  events are expected in the signal region.

When the cut for the muon hit association to the jet chamber track is relaxed from  $2 \sigma$  to

$5 \sigma$ , we would expect a corresponding increase in the number of muons in the signal region, if they were due to random hits. Above 46.3 GeV no additional event was found in the signal region after relaxing the muon cut.

To summarize, though some of the muons do not belong to the cleanest sample, no indication was found that the five muon events are due to enhanced hadron contamination at  $\sqrt{s} \geq 46.3$  GeV.

### 6.4 Inclusive electrons

The same multihadron data sample was searched for an excess of similar events with isolated electrons. For the electrons with  $p > 1.8$  GeV we do not observe any event in the signal region ( $T < 0.8$  and  $|\cos \delta| < 0.7$ ) above 46.3 GeV, and expect 0.7 events from an extrapolation of the lower energy data (see fig. 15). The electron in event E does not fulfill the quality requirements on the  $dE/dx$  measurement due to a nearby track. Another electron in event A is removed by the photon conversion cuts.




Figure 15:  $T$  and  $|\cos \delta|$  for electron inclusive events

In ordinary multihadron events the detection efficiencies for electrons and muons coming from  $b$ - and  $c$ - decays are similar within the acceptance cut  $|\cos \theta| < 0.76$ . However, Monte Carlo studies have shown that multiparticle effects are more important for the detection of electrons than for muons. In particular for close tracks it may happen that an electromagnetic shower cannot be uniquely assigned to one track. Such overlap problems are less severe in muon identification, where the presence of a muon would still be recognized in the muon filter. Therefore the electron

detection efficiency is more strongly influenced by the special event topology caused by a possible new process.

In order to roughly estimate how many events with isolated electrons in a low thrust topology would correspond to the five observed muon events, the muon tracks in these events were replaced by electrons, and the electron selection criteria were applied. Two of the replaced muons are outside the electron acceptance  $|\cos\theta| < 0.76$ , two others have a high probability of escaping detection due to nearby particles, and one event has a similar detection probability as either an electron or a muon event. These studies imply that the absence of electron events in our present data does not contradict lepton universality.

## 6.5 Comparison with other experiments

The cross section for inclusive muon events with  $T < 0.8$  and  $|\cos\delta| < 0.7$  is plotted as a function of the centre of mass energy in fig. 16 and compared with the result of Mark J [21]. The data are not corrected for acceptance. Both experiments, Mark J and JADE, observe an excess in the highest energy bin. Mark J sees seven events and expects  $0.5$  in its data sample of  $2.8 \text{ pb}^{-1}$ .




Figure 16: Cross section for muon events with  $T < 0.8$  and  $|\cos\delta| < 0.7$ . The error bars are shown to give a feeling for possible fluctuations; for such low statistics they cannot be used to calculate significances in terms of standard deviations.

CELLO [23] does not confirm this observation, observing one event at  $\sqrt{s} \geq 46.3 \text{ GeV}$  with an integrated luminosity of  $2.1 \text{ pb}^{-1}$ . Below  $46.3 \text{ GeV}$  the rates measured by CELLO agree with those of Mark J and JADE. CELLO's detection efficiency for the events observed by Mark J and JADE was found to be comparable to that of JADE and Mark J [23], [20].

## 7 Summary

A search for rare processes leading to events containing hadrons and muons or missing energy was conducted. Three events of the type two isolated muons – hadrons have been observed. The rate and their kinematic configurations agree with the QED expectation for the reaction  $e^+e^- \rightarrow q\bar{q}\mu^+\mu^-$ .

Leptoquarks of the second generation are excluded in the mass range  $5 \text{ GeV}$  up to  $20.8 \text{ GeV}$ , color octet muons up to  $23.2 \text{ GeV}$  and color octet neutrinos from  $9 \text{ GeV}$  up to  $21.9 \text{ GeV}$ .

At the highest PETRA energies,  $46.30 \text{ GeV} \leq \sqrt{s} \leq 47.78 \text{ GeV}$ , five low thrust events with isolated muons were observed, whereas only  $0.56 \pm 0.18$  events were expected from an extrapolation from lower energies. No satisfactory explanation for this excess in terms of background could be found. Similar observations were made by Mark J, while CELLO does not observe this effect. Events from a new heavy quark, e.g. a charged  $1/3$  quark of the fourth generation [24], which cannot yet be excluded by PETRA, are expected to be more spherical than most of the five observed events. With the close down of PETRA, it is left to experiments at TRISTAN to determine whether these events indicate more than a statistical fluctuation.

## Acknowledgements

We are indebted to the PETRA machine group and the DESY computer centre staff for their excellent support during the experiment and to all the engineers and technicians of the collaborating institutions who have participated in the construction and maintenance of the apparatus. We thank J. Fuster of the CELLO group and B. Wyslouch of the Mark J Group for their cooperation, and B. Schrempp, F. Schrempp and K. Hagiwara for helpful discussions. This experiment was supported by the Bundesministerium für Forschung und Technologie, by the Ministry of Education, Science and Culture of Japan, by the UK Science and Engineering Research Council through the Rutherford Appleton Laboratory and by the US Department of Energy. The visiting groups at DESY wish to thank the DESY directorate for the hospitality extended to them.

## References

- [1] Mark J Collaboration, B. Adeva et al., Phys. Rev. D34 (1986) 681
- [2] JADE Collaboration, W. Bartel et al., Phys. Lett. 88B (1979) 171  
B. Naroska, Phys. Rep. 148 (1987) 67
- [3] JADE Collaboration, W. Bartel et al., Phys. Lett. 129B (1983) 145
- [4] J. Allison et al., Nucl. Instr. Meth. A238 (1985) 220
- [5] J. Allison et al., Nucl. Instr. Meth. A238 (1985) 230
- [6] JADE Collaboration, W. Bartel et al., Phys. Lett. 160B (1985) 337
- [7] JADE Collaboration, W. Bartel et al., Z. Phys. C 33 (1987) 339
- [8] F. A. Berends, P. H. Daverfeldt, R. Kleiss, Phys. Lett. 148B (1984) 489, Nucl. Phys. B253 (1985) 441
- [9] B. Anderson, G. Gustafson und T. Sjöstrand, Phys. Lett. 94B (1980) 243  
T. Sjöstrand, Computer Phys. Comm. 27 (1982) 105
- [10] B. Schrempp, F. Schrempp, Phys. Lett. 153B (1985) 101
- [11] R. N. Mohapatra, G. Segrè, L. Wolfenstein, Phys. Lett. 145B (1984) 433
- [12] C. Peterson et al., Phys. Rev. D27 (1983) 105
- [13] JADE Collaboration, W. Bartel et al., Phys. Lett. 146B (1984) 126
- [14] JADE Collaboration, W. Bartel et al., Phys. Lett. 155B (1985) 288
- [15] CELLO Collaboration, H. - J. Behrend et al., Phys. Lett. 141B (1984) 217
- [16] CELLO Collaboration, H. - J. Behrend et al., Phys. Lett. 178B (1986) 452
- [17] H. Harari, Phys. Lett. 156B (1985) 250  
H. Harari, report WIS-85/6 Ph-Feb, WIS-85/16 Ph-March and WIS-85/23 Ph-June, Weizman Institute of Science, Rehovot 1985
- [18] S. F. King, S. R. Sharpe, Nucl. Phys. B253 (1985) 1
- [19] S. Komamiya, Proc. of the 1985 International Symposium on Lepton and Photon Interactions, Kyoto-1985
- [20] Mark J Collaboration, B. Wyslouch, private communication
- [21] A. Böhm, Proc. of the XXth Rencontre de Moriond, Les Arcs 1985/1, 559
- [22] JADE Collaboration, W. Bartel et al., Z. Phys. C 21 (1983) 37
- [23] CELLO Collaboration, H. - J. Behrend et al., DESY 87-016, Hamburg 1987
- [24] F. Cornet, E. W. N. Glover, K. Hagiwara, A. D. Martin and D. Zeppenfeld, Phys. Lett. 174B (1986) 224  
V. Barger, R. J. N. Phillips, A. Soni, Phys. Rev. Lett. 57 (1986) 1518