

DEUTSCHES ELEKTRONEN-SYNCHROTRON **DESY**

DESY 87-053
May 1987



THE CHIRAL ANOMALY OF ANTISYMMETRIC TENSOR FIELDS

by

M. Reuter

Deutsches Elektronen-Synchrotron DESY, Hamburg

ISSN 0418-9833

NOTKESTRASSE 85 · 2 HAMBURG 52

DESY behält sich alle Rechte für den Fall der Schutzrechtserteilung und für die wirtschaftliche Verwertung der in diesem Bericht enthaltenen Informationen vor.

DESY reserves all rights for commercial use of information included in this report, especially in case of filing application for or grant of patents.

**To be sure that your preprints are promptly included in the
HIGH ENERGY PHYSICS INDEX ,
send them to the following address (if possible by air mail) :**

**DESY
Bibliothek
Notkestrasse 85
2 Hamburg 52
Germany**

The Chiral Anomaly of Antisymmetric
Tensor Fields

M. Reuter

Deutsches Elektronen-Synchrotron DESY
Notkestrasse 85
2000 Hamburg 52

Abstract

For antisymmetric tensor gauge fields of rank $2n-1$ coupled to gravity in $4n$ dimensions it is shown that the symmetry under duality rotations is broken by quantum effects. The anomaly is related to a local version of the signature index theorem. The zeta-function technique, Fujikawa's method and the stochastic regularization scheme are discussed.

(I) Introduction

As was first pointed out by Alvarez-Gaumé and Witten [1], (anti-) self-dual antisymmetric tensor fields in $4n-2$ dimensions have anomalies in their coupling to gravity. Similar to the case of fermions, these gravitational anomalies are related to chiral anomalies in $4n$ dimensions [2]. For the antisymmetric tensor fields these chiral transformations are realized as duality rotations of a tensor field of rank $2n$. On the other hand, considering this field $F_{\mu_1 \dots \mu_{2n}} \equiv 2n \partial_{[\mu_1} A_{\mu_2 \dots \mu_{2n}]}$ as being the field strength of a gauge potential $A_{\mu_1 \dots \mu_{2n-1}}$ one is led to study the "duality anomaly" of a $U(1)$ gauge theory where the gauge fields are antisymmetric tensor fields of rank $2n-1$. For $n=1$, say, this means that there is an anomaly associated with duality transformations of the field strength $F_{\mu\nu}$ belonging to the photon A_μ .

Along a different line of investigation, Zakharov [3] recently used dispersion relation techniques to show that if the photon field is quantized in a 4-dimensional curved space-time (with Minkowski signature) for which the pseudo-scalar $\epsilon_{\mu\nu\rho\sigma} R^{\mu\nu}{}_{\alpha\beta} R^{\alpha\beta\rho\sigma}$ does not vanish, the vacuum expectation value of the Pauli-Ljubanski vector

$$K^\mu = g^{-\frac{1}{2}} \epsilon^{\mu\nu\rho\sigma} A_\nu \partial_\rho A_\sigma \quad (1.1)$$

is not conserved. This means that the pseudo-scalar $F_{\mu\nu} {}^* F^{\mu\nu}$ acquires a vacuum expectation value:

$$\nabla_\mu \langle K^\mu \rangle \equiv \frac{1}{2} \langle F_{\mu\nu} {}^* F^{\mu\nu} \rangle = \frac{1}{192 \pi^2} g^{-\frac{1}{2}} \epsilon_{\mu\nu\rho\sigma} R^{\mu\nu}{}_{\alpha\beta} R^{\alpha\beta\rho\sigma} \quad (1.2)$$

As we shall see below, this equation expresses the fact that quantum effects spoil the invariance of the classical theory under duality transformations. For a free electromagnetic field this symmetry causes the difference of the numbers of right and

left circularly polarized photons to be conserved [4]. Hence, if the RHS of (1.2) is non-zero, the gravitational field continuously produces (chiral) photons from the vacuum. This is analogous to the anomalous fermion pair creation by Yang-Mills fields as expressed by the famous relation

$$\Delta Q_5 = \frac{1}{8\pi^2} \int d^4x \operatorname{tr} (F_{\mu\nu} * F^{\mu\nu}) \quad (1.3)$$

where ΔQ_5 is the change of the chiral charge.

The purpose of this paper is to show how Zakharov's result is related to an anomalous breaking of the duality symmetry. Furthermore, we shall see that a similar effect exists in all $4n$ -dimensional theories containing antisymmetric tensor gauge fields of rank $2n-1$ coupled to gravity. Working in Euclidean space, we will relate the generalization of (1.2) to a local version of the signature index theorem so that for all n the anomaly can be expressed by the Hirzebruch L-polynomial [5,6]. In section (II) this result is derived using the zeta-function method for regularizing infinite dimensional determinants. Then, in section III, we show that, similar to the fermionic case, the chiral anomaly of antisymmetric tensor fields is associated with a non-trivial Jacobian of the path integral measure [7]. Finally, in section (IV) it is briefly described how the anomaly is obtained in the framework of stochastic quantization [8].

(II) Zeta-function regularization

We are considering a $4n$ -dimensional oriented Riemannian manifold \mathcal{M} of Euclidean signature which has no boundary: $\partial\mathcal{M} = \emptyset$. We define totally antisymmetric tensor fields $A_{\mu_1 \dots \mu_{2n-1}}(x)$ on \mathcal{M} which we frequently will write as differential forms:

$$A(x) = \frac{1}{(2n-1)!} A_{\mu_1 \dots \mu_{2n-1}}(x) dx^{\mu_1} \dots dx^{\mu_{2n-1}} \quad (2.1)$$

We associate to A a field strength $2n$ -form in the usual way [9]

$$F = dA \quad (2.2a)$$

where

$$F_{\mu_1 \dots \mu_{2n}} = 2n \partial_{[\mu_1} A_{\mu_2 \dots \mu_{2n}]} \quad (2.2b)$$

Obviously F is invariant under gauge transformations

$$A \rightarrow A + dx \quad (2.3)$$

for any $(2n-1)$ -form \mathcal{A} , so that for $n=1$ ordinary (Euclidean) electrodynamics is recovered. For our purposes it is convenient to introduce the scalar product

$$\begin{aligned} (\alpha, \beta) &\equiv *[\alpha \wedge * \beta] \\ &= \frac{1}{p!} \alpha_{\mu_1 \dots \mu_p} \beta^{\mu_1 \dots \mu_p} \end{aligned} \quad (2.4)$$

for all p -forms α and β . Here we used the Hodge operator $*$ defined as

$$* dx^{\mu_1} \dots dx^{\mu_p} = \frac{\sqrt{g}}{(4n-p)!} \varepsilon^{\mu_1 \dots \mu_p \mu_{p+1} \dots \mu_{4n}} dx^{\mu_{p+1}} \dots dx^{\mu_{4n}} \quad (2.5)$$

The action for A is the following generalization of the Maxwell action (ε denotes the volume form $\sqrt{g} dx^1 \dots dx^{4n}$):

$$S = \frac{1}{2} \int (F, F) \varepsilon = \frac{1}{2} \int (A, \delta d A) \varepsilon \quad (2.6)$$

The second equality follows from the fact that the co-derivative δ is the adjoint of d . We will perform all calculation in a generalized Lorentz gauge defined by

$$\delta A = 0 \quad (2.7)$$

For $n=1$ this reduces to the ordinary Lorentz condition $\nabla_\mu A^\mu = 0$,

where ∇_μ is the covariant derivative constructed from the metric of \mathcal{M} . For fields fulfilling (2.7) the action can be written as

$$S' = \frac{1}{2} \int (A, \Delta A) \varepsilon \quad (2.8)$$

where $\Delta = d\delta + \delta d$ is the Laplacian. In its original form (2.6), S is invariant under global duality rotations (or "chiral transformations") of the form

$$F \rightarrow F \cos \alpha + *F \sin \alpha, \quad (2.9)$$

or infinitesimally:

$$\delta_\alpha F = \alpha *F \quad (2.10)$$

Now we turn to our main task, namely the calculation of the vacuum expectation value of

$$\nabla_\mu K^\mu \equiv (F, *F) = \frac{1}{(2n)!} F_{\mu_1 \dots \mu_{2n}} *F^{\mu_1 \dots \mu_{2n}} \quad (2.11)$$

for the prescribed background gravitational field given by the metric $g_{\mu\nu}(x)$ of \mathcal{M} . (The generalization of (1.1) for arbitrary n can be read off from $(F, *F) = \delta K$ where $K = *(AdA)$.) We define the generating functional

$$Z[\eta] = \int [DA]_{LG} \exp \left\{ -\frac{1}{2} \int (A, \Delta A) \varepsilon + \int (dA, \eta *dA) \right\} \quad (2.12)$$

for any real, scalar function η because then the expectation value of (2.11) reads:

$$\begin{aligned} \langle (F(x), *F(x)) \rangle &\equiv \frac{\langle 0 | (F(x), *F(x)) | 0 \rangle}{\langle 0 | 0 \rangle} \\ &= \frac{\delta}{\delta \eta(x)} \ln Z[\eta] \Big|_{\eta=0} \end{aligned} \quad (2.13)$$

The subscript "LG" at the path integral measure is to indicate that the integration has to be performed only over fields obeying the Lorentz gauge condition (2.7). We are not going to exponentiate this constraint since it is much simpler to explicitly take it into account in doing the Gaussian integration (see below). In particular, any complication due to the necessity of introducing ghosts for the ghosts is avoided [10].

In the representation (2.13) we can evaluate $\langle (F, *F) \rangle$ using the same technique as developed in ref. [11] for fermionic chiral anomalies and subsequently used for the evaluation of various other types of anomalies [12]. Performing the integral for Z one formally obtains

$$\langle (F(x), *F(x)) \rangle = -\frac{1}{2} \frac{\delta}{\delta \eta(x)} \ln \det \Omega \Big|_{\eta=0} \quad (2.14)$$

where the operator Ω is given by

$$\Omega = \Delta + \mu^2 + 2 * (d\eta) \wedge d \quad (2.15)$$

To control the usual IR divergences associated with massless fields we have introduced a small mass parameter μ . For $\eta(x) \equiv 0$ the operator Ω is positive and hermitian. Because we may consider η infinitesimal, this is sufficient for the zeta-function method to be applicable, i.e. we can define the determinant as $\exp \{-f'(\Omega|0)\}$, where $f(\Omega|s)$ is the zeta-function associated with Ω [13]. Hence one has

$$\langle (F(x), *F(x)) \rangle = \frac{1}{2} \frac{\delta}{\delta \eta(x)} \frac{d}{ds} \Big|_0 f(\Omega|s) \quad (2.17)$$

To evaluate the RHS of (2.17) we introduce a complete set of normalized eigenfunctions of the operator $\Delta + \mu^2$ acting on $(2n-1)$ -forms:

$$(\Delta + \mu^2) a_i(x) = \lambda_i a_i(x), \quad \lambda_i > 0 \quad (2.18a)$$

$$\int (a_i, a_j) \varepsilon = \delta_{ij} \quad (2.18b)$$

It has been shown in ref. 11, for instance, that in terms of the a_i 's the functional derivative of the zeta-function can be written as

$$\frac{\delta \zeta(\Omega|s)}{\delta \eta(x)} = -s \sum_i \lambda_i^{-(1+s)} \int (a_i, \frac{\delta \Omega}{\delta \eta(x)} a_i) \varepsilon \quad (2.19)$$

As was already mentioned, the path integral (2.12) has to be performed only over fields satisfying $\delta A = 0$. Therefore it is only their eigenvalues which contribute to the determinant in (2.14). Consequently, in (2.19) the sum runs only over eigenvectors a_i fulfilling $\delta a_i = 0$. Inserting (2.15) yields

$$\langle (F(x), * F(x)) \rangle = \frac{d}{ds} \Big|_0 s \sum_i \lambda_i^{-(1+s)} (da_i(x), * da_i(x)) \quad (2.20)$$

At this point it is advantageous to change the normalization of the basis fields; we introduce

$$\alpha_i(x) = \lambda_i^{-\frac{1}{2}} a_i(x) \quad (2.21)$$

For $\mu \rightarrow 0$ the α_i are normalized according to

$$\int (d\alpha_i, d\alpha_j) = \delta_{ij} \quad (2.22)$$

Exploiting the identity

$$x^{-s} = \frac{1}{\Gamma(s)} \int_0^\infty dt t^{s-1} e^{-xt}, \quad x > 0 \quad (2.23)$$

we obtain

$$\langle (F(x), * F(x)) \rangle = \frac{d}{ds} \Big|_0 \frac{s}{\Gamma(s)} \int_0^\infty dt t^{s-1} e^{-\mu^2 t} \cdot \sum_i (d\alpha_i(x), * d\alpha_i(x)) e^{-\lambda_i t} \quad (2.24)$$

In this representation we can relate the expectation value of $(F, *F)$ to the index theorem for the signature complex [5,6]. Let us recall that the signature τ of a $4n$ dimensional manifold \mathcal{M} , i.e. the index of the signature complex is defined by

$$\tau(\mathcal{M}) = \text{Tr}_+ [e^{-t\Delta}] - \text{Tr}_- [e^{-t\Delta}] \quad (2.25)$$

The traces Tr_\pm refer to the space of self-dual and anti-self-dual $2n$ -forms, respectively. Note that because the projectors on these spaces are $P_\pm = \frac{1}{2} (1 \pm *)$ this also could be written as

$$\tau(\mathcal{M}) = \text{Tr} [* e^{-t\Delta}], \quad (2.26)$$

where the trace is over all $2n$ -forms now. (One even could perform the trace with respect to the whole exterior algebra; all additional contributions would cancel between the p - and the $(2n-p)$ -forms.) A standard argument shows that only the zero-modes of Δ contribute to the signature. Hence $\tau(\mathcal{M})$ is the difference of the number of self-dual and anti-self-dual zero modes of the Laplacian. The signature can be explicitly calculated from the asymptotic expansion of the relevant heat-kernels [5,14,15]:

$$k_\pm(x;t) \equiv \text{tr} \langle x | e^{-\Delta_\pm t} | x \rangle = \frac{1}{t^{2n}} \sum_{k=0}^\infty B_{2k}^\pm(x) t^k \quad (2.27)$$

The trace tr refers to the tensor indices only and $\Delta_\pm = \Delta P_\pm$ is the Laplacian restricted to the space of (anti-) self-dual fields. One finds

$$\tau(\mathcal{M}) = \int d^{4n}x \sqrt{g} B_{4n}(x), \quad (2.28)$$

where $B_{4n} = B_{4n}^+ - B_{4n}^-$. One possible strategy to evaluate the coefficients $B_{2k}(x)$ is to use a method similar to Fujikawa's computation of spinorial chiral anomalies [7]. This has recently been done by Endo and Takao [2]. They add to the kernel (2.27) additional tensor fields (cf. the remarks above) to form a Dirac-

Kähler fermion [1,16]; the computation is then similar to the evaluation of the anomaly for the Rarita-Schwinger field [17]. In accordance with the mathematical literature their result can be represented as

$$\tau(\mathcal{M}) = \int \det \left[\frac{(\Omega/2\pi)}{\tanh(\Omega/2\pi)} \right]^{1/2} \quad (2.29)$$

The integrand is the Hirzebruch L-polynomial for the curvature 2-forms

$$\Omega^{\lambda}_{\nu} = \frac{1}{2} R^{\lambda}_{\nu\rho\sigma} dx^{\rho} dx^{\sigma} \quad (2.30)$$

constructed from the metric of \mathcal{M} . This notation means that we have to expand the integrand in a power series in Ω and to keep only those terms which have the correct 4n-dimensional volume form.

To make contact with equation (2.24) we consider the kernel

$$\begin{aligned} K(x;t) &= K_+(x;t) - K_-(x;t) \\ &= \sum_i (f_i(x), *f_i(x)) e^{-\lambda_i t} \end{aligned} \quad (2.31)$$

The sum runs over a complete set of 2n-forms f_i with

$$\Delta f_i = \lambda_i f_i, \quad \lambda_i \geq 0, \quad (2.32)$$

and

$$\int (f_i, f_j) \varepsilon = \delta_{ij}. \quad (2.33)$$

According to the Hodge decomposition theorem, our 2n-forms f can be uniquely decomposed as a sum of an exact form (the derivative of a (2n-1)-form α), a co-exact form (the co-derivative of a (2n+1)-form β) and a harmonic form:

$$f = d\alpha + \delta\beta + \phi, \quad \Delta\phi = 0. \quad (2.34)$$

The three pieces are mutually orthogonal and fulfil the eigenvalue equation (2.32) separately. This means that we can divide the f_i 's into three classes: $f_i^{(1)} = d\alpha_i$, $f_i^{(2)} = \delta\beta_i$ and $f_i^{(3)} = \phi_i$. Hence (2.31) decomposes according to

$$\begin{aligned} K(x;t) &= \sum_{\alpha} (d\alpha_i, *d\alpha_i) e^{-\lambda_i t} + \sum_{\beta} (\delta\beta_i, *\delta\beta_i) e^{-\lambda_i t} \\ &\quad + \sum_i (\phi_i, *\phi_i). \end{aligned} \quad (2.35)$$

It is important to note that the first two sums in (2.35) are equal. This follows from the identity

$$(\delta\beta_i, *\delta\beta_i) = (d[*\beta_i], *d[*\beta_i])$$

and the fact that the Hodge operator provides an isomorphism between the space of the α 's and the β 's. Furthermore, this sum coincides with the one appearing in equation (2.24) since for $\lambda \neq 0$ the spectrum of Δ acting on α and $d\alpha$ coincides and the normalization (2.22) is the same as in (2.33).

Returning to (2.24) we may write

$$\langle (F(x), *F(x)) \rangle = \frac{1}{2} \frac{d}{ds} \Big|_0 \frac{s}{\Gamma(s)} \int_0^{\infty} dt t^{s-1} e^{-\mu^2 t}. \quad (2.36)$$

$$\cdot \left\{ K(x;t) - \sum_i (\phi_i(x), *\phi_i(x)) \right\}.$$

Inserting the expansion (2.27) one easily finds the μ -independent relation

$$\nabla_{\mu} \langle K^{\mu} \rangle = \langle (F(x), *F(x)) \rangle = \frac{1}{2} B_{4n}(x) - \frac{1}{2} \sum_i (\phi_i(x), *\phi_i(x)) \quad (2.37)$$

This is the desired result. It states that apart from the 2n-form zero modes of the Laplacian the vacuum expectation value is given by the well known Seeley coefficients $B_{4n}(x)$ [5,6,14]. For $n=1$ and 2 they are explicitly given by

$$B_4(x) = \frac{1}{96\pi^2} g^{-1/2} \varepsilon_{\mu\nu\rho\sigma} R^{\mu\nu}{}_{\alpha\beta} R^{\alpha\beta\rho\sigma}$$

$$B_8(x) = \quad (2.38)$$

$$\frac{1}{4608\pi^4} g^{-1/2} \varepsilon^{\mu_1 \dots \mu_8} R^{\nu_1}_{\nu_2 \mu_1 \mu_2} R^{\nu_2}_{\nu_3 \mu_3 \mu_4} R^{\nu_3}_{\nu_4 \mu_5 \mu_6} R^{\nu_4}_{\nu_5 \mu_7 \mu_8} \\ - \frac{7}{11520\pi^4} g^{-1/2} \varepsilon^{\mu_1 \dots \mu_8} R^{\nu_1}_{\nu_2 \mu_1 \mu_2} R^{\nu_2}_{\nu_3 \mu_3 \mu_4} R^{\nu_3}_{\nu_4 \mu_5 \mu_6} R^{\nu_4}_{\nu_5 \mu_7 \mu_8} .$$

If we integrate (2.37) over a manifold which has no boundary so that the $\nabla_\mu \langle K^\mu \rangle$ -term does not contribute, we recover the signature index theorem

$$n_+ - n_- = \int d^{4n}x \sqrt{g} B_{4n}(x) \quad , \quad (2.39)$$

where n_+ and n_- denotes the number of self-dual and anti-self-dual zero modes. For $n=1$ we have found

$$\nabla_\mu \langle K^\mu \rangle = \frac{1}{2} \langle F_{\mu\nu} * F^{\mu\nu} \rangle \\ = \frac{1}{192\pi^2} g^{-1/2} \varepsilon_{\mu\nu\rho\sigma} R^{\mu\nu}{}_{\alpha\beta} R^{\alpha\beta\rho\sigma} - \frac{1}{2} \sum_i (\phi_i * \phi_i) \quad (2.40)$$

This differs from Zakharov's result (1.2) for Minkowski space by the zero mode term. The situation is similar to the case of the fermionic anomaly. In Euclidean space the complete form of the axial vector divergence reads [18]

$$\nabla_\mu \langle j_5^\mu \rangle = -\frac{1}{384\pi^2} g^{-1/2} \varepsilon_{\mu\nu\rho\sigma} R^{\mu\nu}{}_{\rho\sigma} R^{\alpha\beta\rho\sigma} \\ + \frac{1}{8\pi^2} \text{tr}(F_{\mu\nu} * F^{\mu\nu}) + 2 \sum_i \psi_{0i} \gamma_5 \psi_{0i} \quad , \quad (2.41)$$

where the ψ_{0i} 's are the zero modes of the Dirac operator \not{D} . Integrating equation (2.41) over a manifold with $\partial M = \emptyset$ one reproduces the index theorem for the spin complex. In analogy we therefore expect that in Minkowski space $\nabla_\mu \langle K^\mu \rangle$ is given by $\frac{1}{2} B_{4n}(x)$ for all n .

III. The anomalous Jacobian

In this section we show that the anomaly (2.40) can be understood as arising from a non-trivial Jacobian of the path integral measure. The problem one encounters in this approach is that the relevant transformations, the dual rotations (2.9) or (2.10), are defined in terms of F rather than in terms of the integration variable A . One possibility to avoid this complication is to use the first order formalism [19]. Here we use a different approach. We define a chiral transformation of A by the requirement

$$d(\delta_\alpha A) = \alpha * dA \quad . \quad (3.1)$$

This transformation law guarantees that (2.10) is fulfilled by $F = dA$. As we shall see below, for our purposes it is not necessary to solve this equation for $\delta_\alpha A$. Following Fujikawa [7], the derivation of the anomaly proceeds as follows. Consider the path integral

$$Z \equiv \int [dA]_{L_G} e^{-\frac{1}{2} \int (\alpha A, \alpha A) \varepsilon} \quad (3.2)$$

and change the integration variable from A to $A' = A + \delta_\alpha A$, where the parameter α is allowed to depend on the space time point x . Then the classical action changes by $\int \alpha (F, * F) \varepsilon$. For the Jacobian we make the ansatz

$$[dA]_{L_G}' = \exp \left\{ \int d^{4n}x \sqrt{g} \alpha(x) \mathcal{A}_{4n}(x) \right\} [dA]_{L_G} \quad (3.3)$$

Because Z is independent of α , one finds the "ward identity"

$$\int [dA]_{L_G} \left\{ (F(x), * F(x)) - \mathcal{A}_{4n}(x) \right\} e^{-S} = 0 \quad (3.4)$$

If \mathcal{A}_{4n} would vanish, K_μ would be conserved and the duality rotations would be an intact symmetry even at the quantum level. To show that this is not the case in general, we first expand the gauge field as

$$A(x) = \sum_i c_i a_i(x), \quad (3.5)$$

where the $(2n-1)$ -forms a_i are a complete set of basis vectors which are orthonormalized as in (2.18b). For convenience we choose them as eigenvectors of the Laplacian. Furthermore, because the integral (3.2) is only over fields fulfilling the Lorentz gauge condition, they are constrained to satisfy $\delta a_i = 0$. Thus one has

$$[\mathcal{D}A]_{LG} = \prod_i dc_i \quad (3.6)$$

Under $A \rightarrow A'$ the coefficients c_i change according to

$$c_i \rightarrow c_i' = \sum_j \int (a_i, [1 + \delta_\alpha] a_j) \varepsilon \equiv \sum_j (\delta_{ij} + M_{ij}) c_j, \quad (3.7)$$

i.e. the Jacobian for an infinitesimal transformation is given by

$$\begin{aligned} \det M &= \exp \left\{ \text{Tr} \ln(1 + M) \right\} \\ &= \exp \left\{ \sum_i \int (a_i, \delta_\alpha a_i) \varepsilon \right\} \end{aligned} \quad (3.8)$$

To be able to apply (3.1), we exploit the fact that the a_i 's are eigenfunctions of Δ (with eigenvalues λ_i) and satisfy $\delta a_i = 0$ to write

$$a_i = \lambda_i^{-1} (d\delta + \delta d) a_i = \lambda_i^{-1} \delta d a_i$$

Inserting this in (3.8), integrating by parts and using (3.1) one obtains

$$\begin{aligned} \sum_i \int (a_i, \delta_\alpha a_i) \varepsilon &= \sum_i \lambda_i^{-1} \int (\delta d a_i, \delta_\alpha a_i) \varepsilon \\ &= \sum_i \lambda_i^{-1} \int (d a_i, d \delta_\alpha a_i) \varepsilon \\ &= \sum_i \lambda_i^{-1} \int (d a_i, \alpha * d a_i) \varepsilon \end{aligned} \quad (3.9)$$

Again introducing the differently normalized function α_i of (2.21), we find by comparison with (3.3):

$$\mathcal{A}_{4n}(x) = \sum_i (d \alpha_i(x), * d \alpha_i(x)) \quad (3.10)$$

Because of the completeness of the α_i 's this is an ill defined quantity of the form $0 \cdot \infty$. (This is analogous to the sum $\sum_i \varphi_i^+ \delta_5 \varphi_i$ for the fermionic case.) We regularize it by introducing a Gaussian cutoff:

$$\mathcal{A}_{4n}(x) = \lim_{M \rightarrow \infty} \sum_i (d \alpha_i, * e^{-\lambda_i/M^2} d \alpha_i) \quad (3.11)$$

According to the discussion which led to (2.36) this is the same as

$$\begin{aligned} \mathcal{A}_{4n}(x) &= \lim_{M \rightarrow \infty} \frac{1}{2} \left\{ K(x, \frac{1}{M^2}) - \sum_i (\phi_i(x), * \phi_i(x)) \right\} \\ &= \frac{1}{2} \mathcal{B}_{4n}(x) - \frac{1}{2} \sum_i (\phi_i(x), * \phi_i(x)) \end{aligned} \quad (3.12)$$

This shows that, contrary to the claims in ref. 2, Fujikawa's method is very well applicable even in a first order formulation. The result coincides with that of the zeta-function method and the calculation clearly displays the analogy with the fermionic anomaly.

IV. Stochastic quantization

Recently stochastic quantization [8] received much attention as an alternative to the usual canonical or path integral quantization. One of the reasons might have been that this scheme provides a new type of invariant regularization which was hoped to respect simultaneously all symmetries of the field theory model under consideration. Later on it turned and, however, that the anomalies associated with continuous symmetries also appear within stochastic quantization [20,21]. It is only in the case of the parity violating anomaly in $2n+1$ dimensions that an ambiguity has been observed [22]. In this section we are going to explicitly show that the (continuous) duality anomalies, too, are unambiguously reproduced in the framework of stochastic quantization. Our essential tool will be the stochastic regulator function which was first introduced by Breit, Gupta and Zaks [23].

The basic ingredients to start with are the Langevin equation derived from the action (2.6),

$$\frac{\partial}{\partial \tau} A(x, \tau) = -\delta \alpha A(x, \tau) + \eta(x, \tau) \quad (4.1)$$

and the correlation function for the random source $\eta(x, \tau)$:

$$\begin{aligned} & \langle \eta_{\mu_1 \dots \mu_{2n-1}}(x, \tau) \eta^{\nu_1 \dots \nu_{2n-1}}(x', \tau') \rangle_{\eta} \\ & = 2 g^{-1/2}(x) g_{[\mu_1}^{\nu_1} g_{\mu_2}^{\nu_2} \dots g_{\mu_{2n-1}] }^{\nu_{2n-1}} \delta(x-x') \alpha_{\lambda}(\tau-\tau') \end{aligned} \quad (4.2)$$

Being a source for $A(x, \tau)$, the noise $\eta(x, \tau)$ is a $(2n-1)$ -form, too. In (4.2) we introduced the Breit, Gupta, Zaks regulator $\alpha_{\lambda}(\tau-\tau')$ defined by the properties [23]

$$\begin{aligned} \lim_{\lambda \rightarrow \infty} \alpha_{\lambda}(\tau-\tau') &= \delta(\tau-\tau') \ , \\ \alpha_{\lambda}(\tau-\tau') &= \alpha_{\lambda}(\tau'-\tau) \ , \\ \int d\tau' \alpha_{\lambda}(\tau-\tau') &= 1 \end{aligned} \quad (4.3)$$

The limit $\lambda \rightarrow \infty$ will be performed after all calculations have been done. As is well known, the stochastic quantization of gauge theories does not necessarily require the fixing of a gauge [24], but, nevertheless, it can be computationally advantageous to fix a gauge. In our case we would like to do this in a way so that the Langevin equation contains the complete Laplacian rather than the $\delta \alpha$ -operator only. To achieve this we perform a gauge transformation which depends on the stochastic time τ :

$$A'(x, \tau) = A(x, \tau) + d\chi(x, \tau) \quad (4.4)$$

This transformation changes the form of the Langevin equation, but it does not change any gauge invariant expectation value calculated from it [8,24]. Choosing χ to satisfy

$$\frac{\partial}{\partial \tau} d\chi + d\delta(A+d\chi) = 0 \quad (4.5)$$

the new Langevin equation has the desired form:

$$\frac{\partial}{\partial \tau} A'(x, \tau) = -\Delta A'(x, \tau) + \eta(x, \tau) \quad (4.6)$$

A possible solution of (4.5) is

$$\chi(x, \tau) = -\int_0^{\tau} d\tau' e^{-\Delta(\tau-\tau')} \delta A(x, \tau') \quad (4.7)$$

For notational simplicity we do the following calculations for $n=1$; the generalization will be obvious. To solve equation (4.6), we define

$$G_{\mu}^{\nu}(x, x'; \tau) = \langle \mu, x | e^{-\tau \Delta} | \nu, x' \rangle \quad (4.8)$$

The solution of (4.6) with the initial condition $A(x, 0)=0$ then reads

$$A_{\mu}^{\nu}(x, \tau) = \int_0^{\tau} d\tau' \int d^4x' \sqrt{g} G_{\mu\nu}(x, x'; \tau-\tau') \eta^{\nu}(x', \tau') \quad (4.9)$$

(We omit the prime from A again.) Employing this solution, we can compute

$$\langle (F(x), *F(x)) \rangle = \frac{1}{4} g^{-1/2} \varepsilon^{\mu\nu\sigma\rho}$$

$$\lim_{x' \rightarrow x} \frac{\partial}{\partial x'^{\mu}} \frac{\partial}{\partial x^{\rho}} \langle A_{\nu}(x') A_{\sigma}(x) \rangle \quad (4.10)$$

by straightforwardly evaluating the expectation value:

$$\langle A_{\nu}(x') A_{\sigma}(x) \rangle = \lim_{\lambda \rightarrow \infty} \lim_{\tau \rightarrow \infty} \langle A_{\nu}^{\eta}(x', \tau) A_{\sigma}^{\eta}(x, \tau) \rangle_{\eta} \quad (4.11)$$

$$\begin{aligned} &= \lim_{\lambda \rightarrow \infty} \lim_{\tau \rightarrow \infty} \int_0^{\tau} d\tau_1 \int_0^{\tau} d\tau_2 \int d^4 y_1 \sqrt{g(y_1)} \int d^4 y_2 \sqrt{g(y_2)} \\ &\quad \cdot G_{\nu\alpha}(x', y_1; \tau - \tau_1) G_{\sigma\beta}(x, y_2; \tau - \tau_2) \langle \eta^{\alpha}(y_1, \tau_1) \eta^{\beta}(y_2, \tau_2) \rangle_{\eta} \\ &= \lim_{\lambda \rightarrow \infty} \lim_{\tau \rightarrow \infty} \int_0^{\tau} d\tau_1 \int_0^{\tau} d\tau_2 2\alpha_{\lambda}(\tau_1 - \tau_2) G_{\nu\sigma}(x', x; 2\tau - \tau_1 - \tau_2). \end{aligned}$$

Inserting into (4.11) yields

$$\begin{aligned} \langle (F(x), *F(x)) \rangle &= \lim_{\lambda \rightarrow \infty} \lim_{\tau \rightarrow \infty} \int_0^{\tau} d\tau_1 \int_0^{\tau} d\tau_2 2\alpha_{\lambda}(\tau_1 - \tau_2) \\ &\quad \cdot \sum_i (da_i(x), *da_i(x)) e^{-\lambda_i(2\tau - \tau_1 - \tau_2)} \quad (4.12) \end{aligned}$$

Here the heat-kernel (4.8) has been represented in terms of a complete set of orthonormalized eigenfunctions of the Laplacian:

$$G^{\mu\nu}(x, x'; \tau) = \sum_i a_i^{\mu}(x) a_i^{\nu}(x') e^{-\lambda_i \tau} \quad (4.13)$$

Equation (4.12) holds for all values of $n \geq 1$ again. The following steps parallel Tzani's [21] treatment of the fermionic case.

Introducing

$$t = \tau_1 - \tau_2$$

$$T = \frac{1}{2}(\tau_1 + \tau_2)$$

we have

$$\begin{aligned} \langle (F, *F) \rangle &= \lim_{\lambda \rightarrow \infty} \lim_{\tau \rightarrow \infty} \left(\int_0^{\tau/2} dT \int_{-2T}^{+2T} dt + \int_{\tau/2}^{\tau} dT \int_{-2(\tau-T)}^{+\tau} dt \right) 2\alpha_{\lambda}(t) \\ &\quad \cdot \sum_i (da_i, *da_i) e^{-\lambda_i(2\tau - \tau_1 - \tau_2)} \quad (4.14) \end{aligned}$$

The properties (4.3) imply for the regulator function

$$\begin{aligned} \int_{-2T}^{2T} dt \alpha_{\lambda}(t) &= \Theta\left(T - \frac{1}{2\lambda^2}\right) + O\left(\frac{1}{\lambda^2}\right) \\ \int_{-2(\tau-T)}^{+2(\tau-T)} dt \alpha_{\lambda}(t) &= \Theta\left(\tau - \frac{1}{2\lambda^2} - T\right) + O\left(\frac{1}{\lambda^2}\right) \quad (4.15) \end{aligned}$$

Inserting this into (4.15) and doing the T-integration yields

$$\begin{aligned} \langle (F, *F) \rangle &= \lim_{\lambda \rightarrow \infty} \lim_{\tau \rightarrow \infty} \sum_i \lambda_i^{-1} (da_i, *da_i) \left\{ e^{\lambda_i(2\tau - \frac{1}{\lambda^2})} - e^{\lambda_i/\lambda^2} \right\} e^{-2\lambda_i \tau} \quad (4.16) \end{aligned}$$

At this point the τ -limit may be performed:

$$\langle (F, *F) \rangle = \lim_{\Lambda \rightarrow \infty} \sum_i \lambda_i^{-1} (d\alpha_i, *d\alpha_i) e^{-\lambda_i/\Lambda^2} \quad (4.17)$$

(To obtain strictly positive eigenvalues λ_i one could introduce a small "photon" mass as in section II.) Finally we can change the normalization of the a_i 's according to (2.21) and (2.22). What we find is

$$\langle (F, *F) \rangle = \lim_{\Lambda \rightarrow \infty} \sum_i (d\alpha_i, *e^{-\lambda_i/\Lambda^2} d\alpha_i) \quad (4.18)$$

i.e. the same expression as in Fujikawa's approach, cf. equation (3.11). Obviously, Fujikawa's prescription for cutting off the large eigenvalues is equivalent to using the Breit-Gupta-Zaks regularized version of the noise-noise correlation function. Hence the stochastic quantization scheme leads to the same anomaly as the path integral or zeta-function method. This gives further support to the assumption that the stochastic method correctly reproduces all anomalies associated with continuous symmetries.

V. Conclusions

Using various independent methods, we have established the Euclidean analogue of Zakharov's formula (1.2) for an arbitrary $4n$ -dimensional space-time. The result (2.37) differs from the Minkowski space version of the anomaly equation by the zero mode terms. Their presence allows the anomalous divergence equation to be interpreted as a local version of the signature index theorem. In the path integral formalism the anomaly manifests itself in a non-trivial Jacobian for the duality transformations. This result has been established within the second order formulation of the quantum theory, where duality transformations are defined by (3.1). We also found that the anomaly is correctly reproduced by the stochastic quantization procedure and that the stochastic regulator a_Λ is equivalent to Fujikawa's prescription. All these

properties of the duality, or chiral anomalies of antisymmetric tensor fields are very similar to those of spinor fields.

To close, let us ask for possible physical applications of (1.2) or its generalizations. An example of a metric for which the pseudo-scalar $\epsilon_{\mu\nu\rho\sigma} R^{\mu\nu}{}_{\alpha\beta} R^{\alpha\beta}{}_{\rho\sigma}$ does not vanish is the metric of a rotating mass distribution, a rotating star, say. This means that such a star permanently creates photons and thereby reduces its angular momentum in very much the same way as a dyon produces fermion pairs via (1.3) and thereby reduces its electric charge [3]. It is important to note that this radiation has nothing to do with the familiar Hawking radiation or the gravitational particle creation in expanding universes [25]. These phenomena have their origin in a non-trivial Bogolubov transformation between the creation and annihilation operators used by different observers: the vacuum of one observer can correspond to a state with non-zero particle number for another observer. On the other hand, the (pseudo) scalar $\epsilon_{\mu\nu\rho\sigma} R^{\mu\nu}{}_{\alpha\beta} R^{\alpha\beta}{}_{\rho\sigma}$ which is responsible for the anomalous photon creation, cannot be arranged to vanish by employing a particular vacuum state; it is the same for any observer.

I would like to thank Professor V.I. Zakharov (ITEP) for bringing these problems to my attention and for a helpful discussion.

References

- [1] L. Alvarez-Gaumé, E. Witten, Nucl. Phys. B234, 269 (1984)
- [2] R. Endo, M. Takao, Osaka University report OU-HET 101 (1987), unpublished
- [3] V.I. Zakharov, to be published
- [4] M.G. Calkin, Am. J. Phys. 33, 958 (1965);
D.M. Lipkin, J. Math. Phys. 5, 696 (1964)
- [5] P.B. Gilkey, Invariance Theory, the Heat Equation, and the Atiyah-Singer Index Theorem, Publish or Perish, Wilmington (1984)
- [6] T. Eguchi, P.B. Gilkey, A.J. Hanson, Phys. Rep. 66, 213 (1980)
- [7] K. Fujikawa, Phys. Rev. Lett. 42, 1195 (1979); Phys. Rev. D21, 2848 (1980)
- [8] G. Parisi, Wu-yong-shi, Sci. Sin. 24, 484 (1981);
B. Sakita, in 7th Johns Hopkins Workshop, edited by
G. Domokos and S. Kovesi-Domokos, World Scientific,
Singapore (1983);
P.H. Damgaard, H. Hüffel, to appear in Physics Reports
- [9] Our conventions are those of reference 6
- [10] P.K. Townsend, Phys. Lett. 88B, 97 (1979);
W. Siegel, Phys. Lett. 93B, 170 (1980);
H. Hata, T. Kugo, N. Ohta, Nucl. Phys. B178, 527 (1981)
- [11] M. Reuter, Phys. Rev. D31, 1374 (1985)
- [12] M. Reuter, W. Dittrich, Phys. Rev. D32, 513 (1985);
Phys. Rev. D33, 601 (1986);
W. Dittrich, M. Reuter, Selected Topics in Gauge Theories,
Lecture Notes in Physics, Vol. 244, Springer, New York
(1986)
- [13] J.S. Dowker, R. Critchley, Phys. Rev. D13, 3224 (1976);
S.W. Hawking, Commun. Math. Phys. 55, 133 (1977)
- [14] J.S. Dowker, J. Phys. A11, 347 (1978)
- [15] V.N. Romanov, A.S. Schwarz, Teor. Mat. Fiz. 41, 190 (1979);
A.S. Schwarz, Commun. Math. Phys. 64, 233 (1979)
- [16] P. Becher, H. Joos, Z. Phys. C15, 343 (1982);
T. Banks, Y. Dothan, D. Horn, Phys. Lett. 117B, 413 (1982)
- [17] R. Endo, M. Takao, Prog. Theor. Phys. 73, 803 (1985)
- [18] N.K. Nielsen, H. Römer, B. Schroer, Phys. Lett. 70B, 445 (1977);
B. Schroer, Acta Phys. Austr., Suppl. XIX,155 (1978) and
references therein
- [19] This has been done by the authors of reference 2. However,
their anomaly equation (2.10) has no simple interpretation
in the context of ordinary electrodynamics, say, since the
standard relation between F and A is lost in this approach.
- [20] J. Alfaro, M.B. Gavela, Phys. Lett. 158B, 473 (1985);
M.B. Gavela, N. Parga, Phys. Lett. 174B, 319 (1986);
R. Kirschner, E.R. Nissimov, S.J. Pacheva, Phys. Lett. 174B,
324 (1986); J. Magpantay, M. Reuter, DESY report 87-004
- [21] R. Tzani, Phys. Rev. D33, 1146 (1986)
- [22] E.R. Nissimov, S.J. Pacheva, CERN report TH4435/86,
unpublished;
M. Reuter, DESY report 86-153, to appear in Phys. Rev. D

- [23] J.D. Breit, S. Gupta, A. Zaks, Nucl. Phys. B233, 61 (1984)
- [24] D. Zwanziger, Nucl. Phys. B192, 259 (1981);
L. Baulien, D. Zwanziger, Nucl. Phys. B193, 163 (1981);
M. Namiki, I. Ohba, K. Okano, Y. Yamanaka, Prog. Theor.
Phys. 69, 1580 (1983)
- [25] For a review see N.D. Birrell, P.C.W. Davies, Quantum fields
in curved space, Cambridge University press (1982)