DEUTSCHES ELEKTRONEN-SYNCHROTRON DESY

DESY 87-078 July 1987



MIRROR FAMILIES AND RADIATIVE SU2 × U1-BREAKING

by

B. Lampe

Institut für Theoretische Physik, Universität Hannover

ISSN 0418-9833

NOTKESTRASSE 85 · 2 HAMBURG 52

DESY behält sich alle Rechte für den Fall der Schutzrechtserteilung und für die wirtschaftliche Verwertung der in diesem Bericht enthaltenen Informationen vor.

DESY reserves all rights for commercial use of information included in this report, especially in case of filing application for or grant of patents.

To be sure that your preprints are promptly included in the HIGH ENERGY PHYSICS INDEX , send them to the following address (if possible by air mail) :



DESY 87-078 July 1987

ITP-UH 11/87

Mirror Families and Radiative $SU_2 \ge U_1$ -Breaking

B.Lampe

Institut für Theoretische Physik der Universität Hannover

Abstract

We examine the possibility that a heavy fourth family with mirror quantum numbers is the driving force for radiative $SU_2 \times M_4$ -breaking in the framework of supersymmetric GUT's coupled to N=1 supergravity. We compare the results to the case of a sequential fourth family.

1. Introduction

We consider a supersymmetric grand unified theory coupled to N=1 supergravity with four generations of quarks and leptons. It is conceivable that the fourth generation is a mirror family which is the case, for instance, in supersymmetric σ -models based on the exceptional group $E_{\delta}[1]$. Mirror fermions [2] are fermions with $SU_{3} \times SU_{2} \times M_{4}$ quantum numbers identical to those of the known quarks and leptons but with opposite handedness.

We assume that supergravity is broken and that the matter fields of the GUT feel the supersymmetric breaking at the scale M_W (=the weak boson mass [3-7]). Formally this can be arranged as follows: The effects of the breaking appear in the effective Langrangian as explicit soft breaking terms with coefficient of $O(m_{3/2})$, where $m_{3/2}$ is the gravitino mass [6]. Therefore one simply has to assume $m_{3/2}$ to be of the order of M_W .

By looking at the oneloop renormalization group equations evolving from the Planck scale (M_P) to M_W several groups have shown that with this assumption the $SU_2 \times U_4$ - invariance of the Weinberg-Salam theory can be broken radiatively to electromagnetic U_1 [3,7]. In fact they have shown that M_W can be fitted to its experimental value by considering $m_{3/2}$ as a free parameter and choosing it adequately. Those groups work with three families. In general the $SU_2 \times U_4$ -breaking is driven by the biggest Yukawa coupling. Therefore lower limits on the possible value of the t-quark mass could be "derived".

With a fourth family that is not possible any more. A reasonable assumption might seem that now the top of the fourth generation should drive the $Su_2 \times u_4$ -breaking [4]. If the fourth family is a mirror family even this is questionable. Therefore we have scanned through all initial values for the Yukawa couplings of the fourth generation. We also considered the possibility that these are not too far from the corresponding value for the top quark, i.e. we did not neglect the influence of the third family on the breaking.

2. Renormalization Group Equations

Besides the four families we work with a minimal set of other fields. This means just two additional Higgs doublets H and H' with opposite U_1 -charge. For the case that all four families are ordinary families the renormalization group equations have been given in appendix B of ref.[5]. We call this "case N". Let u_3 be the Yukawa coupling of the t-quark and u_4, d_4 be the quark Yukawas of the fourth family. We assume that all other Yukawa couplings have no influence on the $SU_2 \times M_4$ -breaking. Then the superpotential is [5]

$$f_N \cong u_3 \mathcal{U}_3 \mathcal{Q}_3 \mathcal{H}' + \mu \mathcal{H} \mathcal{H}' + u_4 \mathcal{U}_4 \mathcal{Q}_4 \mathcal{H}' + d_4 \mathcal{D}_4 \mathcal{Q}_4 \mathcal{H}$$
(1)

Here Q_n are the left handed quark doublets of the n-th family and U_n, D_n the corresponding right handed singlets. Now we assume the fourth family to be a mirror family ("case M"). One has

$$f_{M} \cong \mathcal{M}_{3}\mathcal{M}_{3}\mathcal{Q}_{3}H' + \mu HH' + \mathcal{M}_{4}\mathcal{M}_{4}\mathcal{Q}_{4}H + \mathcal{M}_{4}\mathcal{D}_{4}\mathcal{Q}_{4}H' \qquad (1')$$

since the only difference between the two cases are the opposite U_1 -charges. In (1') we have excluded a term $\sim Q_3 Q_4$, , which would give rise to a pathological phenomenology of the mirror fermions [2]. The renormalization group equations depend only on the squares of the charges and ask only for what types of particle couple to what. Therefore the renormalization group equations for case M can be almost read off from those for case N and will not be written down here in full. We give only one characteristic example: Consider the mass m_{in} of the Higgs H' in the soft breaking terms. For case N we have

$$8\pi^{2} \frac{dm_{H}^{2}}{dt} = 3\left(m_{H^{1}}^{2} + m_{H^{3}}^{2} + m_{H^{3}}^{2} + \eta_{H^{3}}^{2}\right)M_{3}^{2} \qquad (2)$$
$$+ 3\left(m_{H^{1}}^{2} + m_{H^{4}}^{2} + m_{H^{4}}^{2} + \eta_{H^{4}}^{2}\right)M_{4}^{2} - g_{1}^{2}M_{4}^{2} - 3g_{2}^{2}M_{2}^{2}$$

Here $g_{1,2,3}$ are the coupling constants of the $SU_3 \times SU_2 \times U_1$ gauge group and $M_{4,2,3}$ the corresponding gaugino masses. t is related to the renormalization mass scale S via $t = M_3 \times M_p$. M_i are the soft masses of the fields i and γ_i are the coefficients of the corresponding trilinear soft terms.

For case M we have

$$8\pi^{2} \frac{dm_{H'}^{2}}{dt} = 3 \left(m_{H'}^{2} + m_{u3}^{2} + m_{q3}^{2} + \gamma_{u3}^{2} \right) u_{3}^{2}$$

$$+ 3 \left(m_{H'}^{2} + m_{p_{4}}^{2} + m_{q_{4}}^{2} + \gamma_{D_{4}}^{2} \right) d_{4}^{2} - g_{4}^{2} \mu_{4}^{2} - 3g_{2}^{2} \mu_{2}^{2}$$
(3)

The reason is simply that one should sum over all contributions from particles which couple to H' in the superpotential. Some remarks are in order.

i) For $u_3 = 0$ it is appropriate to say that the two cases differ from each other only by the interchange of H and H'. The reason is that in this case all the renormalization group equations are identical up to $m_H \leftrightarrow m_{H'}$. For $u_3=0$ there is no real difference between the two cases from the standpoint of the renormalization group equations. One must include the top to see a difference. The influence of the top quark will be the bigger the smaller the mass difference between the third and fourth family is. We shall discuss this in detail in section 4.

The equivalence of the two cases for $u_3 \equiv 0$ extends to the masses of U_4 and D_4 . This is obvious, since in case N the mass of U_4 is proportional to $\langle \mu^{*} \rangle$, while in case M it is proportional to $\langle \mu \rangle$ (c.f. equation (1)). (Exchanging m_{μ} and m_{μ} means exchanging the values of $\langle \mu \rangle$ and $\langle \mu^{*} \rangle$). One may convert these considerations into an argument that in case M $m(U_4) \langle m(D_4)$, whereas in case N $m(U_4) \rangle m(D_5)$ is to be expected. For this one needs the <u>assumption</u> that the mass relation $m(U_3) \gg m(D_3)$ for the third family is a hint for a hierachy $|\langle \mu^{*} \rangle| > |\langle \mu^{*} \rangle|$ and is only partly due to a hierachy of the top- and bottom-Yukawas at the Planck scale.

For $u_3 \neq 0$ a difference between case M and case N arises. For one u_3 is driven by d_4 in case M and by u_4 in case N and u_4 and d_4 behave differently because of their different charges. Second u_3 contributes to H' in both cases, so that one cannot exchange the role of H and H' any more. Therefore the quark masses are not correlated any more in the two cases.

ii) We have excluded a direct mass term between the two heavy families in case M. Therefore there is no mixing allowed between the third and the fourth family. Since we are interested in case N only for reasons of comparison, we neglect mixing also in case N. iii) Let us collect the free parameters that can be varied. At the Planck scale we put all soft masses equal to $m_{3/2}$. The trilinear soft breaking terms are all assumed to have coefficients $\gamma_i(M_p) = a_0 m_{3/2}$, where a_0 is a free parameter in the range $0.5 < a_0 < 3$. Similarly for the bilinear soft term one assumes a coefficient $\beta = b_0 m_{3/2}$ with $0.5 < b_0 < 2$ at M_P . μ and the μ_i are also assumed to be of the order of the gravitino mass:

μ(Mp) = co m312, μ; (Mp) = do m312 with 0.4 < co, do < 1.5

and all μ_i being equal at M_P . These assumptions can be justified in the simplest of supergravity models [6]. Besides that there are the three Yukawa-couplings, which in principle may take any value between 0 and, say, 5.

The procedure is now as follows: One picks up any of the possible values of a_0, b_0, c_0 and d_0 and keeps them fixed. Now one looks for values of $u_3(M_p), u_4(M_p)$ and $d_4(M_p)$, which realize the breaking of $SU_2 \times AU_1$ (c.f. section 3). This is done numerically. At this stage m_{312} is chosen to be 100 GeV. This is no restriction, because the symmetry breaking does not depend on m_{312} . Having found appropriate values of $u_3(M_p), u_4(M_p)$ and $d_4(M_p)$ one can change m_{312} in such a way that the vacuum expectation values come out as they should $(\langle H \rangle^2 + \langle H' \rangle^2 = U^2 = (A^2 + GeV)^2)$.

In general the range of values of $u_3(M_p)_1 u_4(M_p)$ and $d_4(M_p)$, that give the desired symmetry breaking, is quite restricted. From that one can deduce restrictions on the possible masses of $U_3 - U_4 - U_4$ and D_4 -quark. The restrictions are weakened, however, as soon as one also varies $a_{o_1}b_{o_1}c_{o_1}$ and d_0 . The results will be discussed in detail in section 4.

3. Spontaneous Symmetry Breaking

We have the Higgs potential as usually assumed in the literature [7]. Spontaneous symmetry breaking sets in at that value of t, where

$$S := (m_{H}^{2} + \mu^{2})(m_{H'}^{2} + \mu^{2}) - \beta^{2} \mu^{2}$$
(4)

becomes negative. For the potential to be bounded from below

$$C := m_{H}^{2} + m_{H}^{2} + 2\mu^{2} - 2|\beta\mu|$$
 (5)

must remain positive over the whole range $M_w < \zeta < M_p$. This usually implies that the point where S becomes negative is not far above M_W . It also implies that

$$\sin 2\theta = \frac{12\beta\mu 1}{(m_{H}^{2} + m_{H'}^{2} + 2\mu^{2})}$$
(6)

defines an angle θ . θ should be choosen in such a way that $\cos 2\theta < 0$, if $m_{\mu}^2 < m_{\mu'}^2$, v can be given by means of this angle.

$$\nabla^{2} = \frac{2}{(g_{1}^{2} + g_{2}^{2}) | \cos 2\theta |} \left[|m_{H}^{2} - m_{H'}^{2}| - (m_{H}^{2} + m_{H'}^{2} + 2\mu^{2}) | \cos 2\theta | \right]$$
(7)

In fact Θ parametrizes the relative strenth of the vacuum expectation values of II and H':

<h> = u sin 0</h>	(80)
< H'> = v cos O	(85)

Because of (1) and (8) the quark masses are

$$m_{\mu\nu} = \mu_3 v |\omega_s \Theta| \qquad (9a)$$

 $m_{\mu\nu} = \mu_{\mu} v \left[\cos \theta \right] \tag{9b}$

 $m_{D+} = d_{+} v |\sin \Theta| \qquad (9c)$

in case N. To get case M one has to interchange cos and sin in (9b) and (9c).

From equation (3) and the initial conditions (iii) one can deduce that $m_{H'}$ changes linearly with $m_{3/2}$. The same is true for $m_{H'}$ and μ . Therefore according to equation (7) it is also true for $v - a_s$ was anticipated at the end of section 2.

4. Results

First we discuss the features that are independent of whether one considers case N or M. a) The values of the Yukawa couplings at M_W usually are rather independent from the choice of their initial values at M_P (c.f. fig. 1 of ref. [4]).

b) The point, at which the symmetry breaking sets in, is mainly determined by the maximum of these initial values. For initial values above 0.5 S becomes negative almady at high energies. so that at M_W the consistency condition C>0 is violated. (In such a case one has to calculate one loop corrections to decide which is the true vacuum [4,8]. We will not pursue this possibility here, but stick to smaller values of the Yukawa couplings, where the symmetry is broken at the tree level.) c) For initial values below 0.05 S never becomes negative.

The effect b produces an upper limit for the masses of the heavy quarks. If one scans through the $a_0-b_0-c_0-d_0$ -parameter space one does not find bigger masses than 200 GeV [8].

The combined effects a and b restrict the values of masses for a fixed set of parameters $a_{o_1}b_{o_1}c_o$ and d_o . If these are also varied, however, no quantitative predictions are possible any more. Therefore we will discuss only the qualitative features of our result. As a characteristic example we may choose $a_o=A$, $b_o=o.8$, $c_o=o.5$, $d_o=0.8$, $m_{3/2}=100$ GeV.

We know that cases M and N are equivalent for us o up to a trivial interchange sin **9** . Varying the ratio $u_{\mu}(M_{p})/d_{\mu}(M_{p})$ one can accomodate any mass cos **O** ratio $m(\mathcal{M}_{u})/m(\mathcal{P}_{u})$ in both cases. There is no strict proportionality between the two is always such that it makes $\max\left\{\frac{m(U_n)}{m(D_n)}, \frac{m(D_n)}{m(U_n)}\right\}$ larger , i.e. Θ always supports the higher mass quark. If one ratios. In fact the value of Θ than $\max\left(\frac{u_{\mu}(M_{p})}{d_{\mu}(M_{p})}, \frac{d_{\mu}(M_{p})}{u_{\mu}(M_{p})}\right)$, i.e. Θ always supports the higher mass quark. If on switches on a small $u_{\lambda}(M_{p})$, this produces not only a top mass, but also increases $m(\mathcal{U}_{\mu})$ by an amount of the order of the top mass. To examine the differences and w (D.) between the two cases M and N we consider certain specific cases. In case N we think it is an interesting phenomenological possibility that $m(\mathcal{A}_{ij}) > m(\mathcal{A}_{jj}) > m(\mathcal{A}_{ij})$. Therefore we examined the extreme case $W_{i_{p}}(H_{p}) = M_{3}(H_{p}) = :C_{N,j}d_{i_{p}} \equiv 0$. We found that there is a broad window (0.42 $\leq c_N \leq 0.48$) which allows for spontaneous symmetry breaking. $m(\mathcal{U}_3)$ come out the same and vary in the range 100 GeV $\leq m(\mathcal{M}_{3,*}) \leq 125$ GeV $\stackrel{\text{def}}{\longrightarrow}$ It is and m(U_) interesting that under the above assumptions only a very tiny SSB window exists in case M. In fact for case M the interesting phenomenological possibility is $m(D_{4}) \ge m(\mathcal{U}_{3}) > m(\mathcal{U}_{4})$. The associated extreme case is $d_{\psi}(M_p) = c_{\psi}(M_p) = : c_{M_1} \cup u_{\psi} \equiv 0$. This time there is a window $(0.12 \le c_m \le 0.18)$ only in case M, but not in case N. The masses vary in the range 99 GeVem (U3) \$ 125 wey *) . 36 GeVem (D4) = 121 GeV * (The D4 - mass always comes out slightly smaller than the top mass. This is not a significant effect. It can be reversed easily by choosing $d_4(M_P)$ slightly bigger than $u_3(M_P)$.)

In table 1 we have listed the width of the SSB window for some interesting cases, among them also cases where M and N do not behave differently. (We know already from section 2 that the rows a.b and c of table 1 must be symmetric under exchange of M and N. Now we see that the same is true for the row d.)

5. Conclusions

We have examined the effect of a heavy mirror family on supergravity induced breaking of $SU_{2x}M_{4}$. As compared to a sequential family the role of fourth up- and down-type quarks in the alignment of the vaccuum is reversed. Therefore naively one expects the mass of the fourth up-type quark to be lower than that of the fourth down-type quark. We have discussed some renormalization group arguments in favour of this expectation. We have also examined the influence the top quark has on it. In particular we have elaborated on the possibility that the mass of the top is higher than that of the fourth up-type quark. Our results should not be taken as quantitative predictions, because there are two many unknowns in the game. For instance we did not discuss the electron of the fourth family which may or may not have a bigger influence on the symmetry breaking than the top. Also we did not discuss in detail the effect of other values a_0 , b_0 , c_0 and d_0 . We only note in passing that for all of them a similar picture arises. Varying them is only of use, if one wants to fit the biggest quark masses to some future experimental value.

Acknowledgements

I would like to thank W. Buchmüller and N. Falck for many helpful and stimulating discussions and N. Dragon for some suggestions on the final version of the manuscript.

Footnote

This range remains true even for other values of a_0, b_0, c_0 and d_0 like, for instance, the supergravity inspired [5] combination $a_0 = 3, b_0 = 2, c_0 = 4$. Note that for these values the width of the windows is in general smaller than for those discussed in the main text.

	۰.	· · · ·			
		$a_0 = 1$, $b_0 = .8$ $c_0 = .5$ $d_0 = .8$		$a_0 = 3$ $b_0 = 2$ $c_0 = 1$ $d_0 = 1$	
	Yukawas at M p	case M	case N	case M	case N
.)	$u_{4} >> d_{4}, u_{3}$ ($\Leftrightarrow u_{3} >> d_{4}, u_{4}$)	0.174 - 0.362	0.174 - 0.362	0.226 - 0.343	0.22 6 - 0.343
,)	a ₄ >> u ₃ , u ₄	0.172 - 0.384	0.172 - 0.384	0.243 - 0.357	0.243 - 0.357
•)	$u_{1} = d_{1} >> u_{3}$	0.168 - 0.172	0.168 - 0.172	0.178	0.178
.)	$u_3 = u_4 = d_4$	0.111 - 0.120	. 0.111 - 0.120	0.135 - 0.136	0.135 - 0.136
·)	$u_3 = u_4 >> d_4$	0.160	0.110 - 0.196	no window within the numerical error	0.152 - 0.189
•)	$u_3 = d_4 >> u_4$	0.110 - 0.196	0.160	0.152 - 0.189	no window within the numerical error

and the and the second s

.....

Table 1: SSB windows; the error is always 1 in the last digit

. .. - ----- -

References

- C.L.Ong, Phys.Rev.D31 (1985) 3271
 K.Itoh, T.Kugo, H.Kunimoto, Progr.Theor.Phys.75(1986)386
 W.Buchmüller, O.Napoly, Phys.Lett.163B(1985)161
 S.Irie, Y.Yasue, Z.Phys.C29(1985)123
- [2] G.Senjanovic, F.Wilczek, A.Zee, Phys. Lett.141B(1984)389 and references therein
- [3] L.Alvarez-Gaume, J.Polchinski, M.B.Wise, Nucl. Phys. B221(1983)495
- [4] M.Cvetic, C.R. Preitschopf, Nucl. Phys. B272 (1986)490
- [5] N.K.Falck, Z.Phys.C30(1986)247
- [6] R.Barbieri, S.Ferrara, C.A.Savoy, Phys.Lett. 119B(1982)343
- [7] K.Inoue, A.Kakuto, H.Komatsu, H.Takeshita, Progr. Theor. Phys. 67(1982)1889, 71(1984)413
- [8] H.Goldberg, Northeastern University preprint NUB 2680 (1985)

Average a second s

[9] K.Enquist, M.Mursula, M.Ross, Nucl. Phys. B226(1983)121 and references therein