DEUTSCHES ELEKTRONEN-SYNCHROTRON DESY

DESY 85-142 December 1985 受入 86-3--33 島王研図書室

TOPOLOGICAL CHARGE OF (LATTICE) GAUGE FIELDS

by

M. Göckeler

Institut f. Theoretische Physik, Universität Heidelberg

M.L. Laursen

NORDITA, Copenhagen

G. Schierholz

Institut 6. Theoretische Physik, Universität Kiel

and

Deutsches Elektronen-Synchrotron DESY, Hamburg

U.-J. Wiese

Institut 6. Theoretische Physik, Universität Hannover

ISSN 0418-9833

NOTKESTRASSE 85 · 2 HAMBURG 52

DESY behält sich alle Rechte für den Fall der Schutzrechtserteilung und für die wirtschaftliche Verwertung der in diesem Bericht enthaltenen Informationen vor.

ی در ۲۰۰۳ این در ۱۹۹۱ در به دولار سر ۲۰ ۱۹۹۲ در ۲۰۱۰ در ۲۰۰۰ ۱۹۹۵ در ۲۰۱۰ در ۲۰۰۱

n an Central Control of Central Centra

. An est Sect.

and the second second

Ύ.

ni sedi ding

Second Second

i agenti i s

14

y an film la in shi 12

. na i di s

હેસ

2014

net.

19 1. B. 19 1.

ad her

294

DESY reserves all rights for commercial use of information included in this report, especially in case of filing application for or grant of patents.

> To be sure that your preprints are promptly included in the HIGH ENERGY PHYSICS INDEX ,

send them to the following address (if possible by air mail) :

DESY Bibliothek Notkestrasse 85 2 Hamburg 52 Germany

DESY 85-142 December 1985 ISSN 0418-9833

I. Introduction

Differentiable SU(N) gauge fields on a compact manifold **M** carry a topological charge¹⁾

$$Q = -\frac{1}{32\pi^2} \int d^4x \, \epsilon_{\mu\nu\rho\sigma} \, Tr \left[\overline{T}_{\mu\nu} \, \overline{T}_{\rho\sigma} \right], \qquad (1)$$

where

$$\overline{T}_{\mu\nu} = \partial_{\mu}A_{\nu} - \partial_{\nu}A_{\mu} + [A_{\mu}, A_{\nu}].$$
⁽²⁾

The charge Q is a measure for topologically non-trivial properties of the gauge fields, which have been argued to play an important role in the physics of the vacuum of QCD and SU(N) gauge theories.

Preliminary results of calculations of Q in SU(2) gauge theory on the lattice^{2,3)} hold out hope for a quantitative resolution of the U(1) problem⁴⁾. The recent finding⁵⁾ that the vacuum of the quantized (pure) SU(2) gauge theory possesses an underlying instanton structure brings us furthermore in touch with semi-classical ideas of the QCD vacuum⁶⁾ and a possible mechanism for chiral symmetry breaking⁷⁾, which could be the beginning of a better understanding of the non-perturbative phenomena of these theories.

So far these investigations have been limited to smaller lattices and hence to smaller values of β . The difficulty involved is that a meaningful transcription of the topological charge (1), as it stands, onto the (periodic) lattice⁸ is mathematically voluminous and very time consuming to compute, since the transition functions and their derivatives must be constructed everywhere on the boundary of each hypercube.

We realize, however, that the 4-dimensional integral in eq. (1) can be performed analytically with the help of the cochain reduction given in ref.9. The outcome is a local algebraic expression for the topological charge (1) in the continuum, which is relatively easy to implement on the lattice and fast to compute. The derivation of this result, which we believe is also of interest beyond the scope of lattice gauge theory applications as it allows a simple geometrical interpretation of topology, will be given in the present paper.

M. Göckeler¹⁾, M.L. Laursen²⁾, G. Schierholz^{3,4)} and U.-J. Wiese⁵⁾

Topological charge of (lattice) gauge fields

1) Institut für Theoretische Physik der Universität, Heidelberg

²⁾ NORDITA, Copenhagen

³⁾ Institut für Theoretische Physik der Universität, Kiel

4) Deutsches Elektronen-Synchrotron DESY, Hamburg

⁵⁾ Institut für Theoretische Physik der Universität, Hannover

<u>Abstract:</u> Using recently derived explicit formulae for the 2- and 3-cochains in SU(2) gauge theory, we are able to integrate the Chern-Simons density analytically. We arrive - in SU(2) - at a local algebraic expression for the topological charge, which is the sum of local winding numbers associated with the corners (lattice points) of the cells covering the manifold plus contributions from possible isolated gauge singularities which manifest themselves as "vortices" in the 1-, 2- or 3-cochains. Among others we consider hypercubic geometry - i.e. covering the manifold by hypercubes - which is of particular interest to lattice Monte Carlo applications. Finally, we extend our results to SU(3) gauge theory. The paper is organized as follows. Sections II and III deal with the continuum: in section II we do the integral for SU(2) gauge fields covering the manifold M by a "generic" set of cells, while in section III we take M to be the 4-torus and cover it by hypercubes. In section IV we briefly outline what it involves to evaluate the new expression for Q on the lattice. In section V we extend our results to gauge group SU(3). We finish with some concluding remarks in section VI.

II. Continuum SU(2) gauge fields and generic geometry

The derivation in this section will follow the cochain reduction of the Chern-Simons density given in ref.9.

We cover the compact manifold M by a set of cells c_i , i.e.

 $M = \bigcup_{i} C_{i}$

with

$$C_i \wedge C_j = \Im C_i \wedge \Im C_j \quad (4)$$

(3)

.

The cells are chosen such that 5-n cells overlap in an n-dimensional intersection. This we refer to as generic geometry. The topological charge (1) can then be written

$$Q = -\frac{1}{32\pi^2} \sum_{i} \int_{C_i} d^4 \times \epsilon_{\mu\nu\rho\sigma} \operatorname{Tr} \left[\mathcal{F}_{\mu\nu}^{i} \mathcal{F}_{\rho\sigma}^{i} \right], \qquad (5)$$

where the index i on the fields refers to the gauge in the cell c_i . In each cell we may gauge transform the gauge fields into a complete axial gauge by

$$\hat{A}_{\mu}^{i} = g_{i}^{-\prime} (A_{\mu}^{i} + \partial_{\mu}) g_{i} .$$
 (6)

Making use of the fact that the Chern-Pontryagin density is a total divergence, i.e.

the second second second second second

$$P = -\frac{1}{32\pi^2} \epsilon_{\mu\nu\rho\sigma} Tr \left[\mathcal{F}_{\mu\nu} \mathcal{F}_{\rho\sigma} \right] = \mathcal{D}_{\mu} \mathcal{Q}_{\mu}^{(o)}(i)$$
(7)

with

$$\Omega_{\mu}^{(0)}(i) = -\frac{1}{3\pi^2} \epsilon_{\mu\nu\rho\sigma} Tr \left[\hat{A}_{\nu}^{i} \left(\partial_{\rho} \hat{A}_{\sigma}^{i} + \frac{2}{3} \hat{A}_{\rho}^{j} \hat{A}_{\sigma}^{i} \right], \qquad (8)$$

we obtain

$$Q = \sum_{i} \int d^{3} \sigma_{\mu} \Omega_{\mu}^{(0)} = \frac{1}{2!} \sum_{i,j} \int d^{3} \sigma_{\mu} (\Omega_{\mu}^{(0)} - \Omega_{\mu}^{(j)}), \quad (9)$$

where $c_i n c_i$ has the orientation of $\Im c_i$.

We introduce transition functions relating the gauges i,j in the cells $\mathbf{c}_i,\,\mathbf{c}_i$ by

$$\widetilde{A}_{\mu}^{a} = \widetilde{v_{ij}} \left(\widetilde{A}_{\mu}^{a} + \widetilde{v}_{\mu} \right) \widetilde{v_{ij}} .$$
⁽¹⁰⁾

Comparing this with eq. (6) one finds

$$\mathcal{V}_{\gamma j} = g_{\gamma}^{-\prime} g_{j}^{\prime} . \tag{11}$$

We recognize now that the integrand on the r.h.s. of eq. (9) is the coboundary operation

$$\Omega_{\mu}^{(0)} - \Omega_{\mu}^{(0)}(\dot{q}) = \Delta \Omega_{\mu}^{(0)}(\dot{q}, \dot{q}) = \mathcal{D}_{\mu\nu}^{(0)}(\dot{q}, \dot{q}), \quad (12)$$

which is again a total divergence. This excludes singular points

$$\times \in (c_i \cap c_j) \setminus \mathcal{I}(c_i \cap c_j), \qquad (13)$$

at which

$$\overline{v_{ij}} = exp \ i \ \vec{\alpha} \ \vec{t} = \cos \alpha + i \sin \alpha \ \vec{e_x} \ \vec{t}, \ \alpha \in [0, \pi]$$
(14)

becomes -1, i.e.
$$\propto = \pi$$
, where⁹

has a "vortex"

$$-\frac{1}{8\pi} \epsilon_{\mu\nu\rho\sigma} \vec{e}_{\alpha} \cdot (\partial_{\rho} \vec{e}_{\mu} * \partial_{\sigma} \vec{e}_{\mu}). \tag{16}$$

Applying Gauss' theorem, eq. (9) then reduces to

$$Q = Q^{(\prime)} + Q^{(\prime)}_{\Sigma}$$
⁽¹⁷⁾

$$Q_{Z}^{(i)} = \frac{1}{2!} \sum_{i,j \neq 0} \int d^{2} \sigma_{\mu\nu} \Omega_{\mu\nu}^{(i)} (i,j)$$

$$= \frac{1}{3!} \sum_{i,j,k} \int d^{2} \sigma_{\mu\nu} \left[\Omega_{\mu\nu}^{(i)} (i,j) - \Omega_{\mu\nu} (i,k) + \Omega_{\mu\nu}^{(i)} (j,k) \right]^{(18)}$$

1.5

- 5 -

and

$$Q^{(1)} = \frac{1}{2!} \sum_{i,j} \sum_{x \in (C_i, n \subset_j) \setminus V(C_i, n \subset_j)} \int_{\varepsilon_e} d^2 \sigma_{\mu\nu} \frac{1}{\delta \pi} \epsilon_{\mu\nuj\sigma} \overline{e_x} (\mathcal{O}_j \overline{e_x} \times \mathcal{O}_{e_x})_{(19)}$$

where $S_e^2(x)$ is a sphere of radius ϵ around x and where $c_i \wedge c_j \wedge c_k$ and $S_e^2(x)$ have the orientation of $\Im(c_i \wedge c_j)$.

The integral in eq. (19) gives

$$\int_{\mathbf{c}} d^{2} \sigma_{\mu\nu} f_{\pi} \epsilon_{\mu\nu\rho\sigma} \vec{e}_{\chi} \left(\partial_{\rho} \vec{e}_{\chi} \times \partial_{\sigma} \vec{e}_{\chi} \right) = n^{(\prime)} (x; i, j) , \qquad (20)$$

where $n^{(1)}$ is the S² winding number associated with the singularity at x. In the vicinity of x we can write

$$\mathcal{T}_{ij} = -1 + i M_{ajs} \, dx_a \, \overline{\iota}_{js} + \cdots , \qquad (21)$$

which for det $M \neq 0$ results in

$$n^{(1)}(x_j, i, j) = sign(det M) = \pm 1.$$
 (22).

In total we then obtain

$$Q^{(i)} = \sum_{\Lambda^{(i)}} n^{(i)}(x_{j}, i, j), \qquad (23)$$

where the sum is over all singular points x, i.e.

$$\Lambda^{(1)} = \bigcup_{i,j} \left\{ x \in (c_i, nc_j) \setminus \mathcal{D}(c_i, nc_j) \right\}.$$
(24)

From eq. (11) we derive the cocycle condition

which, writing

$$\mathcal{V}_{ij} = \exp(i\vec{\alpha}\vec{\tau}), \quad \mathcal{V}_{jk} = \exp(i\vec{\beta}\vec{\tau}), \quad \mathcal{V}_{ik} = \exp(i\vec{\sigma}\vec{\tau}), \quad (26)$$

defines a spherical triangle as has been shown in ref.9. The integrand on the r.h.s. of eq. (18) is the coboundary operation

$$\mathcal{Q}_{\mu\nu}(i,j) - \mathcal{Q}_{\mu\nu}(i,k) + \mathcal{Q}_{\mu\nu}(j,k) = \Delta \mathcal{Q}_{\mu\nu}(i,j,k) = \mathcal{Q}_{\mu\nu}(i,j,k), (27)$$

which again is a total divergence. This excludes singular points

$$\times \in (c_i \land c_j \land c_k) \setminus \Im(c_i \land c_j \land c_k), \qquad (28)$$

at which

$$(\vec{\alpha}+\vec{\beta}-\vec{\epsilon})\vec{e}_{\chi} = (\vec{\alpha}+\vec{\beta}-\vec{\epsilon})\vec{e}_{\chi} = -(\vec{\alpha}+\vec{\beta}-\vec{\epsilon})\vec{e}_{\chi} = -2\pi$$
, (29)

i.e. where the spherical triangle defined by the transition functions (26) degenerates to a circle, where $^{9)}$

$$\begin{aligned} \mathcal{L}_{\mu\nu\rho}^{(2)}(i,j,k) &= -\frac{1}{8\pi^2} \mathcal{L}_{\mu\nu\rho\sigma}^{(1+2\cos\alpha\beta\beta\beta\beta\gamma\delta\delta\delta\delta\delta} - \cos^2\alpha - \cos^2\beta - \cos^2\delta)^{-1} \\ &\cdot \left\{ (\vec{\alpha}, +\vec{p}, -\vec{s}) \cdot (\sin\alpha\vec{e}_{\mu}) \right\} \left[\mathcal{D}_{\mu}^{(4)}(\vec{e}_{\mu}) \cdot (\sin\beta\vec{e}_{\mu}) - (\sin\beta\vec{e}_{\mu}) \cdot \mathcal{D}_{\mu}^{(5)}(\vec{s}_{\mu}) \right] \\ &+ (\vec{\alpha}, +\vec{p}, -\vec{s}) \cdot (\sin\beta\vec{e}_{\mu}) \left[\mathcal{D}_{\mu}^{(5)}(\sin\beta\vec{e}_{\mu}) - (\sin\beta\vec{e}_{\mu}) \cdot \mathcal{D}_{\mu}^{(5)}(\sin\beta\vec{e}_{\mu}) \right] \\ &+ (\vec{\alpha}, +\vec{p}, -\vec{s}) \cdot (\sin\beta\vec{e}_{\mu}) \left[\mathcal{D}_{\mu}^{(5)}(\sin\beta\vec{e}_{\mu}) \cdot (\sin\beta\vec{e}_{\mu}) - (\sin\beta\vec{e}_{\mu}) \cdot \mathcal{D}_{\mu}^{(5)}(\sin\beta\vec{e}_{\mu}) \right] \right] \end{aligned}$$

has a "vortex"

$$-\frac{1}{4\pi^2} \epsilon_{\mu\nu\rho\sigma} \frac{(\vec{x}+\vec{s}-\vec{\sigma})\cdot(\partial_{\sigma}\vec{S}\times\vec{S})}{S^2} = -\frac{1}{2\pi} \epsilon_{\mu\nu\rho\sigma} \frac{\vec{e}_{\mu}\cdot(\partial_{\sigma}\vec{S}\times\vec{S})}{S^2}, \quad (31)$$

$$\vec{S} = \cos \alpha \sin \beta \sin \beta \vec{e}_{p} \times \vec{e}_{p} + \cos \beta \sin \beta \sin \beta \vec{e}_{p} \times \vec{e}_{p} + \cos \beta \sin \alpha \sin \beta \vec{e}_{p} \times \vec{e}_{p}, \vec{S} = \vec{e}_{p} - \vec{e}_{p}$$

The vector \vec{S} defines the orientation of the great sphere (in group space) spanned by the transition functions (26) and winds around by 2π as one runs round the singularity.

Applying Gauss' theorem, eq.(18) then reduces to

$$Q_{\Sigma}^{(\prime)} = Q_{\Sigma}^{(2)} + Q_{\Sigma}^{(2)}$$
 (32)

with

$$Q_{Z}^{(2)} = \frac{1}{3!} \sum_{i,j,k} \int d\sigma_{\mu\nu\rho} \Omega_{\mu\nu\rho} (i,j,k)$$

$$= \frac{1}{4!} \sum_{i,j,k} \int d\sigma_{\mu\nu\rho} [\Omega_{\mu\nu\rho} (i,j,k) - \Omega_{\mu\nu\rho} (i,j,l)$$

$$+ \Omega_{\mu\nu\rho} (i,k,l) - \Omega_{\mu\nu\rho} (j,k,l)]$$
(33)

.....

and

$$Q^{(2)} = \frac{1}{3!} \sum_{i,j,k} \sum_{x \in (c_i, n_{c_j}, n_{c_k}) \setminus \Im(c_i, n_{c_j}, n_{c_k})} \int_{S_{e}(x)} d\sigma_{\mu\nu\rho} \frac{1}{2\pi} \epsilon_{\mu\nu\rho\sigma} \frac{1}{\delta^2} \frac{1}{\delta$$

where $S_e(x)$ is a circle of radius ϵ around x and where $c_i \wedge c_j \wedge c_k \wedge c_1$ and $S_e(x)$ have the orientation of $\Im(c_i \wedge c_k)$.

One realizes that the integral in eq. (34) gives

$$\int_{\mathbf{s}_{\ell}(x)} d\sigma_{\mu\nu g} \frac{1}{2\pi} \epsilon_{\mu\nu g\sigma} \frac{\vec{e}_{\alpha} \cdot (\partial_{\sigma} \vec{s} \times \vec{s})}{\delta^{2}} = n^{(2)} (x; i, j, k), \quad (35)$$

where $n^{(2)}$ is the S¹ winding number associated with the singularity x which assumes the values

$$n^{(2)}(x_j, i, j, k) = \pm 1$$
 (36)

ه ه

depending on whether $\vec{s}, \vec{\delta}$ wind around in the positive or negative sense. We then obtain

$$Q^{(2)} = \sum_{\Lambda^{(2)}} n^{(2)}(x_{j}, i, j, k)$$
(37)

with

$$\Lambda^{(2)} = \bigcup_{i,j,k} \left\{ x \in (c_i n c_j n c_k) \setminus \Im(c_i n c_j n c_k) \right\}.$$
(38)

Expression (33) brings together 4 spherical triangles, which build a spherical tetrahedron as illustrated in ref.9.

The descent continues:

In ref.9 we have shown that

$$\Omega_{\mu\nu\rho\tau}(i,j,k,l) = \frac{1}{2\pi^2} \epsilon_{\mu\nu\rho\tau} V(i,j,k,l), \qquad (40)$$

where V(i,j,k,l) is the volume of the spherical tetrahedron defined by the transition functions relating the gauges i,j,k,l. We realize that V(i,j,k,l) has a discontinuity at points

$$x \in (C_i \cap C_k \cap C_\ell) \setminus \mathcal{I}(C_i \cap C_j \cap C_k \cap C_\ell), \qquad (41)$$

where the spherical tetrahedron degenerates to a sphere. Applying Gauss' theorem we then obtain two contributions to eq. (33):

$$Q_{\Sigma}^{(2)} = Q + Q_{\Sigma}^{(3)}$$
 (42)

The first contributions gives

$$Q^{(3)} = \sum_{\Lambda^{(3)}} n^{(3)}(x; i; j, h, \ell)$$
(43)

with

$$\Lambda^{(3)} = \bigcup \left\{ X \in (C_i \cap C_j \cap C_k \cap C_\ell) \setminus \mathcal{D}(C_i \cap C_j \cap C_k \cap C_\ell) \right\}$$

$$(44)$$

where $n^{(3)}$ accounts for the change of the orientation of the spherical tetrahedron at the point x, i.e.

$$n^{(3)}(x;i,j,k,l) = \frac{1}{2\pi^2} \epsilon_{\mu\nu\rho\sigma} \Delta V(i,j,k,l) / = \pm 1.$$
(45)

For the second contribution we obtain

$$Q_{\Sigma}^{(3)} = \frac{1}{4!} \sum_{\substack{Q(c_i, c_j, nc_k, nc_k) \in \mathbb{Z}}} \sigma_{\frac{1}{2\pi^2}} V(i_{jj}, k, \ell)$$

$$= \frac{1}{5!} \sum_{\substack{C_i, nc_j, nc_k, nc_k, nc_k}} \sigma_{\frac{1}{2\pi^2}} [V(i_{jj}, k, \ell) - V(i_{jj}, k, m)] \quad (46)$$

where $\sigma = \pm 1$ is a sign factor, which denotes the orientation of $\Im(c_i n c_j n c$

The intersection of 5 4-dimensional cells defines a point which naturally leads to the notation of the "lattice":

$$\Lambda = \bigcup C_i \cap C_j \cap C_k \cap C_k \cap C_m .$$

$$i, j, k, l, m$$
(47)

Thus we can rewrite eq. (46) in the form

$$Q_{\Sigma}^{(3)} = \sum_{\Lambda} n(i,j,k,\ell,m)$$
⁽⁴⁸⁾

with

$$n(i,j,k,l,m) = \sum_{2\pi^2} \left[V(i,j,k,l) - V(i,j,k,m) + V(i,j,l,m) - V(i,k,l,m) + V(j,k,l,m) \right].$$
(49)

Note that the 1/5! factor in eq. (46) has disappeared because of the implicit summation over all permutations of cells. As we have discussed in ref.9, n combines 5 spherical tetrahedra, which wind around S^3 , the group space of SU(2). Since the volume of S^3 is $2\pi^2$ and the 5 spherical tetrahedra together are compact and so cover S^3 (but at most once), we find

$$\mathcal{N} = O_{1} \pm I \,. \tag{50}$$

The calculation of n(i,j,k,l,m) proceeds as follows. We take the transition functions $\mathcal{F}_{ij}, \mathcal{F}_{jk}, \ldots$ and determine the angles between adjacent spherical triangles intersecting along a "hinge" of the spherical tetrahedra (cf. ref.9). In Fig. 1a we have illustrated this for a particular "hinge" with which are associated 3 angles, A_1 , A_2 and A_3 , belonging to 3 different spherical tetrahedra. For geometrical reasons

$$\frac{(A_1 + A_2 + A_3)}{A_{\overline{L}}} \equiv p = 0, \pm 1, \qquad (51)$$

a special example of which is given in Fig. 1b. If one of the 10 p's is zero, then evidently n = 0. If all p's are + 1, then n = +1, and if all p's are - 1, we will have n = -1.

Collecting our results now, we obtain for the topological charge

$$Q = \sum_{\Lambda} n(i,j,k,l,m) + \sum_{\Lambda^{(1)}} n^{(1)}(x_j,i,j)$$

$$+ \sum_{\Lambda^{(2)}} n^{(2)}(x_j,i,j,k) + \sum_{\Lambda^{(3)}} n^{(3)}(x_j,i,j,k,l) ,$$
(52)

which is the sum of local winding numbers and, consequently, assumes integer values. Expression (52) is gauge invariant by construction. It furthermore has a simple, geometrical interpretation.

In the continuum we can arrange that neither $\mathcal{T}_{i,j} = -1$ in $(c_i c_j) \setminus \partial (c_i c_j)$, $(\vec{\alpha} + \vec{\beta} - \vec{\sigma}) \vec{e}_x = 2\pi$ in $(c_i c_j c_k) \setminus \partial (c_i c_j c_k)$ nor $\Delta V(i, j, k, l) l_x = \pm 2\pi^2$ at $x \in (c_i c_j c_k c_k) \setminus \partial (c_i c_j c_k c_k)$, in which case the topological charge takes the simple form

$$Q = \sum_{\Lambda} n(i,j,k,l,m).$$
 (53)

For a further discussion see section IV.

III. Hypercubic geometry

So far we have considered a generic geometry of cells covering M. Now we come to the case of a hypercubic geometry, which is of special interest to lattice gauge theory calculations, and where M is the 4-torus T⁴.

The "lattice" is now defined by

$$\Lambda = \{ s \in M \mid s_{p} \in \mathbb{Z}, \ \mu = 0, 1, 2, 3 \}.$$
 (54)

The cells covering M are the hypercubes (of length 1),

$$M = \bigcup_{s \in \Lambda} C(s)$$
(55)

with

$$C(s) = \{ g \in M \mid s_{\mu} \leq g_{\mu} \leq s_{\mu} + 1 \}.$$
(56)

The topological charge (1) can then be written

$$Q = \sum_{s \in \Lambda} \int d^{4}x \mathcal{P}, \qquad (57)$$

where P is the Chern-Pontryagin density given in eq. (7). Using

$$\mathcal{P} = \mathcal{D}_{\mu} \cdot \mathcal{Q}_{\mu}^{(0)}(\mathbf{s}) \tag{58}$$

we can perform one integration as before and obtain

$$Q = \sum_{s \in \Lambda} \sum_{p} \int d^{3} \sigma_{p} \left[\Omega_{p}^{(0)}(s) - \Omega_{p}^{(0)}(s, p) \right], \quad (59)$$

where $f(s,\mu)$ are the faces

$$f(s,\mu) = C(s) \cap C(s-\mu), \qquad (60)$$

which have the orientation of $\Im \mathcal{C}(\mathcal{S})$, and $\hat{\mu}$ is the unit vector in the μ -direction.

Following the discussion in the previous section we can write

$$(a) = (b) = (a) = (a)$$

except for a set of singular points x $\in f(s,\mu) \setminus \Im f(s,\mu)$. As before the topological charge receives contributions from two terms, $Q = Q^{(1)} + Q^{(1)}_{\Sigma}$, where now

$$Q_{\Sigma}^{(1)} = \sum \sum \int d^{2}\sigma_{\mu\nu} \left[\Omega_{\mu\nu}^{(1)}(s,s\hat{\mu}) - \Omega_{\mu\nu}^{(1)}(s-\hat{\nu},s-\hat{\mu}-\hat{\nu}) \right]$$
(62)

with

$$p(s,\mu,\nu) = C(s) \cap C(s-\hat{\mu}) \cap C(s-\hat{\nu})$$
(63)

being a plaquette, which was the orientation of $\Im f(s,\mu)$. The contribution $Q_{ij}^{(\prime)}$, which is independent of the particular geometry, is precisely the same as in the previous case,

$$Q^{(n)} = \sum_{\Lambda^{(n)}} n^{(n)}(x; s, s-\hat{p}), \qquad (64)$$

where the sum is over all singular points

$$\Lambda^{(\prime)} = \bigcup \left\{ x \in f(s, \mu) \cap \mathcal{J}(s, \mu) \right\}. \tag{65}$$

Noticing that (cf. eq. (15))

•

and the second second

- 12 -

$$Q_{\Sigma}^{(1)} \text{ can be rewritten}$$

$$Q_{\Sigma}^{(1)} = \sum_{s \in \Lambda} \sum_{\mu,\nu} \int_{\rho(s_{\mu},\nu)} \partial_{r}^{2} \sigma_{\mu\nu} \left[\Omega_{\mu\nu}^{(1)} (s_{\rho} s_{\mu}^{-1} \cdot \nu) - \Omega_{\mu\nu}^{(1)} (s_{\rho} s_{\mu}^{-1} \cdot \nu) \right]$$

$$= \sum_{s \in \Lambda} \sum_{\mu,\nu} \int_{\rho(s_{\mu},\nu)} \partial_{r}^{2} \sigma_{\mu\nu} \Delta \Omega_{\mu\nu}^{(1)} (s_{\rho} s_{\mu}^{-1} \cdot \nu) + \Omega_{\mu\nu}^{(1)} (s_{\rho} s_{\mu}^{-1} \cdot \nu)$$

$$= \sum_{s \in \Lambda} \sum_{\mu,\nu} \int_{\rho(s_{\mu},\nu)} \partial_{r}^{2} \sigma_{\mu\nu} \partial_{\rho} \Omega_{\mu\nu\rho}^{(2)} (s_{\rho} s_{\mu}^{-1} \cdot s_{\mu}^{-1} \cdot \nu)$$

$$= \sum_{s \in \Lambda} \sum_{\mu,\nu} \int_{\rho(s_{\mu},\nu)} \partial_{r}^{2} \sigma_{\mu\nu} \partial_{\rho} \Omega_{\mu\nu\rho}^{(2)} (s_{\rho} s_{\mu}^{-1} \cdot s_{\mu}^{-1} \cdot \nu) ,$$

$$\text{which gives as before } Q_{\Sigma}^{(1)} = Q^{(2)} + Q_{\Sigma}^{(2)}, \text{ where}$$

$$Q_{\Sigma}^{(2)} = \sum_{s \in \Lambda} \sum_{\mu,\nu,\rho} \int_{\rho(s_{\mu},\nu,\rho)} \partial_{r} \sigma_{\mu\nu\rho}^{(s,\mu,\rho,\mu,\nu)} (s_{\rho} s_{\mu}^{-1} \cdot \nu) - \Omega_{\mu\nu\rho}^{(2)} (s_{\rho} s_{\mu} \cdot s_{\mu}^{-1} \cdot \nu) \int_{\mu\nu\rho}^{(2)} (s_{\rho} s_{\mu} \cdot s_{\mu}^{-1} \cdot \nu) \partial_{\rho}^{(2)} (s_{\rho} \cdot \nu) \partial$$

$$\mathcal{L}(s,\mu,\nu,p) = \mathcal{C}(s) \cap \mathcal{C}(s,\mu) \cap \mathcal{C}(s,\mu) \qquad (69)$$

being a link, which has the orientation of $\Im p(s, \mu, v)$. The contribution $Q^{(2)}$ has the form

$$Q^{(2)} = \sum_{\Lambda^{(2)}} n^{(2)}(x; s, s, \tilde{\mu}, s, \tilde{\mu}, s, \tilde{\mu}), \qquad (70)$$

where

Q

with

(n)

 Q_{z}

$$\Lambda^{(2)} = \bigcup \left\{ x \in p(s, \mu, v) \setminus \partial p(s, \mu, v) \right\}.$$
(71)

Making use of the fact that⁹⁾

$$\begin{array}{l}
\mathcal{Q}_{\mu\nu\rho}^{(2)}(s,s-\hat{\mu},s-\hat{\mu}-\hat{\nu}-\hat{\rho}) = -\mathcal{Q}_{\mu\rho\nu}^{(2)}(s,s-\hat{\mu},s-\hat{\mu}-\hat{\nu}-\hat{\rho}), \\
\mathcal{Q}_{\mu\nu\rho}^{(2)}(s,s-\hat{\mu}-\hat{\nu},s-\hat{\mu}-\hat{\nu}-\hat{\rho}) = -\mathcal{Q}_{\nu\mu\rho}^{(2)}(s,s-\hat{\mu}-\hat{\nu},s-\hat{\mu}-\hat{\nu}-\hat{\rho}), \quad (72) \\
\mathcal{Q}_{\mu\nu\rho}^{(2)}(s-\hat{\rho},s-\hat{\mu}-\hat{\rho},s-\hat{\mu}-\hat{\nu}-\hat{\rho}) = -\mathcal{Q}_{\rho\mu\nu}^{(2)}(s-\hat{\rho},s-\hat{\mu}-\hat{\rho},s-\hat{\mu}-\hat{\nu}-\hat{\rho}), \\
\end{array}$$

5 . . . **.** .

- 13 -

eq. (68) can be written

$$Q_{\Sigma}^{(2)} = \sum_{s \in \Lambda} \sum_{p_1 \vee j} \int d\sigma_{p_1 \vee p} \left[\Omega_{p_1 \vee p}(s, s \cdot p_1, s \cdot p_1 \cdot \hat{\nu}) - \Omega_{p_1 \vee p}(s, s \cdot p_1, s \cdot p_1 \cdot \hat{\nu}, \hat{p}) \right] + \Omega_{p_1 \vee p}^{(2)}(s, s \cdot p_1 \cdot \hat{\nu}, s \cdot p_1 \cdot \hat{\nu}, \hat{p}) - \Omega_{p_1 \vee p}^{(2)}(s, s \cdot p_1 \cdot \hat{\nu}, s \cdot p_1$$

$$Q^{(3)} = \sum_{\Lambda^{(3)}} n^{(3)}(x; s, s-f, s-f-\hat{v}, s-f-\hat{v}-\hat{f})$$

with

$$\begin{cases} {}^{(3)} = \bigcup \{ x \in \ell(s, \mu, v, g) \mid \partial \ell(s, \mu, v, g) \} \\ s, \nu, v, g \end{cases}$$

$$(75)$$

For $Q_{\frac{1}{2}}^{(3)}$ we obtain, together with eq. (40),

$$Q_{\Sigma}^{(3)} = \sum_{s \in \Lambda} \sum_{\mu, \nu, \beta, \sigma} \epsilon_{\mu \nu \beta \sigma} \frac{1}{2\pi^{2}} \left[V(s, s \hat{\mu}, s - \hat{\mu} - \hat{\nu}, s - \hat{\mu} - \hat{\nu} - \hat{g}) \right]$$

$$- V(s - \hat{\sigma}, s - \hat{\mu} - \hat{\sigma}, s - \hat{\mu} - \hat{\nu} - \hat{\sigma}, s - \hat{\mu} - \hat{\nu} - \hat{g} - \hat{\sigma}) \right],$$
(76)

where $V(s, s, \hat{\mu}, \hat{s}, \hat{\mu}, \hat{s}, \hat{\mu}, \hat{s}, \hat{s}, \hat{\mu}, \hat{s}, \hat{s}, \hat{s}, \hat{s}, \hat{s}, \hat{\mu}, \dots$ defined by the transition functions $v_{s,s-\hat{\mu}}, \dots$. Again, making use of the symmetry/antisymmetry properties of $\epsilon_{\mu\nu\rho\sigma}$, eq. (76) can be written

$$Q_{\Sigma}^{(3)} = \sum_{s \in \Lambda} \sum_{p, v, \beta, \tau} \epsilon_{p, v, \beta \tau} \frac{1}{2\pi^2} \left[V(s, s - \hat{p}, s - \hat{p} - \hat{v}, s - \hat{p} - \hat{v} - \hat{g}) - V(s, s - \hat{p}, s - \hat{p} - \hat{v} - \hat{g}, s - \hat{p} - \hat{v} - \hat{g}) - V(s, s - \hat{p}, s - \hat{p} - \hat{v} - \hat{g}, s - \hat{p} - \hat{v} - \hat{g} - \hat{\sigma}) - V(s, s - \hat{p} - \hat{v} - \hat{g}, s - \hat{p} - \hat{v} - \hat{g} - \hat{\sigma}) - V(s, s - \hat{p} - \hat{v} - \hat{g} - \hat{v} - \hat{g} - \hat{\sigma}) - V(s, s - \hat{p} - \hat{v} - \hat{g} - \hat{\sigma}) + V(s - \hat{p}, s - \hat{p} - \hat{v} - \hat{g} - \hat{\sigma}) - V(s, s - \hat{p} - \hat{v} - \hat{g} - \hat{\sigma}) - \hat{v} - \hat{g} - \hat{\sigma}) - V(s, s - \hat{p} - \hat{v} - \hat{g} - \hat{s}) - \hat{v} - \hat{g} - \hat{s} - \hat{s}) - \hat{v} - \hat{s} - \hat{s} - \hat{s} - \hat{s}) - \hat{v} - \hat{s} - \hat{s} - \hat{s} - \hat{s} - \hat{s}) - \hat{v} - \hat{s} -$$

so that finally we obtain

$$Q_{Z}^{(3)} = \sum_{s \in \Lambda} \sum_{\mu, \nu, \beta, \sigma} n(s, s - \hat{\mu}, s - \hat{\mu} - \hat{\nu}, s - \hat{\mu} - \hat{\nu} - \hat{\beta} - \hat{\sigma}), \quad (78)$$

where n is the winding number associated with the 5 spherical tetrahedra covering S^3 as introduced before with values 0, $\pm 1.$

In total, the topological charge then reads

$$Q = \sum_{s \in \Lambda} \sum_{\mu, \nu, \beta, \sigma} n(s, s - \hat{\mu}, s - \hat{\mu} - \hat{\nu}, s - \hat{\mu} - \hat{\nu} - \hat{\beta}, s - \hat{\mu} - \hat{\nu} - \hat{\beta} - \hat{\sigma}) + \sum_{\Lambda^{(1)}} n^{(1)}(x; s, s - \hat{\mu}) + \sum_{\Lambda^{(2)}} n^{(2)}(x; s, s - \hat{\mu}, s - \hat{\mu} - \hat{\nu})$$
(79)
+
$$\sum_{\Lambda^{(3)}} n^{(3)}(x; s, s - \hat{\mu}, s - \hat{\mu} - \hat{\nu} - \hat{\beta}) .$$

Again, it can be arranged that

$$Q = \sum_{s \in \Lambda} \sum_{\mu, \nu, \beta, \sigma} n(s, s - \hat{\mu}, s - \hat{\mu} - \hat{\nu}, s - \hat{\mu} - \hat{\nu} - \hat{\beta}, s - \hat{\mu} - \hat{\nu} - \hat{\beta} - \hat{\sigma}).$$
(80)

The virtue of this construction of the topological charge Q is that it can easily be adopted to any geometry of cells.

IV. Lattice gauge fields

In sections II and III we have assumed that the gauge fields are known in the continuum (of M). This is no longer true if the theory is regularized by formulating it on a discrete lattice of points in space-time – usually a hypercubic lattice, which we shall identify with Λ (eq. (54)). In this case the transition functions $\mathcal{T}_{s,s-\hat{\mu}}$ are a priori only given at the corners of the faces f(s, μ), i.e. the lattice points.

For the computation of $Q_{3}^{(3)}$ this is all that is required. For the computation of $Q_{1}^{(1)}$, $Q_{2}^{(2)}$ and $Q_{3}^{(3)}$ we need - if no further provisions are made to know whether $\Delta \Omega_{\mu\nu}^{(0)}(s,s-\hat{\mu})$, $\Delta \Omega_{\mu\nu}^{(1)}(s,s-\hat{\mu},s-\hat{\mu}-\hat{\nu})$ and $\Delta \Omega_{\mu\nu}^{(2)}(s,s-\hat{\mu},s-\hat{\mu}-\hat{\nu},\hat{s})$ are singular in $f(s,\mu)$, $\Im f(s,\mu)$, $p(s,\mu,\nu)$, $\Im p(s,\mu,\nu)$ and $\ell(s,\mu,\nu,p)$, $\Im \ell(s,\mu,\nu,p)$, respectively. The exact positions of the singularities ($x \in \Lambda^{(1)}$, $\Lambda^{(2)}$ and $\Lambda^{(3)}$ respectively) are, however, not necessary to know. This requires (only) an interpolation of $\sigma_{s,s-\hat{\mu}}$ throughout the plaquettes $p(s,\mu,\nu)$ and, possibly, to a single (but arbitrary) point in the interior of $f(s,\mu)$. A potential interpolation (which satisfies the cocycle condition and maintains gauge invariance) has been given in the literature⁸.

In the "continuum region", where the lattice spacing (which we have set to 1) becomes small in physical units and the gauge fields contributing to the functional integral have a small (lattice) action density,

$$\mathcal{T}_{\mathbf{r}}\left[I - U(\partial_{\mathbf{p}})\right] < \epsilon \tag{81}$$

(U(\Im p): parallel transporter around the plaquette p), the computation of Q = = $Q_{\Sigma}^{(3)} + Q_{\Gamma}^{(1)} + Q_{\Gamma}^{(2)} + Q_{\Gamma}^{(3)}$ simplifies greatly. Using Lüscher's interpolation⁸ of the transition functions, one derives the following (sufficient) conditions under which $\Delta \Omega_{\mu\nu}^{(3)}(s,s_{\mu}), \Delta \Omega_{\mu\nu}^{(3)}(s,s_{\mu}), s_{\mu} \cdot s_{\nu}$ and $\Delta \Omega_{\mu\nu}^{(3)}(s,s_{\mu}), s_{\mu} \cdot s_{\nu}$ have no gauge singularities:

$$\begin{aligned} d(v_{s,s,\hat{p}}^{(s)}, 1) + & \text{Harccos}\left(I - \frac{1}{2}\epsilon\right) \leq \pi, \\ d(v_{s,s,\hat{p}}^{(s)}, 2) + d(v_{s,\hat{p},\hat{s},\hat{p}}^{(s)}, 2) + d(v_{s,\hat{s},\hat{p}}^{(s)}, 1) + \delta \operatorname{arccos}(I - \frac{1}{2}\epsilon) \leq 2\pi, \\ o(v_{s,\hat{s},\hat{p}}^{(s)}, 1) + \max \left\{ d(v_{s,\hat{p},\hat{s},\hat{p}}^{(s)}, 2) \left[v_{s\hat{p},\hat{s},\hat{p}}^{(s)}, v_{s\hat{p},\hat{s},\hat{p}}^{(s)}, 2 \right]^{\lambda}, 1 \right\} \end{aligned} \tag{82} \\ + d(v_{s,\hat{s},\hat{p}}^{(s)}, v_{s\hat{p},\hat{s},\hat{p}}^{(s)}, 3(s) \left[v_{s\hat{p},\hat{s},\hat{p}}^{(s)}, v_{s\hat{p},\hat{s},\hat{p}}^{(s)}, 2 \right]^{\lambda}, 1) \right\} + \mathcal{B}\operatorname{arccos}(I - \frac{1}{2}\epsilon) \leq 2\pi. \end{aligned}$$

for $\mu, \nu, g \in \{2, 3, 4\}$ (for μ, ν or g = 1 the fields that give rise to gauge singularities are of measure zero in the functional integral), where $d(\mu, \nu)$ is the metric of SU(2) = S³, i.e.

$$d(n,1) = \arccos\left(\frac{1}{2}Trn\right). \tag{83}$$

This means that for small ϵ it is possible to gauge transform the transition functions - if necessary - such that the bounds (82) are satisfied. The computation of Q then reduces to the computation of $Q_{\underline{x}}^{(3)}$, and the charge can be viewed as the sum over local SU(2) winding numbers.

For gauge field configurations with larger action density it is also not necessary to compute all of the Q's. One finds that it is always possible to (explicitly) gauge transform the transition functions such that $Q_{\Sigma}^{(3)} = Q^{(2)} = Q^{(1)} = 0$. This then reduces the problem to determine whether $\overline{\sigma}_{s,s-\hat{\mu}} = -1$ in $f(s,\mu)\setminus \partial f(s,\mu)$, which can be done on purely geometrical grounds.

It is not the purpose of the present paper to go into further details of the numerical evaluation of Q on the lattice. This will be the subject of a forthcoming publication, where we will also present results of a Monte Carlo calculation of $\langle Q^2 \rangle$ on large lattices.

<u>V. SU(3)</u>

It is not trivial to extend the results of sections II and III to the physically more interesting case of gauge group SU(3). Therefore one is led to ask the question: can we reduce the problem of computing the topological charge for SU(3) gauge fields to the case of $SU(2)^{10}$?

In the continuum this is possible by means of the so-called reduction of the structure group¹¹⁾. This means the following. A fibre bundle, which has structure group G and is therefore in general given by G-valued transition functions, may under certain circumstances be described by transition functions with values in a subgroup of G. By means of the theorems proved in ref.11 it is easily shown that the structure group of any SU(3) bundle over T^4 can be reduced to SU(2).

The explicit construction makes use of the fact that SU(3)/SU(2) is equivalent to S^5 . Actually, SU(3) is a SU(2) principal bundle over S^5 (cf. ref.12 for the analogous considerations in case of the unitary groups). The projection

$$SU(3) \rightarrow SU(3)/SU(2) \cong S^{S}$$
 (84)

simply maps a SU(3) matrix onto its first column. Here and in the following we describe points of S^5 by a triplet of complex numbers,

$$(a, b, c) \in \mathcal{C}^3$$
, $|a|^2 + |b|^2 + |c|^2 = 1$. (85)

As S⁵ minus one point is a contractible space, SU(3) minus the fibre over the removed point is a trivial SU(2) bundle. For example, any $\mathcal{M} \in$ SU(3) with $\alpha_{\prime\prime} \neq -1$ may be decomposed as

$$\mathcal{U} = \omega(\mathcal{U}_{\prime\prime}, \mathcal{U}_{2\prime}, \mathcal{U}_{3\prime}) \begin{pmatrix} \prime & o & o \\ o & \alpha & -\beta^* \\ o & \beta & \alpha^* \end{pmatrix}, \quad (86)$$

where $|\alpha|^2 + |\beta|^2 = 1$ and

$$\omega(a,b,c) = \begin{pmatrix} a & -b^{*} & -c^{*} \frac{1+a}{1+a^{*}} \\ b & \frac{1}{1+a} (1+a^{*}-1b)^{2}) & -\frac{bc^{*}}{1+a^{*}} \\ c & -\frac{cb^{*}}{1+a} \frac{1}{1+a^{*}} (1+a^{*}-1c)^{2} \end{pmatrix}$$
(87)

for $|a|^2 + |b|^2 + |c|^2 = 1$, $a \neq -1$. The first factor in (86) represents a point of the base space S⁵ in SU(3) while the second one describes the fibre SU(2).

Now the reduction procedure as given in ref.11 goes as follows. Let \mathcal{P} be an SU(3) bundle over T⁴ specified by transition functions (we shall restrict ourselves here to hypercubic geometry)

$$\overline{v}_{s,s-\mu}(z) \in SU(3), \ z \in f(s,\mu)$$
⁽⁸⁸⁾

analogous to the case of SU(2). We first construct a section of the bundle $\mathcal{P}/SU(2)$ which has T⁴ as base space and S⁵ as fibre. Such a section is described by smooth functions

- 18 -

$$\mathbf{z}_{\mathbf{s}} : \mathbf{C}(\mathbf{s}) \to \mathbf{S}^{\mathbf{s}}$$
(89)

satisfying

$$z_{s}(q) = \overline{v}_{s,s-\beta}(q) \, z_{s-\beta}(q) \tag{90}$$

for $\mathcal{F} \in f(s,\mu)$. The fact that $\mathcal{P}/SU(2)$ admits a section follows from

$$\overline{\mathcal{T}}_{1}(S^{5}) = \overline{\mathcal{T}}_{2}(S^{5}) = \overline{\mathcal{T}}_{3}(S^{5}) = 0.$$
 (91)

If $\mathbf{E}_{\mathcal{S}}$ is not surjective - this condition is generically fulfilled - we can find a smooth map

$$\hat{\hat{z}}_{s}: C(s) \to SU(3) \tag{92}$$

such that the first column of $\hat{e}_{s}(y)$ coincides with $e_{s}(y)$ for all $y \in c(s)$:

$$\hat{\vec{z}}_{s}(\vec{z}) \begin{pmatrix} \prime \\ o \\ o \end{pmatrix} = \vec{z}_{s}(\vec{z}).$$
 (93)

For example, if $z_{e}(y) \neq (-1,0,0)$ for all $y \in c(s)$ we can define

$$\hat{\boldsymbol{\varepsilon}}_{\boldsymbol{s}}(\boldsymbol{\eta}) = \boldsymbol{\omega} \left(\boldsymbol{z}_{\boldsymbol{s}}(\boldsymbol{\eta}) \right). \tag{94}$$

For y € f(s,µ) we have

$$\hat{z}_{s}(q) \begin{pmatrix} 0 \\ 0 \end{pmatrix} = \hat{z}_{s}(q) = \overline{\sigma}_{s,s-\beta}(q) \hat{z}_{s-\beta}(q) = \overline{\sigma}_{s,s-\beta}(q) \hat{z}_{s-\beta}(q) \begin{pmatrix} 0 \\ 0 \end{pmatrix}.$$
(95)

Consequently, if we use the functions \hat{z}_s to gauge transform the $\overline{v}_{s,s-\hat{\mu}}$, we arrive at transition functions with values in SU(2):

$$\hat{\boldsymbol{z}}_{s}(\boldsymbol{z})^{-1} \boldsymbol{\overline{v}}_{s,s-\hat{\boldsymbol{\mu}}}(\boldsymbol{z}) \hat{\boldsymbol{z}}_{s-\hat{\boldsymbol{\mu}}}(\boldsymbol{z}) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \boldsymbol{\overline{v}}_{s,s-\hat{\boldsymbol{\mu}}} \end{pmatrix}, \quad \boldsymbol{\overline{v}}_{s,s-\hat{\boldsymbol{\mu}}}(\boldsymbol{z}) \in SU(\boldsymbol{z}). \quad (96)$$

As the topological charge is invariant under gauge transformations like (96), it can be computed from the SU(2)-valued transition functions $\mathfrak{T}_{S-S-\hat{\mu}}(y)$.

To carry out the reduction explicitly we need to know $\geq_s(y)$. In the continuum, where the SU(3)-valued transition functions $\bar{v}_{s,s-\bar{\mu}}$ are explicitly given in the form

$$\overline{\upsilon}_{s,s-\hat{\mu}}(\mathfrak{z}) = \overline{\mathfrak{g}}_{s}(\mathfrak{z})^{-} \overline{\mathfrak{g}}_{s-\hat{\mu}}(\mathfrak{z}) \tag{97}$$

(cf. eq.(11)) for all $y \in f(s,\mu)$, we may take

$$\mathcal{Z}_{3}(\mathfrak{z}) = \overline{\mathfrak{g}}_{3}(\mathfrak{z})^{-1}(\mathfrak{z}), \qquad (98)$$

which, as is easily seen, satisfies eq.(90)(and, as a result of (91), can be extended to c(s)). On the lattice we follow ref.8 and construct transition functions at the corners of $f(s,\mu)$ with help of the standard parallel transporters $\widehat{\phi}^{s}$:

$$\overline{v}_{s,s-\mu}(g) = \overline{v}_{s}(g) \cdot \overline{v}_{s}(g)^{-\prime}.$$
(99)

To extend the $\overline{v}_{s,s-\hat{\mu}}$'s to the whole of $f(s,\mu)$ we may take Lüscher's interpolation⁸) (which is trivial to extend to the case of gauge group SU(3)). It has been shown in ref.13 that this can be written for all $y \in f(s,\mu)$ in the form (99). The explicit expressions for the parallel transporters $\overline{v}(y), y \in \partial c \omega$ are also given in this reference. We then may define

$$\mathcal{Z}_{s}(\boldsymbol{z}) = \overline{\boldsymbol{v}}^{s}(\boldsymbol{z}) \begin{pmatrix} \boldsymbol{z} \\ \boldsymbol{z} \end{pmatrix}, \qquad (100)$$

which satisfies eq.(90) and provides us with SU(2)-valued transition functions $\sigma_{S,S-\hat{\mu}}(y)$ for all $y \in f(s,\mu)$.

V. Conclusions

We have integrated the Chern-Pontryagin density completely and obtained a closed expression for the topological charge Q - which requires to know the gauge singularities of $\Delta \Omega_{\mu\nu}^{(0)}$, $\Delta \Omega_{\mu\nu}^{(1)}$ and $\Delta \Omega_{\mu\nu}^{(2)}$ though. This result applies as well for gauge group SU(3), whose transition functions can be gauge transformed into transition functions with values in SU(2). The practical use of the expression is that it allows us to calculate Q, in the continuum and on the lattice, without resort to numerical integration.

We have seen that eq.(79) leaves considerable room for the practical evaluation of Q on the lattice - including the choice of an interpolation of the transition functions. The by far fastest procedure is the first method described in section IV, in which it is sufficient to compute $Q_{\mathcal{L}}^{(3)}$. This will be applicable in the "continuum region" only. But notice that alone in the "continuum region" a lattice gauge field configuration can be assigned a unique topological charge⁸.

While we were preparing this paper we received a preprint by Phillips and Stone¹⁴⁾, who also were able to compute the topological charge of SU(2) gauge fields by analytical means. The amount of algebra to do in their algorithm is similar to that of the second method sketched in section IV, where one has to compute $Q^{(1)}$ only. It will be interesting to compare their charge with, e.g., Lüscher's interpolation of the transition functions on individual gauge field configurations at various values of the coupling constant to test for uniqueness of the results.

References

- A.A. Belavin, A.M. Polyakov, A.S. Schwartz, Yu.S. Tyupkin: Phys. Lett. <u>59B</u>, 85 (1975)
- I.A. Fox, J.P. Gilchrist, M.L. Laursen, G. Schierholz: Phys. Rev. Lett. <u>54</u>, 749 (1985)
- 3. Y. Arian, P. Woit: Stony Brook preprint ITP-SB-85-60 (1985)
- 4. G. 't Hooft: Phys. Rev. Lett. 37, 8 (1976); Phys. Rev. D14, 3432 (1976)
- 5. E.-M. Ilgenfritz, M.L. Laursen, M. Müller-Preußker, G. Schierholz, H. Schiller: DESY preprint DESY 85-108 (1985), to be published in Nucl. Phys. <u>B</u>
- C. Callan, R. Dashen, D. Gross: Phys. Rev. <u>D19</u>, 1826 (1979);
 E.V. Shuryak: Phys. Rep. <u>115</u>, 151 (1984)
- 7. C. Callan, R. Dashen, D. Gross: Phys.Rev. <u>D17</u>, 2717 (1978)
- 8. M. Lüscher: Comm. Math. Phys. 85, 39 (1982)
- M.L. Laursen, G. Schierholz, U.-J. Wiese: DESY preprint DESY 85-062 (1985), to be published in Comm. Math. Phys.
- 10. G. Parisi, F. Rapuano: Phys. Lett. <u>152B</u>, 218 (1985)
- Y. Choquet-Bruhat, C. DeWitt-Morette, M. Dillard-Bleick: Analysis, Manifolds and Physics (North-Holland, Amsterdam, 1982)
- 12. N. Steenrod: The Topology of Fibre Bundles (Princeton University Press, Princeton, 1951)
- 13. U.-J. Wiese: Thesis, to be published
- 14. A. Phillips, D. Stone: Stony Brook preprint (1985)



Fig.1 The 5 spherical tetrahedra covering S^3 . A_1 is the angle between the spherical triangles (i,j,k) and (i,j,1), A_2 is the angle between the spherical triangles (i,j,k) and (i,j,m) and A_3 is the angle between the spherical triangles (i,j,m) and (i,j,1).





Fig.1

11 į,

ŝ

, t

1999 - K

с 1 4

× ε